



UNIVERSITÀ  
DEGLI STUDI  
DI UDINE

## Università degli studi di Udine

Hydrolysis of new Pt(II) complexes with ligands of natural origin: a DFT study

*Original*

*Availability:*

This version is available <http://hdl.handle.net/11390/1097933> since 2016-12-02T10:09:43Z

*Publisher:*

*Published*

DOI:

*Terms of use:*

The institutional repository of the University of Udine (<http://air.uniud.it>) is provided by ARIC services. The aim is to enable open access to all the world.

*Publisher copyright*

(Article begins on next page)

## Hydrolysis of new Pt(II) complexes with ligands of natural origin: a DFT study

Daniele Veclani, Andrea Melchior, Marilena Tolazzi

*Dipartimento di Chimica Fisica e Ambiente, Università di Udine, Italy*

Since the discovery of the first complex which exhibited anti-tumor properties (cis-diamminedichloro-Pt(II), cisplatin) [1] numerous Pt(II) complexes have been used as anti-cancer drugs, but only few have been approved for clinical use. The accepted mechanism of action of cisplatin consist in two stages: intracellular activation by the hydrolysis of a ligand (for example Cl<sup>-</sup> in cisplatin) and formation of intra-strand cross-links in DNA which produces a bending that causes the generation of defective proteins and ultimately the cell death [2]. Unfortunately, its therapeutic efficacy is somewhat compromised by occurrence of serious side effects [3]. In order to overcome these limitations, many studies have been done to find other platinum drugs with an equivalent or improved range of activity but with less toxic side effects [4]. Hydrolysis reactions are the rate determining step of the whole process and therefore many experimental and theoretical studies [5] have been carried out to obtain structure-reactivity relationships.

In this work we report the theoretical study of the hydrolysis reaction processes of Pt(II)[(NH<sub>3</sub>)<sub>2</sub>(C<sub>9</sub>O<sub>5</sub>H<sub>8</sub>)] (A), Pt(II)[(PH<sub>3</sub>)<sub>2</sub>(C<sub>9</sub>O<sub>5</sub>H<sub>8</sub>)] (B) and Pt(II)[(PH<sub>3</sub>)<sub>2</sub>(Cl)<sub>2</sub>] (C) shown in figure 1. The complexes A and B once hydrolyzed within the cell, release ethylene gallate (C<sub>9</sub>O<sub>5</sub>H<sub>8</sub>), compound which has anti-oxidant and anti-inflammatory and antiproliferative properties [6]. Complex C has been studied for comparison.




Figure 1: complexes based on platinum studied in this work.

These complexes have been studied using density functional theory (DFT) in order to obtain detailed data on its mechanism of action. DFT calculations were performed with the Gaussian 09 program, using the mPW1PW91 functional. Geometry Optimizations were carried out in vacuum and PCM with a 6-31+G(d) basis set for all atoms except the platinum atom, which was described by the quasi-relativistic Stuttgart-Dresden pseudopotential (SDD). Solvent effect have been introduced by the PCM approach. The results were compared with those present in the literature for several anti-cancer drugs [5,7-8].

The computed potential energy surface shows that the lower activation energy is found for [Pt(II)[(PH<sub>3</sub>)<sub>2</sub>(Cl)<sub>2</sub>]. For complex Pt(II)[(PH<sub>3</sub>)<sub>2</sub>(C<sub>9</sub>O<sub>5</sub>H<sub>8</sub>)] it can be observed that activation energy is comparable to cisplatin. Pt(II)[(NH<sub>3</sub>)<sub>2</sub>(C<sub>9</sub>O<sub>5</sub>H<sub>8</sub>)], instead, shows the highest energy barrier, this indicates that the hydrolysis reaction is disfavored in this complex.

[1] B. Rosenberg, L. Camp, T. Krigas, *Nature*, 1965, **205**, 698.

[2] M.J. Bloemink, J. Reedijk, *Met Ions Biol Syst*, 1999, **32**, 641.

[3] C. Bokemeyer, C. Berger, B. Kynast, H.J. Schmoll, H. Poliwada, *Eur. J. Cancer*, 1993, **29**, S241.

[4] D. Wang, S. Lippard, *J. Nat. Rev. Drug Discovery*, 2005, **4**, 307.

[5] A. Melchior, E. S. Marcos, R. R. Pappalardo, J. M. Martinez, *Theor Chem Acc*, 2011, **128**, 627

[6] S. H. Lee, J. K. Kim, D. W. Kim, H. S. Hwang, W. S. Eum, J. Park, K. H. Han, J. S. Oh, S. J. Choi, *Biochim. Biophys. Acta*, 2013, **1830**, 4017

[7] M. F. A. Lucas, M. Pavelka, M. E. Alberto, N. Russo, *J. Phys. Chem.*, 2009, **B113**, 831.

[8] M. E. Alberto, M. F. A. Lucas, M. Pavelka, N. Russo, *J. Phys. Chem.*, 2009, **B113**, 14473.