



UNIVERSITÀ
DEGLI STUDI
DI UDINE

Università degli studi di Udine

Exploitation of lettuce waste flour to increase bread functionality: effect on physical, nutritional, sensory properties and on consumer response

Original

Availability:

This version is available <http://hdl.handle.net/11390/1138467> since 2020-02-27T15:26:23Z

Publisher:

Published

DOI:10.1111/ijfs.13820

Terms of use:

The institutional repository of the University of Udine (<http://air.uniud.it>) is provided by ARIC services. The aim is to enable open access to all the world.

Publisher copyright

(Article begins on next page)

WILEY

Online Proofing System Instructions

The Wiley Online Proofing System allows proof reviewers to review PDF proofs, mark corrections, respond to queries, upload replacement figures, and submit these changes directly from the locally saved PDF proof.

1. For the best experience reviewing your proof in the Wiley Online Proofing System ensure you are connected to the internet. This will allow the PDF proof to connect to the central Wiley Online Proofing System server. If you are connected to the Wiley Online Proofing System server you should see a green check mark icon above in the yellow banner.



Connected



Disconnected

2. Please review the article proof on the following pages and mark any corrections, changes, and query responses using the Annotation Tools outlined on the next 2 pages.



3. Save your proof corrections by clicking the “Publish Comments” button in the yellow banner above. Corrections don’t have to be marked in one sitting. You can publish comments and log back in at a later time to add and publish more comments before you click the “Complete Proof Review” button below.



4. If you need to supply additional or replacement files bigger than 5 Megabytes (MB) do not attach them directly to the PDF Proof, please click the “Upload Files” button to upload files:

5. When your proof review is complete and all corrections have been published to the server by clicking the “Publish Comments” button, please click the “Complete Proof Review” button below:

IMPORTANT: Did you reply to all queries listed on the Author Query Form appearing before your proof?

IMPORTANT: Did you click the “Publish Comments” button to save all your corrections? Any unpublished comments will be lost.


IMPORTANT: Once you click “Complete Proof Review” you will not be able to add or publish additional corrections.

Required software to e-Annotate PDFs: **Adobe Acrobat Professional** or **Adobe Reader** (version 11 or above). (Note that this document uses screenshots from **Adobe Reader DC**.)

The latest version of Acrobat Reader can be downloaded for free at: <http://get.adobe.com/reader/>

Once you have Acrobat Reader open on your computer, click on the **Comment** tab (right-hand panel or under the Tools menu).

This will open up a ribbon panel at the top of the document. Using a tool will place a comment in the right-hand panel. The tools you will use for annotating your proof are shown below:





1. Replace (Ins) Tool – for replacing text.



Strikes a line through text and opens up a text box where replacement text can be entered.

How to use it:

- Highlight a word or sentence.
- Click on .
- Type the replacement text into the blue box that appears.




2. Strikethrough (Del) Tool – for deleting text.



Strikes a red line through text that is to be deleted.

How to use it:

- Highlight a word or sentence.
- Click on .
- The text will be struck out in red.

experimental data if available. For ORFs to be had to meet all of the following criteria:



1. Small size (35-250 amino acids).
2. Absence of similarity to known proteins.
3. Absence of functional data which could not be the real overlapping gene.
4. Greater than 25% overlap at the N-terminus terminus with another coding feature; over both ends; or ORF containing a tRNA.


3. Commenting Tool – for highlighting a section to be changed to bold or italic or for general comments.



Use these 2 tools to highlight the text where a comment is then made.

How to use it:

- Click on .
- Click and drag over the text you need to highlight for the comment you will add.
- Click on .
- Click close to the text you just highlighted.
- Type any instructions regarding the text to be altered into the box that appears.

informal invariance:  for
A: Math. Gen., Vol. 12, N

lified theory for a matrix
'ol. 8, 1984, pp. 305-323
d manuscript, 1984.

ching fractions for $D_0 \rightarrow K+K$
relation in D_0 decays' Phys





4. Insert Tool – for inserting missing text at specific points in the text.



Marks an insertion point in the text and opens up a text box where comments can be entered.

How to use it:

- Click on .
- Click at the point in the proof where the comment should be inserted.
- Type the comment into the box that appears.




5. Attach File Tool – for inserting large amounts of text or replacement figures.



Inserts an icon linking to the attached file in the appropriate place in the text.

How to use it:

- Click on  .
- Click on the proof to where you'd like the attached file to be linked.
- Select the file to be attached from your computer or network.
- Select the colour and type of icon that will appear in the proof. Click OK.

The attachment appears in the right-hand panel.


chondrial preparator
ative damage injury
the extent of membra
malondialdehyde (TBARS) formation.
used by high perform

6. Add stamp Tool – for approving a proof if no corrections are required.



Inserts a selected stamp onto an appropriate place in the proof.

How to use it:


- Click on  .
- Select the stamp you want to use. (The [Approved](#) stamp is usually available directly in the menu that appears. Others are shown under *Dynamic*, *Sign Here*, *Standard Business*).
- Fill in any details and then click on the proof where you'd like the stamp to appear. (Where a proof is to be approved as it is, this would normally be on the first page).

of the business cycle, starting with the
on perfect competition, constant ret
production. In this environment goods
extra costs are incurred to make marks
he total cost of production is deter
etermined by the model. The New-Key
otaki (1987), has introduced produc
general equilibrium models with nomin
and real variables. Most of this litera

APPROVED




Drawing tools available on comment ribbon




7. Drawing Markups Tools – for drawing shapes, lines, and freeform annotations on proofs and commenting on these marks.

Allows shapes, lines, and freeform annotations to be drawn on proofs and for comments to be made on these marks.

- Click on one of the shapes in the [Drawing Markups](#) section.
- Click on the proof at the relevant point and draw the selected shape with the cursor.
- To add a comment to the drawn shape, right-click on shape and select *Open Pop-up Note*.
- Type any text in the red box that appears.



For further information on how to annotate proofs, click on the [Help](#) menu to reveal a list of further options:



Author Query Form

WILEY

Journal: IJFS

Article: 13820

Dear Author,

During the copyediting of your manuscript the following queries arose.

Please refer to the query reference callout numbers in the page proofs and respond to each by marking the necessary comments using the PDF annotation tools.

Please remember illegible or unclear comments and corrections may delay publication.

Many thanks for your assistance.

Query reference	Query	Remarks
A	AUTHOR: Please note that this proof exceeds the journal's free pages allocation (7pp) and will be subject to an excess page charge of £100 for each typeset page exceeding the first 7 pages. Please download the page charge form for your paper is more than 7 proof pages from http://onlinelibrary.wiley.com/store/10.1111/(ISSN)1365-2621/asset/homepages/Page_Charge_Form.pdf?v=1&s=c0fd894e13c9-fa7ae37a4b2149dec48a4f8b162f&isAguDoi=false . Please return the completed form to the Production Editor (ijfs@wiley.com , fax: +44 4219 7754). A hardcopy via regular mail would not be necessary. Please pay excess page charges when the article is published in an issue of the journal.	
1	AUTHOR: Please verify that the linked ORCID identifier is correct for the author "Lara Manzocco".	
2	AUTHOR: Please confirm that given names (red) and surnames/family names (green) have been identified correctly.	
3	AUTHOR: Please provide blurb content for graphical abstract.	
4	AUTHOR: Mastromatteo <i>et al.</i> , 2014 has not been included in the Reference List, please supply full publication details.	
5	AUTHOR: Ortolan <i>et al.</i> , 2016 has been changed to Ortolan <i>et al.</i> , 2015 so that this citation matches the Reference List. Please confirm that this is correct.	
6	AUTHOR: Ajila <i>et al.</i> , 2008 has not been included in the Reference List, please supply full publication details.	
7	AUTHOR: Lattimer and Haub, 2014 has been changed to Lattimer & Haub, 2010 so that this citation matches the Reference List. Please confirm that this is correct.	

Original article

Exploitation of lettuce waste flour to increase bread functionality: effect on physical, nutritional, sensory properties and on consumer responseStella Plazzotta, Sandro Sillani & Lara Manzocco* 

Department of Agricultural, Food, Environmental and Animal Sciences, University of Udine, Via Sondrio 2/A, 33100 Udine, Italy

(Received 28 March 2018; Accepted in revised form 24 April 2018)

Summary Phenol and fibre-rich flour obtained by air-drying and grinding of lettuce waste was partially substituted (26, 53, 170 and 575 g kg⁻¹) with wheat flour to produce functional bread. The addition of flour progressively decreased dough leavening capacity while increased bread moisture and firmness. Lettuce waste flour significantly increased the polyphenolic content (up to 3.4 g GAE kg⁻¹) of bread samples and enhanced their antioxidant activity by 200%. Bread containing 170 and 575 g kg⁻¹ of lettuce waste flour presented sensory properties and consumer acceptability comparable to those of commercial wholemeal bread with similar rye bran content. Bread containing at least 170 g kg⁻¹ of lettuce waste flour could be associated to nutritional claims related to its enhanced fibre content (>30 g kg⁻¹). Data obtained by conjoint analysis demonstrate the possibility of increasing consumer preference for lettuce waste flour bread by proper nutritional (fibre content) and sustainability (lettuce waste valorisation) claims.

Keywords Acceptability, consumer response, functional bread, lettuce waste, sensory properties.

Introduction

The market of functional foods has been constantly growing, following consumer awareness of their potential in maintaining a healthy state (Sikand *et al.*, 2015; Gul *et al.*, 2016). Being a staple food in several countries, bread is an optimal candidate for functionalisation (Akhtar *et al.*, 2011). To this aim, wholemeal flour is traditionally used due to its content in antioxidants and fibres from bran and aleurone (Dewettinck *et al.*, 2008; Dziki *et al.*, 2014). This goal could be equally reached using flours from fruit and vegetables or from their wastes, which are often richer in nutritional compounds (Nilnakara *et al.*, 2009; Mastromatteo *et al.*, 2014). Functional bread has been produced using flours from tomato, cabbage and pineapple waste (Nilnakara *et al.*, 2009; Nour *et al.*, 2015; Wu & Shiau, 2015; Chareonthaikij *et al.*, 2016).

Fresh-cut processing of lettuce heads generates huge amounts of waste, due to removal of external leaves and core. Waste amounts up to 40% of the initial lettuce weight, leading to high management costs (Plazzotta *et al.*, 2017). This waste can be air-dried and ground to obtain flour with 3.05 mg GAE g⁻¹ dw polyphenols, similar to that of cabbage and pumpkin

functional flour (Que *et al.*, 2008; Nilnakara *et al.*, 2009), and fibre content (260 g kg⁻¹) comparable to that of rice and oat bran (USDA, 2016). Lettuce waste flour could thus represent a suitable ingredient for functional bread. However, its use is expected to strongly affect product processability as well as physical, sensory and nutritional properties.

Functional bread containing lettuce waste flour could represent a value-added food derived from a cheap and always available source, associated to an eco-friendly image appreciated by consumers (Simoes *et al.*, 2015). Reversely, the presence of a waste derivative in bread could negatively affect consumers' reaction (Pickett-Baker & Ozaki, 2008; Simoes *et al.*, 2015).

The aim of this study was to evaluate the potential of flour from *Iceberg* lettuce waste in functional bread production. Bread containing increasing amounts of lettuce waste flour were characterised for colour, moisture, phenol and fibre content, antioxidant activity, firmness, specific volume and sensory properties. Bread acceptability was evaluated and compared to that of commercial bread containing rye bran. Consumer response towards claims associable to bread containing lettuce waste flour was investigated by conjoint analysis.

*Correspondent: E-mail: lara.manzocco@uniud.it

Dispatch: 10.5.18	CE: Menaka T	PE: Eswari M.
No. of pages: 8		
WILEY		
13820	Manuscript No.	
I J F S	Journal Code	
		

Materials and methods

Lettuce waste flour

Lettuce waste was prepared by removing external leaves and core from heads, simulating fresh-cut processing. One kg waste was then air-dried (15 h) in single layers (UM100; Memmert, Schwabach, Germany) at 70 ± 0.5 °C and a 55–65 ERH%, ground into flour (MC3001; Moulinex, Milan, Italy), sieved using a 125 µm sieve (Endecotts Ltd, London, UK) and stored at 20 °C in sealed aluminised aseptic bags until use (Plazzotta *et al.*, 2018). Flour presented a moisture amount of 42 ± 2 g kg⁻¹.

Bread

Bread was obtained substituting Manitoba type '0' wheat flour (Molino Spadoni, Coccolia, Italy; 141 g kg⁻¹ moisture, 12 g kg⁻¹ fat, 675 g kg⁻¹ carbohydrates of which 15 g kg⁻¹ sugars, 22 g kg⁻¹ fibres and 135 g kg⁻¹ proteins) with increasing amounts of lettuce waste flour (0, 10, 25, 70, 225 g kg⁻¹ of dough, corresponding to a replacement of wheat flour of 2%, 4%, 12% and 40% w/w), while maintaining a constant ratio among the other ingredients (Table 1). Water (at 30 ± 0.5 °C), sugar and fresh yeast were premixed 3 min (KM285; Kenwood, Milan, Italy), added with flour and salt and mixed 15 min. Subsequently, dough portions (250 g) were manually rounded, leavened on a tray at 37 °C and 80 ERH% (ST500; Pol-Eko-Aparatura S.P.J., Wodzslaw, Poland) for 60 min, baked (170 °C, 20 min; 10GN1/1; Air-O-Steam Touchline, Electrolux, Porcia, Italy) and cooled at room temperature for 1 h (Mastromatteo *et al.*, 2012).

Commercial wholegrain bread samples (Gilli srl, Laives, Italy) containing 180 and 510 g kg⁻¹ rye bran (fibre content of 55 and 93 g kg⁻¹, respectively, as declared on the product label) were also purchased on the local market.

Image acquisition

Images were acquired by a EOS-550D camera (Canon, Milan, Italy) placed 45 cm above a black cardboard base where samples were positioned and enlightened by 4 100 W-frosted photographic floodlights, in a position allowing minimum shadow and glare (Immagini & Computer, Bareggio, Italy).

Colour

A tristimulus Chromameter-2-Reflectance colorimeter (Minolta, Osaka, Japan) with a CR-300 measuring head, standardised against a white tile, was used and data were expressed in L*, a* and b* Hunter-scale

Table 1 Formulation of dough samples containing increasing amounts of lettuce waste flour

Dough ingredient (g kg ⁻¹)					
Lettuce waste flour	Wheat flour	Water	Sugar	Salt	Yeast
0	561	404	9	13	13
10	551	404	9	13	13
25	536	404	9	13	13
70	491	404	9	13	13
225	336	404	9	13	13

parameters (Ortolan *et al.*, 2015). Samples were positioned on a white cardboard, and the colorimeter head was placed perpendicular to sample surface. At least five measures were taken on different points of bread dough samples and on the crust of bread ones.

Specific volume

Loaf-specific volume (cm³ g⁻¹) was obtained by rapeseed displacement according to AOAC methods (AOAC, 2000).

Moisture

Moisture content was calculated according to AOAC methods (AOAC, 2000). Around 2 g of sample was dried in a vacuum oven (1.32 kPa) at 75 °C until constant weight (12 h).

Firmness

Firmness was measured by uniaxial compression test using an Instron 4301 (Instron LTD., High Wycombe, UK). Samples were tested by a 12.7-mm-diameter cylindrical probe (100 N compression head) at a 5-mm min⁻¹ crosshead speed. Firmness was taken as the maximum force (N) for 5-mm sample penetration. Dough firmness was evaluated by penetrating the surface of the leavened dough. In the case of bread, six slices (20 mm thick) were cut from the central portion of each loaf. Firmness was evaluated by penetrating the crumb of each slice in five different points (Calligaris *et al.*, 2013).

Total dietary fibre

Total dietary fibre was determined according to AOAC methods (AOAC, 2000), using TDF-100A kit (Sigma-Aldrich, St. Louis, MO, USA). Results were expressed as g of fibres per kg of bread.

Polyphenolic extract

Extracts were prepared according to Llorach *et al.* (2004) with some modifications. Bread was extracted

by reflux for 60 min in boiling water (dilution 0.2 g mL⁻¹). Extracts were then added in methanol (1 mL g⁻¹ of bread) and centrifuged (9450 g, 15 min, 20 °C; Mikro 20; Hettich Zentrifugen, Tuttlingen, Germany). The supernatant was used for polyphenols and antioxidant activity analyses.

Total polyphenols

The Folin-Ciocalteu reagent was used (Singleton & Rossi, 1985). The reaction mixture contained 100 µL polyphenolic extract, 500 µL Folin-Ciocalteu reagent, 4 mL water and 2 mL sodium carbonate-water solution (0.15 g mL⁻¹). After 2 h-reaction in the dark at ambient temperature, absorbance was read at 750 nm (UV-2501PC, UV-Vis spectrophotometer; Shimadzu Corporation, Kyoto, Japan). A calibration curve ($R^2 = 0.99$) was made with 0.1–1000 mg L⁻¹ solutions of gallic acid. Results were expressed as mg of gallic acid equivalents (GAE) per kg of bread.

Antioxidant activity

A volume of 1.80 mL of 6.1×10^{-5} M DPPH· methanol solution was added with 150 µL polyphenolic extract. DPPH· bleaching was followed at 515 nm (UV-2501PC; Shimadzu Corporation) at 20 °C for 10 min and resulted proportional to extract concentration. Equation 1 was chosen to obtain the reaction rate of DPPH· bleaching, k (Manzocco *et al.*, 1998):

$$\frac{1}{A^3} - \frac{1}{A_0^3} = 3kt, \quad (1)$$

where A_0 is initial optical density (OD) and A is optical density at increasing time, t . The chain-breaking activity was expressed as k (OD⁻³ min⁻¹) per sample mass.

Sensory attributes

A focus group of ten judges was used to identify sensory attributes of bread containing lettuce flour. Judges were not trained on sensory analysis of bread but were experts in the use of the selected sensory methods. White bread and bread containing 575 g kg⁻¹ of lettuce flour were evaluated. The focus group decided, through consensus and independently on consumer response, which descriptors better discriminated the samples. Judges were then asked to evaluate the intensity of the selected descriptors in bread containing 0, 170 and 575 g kg⁻¹ of lettuce waste flour. Descriptors of bread samples, identified with a three-digit random code, were evaluated on a 1- to 9-point hedonic scale, in which one corresponded to 'extremely low descriptor intensity', and 9 to 'extremely high descriptor intensity'. Three bread samples

were evaluated in each session, and water was used to rinse mouth among samples (Manzocco & Lagazio, 2009).

Consumer acceptability

About eighty bread consumers (thirty-seven men and forty-three women, age 18–55) were recruited at the University of Udine, Italy. Samples, indicated by a three-digit random code, were served in odourless plastic dishes. Consumers were asked to taste samples and score their acceptability on a 1- to 9-hedonic scale anchored with 'highly nonacceptable' (score 1) and 'highly acceptable' (score 9; Peryam & Pilgrim, 1957). Four bread samples containing lettuce flour (170 and 575 g kg⁻¹) and rye bran (180 and 510 g kg⁻¹) were evaluated in each session. Water was used to rinse mouth among samples.

Consumer response

Conjoint analysis was used to evaluate consumer preference towards bread by decomposing total preference in partial preferences relevant to independent product attributes (De Pelsmaecker *et al.*, 2017; Sillani *et al.*, 2017). Five attributes of bread containing lettuce flour were selected as experimental variables and named 'lettuce flour', 'health', 'waste recovery', 'waste reduction' (discrete variables) and 'price' (linear variable). Different levels were associated to each experimental variable (Table 2). For discrete variables, two levels (claim presence or absence) were used. For price, three values were used. Experimental variables were combined according to an orthogonal experimental design, obtaining eleven product profiles, which represent information available to consumer on a possible bread label. A nonprobabilistic sample of 525 bread consumers, equally distributed among men and women (age 18–41), was recruited at the University of Udine, Italy. Consumers were asked to fill up a structured questionnaire, indicating, for each product profile, their preference on a 1–100 scale. No prior information was provided about origin and preparation of lettuce waste flour. In other words, the response of

Table 2 Experimental variables defining bread attributes and relevant levels used for conjoint analysis

Experimental variable	Levels
Lettuce flour	Absent; 'Containing lettuce flour'
Health	Absent; 'Rich in fibre'
Waste recovery	Absent; 'Produced recovering lettuce waste'
Waste reduction	Absent; 'Produced reducing food waste'
Price (€ kg ⁻¹)	3.00; 4.50; 6.00

consumers towards a bread label reporting different information was assessed. A total of 370 responses were valid and analysed.

Data analysis

Determinations were expressed as the mean \pm standard deviation of at least three measurements from three experiment replications. Statistical analysis was performed using R v.2.15.0 (The R foundation for Statistical Computing, Vienna, Austria). Bartlett's test was used to check the homogeneity of variance. One-way ANOVA was carried out, and Tukey's test was used as *post hoc* test to determine statistically significant differences among means ($P < 0.05$). For conjoint analysis, IBM SPSS Statistics 20 (Armonk, NY, USA) was used to calculate partial preference values, their relative importance and model goodness of fit (Pearson's R and Kendall's τ).






Results and discussion

Wheat flour in bread dough was substituted with increasing amounts of lettuce waste flour. Leavened dough was characterised for appearance, colour and firmness (Table 3). The addition of lettuce waste flour decreased luminosity (L^*) and yellowness (b^*), while increased red point (a^*). This can be attributed to the brownish colour of air-dried lettuce flour. Air-drying has actually been reported to promote oxidation of the

main phenolic compounds of *Iceberg* lettuce waste, including 3-O-caffeoylquinic acid, caffeoyltartaric acid, 4-O-caffeoylquinic acid, 5-O-caffeoylquinic acid, caffeic acid derivatives, isochlorogenic acid, chicoric acid, luteolin 7-O-glucuronide and quercetin 3-O-glucuronide (Plazzotta *et al.*, 2018). Images show that lettuce flour hindered dough leavening, leading to progressively firmer and less aerated dough. Lettuce flour reduced gluten concentration in the dough, while increased the presence of water holding fibres. To this regard, Plazzotta *et al.* (2018) demonstrated that 1 g of lettuce flour can hold up to 9 g of water. Water would thus be less available for gluten-starch network formation, reducing dough elasticity, gas entrapment and leavening capacity. The addition of vegetable fibres was demonstrated to affect dough moisture distribution, altering rheological properties and leavening (Ameh *et al.*, 2013; Chareonthaikij *et al.*, 2016).






Increasing amounts of brownish lettuce flour resulted in brown bread samples (Table 4). The changes in luminosity and colorimetric parameters presented a discontinuity point in correspondence of 170 g kg^{-1} lettuce flour concentration. This is probably attributable to the counterbalancing colour effects of the increase in brownish flour and the inhibition of Maillard reaction by water holding fibres, which reduce dehydration rate and extent (Kent-Jones & Amos, 1967). As reported for other vegetable flours (Greene & Bovell-Benjamin, 2004; Marpalle *et al.*, 2014), the water holding capacity of lettuce fibres was

Table 3 Appearance, hunter scale colour parameters (L^* , a^* , b^*) and firmness of leavened dough containing increasing amounts of lettuce waste flour

Lettuce waste flour (g kg^{-1})	Appearance after leavening	Colour			Firmness (N)
		L^*	a^*	b^*	
0		81.8 ± 0.5^a	-0.1 ± 0.3^a	18.6 ± 0.9^c	0.110 ± 0.015^b
10		72.6 ± 1.0^b	2.4 ± 0.3^b	21.7 ± 0.9^b	0.139 ± 0.005^b
25		60.4 ± 0.7^c	7.5 ± 0.4^c	28.1 ± 1.6^a	0.217 ± 0.023^b
70		48.7 ± 0.6^d	10.2 ± 0.3^d	16.2 ± 0.5^d	0.139 ± 0.015^b
225		33.3 ± 1.1^e	11.0 ± 0.4^e	7.3 ± 2.0^e	1.803 ± 0.145^a

Mean values indicated by different superscript letters are statistically different ($P < 0.05$).

Table 4 Visual appearance, crust Hunter scale colour parameters (L^* , a^* , b^*), crumb firmness, specific volume, humidity, total phenolic content (TPC), chain-breaking activity (CBA) and total dietary fibre (TDF) of bread samples containing increasing amount of lettuce waste flour

Lettuce waste flour (g kg ⁻¹)	Dough	Bread	Colour			Firmness (N)	Specific volume (cm ³ g ⁻¹)	Moisture content (g kg ⁻¹)	TPC (mg GAE kg ⁻¹)	CBA (DO ⁻³ min ⁻¹ kg ⁻¹)	TDF (g kg ⁻¹)
			L*	a*	b*						
0			64.9 ± 2.0 ^a	7.5 ± 1.5 ^c	26.4 ± 0.8 ^b	0.573 ± 0.143 ^d	5.595 ± 0.683 ^a	431 ± 1 ^c	438.5 ± 8.3 ^c	3873.3 ± 505.6 ^c	13.2 ± 1.1 ^e
10			52.3 ± 1.1 ^c	11.4 ± 0.4 ^a	24.9 ± 0.9 ^{cd}	0.523 ± 0.033 ^d	3.580 ± 0.288 ^b	445 ± 1 ^c	585.6 ± 4.2 ^c	4486.7 ± 14.1 ^{bc}	16.2 ± 1.2 ^d
25			50.4 ± 3.3 ^c	12.0 ± 0.9 ^a	23.4 ± 1.8 ^c	0.849 ± 0.056 ^c	3.439 ± 0.259 ^b	462 ± 3 ^b	731.2 ± 2.1 ^c	4644.4 ± 937.5 ^{bc}	19.3 ± 1.1 ^c
70			59.1 ± 2.7 ^b	9.9 ± 0.7 ^b	29.3 ± 1.2 ^a	1.351 ± 0.214 ^b	1.625 ± 0.098 ^c	464 ± 2 ^b	1354.7 ± 6.2 ^b	5602.2 ± 288.8 ^b	31.1 ± 2.0 ^b
225			43.9 ± 0.6 ^a	10.9 ± 0.5 ^{ab}	9.4 ± 1.3 ^d	5.212 ± 0.174 ^a	0.886 ± 0.060 ^d	487 ± 8 ^a	3406.2 ± 78.9 ^a	10 290.0 ± 621.6 ^a	71.5 ± 2.2 ^a

Mean values indicated by different superscript letters are statistically different ($P < 0.05$).

reflected in progressively increasing bread loaf moisture (Table 4). This might pose stability issues for lettuce flour bread, due to its altered response to both microbial spoilage and staling (Rosell & Santos, 2010; Ameh *et al.*, 2013). Bread containing increasing amounts of lettuce flour resulted progressively firmer and with lower specific volume (Table 4), due to the water absorption capacity of lettuce fibres and reduced dough leavening (Table 3). The latter should not be regarded as a negative feature, as promoting stomach filling and sense of satiety (Greene & Bovell-Benjamin, 2004; Ameh *et al.*, 2013). The addition of lettuce flour, rich in antioxidant polyphenols (Plazzotta *et al.*, 2018), promoted the increase in bread phenolic content and antioxidant activity (Table 4). Phenolic compounds are mostly located in cereal cell wall, linked to hemicelluloses or other wall constituents, with the highest concentration in the aleurone grain layer. Subsequently, white bread is poor in these compounds. *Iceberg* lettuce flour could be exploited to increase its phenolic content, being rich in these compounds (Llorach *et al.*, 2004; Naczki & Shahidi, 2006). An increase in phenolic content and antioxidant activity of bakery products was also obtained by adding mango peel, dried tomato waste, broccoli, carrot and beetroot (Ajila *et al.*, 2007; Nour *et al.*, 2015; Ranawana *et al.*, 2016). By contrast, no change in total phenolic was detected when apple and lemon fibre was added to cookies (Bilgiçli *et al.*, 2007). Interactions between phenols and wheat proteins/polysaccharides as well as oxidation, isomerisation/epimerisation and degradation of bioactive compounds during dough preparation and baking may account for these contrasting results (Wang & Zhou, 2004). As a result, bread antioxidant properties would depend on phenols naturally occurring in wheat and lettuce flour as well as on thermally induced products and phenol complexes with proteins/polysaccharides (Rupasinghe *et al.*, 2008; Sivam *et al.*, 2010).

As reported for other baked goods containing vegetable derivatives (Bilgiçli *et al.*, 2007; Ajila *et al.*, 2008; Rupasinghe *et al.*, 2008), the addition of lettuce flour also promoted the increase of bread fibre concentration (Table 4). Lettuce mainly contains insoluble dietary fibre, known for its beneficial health effects on intestinal regularity and weight control (Lattimer & Haub, 2010). The fibre content of bread with 170 and 575 g kg⁻¹ lettuce flour could be associated to the nutritional claims 'rich in fibre' and 'source of fibres', respectively (Reg. CE. 1924/2006). As these high levels of fibres and polyphenols may significantly affect bread sensory attributes, these samples were submitted to sensory evaluation, using white bread as control (Mastromatteo *et al.*, 2012). Data reported in Table 5 show that lettuce flour decreased the perceived intensity of yeast odour and flavour while increased silage and herbaceous odour and flavour, dried fruit flavour,

Table 5 Sensory attributes scores of bread samples containing increasing amounts of lettuce waste flour

Lettuce waste flour (g kg ⁻¹)	Sensory attribute			Taste					Flavour					Texture		
	Odour	Bread	Dough	Yeast	Silage	Herbaceous	Acid	Sweet	Salty	Sour	Yeast	Silage	Herbaceous	Dried fruit	Softness	Gumminess
0				6.6 ± 1.7 ^a	1.9 ± 1.9 ^b	1.2 ± 0.4 ^b	3.3 ± 1.9 ^b	3.4 ± 1.2 ^a	6.0 ± 1.6 ^a	1.0 ± 0.5 ^b	6.1 ± 1.5 ^a	1.5 ± 1.3 ^b	1.2 ± 0.4 ^b	1.1 ± 0.3 ^b	6.0 ± 1.9 ^a	2.0 ± 0.8 ^b
70				1.3 ± 0.5 ^b	5.7 ± 1.4 ^a	6.1 ± 1.4 ^a	3.7 ± 1.9 ^{ab}	4.0 ± 1.9 ^a	4.6 ± 1.3 ^a	3.1 ± 0.9 ^{ab}	1.4 ± 0.5 ^b	4.5 ± 1.4 ^{ab}	5.0 ± 1.3 ^{ab}	4.2 ± 1.1 ^a	5.2 ± 1.0 ^a	5.5 ± 1.8 ^a
225				1.8 ± 1.6 ^b	5.9 ± 1.6 ^a	6.1 ± 1.4 ^a	5.8 ± 1.7 ^a	4.3 ± 1.1 ^a	4.3 ± 1.2 ^a	5.2 ± 1.9 ^a	1.7 ± 0.8 ^b	5.6 ± 1.6 ^a	6.4 ± 1.4 ^a	5.3 ± 1.5 ^a	2.5 ± 1.2 ^b	5.4 ± 1.3 ^a

For each sensory attribute, mean values indicated by different superscript letters are statistically different ($P < 0.05$). Sensory attributes were scored by a panel of ten judges.

acid and sour taste. In accordance with increasing bread firmness data (Table 4), lettuce flour promoted an increase in gumminess and a decrease in bread softness. Sensory attributes of bread containing lettuce flour are those typically associated to wholemeal bread (Greene & Bovell-Benjamin, 2004; Ameh *et al.*, 2013). Consumer acceptability of bread containing lettuce waste flour was thus compared to that of commercial rye bran bread (Table 6). To this aim, two commercial products were selected based on rye bran content (180 and 510 g kg⁻¹) comparable to that of lettuce waste flour in the selected samples (170 and 575 g kg⁻¹) as well as on similar colour ($P \geq 0.05$; Table 4). In particular, L*, a* and b* of bread containing 180 and 510 g kg⁻¹ rye bran were 58.1 ± 0.2 , 9.7 ± 1.2 , 30.3 ± 0.2 and 41.9 ± 1.1 , 11.9 ± 1.6 , 10.4 ± 0.9 , respectively.

Bread containing the highest level of rye bran (510 g kg⁻¹) or lettuce waste flour (575 g kg⁻¹) resulted less acceptable, probably due to the peculiar sensory attributes of fibre-rich bread (Table 6; Ameh *et al.*, 2013). Similar concentrations of rye bran or lettuce waste flour were associated to analogous acceptability scores, confirming lettuce flour bread to be just as acceptable as traditional wholemeal bread.

Despite consumer acceptability results, the reaction of consumers towards consumption of bread containing an ingredient deriving from waste could be a critical issue. Conjoint analysis was thus applied to assess

Table 6 Acceptability scores of bread samples containing increasing amounts of lettuce waste flour or rye bran

Sample	Acceptability
Bread with 180 g kg ⁻¹ rye bran	6.4 ± 0.7^a
Bread with 170 g kg ⁻¹ lettuce waste flour	6.6 ± 0.4^a
Bread with 510 g kg ⁻¹ rye bran	4.4 ± 0.9^b
Bread with 575 g kg ⁻¹ lettuce waste flour	4.4 ± 0.4^b

Mean values indicated by different superscript letters are statistically different ($P < 0.05$).

Bread acceptability was scored by eighty consumers.

Table 7 Partial preference coefficients and relative importance of different label information in defining consumer preference

Information	Partial preference coefficient	Relative importance
Price (€ kg ⁻¹) 3.00	-16.057 ± 1.133	32.4 ± 1.1
4.50	-24.086 ± 1.700	
6.00	-32.114 ± 2.267	
Produced recovering lettuce waste	1.353 ± 0.469	20.9 ± 0.8
Rich in fibre	3.466 ± 0.333	17.2 ± 0.6
Produced reducing food waste	2.668 ± 0.344	15.7 ± 0.6
Containing lettuce flour	1.816 ± 0.328	13.9 ± 0.6

Partial preference was scored by 370 consumers.

consumers' response towards a bread label reporting different information relevant to the presence of lettuce waste flour in bread (Table 2). Beside price, claims associated with nutritional value ('containing lettuce flour'; 'rich in fibre') or sustainability issues ('produced recovering lettuce waste'; 'produced reducing food waste') were considered. The obtained model resulted significant with *P*-values of both Pearson's *R* and Kendall's $\tau < 0.0001$. Partial preference coefficients and relative importance of each label information in defining consumer preference are reported in Table 7.

As expected, the increase in price led to a decrease in consumer preference. Price affected consumer preference by more than 30%, resulting the most important variable among those considered. Partial preference coefficients related to the presence of nutritional claims ('rich in fibre'; 'containing lettuce flour') resulted positive, indicating these claims to increase consumer preference. This result is consistent with the increasing consumer awareness of the importance of a diet rich in plant foods containing fibre (Rooney *et al.*, 2017). Sustainability claims ('produced recovering lettuce waste'; 'produced reducing food waste') also promoted a positive consumer reaction, probably due to increasing consumer concern about food sustainability (Grunert *et al.*, 2014; Simoes *et al.*, 2015). Noteworthy, the use of the word 'waste' was not associated with adverse consumer response. By contrast, the claim 'produced recovering lettuce waste' was the most important information after price in defining consumer preference. It can be inferred that waste-related claims could contribute in developing an eco-friendly image of bread containing lettuce waste flour and be strategically exploited to steer consumer preference towards more sustainable bread alternatives.

Conclusions

The use of flour obtained by air-drying of lettuce waste represents a promising strategy to improve bread functionality and sustainability. Such an approach could be easily extended to other vegetable wastes and to baked goods other than bread.

Although this valorisation strategy shows the potential for consumer acceptance, additional studies would be required to assess its feasibility. For instance, the effect of vegetable waste flour addition on bakery product shelf-life could be not negligible. Moreover, the success of this valorisation strategy will depend on the availability of vegetable waste flour with standardised composition and technological performance as well as safety and quality parameters suitable for food production. To this regard, the presence of contaminants deriving from both cultivation practices and process should be carefully assessed.

References

- Ajila, C.M., Leelavathi, K. & Rao, P.U.J.S. (2007). Improvement of dietary fibre content and antioxidant properties in soft dough biscuits with the incorporation of mango peel powder. *Journal of Cereal Science*, **48**, 319–326.
- Akhtar, S., Anjum, F.M. & Anjum, M.A. (2011). Micronutrient fortification of wheat flour: recent development and strategies. *Food Research International*, **44**, 652–659.
- Ameh, M.O., Gernah, D.I. & Igbabul, B.D. (2013). Physico-chemical and sensory evaluation of wheat bread supplemented with stabilized undefatted rice bran. *Food and Nutrition Sciences*, **4**, 43–48.
- AOAC. (2000). *Official Methods of Analysis*. Washington, DC: Association of Official Analytical Chemists.
- Bilgiçli, N., Ibanoglu, S. & Herken, E.N. (2007). Effect of dietary fibre addition on the selected nutritional properties of cookies. *Journal of Food Engineering*, **78**, 86–89.
- Calligaris, S., Manzocco, L., Valoppi, F. & Nicoli, M.C. (2013). Effect of palm oil replacement with monoglyceride organogel and hydrogel on sweet bread properties. *Food Research International*, **51**, 596–602.
- Chareonthaikij, P., Uan-On, T. & Prinyawiwatkul, W. (2016). Effects of pineapple pomace fibre on physicochemical properties of composite flour and dough, and consumer acceptance of fibre-enriched wheat bread. *International Journal of Food Science and Technology*, **51**, 1120–1129.
- De Pelsmaecker, S., Schouteten, J.J., Lagast, S., Dewettinck, K. & Gellynck, X. (2017). Is taste the key driver for consumer preference? A conjoint analysis study. *Food Quality and Preference*, **62**, 323–331.
- Dewettinck, K., Van Bockstaele, F., Kühne, B., Van de Walle, D., Courtens, T.M. & Gellynck, X. (2008). Nutritional value of bread: influence of processing, food interaction and consumer perception. *Journal of Cereal Science*, **48**, 243–257.
- Dziki, D., Rozylo, R., Gawlik-Dziki, U. & Swieca, M. (2014). Current trends in the enhancement of antioxidant activity of wheat bread by the addition of plant materials rich in phenolic compounds. *Trends in Food Science and Technology*, **40**, 48–61.
- Greene, J.L. & Bovell-Benjamin, A.C. (2004). Macroscopic and sensory evaluation of bread supplemented with sweet-potato flour. *Journal of Food Science*, **69**, 167–173.
- Grunert, K.G., Hieke, S. & Wills, J. (2014). Sustainability labels on food products: consumer motivation, understanding and use. *Food Policy*, **44**, 177–189.
- Gul, K., Singh, A.K. & Jabeen, R. (2016). Nutraceuticals and functional foods: the foods for the future world. *Critical Reviews in Food Science and Nutrition*, **56**, 2617–2627.
- Kent-Jones, D.W. & Amos, O.B.E. (1967). *Modern Cereal Chemistry*. London: Food Trade Press Ltd.
- Lattimer, J.M. & Haub, M.D. (2010). Effects of dietary fiber and its components on metabolic health. *Nutrients*, **2**, 1266–1289.
- Llorach, R., Tomás-Barberán, F.A. & Ferreres, F. (2004). Lettuce and chicory byproducts as a source of antioxidant phenolic extracts. *Journal of Agricultural and Food Chemistry*, **52**, 5109–5116.
- Manzocco, L. & Lagazio, C. (2009). Coffee brew shelf life modelling by integration of acceptability and quality data. *Food Quality and Preference*, **20**, 24–29.
- Manzocco, L., Mastrocola, D. & Nicoli, M.C. (1998). Chain-breaking and oxygen scavenging properties of wine as affected by some technological procedures. *Food Research International*, **31**, 673–678.
- Marpalle, P., Sonawane, S.K. & Arya, S.S. (2014). Effect of flaxseed flour addition on physicochemical and sensory properties of functional bread. *LWT – Food Science and Technology*, **58**, 614–619.
- Mastromatteo, M., Danza, A., Guida, M. & Del Nobile, M.A. (2012). Formulation optimisation of vegetable flour-loaded functional bread. Part II: effect of the flour hydration on the bread quality. *International Journal of Food Science and Technology*, **47**, 2109–2116.

- Naczk, M. & Shahidi, F. (2006). Phenolics in cereals, fruits and vegetables: occurrence, extraction and analysis. *Journal of Pharmaceutical and Biomedical Analysis*, **41**, 1523–1542.
- Nilnakara, S., Chiewchan, N. & Devahastin, S. (2009). Production of antioxidant dietary fibre powder from cabbage outer leaves. *Food and Bioprocess Technology*, **87**, 301–307.
- Nour, V., Ionica, M.E. & Trandafir, I. (2015). Bread enriched in lycopene and other bioactive compounds by addition of dry tomato waste. *Journal of Food Science and Technology*, **52**, 8260–8267.
- Ortolan, F., Brites, L.T.G., Montenegro, F.M. et al. (2015). Effect of extruded wheat flour and pre-gelatinized cassava starch on process and quality parameters of French-type bread elaborated from frozen dough. *Food Research International*, **76**, 402–409.
- Peryam, D.R. & Pilgrim, F.J. (1957). Hedonic scale method of measuring food preferences. *Food Technology*, **11**, 9–14.
- Pickett-Baker, J. & Ozaki, R. (2008). Pro-environmental products: marketing influence on consumer purchase decision. *Journal of Consumer Marketing*, **25**, 281–293.
- Plazzotta, S., Manzocco, L. & Nicoli, M.C. (2017). Fruit and vegetable waste management and the challenge of fresh-cut salad. *Trends in Food Science and Technology*, **63**, 51–59.
- Plazzotta, S., Calligaris, S. & Manzocco, L. (2018). Application of different drying techniques to fresh-cut lettuce waste to obtain food ingredients rich in antioxidants and with high solvent loading capacity. *LWT – Food Science and Technology*, **89**, 276–283.
- Que, F., Mao, L., Fang, X. & Wu, T. (2008). Comparison of hot air-drying and freeze-drying on the physicochemical properties and antioxidant activities of pumpkin (*Cucurbita moschata* Duch.) flours. *International Journal of Food Science and Technology*, **43**, 1195–1201.
- Ranawana, V., Raikos, V., Campbell, F. et al. (2016). Breads fortified with freeze-dried vegetables: quality and nutritional attributes. Part II: breads not containing oil as an ingredient. *Foods*, **5**, 62–76.
- Regulation (EC) No 1924/2006 of the European Parliament and of the Council of 20 December 2006 on nutrition and health claims made on foods.
- Rooney, C., McKinley, M.C., Appleton, K.M. et al. (2017). How much is “5-a-day”? A qualitative investigation into consumer understanding of fruit and vegetable intake guidelines. *Journal of Human Nutrition and Dietetics*, **30**, 105–113.
- Rosell, C.M. & Santos, E. (2010). Impact of fibers on physical characteristics of fresh and staled bake off bread. *Journal of Food Engineering*, **98**, 273–281.
- Rupasinghe, H.P.V., Wang, L., Huber, G.M. & Pitts, N.L. (2008). Effect of baking on dietary fibre and phenolics of muffins incorporated with apple skin powder. *Food Chemistry*, **107**, 1217–1224.
- Sikand, G., Kris-Etherton, P. & Boulos, N.M. (2015). Impact of functional foods on prevention of cardiovascular disease and diabetes. *Current Cardiology Reports*, **17**, 17–39.
- Sillani, S., Miccoli, A. & Nassivera, F. (2017). Different preferences for wine communication. *Wine Economics and Policy*, **6**, 28–39.
- Simoes, J.S., Mársico, E.T., da Cruz, A.G., de Freitas, M.Q., Doro, L.H. & Conte-Junior, C.A. (2015). Effect of sustainability information on consumers’ liking of freshwater prawn (*Macrobrachium rosenbergii*). *Journal of the Science of Food and Agriculture*, **95**, 3160–3164.
- Singleton, V.L. & Rossi, J.A.J. (1985). Colorimetry of total phenolics with phosphomolybdic-phosphotungstic acid reagents. *American Journal of Enology and Viticulture*, **16**, 144–158.
- Sivam, A.S., Sun-Waterhouse, D., Quek, S.Y. & Perera, C.O. (2010). Properties of bread dough with added fiber polysaccharides and phenolic antioxidants: a review. *Journal of Food Science*, **75**, R163–R174.
- USDA (2016). Food composition databases. <https://ndb.nal.usda.gov/ndb/>. Accessed 05.01.18.
- Wang, R. & Zhou, W. (2004). Stability of tea catechins in the bread-making process. *Journal of Agricultural and Food Chemistry*, **52**, 8224–8229.
- Wu, M.Y. & Shiau, S.Y. (2015). Effect of the amount and particle size of pineapple peel fiber on dough rheology and steamed bread quality. *Journal of Food Processing and Preservation*, **39**, 549–558.

Graphical Abstract

The contents of this page will be used as part of the graphical abstract of html only. It will not be published as part of main.

