



UNIVERSITÀ  
DEGLI STUDI  
DI UDINE

## Università degli studi di Udine

Compositional characterization of commercial sparkling wines from cv. Ribolla  
Gialla produced in Friuli Venezia Giulia

*Original*

*Availability:*

This version is available <http://hdl.handle.net/11390/1157659> since 2020-03-27T11:15:01Z

*Publisher:*

*Published*

DOI:10.1007/s00217-019-03334-9

*Terms of use:*

The institutional repository of the University of Udine (<http://air.uniud.it>) is provided by ARIC services. The aim is to enable open access to all the world.

*Publisher copyright*

(Article begins on next page)



Draft Manuscript for Review

# **Characterization of the metabolomic profile of Ribolla Gialla commercial sparkling wines**

Journal:	<i>European Food Research and Technology</i>
Manuscript ID	Draft
Manuscript Type:	Original paper
Date Submitted by the Author:	n/a
Complete List of Authors:	Voce, Sabrina; University of Udine, Department of Agricultural, Food, Environmental and Animal Sciences Skrab, Domen; University of Udine, Department of Agricultural, Food, Environmental and Animal Sciences; Fondazione Edmund Mach Istituto Agrario di San Michele all'Adige, Research and Innovation Centre Vrhovsek, Urska; Fondazione Edmund Mach Istituto Agrario di San Michele all'Adige, Research and Innovation Centre Battistutta, Franco; University of Udine, Department of Agricultural, Food, Environmental and Animal Sciences Comuzzo, Piergiorgio; University of Udine, Department of Agricultural, Food, Environmental and Animal Sciences Sivilotti, Paolo; University of Udine, Department of Agricultural, Food, Environmental and Animal Sciences
Keywords:	Ribolla Gialla; sparkling wine; volatile compounds; tryptophan metabolites; lipids

SCHOLARONE™  
Manuscripts

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1     **Characterization of the metabolomic profile of Ribolla Gialla**  
2     **commercial sparkling wines**

3     **Sabrina Voce<sup>1‡</sup>, Domen Skrab<sup>1,2‡</sup>, Urska Vrhovsek<sup>2</sup>, Franco Battistutta<sup>1</sup>, Piergiorgio**  
4     **Comuzzo<sup>1\*</sup>, Paolo Sivilotti<sup>1\*</sup>**

6     <sup>1</sup> University of Udine, Department of Agricultural, Food, Environmental and Animal Sciences,  
7     via delle Scienze 206, 33100 Udine – Italy  
8     <sup>2</sup> Edmund Mach Foundation, Research and Innovation Centre, Department of Food Quality and  
9     Nutrition, via Edmund Mach 1, 38010, San Michele all’Adige, TN – Italy

11    <sup>‡</sup> Equal contribution to the work  
12    \* Corresponding Authors

13    Piergiorgio Comuzzo  
14    Tel: + 39 0432 55 8166  
15    e-mail: [piergiorgio.comuzzo@uniud.it](mailto:piergiorgio.comuzzo@uniud.it)

17    Paolo Sivilotti  
18    Tel: + 39 0432 55 8628  
19    e-mail: [paolo.sivilotti@uniud.it](mailto:paolo.sivilotti@uniud.it)

**Abstract**

Ribolla Gialla (RG) is a white grape variety used in the production of high-quality sparkling wines. It is widespread in North-East Italy, particularly Friuli Venezia Giulia (FVG), as well as Slovenia. Because of its limited area of cultivation, however, there is little information about the composition and chemical characteristics of the sparkling wines produced. This work used different analytical approaches to characterize thirty-three commercial sparkling RG wines from different areas of FVG. The characteristics included the overall volatile profile and content of terpenes, C<sub>13</sub>-norisoprenoids, lipids, and tryptophan metabolites. The aroma profile of RG wines was mainly characterized by fermentative esters and  $\beta$ -damascenone, whereas other norisoprenoids and varietal aromas were below the odor threshold. Appreciable amounts of certain fatty acids were found (e.g., palmitic acid), which could be potentially correlated with greater foam stability. However, high concentrations of tryptophan metabolites highlighted a higher risk of developing atypical ageing defects.

**KEYWORDS:** Ribolla Gialla; sparkling wine; volatile compounds; tryptophan metabolites; lipids

**Introduction**

Ribolla Gialla is a promising white grape variety that has recently been used for the production of premium sparkling wines, which have received good appreciation in international wine markets. Ribolla has been cultivated since ancient times in North-Eastern Italy (in the Friuli Venezia Giulia region), as well as Slovenia and the Ionian Islands (Kefalonia), where it is known as Rebula and Robola, respectively [1]. A certain number of studies have been carried out on the genetic identity and characterization of Ribolla grapes [1–4], but very few studies have dealt with the metabolomic profile of the wines [5, 6]. In addition, none of them have focused on the aromatic characteristics of sparkling wines obtained from Ribolla grapes, including their volatile organic compound (VOC) content, lipids, polyphenols, and tryptophan metabolites. All of these compounds are important because they may be linked with positive and negative aspects of wine quality.

VOCs are fundamental for determining the sensory characteristics of wine and its profile. These compounds are classified into four groups [7]: i) primary grape VOCs, which are present in the cells of grapes; ii) secondary grape aroma, which are formed during the pre-fermentation phases (crushing, pressing, and skin contact) or by thermal, chemical, and enzymatic reactions in must; iii) fermentation bouquet, which includes aroma compounds that form during alcoholic fermentation; iv) and maturation bouquet, which refers to the aroma compounds that develop when ageing the wine. Primary aroma is particularly important for the production of certain wine typologies, such as white sparkling wines. Furthermore, different aroma compounds may be produced after fermentation and ageing that further affect the aroma profile of the final wine, which are mainly due to pre-fermentative processes and yeast metabolism (e.g., C6-compounds, norisoprenoids, fatty acids, alcohols, esters, carbonyl compounds) [8–11].

Moreover, yeast autolysis takes place during the ageing of sparkling wine, which leads to the release of substances in the medium. Lipids represent a wide class of compounds among these substances [8]. Fatty acids may have a significant impact on the sensory properties of the wine.

1  
2  
3 65 A large part of the fatty acids originates from the firm tissues of the grapes, but the greatest  
4  
5 66 amount is formed during alcoholic fermentation. Therefore, fatty acids may be present in wine  
6  
7 67 in free or bound forms as ethyl esters. Both forms directly contribute to the flavor of the wine,  
8  
9 68 while unsaturated fatty acids such as oleic, linolenic, and linoleic acids act as precursors of C6-  
10  
11 69 aldehydes and alcohols with herbaceous notes, and are critical for yeast growth during  
12  
13 70 fermentation [12]. Another part of the sensory assessment of sparkling wine is the relationship  
14  
15 71 between the foaming properties and lipid compounds. Therefore, it has been reported that  
16  
17 72 medium-chain free fatty acids C8, C10, and C12 are negatively correlated with foamability,  
18  
19 73 while the ethyl esters of hexanoic, octanoic, and decanoic acids have a positive effect on the  
20  
21 74 formation and stability of the foam [13].  
22  
23  
24  
25

26 75 The essential amino acid tryptophan (TRP) and its metabolites, especially indole-3-acetic acid  
27  
28 76 (IAA), are considered to be potential precursors of 2-aminoacetophenone (2-AAP), which is an  
29  
30 77 aroma compound that causes an atypical ageing off-flavor (ATA) in *Vitis vinifera* wines. The  
31  
32 78 off-flavor is described with aroma descriptors such as “acacia blossom,” “furniture polish,”  
33  
34 79 “wet wool,” “mothballs,” or “fusel alcohol,” which are combined with a loss of the typical  
35  
36 80 bouquet of the grape variety [14]. Depending on the wine matrix, the detection threshold of 2-  
37  
38 81 AAP varies from 0.5 to 1.5  $\mu\text{g L}^{-1}$ . Strongly aromatic wines are able to integrate more than 1.5  
39  
40 82  $\mu\text{g L}^{-1}$  of 2-AAP, while meager wines might be rejected as tainted by ATA with less than 0.5  
41  
42 83  $\mu\text{g L}^{-1}$ . It is generally accepted that the ultimate cause of ATA development in white wines is a  
43  
44 84 stress reaction in the vineyard triggered by drought, nutritional deficiency, and other viticultural  
45  
46 85 factors, such as the time of harvest and leaf removal [14, 15].  
47  
48  
49  
50

51 86 Most papers on sparkling wines focus on the characterization of internationally known varieties  
52  
53 87 with a worldwide distribution [16–19], but there are few studies on local or less relevant  
54  
55 88 cultivars. Therefore, the aim of this work was to investigate the compositional profile of  
56  
57 89 different Ribolla Gialla sparkling wines produced in different DOC (*Denominazione di Origine*  
58  
59 90 *Controllata*) districts in the Friuli Venezia Giulia region in North-East Italy. The wines were  
60

characterized for their chemical composition (pH, titratable acidity, residual sugars, alcohol, and sulfur dioxide), volatile profile (overall profile of aroma compounds, free and bound terpenes, and norisoprenoids), and content of lipids and tryptophan metabolites. The results are critically discussed with the aim of defining the metabolomic profile of Ribolla Gialla sparkling wines.

## Materials and Methods

### *Reagents and materials*

Hydrogen peroxide (30% w/w), ethanol (96% v/v), ACS grade hydrochloric acid (37%), anhydrous sodium sulfate and citric acid were purchased from Carlo Erba Reagents (Milan, Italy). Sodium chloride (99.5%) was obtained from Honeywell Fluka (Morris Plains, New Jersey). HPLC-grade solvents dichloromethane, *n*-pentane, and methanol, LC-MS grade methanol, acetonitrile, 2-propanol, chloroform, formic acid, ammonium formate, ethyl heptanoate, 1-heptanol, 2-octanol, ethyl hexanoate- $d_{11}$ , 3-(2-hydroxy ethyl)-indole, kynurenic acid, D-tryptophan methyl ester, L-tyrosine ethyl ester, *N*-acetyl tyrosine ethyl ester, and 3,5-di-*tert*-4-butylhydroxytoluene (BHT) were obtained from Sigma-Aldrich (St. Louis, MO, USA).  $C_7$ - $C_{30}$  *n*-alkane solution in *n*-hexane was purchased from Supelco (Bellefonte, PA, USA), while cholesterol- $d_7$  and octadecanoic acid- $d_3$  were obtained from CDN Isotopes (Quebec, Canada). Finally, the glycosidase preparation (Rapidase Revelation Aroma) used for the determination of bound monoterpenes was obtained from Oenobrand (Montellier, France).

### *Wine samples*

Local wineries located in the North-East Italian region of Friuli Venezia Giulia provided thirty-three commercial Ribolla Gialla sparkling wines. The locations of the wineries are shown in Figure 1.

### 114 *Basic analysis of wine samples*

115 The basic quality control parameters (reducing sugars, alcoholic strength, total acidity, volatile  
116 acidity, pH, malic, lactic and tartaric acid) were determined by FTIR spectroscopy with a  
117 Winescan<sup>TM</sup> FT-120 instrument (FOSS, Hillerød, Denmark). All of the samples were analyzed  
118 two times, and the mean value of the two measurements was considered for the data analysis.

### 119 *Determination of free and bound terpenes and norisoprenoids by SPE-GC-MS*

120 Terpenes and C<sub>13</sub>-norisoprenoids in free and bound forms were extracted on Isolute® 500-mg,  
121 6-mL, C<sub>18</sub> SPE cartridges (Biotage, Uppsala, Sweden) according to the method reported by  
122 Comuzzo et al. [20]. A GCMS-QP-2010 system (Shimadzu, Kyoto, Japan) was used for GC-  
123 MS analyses. Volatile compounds were separated on a J&W DB-Wax capillary column (30 m  
124 × 0.25 mm i.d., 0.25 µm film thickness) provided by Agilent Technologies Inc. (Santa Clara,  
125 CA, USA) using the operating conditions described by Loira et al. [21].

126 Electron impact mass spectra were recorded at 70 eV and volatile compounds were  
127 tentatively identified by comparison of their mass spectra and retention times with those of  
128 standard compounds or by comparison of the mass spectrum with those reported in the Wiley  
129 6, NIST 21, and NIST 107 mass spectrum libraries (provided by the manufacturer). Linear  
130 retention indices were calculated according to the retention times of *n*-alkanes and compared  
131 with those reported in literature. The semi-quantitative analysis was based on the internal  
132 standard method using 1-heptanol (312 mg L<sup>-1</sup> in 96 % v/v ethanol) as the internal standard  
133 while considering the response factor to be equal to 1.00.

### 134 *Determination of volatile compounds*

135 The volatile compounds in the commercial wines were determined using two analytical  
136 approaches: liquid-liquid extraction (LLE) and solid-phase microextraction (SPME). Samples



were degassed before analysis by placing them in an ultrasonic bath (Falc Labsonic, Treviglio, Italy) for 2 min.

#### *Liquid-liquid extraction (LLE-GC-MS)*

Aroma compounds were determined as reported by Loira et al. [21] by mixing 5 mL of wine with 5 mL of a 30% (w/v) sodium chloride solution and 200  $\mu$ L of internal standard (ethyl heptanoate, 422 mg L<sup>-1</sup> in 96% v/v ethanol). The mixture was subjected to three extractions using 2.5 mL of pentane:dichloromethane (2:1 v/v) each. The organic phase was collected in a Pyrex tube, dehydrated with anhydrous sodium sulfate, and concentrated under nitrogen flow to a final volume of about 1 mL. The samples obtained were subjected to GC-MS analysis as reported previously[21].

#### *SPME-GC-MS/MS*

Degassed wine (1 mL) was spiked with 50  $\mu$ L of 2-octanol at 2.13 mg L<sup>-1</sup> (IS) in ethanol and placed in a 20-mL headspace vial containing 1.5 g of sodium chloride. Two technical replicates were prepared for each sample, along with a blank sample containing only sodium chloride. GC analysis was performed using a Trace GC Ultra gas chromatograph coupled with a TSQ Quantum Tandem mass spectrometer, which was upgraded to the XLS configuration and equipped with a Triplus autosampler (Thermo Fisher Scientific, Waltham, MA, USA). The method used was adopted by Carlin et al. [22]. Samples were incubated for 5 min at 35°C, and VOCs were extracted for 20 min with a 2-cm-long 50/30- $\mu$ m coated divinylbenzene/carboxen/polydimethylsiloxane (DVB/CAR/PDMS) SPME fiber (Supelco, Sigma Aldrich, Milan, Italy). After solid-phase microextraction, the fiber was desorbed for 3 min at 250°C in the GC system with the injector set in splitless mode. The fiber was reconditioned between each sample at 270°C for 7 min. Helium was used as a carrier gas at a flow rate of 1.2 mL min<sup>-1</sup>.

The GC oven was equipped with a 30-m  $\times$  0.25-mm VF-WAXms column (Agilent Technologies Inc., Santa Clara, CA, USA) with a film thickness of 0.25  $\mu$ m. The oven temperature was held for 2 min at 40°C after the injection, ramped at 6°C min<sup>-1</sup> up to 250°C, and held for 5 min. The ion source was set at 230°C, and electron impact mass spectra were recorded at 70 eV. Data acquisition and analyses were performed using the Xcalibur Workstation software supplied by the manufacturer.

#### *Determination of tryptophan metabolites by UHPLC-MS/MS*

Sparkling wine samples were filtered at 0.22  $\mu$ m by a Millex-GV filtration unit (Merc, Darmstadt, Germany) and directly collected in 2-mL HPLC amber vials. The analyses were performed on an Acquity UHPLC system provided with an autosampler and coupled with a XEVO TQMS mass spectrometer equipped with an electrospray source (Waters Corporation, Milford, MA, USA). Tryptophan metabolites were separated at a flow rate of 0.4 mL min<sup>-1</sup> on a 1.8- $\mu$ m 150  $\times$  2.1-mm Waters Acquity HSS T3 column (Waters Corporation), which was conditioned at 40°C. A linear gradient of water containing 0.1% formic acid (solvent A) and 0.1% formic acid in acetonitrile (solvent B) was used according to the conditions reported by Arapitsas et al.[23]. The injection volume was 10  $\mu$ L, and the wine samples were stored in the autosampler at 6°C during analyses.

The MS conditions included a capillary voltage of 3.5 kV in positive mode and -2.7 kV in negative mode. The ion source temperature was set at 150°C, while the desolvation temperature was 500°C. Nitrogen was used as the cone gas with a flow rate of 50 L h<sup>-1</sup>. The data were processed using Waters MassLynx (version 4.1) and TargetLynx software (Milford, MA, USA).

### 183 *Analysis of lipid molecules by UHPLC-MS/MS*

184 Lipid analysis was performed as described by Della Corte et al. [24] with slight modifications.  
185 An aliquot (3 mL) of degassed wine was introduced in a 50-mL Falcon tube and spiked with  
186 30  $\mu\text{L}$  of a methanolic solution containing cholesterol- $\text{d}_7$  and octadecanoic acid- $\text{d}_3$  (IS) at  
187 concentrations of  $1.01 \text{ mg mL}^{-1}$  and  $1.002 \text{ mg mL}^{-1}$ , respectively. Lipids were extracted two  
188 times in 21 mL of a chloroform–methanol solution (2:1 v/v) containing BHT ( $10 \text{ mg L}^{-1}$ ). The  
189 total lower lipid-rich layer was collected in 100-mL flasks, and the solvent was evaporated to  
190 dryness using a rotary evaporator. The samples were reconstituted in 300  $\mu\text{L}$  of acetonitrile/2-  
191 propanol/water (65:30:5 v/v/v) and filtered (0.22  $\mu\text{m}$ ) into 2-mL HPLC amber vials for  
192 UHPLC-MS/MS analysis.  
193 UHPLC separation was performed on a Dionex 3000 chromatograph (Thermo Fisher Scientific,  
194 Waltham, MA, USA) equipped with an autosampler and coupled with an API 5500 triple-  
195 quadrupole mass spectrometer (Sciex, Concord, Vaughan, ON, Canada), which was provided  
196 with an electrospray ion source (ESI). Lipids were separated on a 2.7- $\mu\text{m}$   $150 \times 2.1\text{-mm}$  RP  
197 Ascentis Express column (Sigma-Aldrich, Milan, Italy) set at  $55^\circ\text{C}$ . The injection volume was  
198 5  $\mu\text{L}$ , and the samples were stored in an autosampler at  $10^\circ\text{C}$  during the analyses. The mobile  
199 phase and chromatographic conditions are those described by Della Corte et al. [24], and the  
200 flow rate was  $0.260 \text{ mL min}^{-1}$ .  
201 The spray voltage of the ESI source was set at 5500 V for positive mode and  $-4500 \text{ V}$  for  
202 negative mode, and the source temperature was  $250^\circ\text{C}$ . The nebulizer (Gas 1) and heater gas  
203 (Gas 2) pressures were set at 40 and 20 psi, respectively. Ultra-high-purity nitrogen (99.999%)  
204 was used as both a curtain gas and collision gas (CAD) at 20 and 9 psi, respectively. Instrument  
205 control and data acquisition were performed by Analyst software (Applera Corporation,  
206 Norwalk, CT, USA), and the data were processed using MultiQuant, version 2.1 (Sciex,  
207 Concord, Vaughan, ON, Canada).

## 208 *Statistical analysis*

209 Statistica for Windows Version 8 (Statsoft, Tulsa, OK, USA) was used to calculate the means,  
210 standard deviations (SDs), coefficients of variation (CVs), minimum values (MIN), and  
211 maximum values (MAX) of the different analytical parameters determined for wine samples.

## 212 **Results and Discussion**

### 213 *Basic quality parameters*

214 Table 1 shows the results related to the basic chemical composition of the wines. The mean  
215 values, SDs, and CVs show that the samples are very homogeneous concerning alcoholic  
216 strength, titratable acidity, tartaric acid, pH, and even volatile acidity, which shows quite low  
217 values despite a slightly higher CV. This relatively homogeneous composition is probably  
218 related to the fact that the sparkling wines analyzed are all DOC productions and were  
219 manufactured in fulfillment of the standards reported in the DOC production guidelines.

220 However, some differences that are more relevant from a practical point of view were observed  
221 for residual sugars (min-max range: 6-19 g L<sup>-1</sup>), malic acid (0.01-3.79 g L<sup>-1</sup>), and lactic acid  
222 (0.00-2.75 g L<sup>-1</sup>). These differences could be related to the application of different winemaking  
223 protocols by the producers, leading to different styles of wine for Ribolla Gialla DOC, even if  
224 they are in agreement to the DOC production guidelines. In regard to sugars, most of the wines  
225 analyzed showed a residual sugar content of less than 12 g L<sup>-1</sup> or between 12 and 17 g L<sup>-1</sup>.  
226 According to the European Commission Regulation (EC) n. 607/2009 [25], these wines are  
227 classified as *brut* and *extra dry*, respectively. Only one sample had higher sugar content (about  
228 19 g L<sup>-1</sup>) and was categorized as *sec* (or *dry*).

229 The variations of malic and lactic acid highlight the different ways of managing malolactic  
230 fermentation (MLF), which are probably connected with the different storage times of the wines  
231 on yeast lees after refermentation. Most of the wines did not contain lactic acid in appreciable

amounts, which was probably due to the lack of malolactic fermentation. MLF generally occurs during ageing on the lees because of the positive effect that they have on promoting the growth of lactic acid bacteria [26]. For this reason, most winemakers would consider Ribolla Gialla sparkling wine as a fresh and young wine according to the data.

Ribolla Gialla is generally refermented via the Martinotti/Charmat short method, which involves refermentation in stainless steel autoclaves without contact with the lees at the end of refermentation. However, six wine samples had a higher content of lactic acid. This accounted for the increase of the CV calculated for lactic acid in Table 1 and was combined with the decrease of malic acid concentration. This evidence may be related to the manufacture of certain Ribolla Gialla wines, with some products being kept in contact with the lees after refermentation in an autoclave. Such *sur-lies* aging may be prolonged for up to several months (2-4 months on average). In addition, certain winemakers also produce Ribolla Gialla by the traditional refermentation method in bottles.

#### *Wine aroma profile*

Table 2 shows the results of the qualitative and quantitative determination of the aroma compounds detected in wine samples by the different analytical techniques. Concerning non-varietal aromas, a total number of fifty-eight volatiles were tentatively identified in the wines, including acids, alcohols, esters, C6 compounds, diols, and carbonyls. Fatty acids are some of the most representative volatiles in the wines analyzed, and generally, they are described as having cheese, rancid, and fatty notes. If their amounts are higher than the odor detection threshold (ODT), they may negatively affect the organoleptic characteristics of the wines.

In the samples analyzed, the average concentration of fatty acid was often close and sometimes higher than their ODT. Nevertheless, their odor activity value (OAV), which is the ratio between the concentration and ODT, was in the range of 1-4 for many of the compounds analyzed. The exception was octanoic acid, which is the most representative fatty acid in the

257 volatile composition of Ribolla wines ( $OAV \approx 30$ ). These results suggest that fatty acids might  
258 have minor relevance in the aromatic characteristics of Ribolla Gialla. Acetic acid (which is  
259 responsible for the typical vinegar off-flavor) was found to be lower than ODT in all samples  
260 analyzed.

261 Interestingly, the variability among the samples (i.e., the CVs) for fatty acids and volatile  
262 compounds in general (Table 2) was higher than that detected for the basic parameters in Table  
263 1. This highlights a certain level of differentiation among Ribolla Gialla DOC wines, which is  
264 probably related to a number of variables, such as those typically connected with the  
265 development of wine aromas. Concerning fatty acids, for instance, the differences among the  
266 samples might have resulted from the different origin of the grapes used, the amount of lipid  
267 substances in the musts, the diverse winemaking conditions, and the yeast strains used for  
268 fermentations [27].

269 Higher alcohols (HA) are also important volatiles among the compounds listed in Table 2. They  
270 are produced by yeasts during alcoholic fermentation as products of amino acid metabolism  
271 [11]. Their contributions to wine aroma vary from honey, rose, and floral characters (2-  
272 phenylethyl alcohol and benzyl alcohol) to pungent and solvent-like smells (1-propanol, 1-  
273 butanol, 2- and 3-methyl-1-butanol) [28], and the effects depend on their concentration [9]. The  
274 average amounts of HA observed in the wines analyzed were generally found to be lower than  
275 their ODT except for 2- and 3-methyl-1-butanol and 2-phenylethanol (ODT: 40 and 10 mg L<sup>-1</sup>,  
276 respectively). This composition may be linked with the yeast strains used, as well as the  
277 winemaking conditions adopted and the amino acidic composition of the grapes. The presence  
278 of significant amounts of 2-phenylethanol is very interesting for its fresh rose-like odor [29],  
279 even if its OAV range was about 1-6.

280 Esters are the most representative class of volatile compounds found in Ribolla Gialla sparkling  
281 wines. Isoamyl acetate (3-methyl-1-butanol acetate), ethyl hexanoate, ethyl lactate, ethyl  
282 octanoate, diethyl succinate, and 2-phenylethyl acetate showed the highest concentrations. The

compounds eluted at higher retention times (e.g., diethyl malate) presumably had a minor impact on the aroma profile because of their low volatility. Some esters, such as ethyl butanoate and ethyl hexanoate, were close to or higher than their ODT (20  $\mu$ g and 65  $\mu$ g) in all the samples analyzed. Others such as isoamyl acetate, 2-phenylethyl acetate, and hexyl acetate were detected in only certain wines with concentrations higher than their ODT. Compounds such as ethyl hexanoate and isoamyl acetate were found to have significant OAVs (e.g., up to 18 for ethyl hexanoate, 85 for hexyl acetate, and approximately 200 for isoamyl acetate). This observation is very interesting from a sensory point of view because esters generally confer floral, fresh, and fruity notes (rose, banana, pear, green apple) to the wines [28, 30, 31], which are generally recognized as typical in Ribolla wines. For this reason, esters appeared to be an important component of the volatile profile of Ribolla Gialla.

Among esters, ethyl lactate and diethyl succinate are considered “ageing esters” (AE), and their content in wines generally increases during ageing and after malolactic fermentation [29, 32]. The same consideration is true for diethyl malate, but this compound is normally detected less frequently in wines. The concentration of AE was generally found to be lower than the ODT in most of the wines analyzed. However, in some samples, ethyl lactate had higher concentrations than its odor threshold (60 mg/L) in accordance with the amount of malic and lactic acid. In fact, the highest concentration of these esters was observed in the same products in which malolactic fermentation was hypothesized to have taken place (i.e., the wines where malic acid was not detected), as shown in Table 1. This may confirm that certain Ribolla Gialla sparkling wines are produced with a more or less prolonged period of *sur-lies* ageing or by refermenting them in bottle by the traditional method.

Carbonyls, diols, and C6 compounds appeared not to have an important contribution to the aromatic characterization of the samples, and most of the cases, they were detected at levels below their ODT. Hexanal was found to overcome its ODT in some samples, but at the low concentrations detected, it may or may not contribute to the fresh vegetal notes of Ribolla wines.



Table 3 reports the detected varietal aromas (terpenes and C<sub>13</sub>-norisoprenoids) in free and bound form in the samples. Terpenes are well known to confer floral odors to wines [29]. In Ribolla, they were especially found in the free form, but in most cases, their average concentrations were below their ODT with a few exceptions for linalool and geraniol. Although it was present below its ODT,  $\alpha$ -terpineol was the most abundant free terpenic alcohol in the wines. Interestingly, even if Ribolla Gialla cannot be considered as an aromatic variety, the concentrations detected for  $\alpha$ -terpineol and linalool were higher than those normally observed in certain international white varieties, such as Pinot blanc, Pinot Gris, Chardonnay, and Sauvignon blanc [19]. Also  $\beta$ -citronellol, was found at lower levels with respect to its ODT (100  $\mu\text{g L}^{-1}$ ) [33], it showed similar average values to those of some Riesling wines [19].

Considering the glycosylated forms, only geraniol was significantly detected as both free and bound terpenol with a prevalence of the latter (combined) form of the molecule. According to other experiments, geraniol (with minor traces of nerol) was the only terpenic alcohol normally found in Ribolla Gialla grapes [34]. For this reason, in Ribolla Gialla, linalool,  $\alpha$ -terpineol, and  $\beta$ -citronellol may be formed from such terpenic alcohols during grape processing. Citronellol, for instance, is reported to be produced by *S. cerevisiae* from geraniol and nerol during alcoholic fermentation, and other formation pathways have also been suggested for linalool (from geraniol) and  $\alpha$ -terpineol (by cyclization of nerol) [35]. For both free and bound geraniol, the average values were generally lower than the ODT reported for the alcohol, but some of the samples showed significant levels of total geraniol (Table 3). Considering that the bound form generally prevails for this terpenol, one interesting way to improve the volatile profile of these wines might be the application of techniques to increase the release of free geraniol from its glycosides in the production process of Ribolla Gialla sparkling wines.

In contrast to terpenes, which did not appear particularly characteristic for the aroma profile of the wines, some C<sub>13</sub>-norisoprenoids were detected in appreciable concentrations (Table 3). TDN and  $\beta$ -damascenone, for instance, overcame their sensory thresholds in different samples.



Norisoprenoids are generally not connected to specific aromatic grapevine varieties and generally have a low odor threshold. The ODT of  $\beta$ -damascenone is 0.04-0.06  $\mu\text{g L}^{-1}$  in a model dilute alcohol solution [29], while a relatively wide range of ODTs was found in wine: Sefton et al. [36] reported 0.14  $\mu\text{g L}^{-1}$  in deodorized white wine, 0.85 – 2.10  $\mu\text{g L}^{-1}$  in deodorized red wine, and 7.00  $\mu\text{g L}^{-1}$  in red wine. The presence of  $\beta$ -damascenone may be particularly interesting for Ribolla wines for not only the relatively high concentrations found but also because it is reported to modify the sensory perception of some esters. Escudero et al.[37] found that the addition of low levels of  $\beta$ -damascenone (0.85  $\mu\text{g L}^{-1}$ ) to a solution of esters increased the fruity notes of the mixture, while higher levels (3.5  $\mu\text{g L}^{-1}$ ) accounted for the development of strong raisin/dry-plum odors.

The amount of norisoprenoids is also dependent on the winemaking conditions. In general,  $\beta$ -damascenone is observed at higher concentrations in young wines [36], whereas for 1,1,6-trimethyl-1,2-dihydronaphthalene (TDN), a positive correlation between its occurrence in wine and ageing was reported.[38] Their odor descriptors range from floral (e.g., vitispiranes) and fruity-honey scents ( $\beta$ -damascenone) to notes of kerosene and petrol (for TDN) [39]. The presence of TDN among the compounds listed in Table 3 may represent further evidence about the ageing of certain Ribolla Gialla sparkling wines after refermentation.

Another interesting aspect related to norisoprenoids is the high concentration observed for some of them in bound form, particularly 3-oxo- $\alpha$ -ionol and 3-hydroxy- $\beta$ -damascone, which are the most representative compounds. As for terpenes, Ribolla Gialla appeared to have an unexpressed aromatic potential for norisoprenoids. Thus, from a technical point of view, the optimization of winemaking process to maximize the release of this bound aromatic potential might be an interesting way to improve the quality of Ribolla Gialla sparkling wines.

### 358 *Metabolomic fingerprint of tryptophan metabolites and indoles*

359 The second group of metabolites analyzed was the amino acid tryptophan and its catabolites,  
360 as shown in Table 4. There were twenty-eight compounds detected, among which the sulfonated  
361 derivates of indole 3-lactic acid, tryptophol, indole 3-lactic acid glucoside, and indole 3-acetic  
362 acid are presented. Sulfonation is the addition of a sulfonic acid group ( $-\text{SO}_3\text{H}$ ) to an organic  
363 compound and is a widespread industrial process used in a diverse range of products. It also  
364 has a major function in modulating biological activities that are known to occur in wine, which  
365 involve several metabolites such as polyphenols and indoles [23].

366 Sparkling wines are well known to have much lower amounts of tryptophan than other wines,  
367 which is probably due to the second fermentation [23]. Tryptophan values for the commercial  
368 samples ranged from 0.01 to 0.39  $\text{mg L}^{-1}$  with a mean value of 0.13  $\text{mg L}^{-1}$ , and which similarly  
369 occurred for its ethyl ester (0.44  $\text{mg L}^{-1}$  mean value). TOL- $\text{SO}_3\text{H}$  had higher amounts among  
370 all the other tryptophan metabolites with a mean value of 3.73  $\text{mg L}^{-1}$ . The unsulfonated form  
371 of tryptophol varied from 0.01 to 1.08  $\text{mg L}^{-1}$  in all samples. This could mean that the  
372 sulfonated/unsulfonated TOL ratio that favored TOL- $\text{SO}_3\text{H}$  may also give rise to further  
373 products similar to 2-AAP.

374 The ILA (0.03  $\text{mg L}^{-1}$ ) and ILA- $\text{SO}_3\text{H}$  (8.53  $\text{mg L}^{-1}$ ) pairing showed similar behavior to the  
375 previously described TOL and TOL- $\text{SO}_3\text{H}$  pairing. Arapitsas et al. [23], found that young red  
376 wines and rosé wines had a much higher concentration of ILA and especially ILA-GLU, which  
377 is synthesized by plants [40]. This was probably due to maceration with the skins during  
378 winemaking. In contrast, white and especially sparkling wines had the lowest concentrations  
379 due to soft pressing and secondary fermentation, which necessitate further nitrogen  
380 consumption.

381 Sulfonated indole 3-acetic acid was detected at concentrations between 0.13 and 0.39  $\text{mg L}^{-1}$ ,  
382 while its parent compounds were detectable at very low concentrations. Hoenicke et al. [41]  
383 showed that the sulfonation of indoles in a model wine solution could be responsible for their

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

degradation and the formation of aromatic aminobenzenes such as 2-aminoacetophenone, which are responsible for some of the heavy aromatic characteristics of white wines. On other hand, the structural similarity of TOL to IAA could also cause the detection of 2-AAP or other similar aromatic compounds in wine. Because of the lack of anthocyanins and the low flavanol content in white wines, the sulfonation of indoles can increase the risk of developing an atypical off-flavor. Thus, the sparkling wines produced from the Ribolla Gialla variety might have a higher tendency to develop atypical ageing defects, which could also be promoted by inappropriate storage temperature.

*Metabolomic fingerprint of lipids*

Table 5 shows the twenty-nine lipid compounds found in the samples of commercial sparkling wines according to UHPLC-MS-MS. Most of the compounds found are saturated long-chain fatty acids (LCFAs, more than 12 carbon atoms), although the mid-chain fatty acids (MCFAs, 4–12 carbons) and their esters have a major influence on the organoleptic properties of wine. In addition to the fact that lipids are an integral part of solid grape tissues, they are also an important building block in wine yeasts, where the majority of LCFAs are esterified with glycerol or glycerophosphate to form mono-, di-, and tri-acylglycerides or glycerophospholipids, respectively [42].

The two most abundant LCFAs detected in the samples were palmitic acid (C16:0), which ranged from 5.21 to 11.49 mg L<sup>-1</sup> with a mean value of 9.24 mg L<sup>-1</sup>, and stearic acid (C18:0), which ranged from 4.43 to 8.40 mg L<sup>-1</sup> (mean value: 6.49 mg L<sup>-1</sup>). Previous studies showed that after the beginning of second fermentation, which is after the inoculation of the medium with yeast, the concentration of saturated fatty acids C16:0 slightly increased, while the proportion of C18:0 remained constant. However, after the end of the growth phase of the yeasts, the distributions of both saturated fatty acids decreased exponentially [43].

Unlike saturated fatty acids (SFA), the values of unsaturated fatty acids (UFA) were lower in the wine samples (*cis*-11-eicosanoic acid 0.11 mg L<sup>-1</sup>, myristoleic acid 0.05 mg L<sup>-1</sup>, linoleic acid 0.04 mg L<sup>-1</sup>, palmitoleic acid 0.02 mg L<sup>-1</sup>, and linolenic acid 0.01 mg L<sup>-1</sup>). UFAs also had a much smaller proportion than SFAs. Because of their membrane fluidity, wine yeasts are able to modify the UFA/SFA ratio in their membrane composition in response to stress conditions, such as the presence of ethanol or cooler fermentation conditions under which fermentation takes place, especially for white wines. Normally, the UFA/SFA ratio is close to 1:1, but a deficiency of UFA leads to impaired biosynthesis of phospholipids and consequently stuck fermentations [12].

Similarly, Pueyo et al. [44] studied the effect of the total contents of linolenic acid and palmitic acid as compounds that best define foam stability in wines and foam height in the cavas, respectively. They found a positive correlation between these compounds and the measured properties, meaning that the wines with a greater amount of these fatty acids in the lipidic fraction could form more foam with high stability than those with lower concentrations. Since the analyzed samples in the present study contained a greater concentration of palmitic acid compared to the linolenic acid, it can be expected that foam formation of the Ribolla Gialla sparkling wines will be increased, which is a key parameter for determining the quality of sparkling wines.

In conclusion, Ribolla Gialla sparkling wines showed homogeneous characteristics concerning basic parameters (e.g., sugar content, alcoholic strength, pH, and titratable acidity) as well as their aroma composition. Sparkling Ribolla wines are generally characterized by low levels of free terpenols, and their aroma seems to be mostly characterized by the development of volatile esters and  $\beta$ -damascenone during fermentations, storage, and processing. The presence of these compounds is probably connected with the fresh and fruity notes that normally characterize the sparkling wines produced from Ribolla grapes.

1  
2  
3 433 A minor amount of malolactic fermentation was found, which reflects that two distinct wine  
4  
5 434 styles coexist for Ribolla sparkling wines. The first characterizes young wines normally  
6  
7 435 produced by refermentation in stainless-steel autoclaves (Martinotti/Charmat short  
8  
9 436 refermentation method), and the second is based on a more or less prolonged ageing period on  
10  
11 437 refermentation lees in an autoclave or bottles (according to the traditional refermentation  
12  
13 438 method). The analysis of lipid molecules, tryptophan, and its metabolites highlighted other  
14  
15 439 interesting features of Ribolla sparkling wines from a practical point of view. Ribolla Gialla  
16  
17 440 shows a high ratio between saturated (e.g., palmitic acid) and unsaturated fatty acids (e.g.,  
18  
19 441 linolenic acid), which may lead to higher foam height in the sparkling wines produced and  
20  
21 442 represents one of the key quality features of sparkling wines in general. In contrast, due to the  
22  
23 443 high amounts of certain tryptophan metabolites, Ribolla Gialla could be prone to the formation  
24  
25 444 of atypical ageing aromas. Considering that this specific Italian product has only recently been  
26  
27 445 appearing in wine markets, further investigations should carefully consider these results to  
28  
29 446 properly address the production practices and techniques towards producing high quality  
30  
31 447 products, thus increasing the local and international competitiveness of Ribolla Gialla sparkling  
32  
33 448 wines.  
34  
35  
36  
37  
38  
39  
40

#### 41 449 **Acknowledgments**

42  
43  
44 450 This research was financially supported by Friuli Venezia Giulia Region (LR 29 Dec 2016, n.  
45  
46 451 25, art. 3, paragraphs 6-10. Project Title: Maturation parameters and optimization of the  
47  
48 452 agronomic and enological techniques for the production of quality sparkling wines, Jul 2017 –  
49  
50 453 Dec 2019) and by the ADP 2017 project and PhD scholarship (D.S.) funded by the Autonomous  
51  
52 454 Province of Trento. The authors are grateful to Dr. Matteo Corradin for his cooperation in aroma  
53  
54 455 analyses.  
55  
56  
57  
58  
59  
60

## References

1. Rusjan D, Jug T, Štajner N (2010) Evaluation of genetic diversity: Which of the varieties can be named “Rebula” (*Vitis vinifera* L.)? *Vitis* 49:189–192
2. De Lorenzis G, Imazio S, Rusjan D, Vouillamoz J F, Nikolaou N, Failla O, Scienza A (2013) Genetic investigation of grapevine varieties ‘Ribolla Gialla’ (Italy), ‘Rebula’ (Slovenia) and ‘Robola’ (Ionian Islands). *Sci Hortic* 150:425–431.
3. Imazio S, Scienza A, Brancadoro L, de Lorenzis G, Failla O (2016) Evidence for a Sympatric Origin of Ribolla gialla, Gouais Blanc and Schiava cultivars (*V. vinifera* L.). *South African J Enol Vitic* 35:149–156.
4. Imazio S, De Lorenzis G, Scienza A, Failla O, Vouillamoz J, Korosec-Koruza Z, Rusjan D, Nikolao N (2014) ‘Ribolla Gialla’ from North Eastern Italy, ‘Rebula’ from Northern Balkans and ‘Robola’ from Ionian Islands; Do They Belong to the Same Population Variety or Are They Genetically Different? *Acta Horti* 1046:645–652.
5. Bavčar D, Baša Česnik H, Čuš F, Vanzo A, Gašperlin L, Košmerl T (2011) Impact of alternative skin contact procedures on the aroma composition of white wine. *South African J Enol Vitic* 32:190–203.
6. Bavčar D, Baša Česnik H, Čuš F, Košmerl T (2011) The influence of skin contact during alcoholic fermentation on the aroma composition of Ribolla Gialla and Malvasia Istriana *Vitis vinifera* (L.) grape wines. *Int J Food Sci Technol* 46:1801–1808.
7. Rapp A, Versini G (1995) Influence of nitrogen compounds in grapes on aroma compounds of wines. *Dev Food Sci* 37:1659–1694
8. Pueyo E, Martínez-Rodríguez A, Polo MC, Santa-María G, Bartolomé B (2000) Release of Lipids during Yeast Autolysis in a Model Wine System. *J Agric Food Chem* 46:116–122.
9. Swiegers JH, Bartowsky EJ, Henschke PA, Pretorius IS (2005) Yeast and bacterial modulation of wine aroma and flavour. *Aust J Grape Wine Res* 11:139–173.

- 483 10. Callejón RM, Margulies B, Hirson GD, Ebeler SE (2012) Dynamic changes in volatile  
484 compounds during fermentation of Cabernet Sauvignon grapes with and without skins.  
485 Am J Enol Vitic 63:301–312.
- 486 11. Hirst MB, Richter CL (2016) Review of Aroma Formation through Metabolic Pathways  
487 of *Saccharomyces cerevisiae* in Beverage Fermentations. Am J Enol Vitic 67:361–370.
- 488 12. Waterhouse AL, Sacks GL, Jeffery DW (2016) Understanding Wine Chemistry. John  
489 Wiley & Sons, Chichester (UK).
- 490 13. Gallart M, Lopez-Tamames E, Suberbiola G, Buxaderas Susana (2002) Influence of  
491 Fatty Acids on Wine Foaming. J Agric Food Chem 50:7042–7045.
- 492 14. Hoenicke K, Simat TJ, Steinhart H, Köhler HJ, Schwab A (2001) Determination of Free  
493 and Conjugated Indole-3-Acetic Acid, Tryptophan, and Tryptophan Metabolites in  
494 Grape Must and Wine. J Agric Food Chem 49:5494–5501.
- 495 15. Schneider V (2014) Atypical Aging Defect: Sensory Discrimination, Viticultural Causes,  
496 and Enological Consequences. A Review. Am J Enol Vitic 65:277–284.
- 497 16. Dziadas M, Jeleń HH (2010) Analysis of terpenes in white wines using SPE-SPME-  
498 GC/MS approach. Anal Chim Acta 677:43–49.
- 499 17. Louw L, Tredoux AGJ, Rensburg PV, Kidd M, Naes T, Nieuwoudt HH (2010)  
500 Fermentation-derived Aroma Compounds in Varietal Young Wines from South Africa.  
501 South African J Enol Vitic 31:213–225.
- 502 18. D’Onofrio C, Matarese F, Cuzzola A (2017) Study of the terpene profile at harvest and  
503 during berry development of *Vitis vinifera* L. aromatic varieties Aleatico, Brachetto,  
504 Malvasia di Candia aromatica and Moscato bianco. J Sci Food Agric 97:2898–2907.
- 505 19. Vilanova M, Genisheva Z, Graña M, Oliveira JM (2013) Determination of Odorants in  
506 Varietal Wines from International Grape Cultivars (*Vitis vinifera*) Grown in NW Spain.  
507 South African J Enol Vitic 34:212–222.
- 508 20. Comuzzo P, Marconi M, Zanella G, Querzè M (2018) Pulsed electric field processing of



- white grapes (cv. Garganega): Effects on wine composition and volatile compounds. Food Chem 264:16–23.
21. Loira I, Morata A, Comuzzo P, Callejo MJ, González C, Calderón F, Suárez-Lepe JA (2015) Use of *Schizosaccharomyces pombe* and *Torulaspora delbrueckii* strains in mixed and sequential fermentations to improve red wine sensory quality. Food Res Int 76:325–333.
22. Carlin S, Vrhovsek U, Franceschi P, Lotti C, Bontempo L, Camin F, Toubiana D, Zottele F, Toller G, Fait A, Mattivi F (2016) Regional features of northern Italian sparkling wines, identified using solid-phase micro extraction and comprehensive two-dimensional gas chromatography coupled with time-of-flight mass spectrometry. Food Chem 208:68–80.
23. Arapitsas P, Guella G, Mattivi F (2018) The impact of SO<sub>2</sub> on wine flavanols and indoles in relation to wine style and age. Sci Rep 8:858.
24. Della Corte A, Chitarrini G, Di Gangi IM, Masuero D, Soini E, Mattivi F, Vrhovsek U (2015) A rapid LC–MS/MS method for quantitative profiling of fatty acids, sterols, glycerolipids, glycerophospholipids and sphingolipids in grapes. Talanta 140:52–61.
25. Commission Regulation (EC) No 607/2009 (2009) Laying down certain detailed rules for the implementation of Council Regulation (EC) No 479/2008 as regards protected designations of origin and geographical indications, traditional terms, labeling and presentation of certain wine sector products. Off J Eur Communities 60–139.
26. Patynowski RJ, Jiranek V, Markides AJ (2002) Yeast viability during fermentation and sur lie ageing of a defined medium and subsequent growth of *Oenococcus oeni*. Aust J Grape Wine Res 8:62–69.
27. Lambrechts MG, Pretorius IS (2000) Yeast and its Importance to Wine Aroma - A Review. South African J Enol Vitic 21:97-129.
28. Fariña L, Villar V, Ares G, Carrau F, Dellacassa E, Boido E (2015) Volatile composition



- 535 and aroma profile of Uruguayan Tannat wines. Food Res Int 69:244–255.
- 536 29. Ribéreau-Gayon P, Glories Y, Maujean A, Dubourdieu D (2006) Handbook of Enology.
- 537 Vol. 2. The Chemistry of Wine Stabilization and Treatments, 2<sup>nd</sup> ed. John Wiley & Sons,
- 538 Chichester (UK).
- 539 30. Ferreira V, Fernandez P, Peiia C, Escudero A (1995) Investigation on the Role Played
- 540 by Fermentation Esters in the Aroma-of Young Spanish Wines by Multivariate Analysis.
- 541 J Sci Food Agric 67:381–392.
- 542 31. Saerens SMG, Delvaux FR, Verstrepen KJ, Thevelein JM (2010) Production and
- 543 biological function of volatile esters in *Saccharomyces cerevisiae*. Microb Biotechnol
- 544 3:165–177.
- 545 32. Boido E, Medina K, Farina L, Carrau F, Versini G, Dellacassa E (2009) The Effect of
- 546 Bacterial Strain and Aging on the Secondary Volatile Metabolites Produced during
- 547 Malolactic Fermentation of Tannat Red Wine. J Agric Food Chem 57:6271–6278.
- 548 33. Guth H (1997) Quantitation and Sensory Studies of Character Impact Odorants of
- 549 Different White Wine Varieties. J. Agric. Food Chem 43:35–37.
- 550 34. Voce S, Pizzamiglio G, Comuzzo P, Sivilotti S, Mosetti D, Bigot G, Lonardi A (2018)
- 551 Defogliazione e aromi della Ribolla Gialla. Vigne, Vini Qual 38–41.
- 552 35. Mateo JJ, Jimenez M (2000) Monoterpenes in grape juice and wines. J Chromatogr A
- 553 881:557–567.
- 554 36. Sefton MA, Skouroumounis GK, Elsey GM, Taylor DK (2011) Occurrence, sensory
- 555 impact, formation, and fate of damascenone in grapes, wines, and other foods and
- 556 beverages. J Agric Food Chem 59:9717–9746.
- 557 37. Escudero A, Campo E, Fariña L, Cacho J, Ferreira V (2007) Analytical Characterization
- 558 of the Aroma of Five Premium Red Wines . Insights into the Role of Odor Families and
- 559 the Concept of Fruitiness of Wines. J Agric Food Chem 55:4501–4510.
- 560 38. Belitz HD, Grosch W, Schieberle P (2004) Food Chemistry, 3<sup>rd</sup> ed. Springer, Berlin.

- 1  
2  
3 561 39. Black CA, Parker M, Siebert TE, Capone DL, Francis IL (2015) Terpenoids and their  
4  
5 562 role in wine flavour: Recent advances. *Aust J Grape Wine Res* 21:582–600.  
6  
7 563 40. Mattivi F, Arapitsas P, Perenzoni D, Guella G (2015) In: Ebeler SB, Sacks G, Vidal S,  
8  
9 564 Winterhalter P (eds) *Advances in Wine Research*. ACS Publications, Washington (USA)  
10  
11 565 41. Hoenicke K, Borchert O, Ning K G, Simat TJ (2002) Untypical Aging Off-Flavor &quot;  
12  
13 566 in Wine: Synthesis of Potential Degradation Compounds of Indole-3-acetic Acid and  
14  
15 567 Kynurenine and Their Evaluation as Precursors of 2-Aminoacetophenone. *J Agric Food*  
16  
17 568 *Chem* 50:4303–4309.  
18  
19 569 42. Ramsay AM, Douglas LJ (1979) Effects of Phosphate Limitation of Growth on the Cell-  
20  
21 570 wall and Lipid Composition of *Saccharomyces cerevisiae*. *J Gen Microbiol* 110:185–  
22  
23 571 191.  
24  
25 572 43. Piton F, Charpentier M, Troton D (1988) Cell Wall and Lipid Changes in *Saccharomyces*  
26  
27 573 *cerevisiae* during Aging of Champagne Wine. *Am J Enol Vitic* 39:221–225.  
28  
29 574 44. Pueyo E, Martin-Alvarez P, Polo MC (1995) Relationship Between Foam Characteristics  
30  
31 575 and Chemical Composition in Wines and Cavas (Sparkling Wines). *Am J Enol Vitic*  
32  
33 576 46:518–524.  
34  
35 577 45. Vilanova M, Sieiro C (2006) Determination of free and bound terpene compounds in  
36  
37 578 Albariño wine. *J Food Compos Anal* 19:694–697.  
38  
39 579 46. Francis IL, Newton JL (2005) Determining wine aroma from compositional data. *Aust J*  
40  
41 580 *Grape Wine Res* 11:114–126  
42  
43 581 47. Sansone-Land A, Takeoka GR, Shoemaker CF (2014) Volatile constituents of  
44  
45 582 commercial imported and domestic black-ripe table olives (*Olea europaea* ). *Food Chem*  
46  
47 583 149:285–295.  
48  
49 584 48. Zhu J, Xiao Z (2018) Characterization of the Major Odor-Active Compounds in Dry  
50  
51 585 Jujube Cultivars by Application of Gas Chromatography–Olfactometry and Odor  
52  
53 586 Activity Value. *J Agric Food Chem* 66:7722–7734.  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

587 49. Ferreira V, Lopez R, Cacho JF (2000) Quantitative determination of the odorants of  
588 young red wines from different grape varieties. J Sci Food Agric 80:1659–1667.  
589 50. Elsharif SA, Buettner A (2018) Structure – Odor Relationship Study on Geraniol, Nerol,  
590 and Their Synthesized Oxygenated Derivatives. J Agric Food Chem 66:2324–2333.  
591  
592

For Peer Review

## Figure Captions

**Figure 1.** Geographical position of Friuli Venezia Giulia region. Grey areas mark the municipalities in which the wineries that supplied wine samples are located. The number of samples/wineries for each municipality are also reported.

For Peer Review

**Table 1. Chemical composition of Ribolla Gialla commercial wines. SD: standard deviation; MIN: minimum value; MAX: maximum value; CV: coefficient of variation.**

Parameter	Mean ± SD	MIN	MAX	CV
alcoholic strength (% v/v)	11.68 ± 0.64	10.40	12.70	0.05
reducing sugars (g L <sup>-1</sup> )	10.64 ± 2.91	6.00	18.87	0.27
titratable acidity (g L <sup>-1</sup> )	5.94 ± 0.48	5.00	7.25	0.08
volatile acidity (g L <sup>-1</sup> )	0.31 ± 0.06	0.20	0.48	0.21
pH	3.22 ± 0.12	3.03	3.51	0.04
malic acid (g L <sup>-1</sup> )	2.08 ± 0.93	n.d. <sup>a</sup>	3.79	0.45
lactic acid (g L <sup>-1</sup> )	0.26 ± 0.63	n.d.	2.75	2.45
tartaric acid (g L <sup>-1</sup> )	2.83 ± 0.36	2.30	3.93	0.13

<sup>a</sup> n.d.: not detected

**Table 2. Non-variatal aroma compounds detected in Ribolla Gialla commercial sparkling wines by different analytical approaches. Concentrations are expressed in  $\mu\text{g L}^{-1}$ .**

Compounds	AM <sup>a</sup>	IM <sup>b</sup>	Mean $\pm$ SD <sup>c</sup>	MIN <sup>d</sup>	MAX <sup>e</sup>	CV <sup>f</sup>	ODT ( $\mu\text{g L}^{-1}$ ) <sup>g</sup>
<b><u>Acids</u></b>							
acetic acid	LLE	MS RI S	2624 $\pm$ 1206	1000	6388	0.46	200000 [33]
2-methylpropanoic acid	LLE	MS RI S	376 $\pm$ 168	n.d. <sup>h</sup>	877	0.45	230 [28]
butanoic acid	LLE	MS RI S	423 $\pm$ 126	218	812	0.30	10000 [33]
3-methylbutanoic acid	LLE	MS RI S	228 $\pm$ 242	n.d.	1103	1.06	250 [28]
hexanoic acid	LLE	MS RI S	4196 $\pm$ 1298	2111	7817	0.31	3000 [33]
heptanoic acid	LLE	MS RI	56 $\pm$ 70	n.d.	224	1.24	-
octanoic acid	LLE	MS RI S	6923 $\pm$ 2390	2878	15070	0.35	500 [45]
nonanoic acid	SPME	MS S	454 $\pm$ 96	239	708	0.21	-
decanoic acid	LLE	MS RI S	872 $\pm$ 736	n.d.	4043	0.84	1000 [45]
benzoic acid	SPME	MS S	5 $\pm$ 3	2	17	0.61	-
dodecanoic acid	SPME	MS S	3 $\pm$ 2	1	9	0.67	-
hexadecanoic acid	LLE	MS RI	2065 $\pm$ 2072	n.d.	7714	1.00	-
<b><u>Alcohols</u></b>							
2-methyl-1-propanol	LLE	MS RI	8784 $\pm$ 2825	5569	18165	0.32	40000 [33]
1-butanol	LLE	MS RI S	210 $\pm$ 103	n.d.	515	0.49	40000 [46]
2- and 3-methyl-1-butanol	LLE	MS RI S	125666 $\pm$ 24151	85297	190584	0.19	40000 [28]
3-ethoxy-1-propanol	LLE	MS RI	172 $\pm$ 349	n.d.	1651	2.03	-
2-phenylethanol	LLE	MS RI S	23645 $\pm$ 12214	9954	63961	0.52	10000 [33]
<b><u>C6 compounds</u></b>							
1-hexanol	LLE	MS RI S	957 $\pm$ 231	603	1443	0.24	2500 [28]

1	<i>trans</i> -3-hexen-1-ol	SPME	MS S	3 ± 1	2	9	0.41	1000	[28]
2	<i>trans</i> -2-hexen-1-ol	SPME	MS S	7 ± 6	n.d.	35	0.84	-	
3	<i>cis</i> -3-hexen-1-ol	LLE	MS RI S	39 ± 118	n.d.	646	3.01	400	[28]
4	hexanal	SPME	MS S	2 ± 7	n.d.	42	3.19	5	[47]
5	<i>trans</i> -2-hexenal	SPME	MS S	4 ± 1	1	8	0.36	82	[48]
6									
7									
8									
9	<b><i>Diols</i></b>								
10	2,3-butanediol	LLE	MS RI	2987 ± 1079	1497	5601	0.36	-	
11	1-2 propandiol	LLE	MS RI	674 ± 250	258	1216	0.37	-	
12									
13									
14	<b><i>Esters</i></b>								
15	2-methyl-1-propanol acetate	SPME	MS S	0 ± 0	n.d.	1	1.07	-	
16	ethyl butanoate	SPME	MS S	44 ± 12	18	64	0.26	20	[33]
17	3-methyl-1-butanol acetate	LLE	MS RI	1182 ± 1168	n.d.	5908	0.99	30	[33]
18	methyl hexanoate	SPME	MS S	1 ± 0	n.d.	2	0.38	-	
19	ethyl hexanoate	LLE	MS RI S	554 ± 268	132	1165	0.48	65	[45]
20	hexyl acetate	LLE	MS RI	8 ± 32	n.d.	171	3.94	2	[47]
21	ethyl lactate	LLE	MS RI	20976 ± 33732	1589	141330	1.61	60000	[28]
22	methyl octanoate	SPME	MS S	3 ± 1	1	6	0.39	200	[47]
23	ethyl octanoate	LLE	MS RI S	433 ± 326	n.d.	982	0.75	580	[45]
24	3-methylbutyl lactate	SPME	MS S	1 ± 0	n.d.	1	0.30	-	
25	ethyl-2-hydroxy-4-methylpentanoate	SPME	MS S	6 ± 3	1	14	0.46	-	
26	methyl decanoate	SPME	MS S	0 ± 0	n.d.	1	0.54	4	[48]
27	ethyl 3-hydroxybutanoate	LLE	MS RI	42 ± 87	n.d.	295	2.04	-	
28	ethyl decanoate	LLE	MS RI S	36 ± 80	n.d.	307	2.25	200	[45]
29	methyl ethyl succinate	SPME	MS S	96 ± 54	25	283	0.56	-	
30	3-methylbutyl octanoate	SPME	MS S	5 ± 2	2	9	0.35	125	[49]
31	diethyl succinate	LLE	MS RI	2555 ± 1849	263	8391	0.72	100000	[28]
32	ethyl 9-decenoate	SPME	MS S	9 ± 11	1	50	1.15	-	
33	methyl salicylate	SPME	MS S	5 ± 6	1	29	1.31	40	[47]
34									
35									
36									
37									
38									
39									
40									
41									
42									
43									
44									
45									
46									

2-phenylethyl acetate	LLE	MS RI S	136 ± 341	n.d.	1707	2.50	250	[33]
ethyl dodecanoate	LLE	MS RI	57 ± 125	n.d.	456	2.20	-	
diethyl malate	LLE	MS RI	5544 ± 4199	679	18639	0.76	760000	[28]
ethyl tetradecanoate	LLE	MS RI	57 ± 140	n.d.	504	2.48	-	
diethyle 2-hydroxypentanedioate	LLE	MS RI	605 ± 406	n.d.	1472	0.67	-	
ethyl hexadecanoate	LLE	MS RI S	1069 ± 1764	n.d.	6185	1.65	-	
ethyl hydrogen succinate	SPME	MS S	57 ± 30	19	125	0.52	1000000	[28]

### Carbonyl compounds

3-hydroxy-2-butanone (acetoin)	LLE	MS RI S	599 ± 311	162	1230	0.52	150000	[49]
furfural	SPME	MS S	29 ± 25	7	138	0.86	770	[28]
benzaldehyde	SPME	MS S	14 ± 28	4	164	2.05	350	[47]
3,4-dimethyl benzaldehyde	SPME	MS S	1 ± 0	n.d.	2	0.34	-	

### Others

dihydro-2-methyl-3(2H)-thiophenone	SPME	MS S	4 ± 3	1	11	0.59	-	
dihydro-2(3H)-furanone (γ-butyrolactone)	LLE	MS RI S	691 ± 204	316	1237	0.30	1000	[28]
methionol	LLE	MS RI	232 ± 145	n.d.	623	0.62	1000	[49]

<sup>a</sup> AM: analytical method; <sup>b</sup> IM: identification method (S comparison of mass spectra and retention time with those of standard compounds; RI comparison of order of elution with those reported in literature; MS comparison of mass spectra with those reported in mass spectrum libraries); <sup>c</sup> SD: standard deviation; <sup>d</sup> MIN: minimum value; <sup>e</sup> MAX: maximum value; <sup>f</sup> CV: coefficient of variation; <sup>g</sup> ODT: odor detection threshold; <sup>h</sup> n.d.: not detected



**Table 3. Free and bound terpenes and norisoprenoids detected in Ribolla Gialla commercial sparkling wines by different analytical approaches.**

**Concentrations are expressed in µg L<sup>-1</sup>.**

Compounds	AM <sup>a</sup>	IM <sup>b</sup>	Mean ±	SD <sup>c</sup>	MIN <sup>d</sup>	MAX <sup>e</sup>	CV <sup>f</sup>	ODT (µg L <sup>-1</sup> ) <sup>g</sup>
<i><b>Free terpenes</b></i>								
β-myrcene	SPME	MS S	0 ± 1		n.d. <sup>h</sup>	7	3.56	-
limonene	SPME	MS S	2 ± 1		1	4	0.33	200 [47]
cis-linalool oxide (furanic)	SPE	MS RI S	4 ± 4		n.d.	12	0.88	6000 [28]
trans-linalool oxide (furanic)	SPE	MS RI S	1 ± 2		n.d.	9	1.61	6000 [28]
linalool	SPE	MS RI S	10 ± 15		n.d.	80	1.51	50 [28]
terpinen-4-ol	SPME	MS S	1 ± 0		1	2	0.34	340 [47]
α-terpineol	SPE	MS RI S	23 ± 34		n.d.	194	1.49	250 [49]
β-citronellol	SPE	MS RI S	2 ± 4		n.d.	18	1.81	100 [33]
nerol	SPME	MS S	1 ± 1		n.d.	6	0.80	60 [50]
geraniol	SPE	MS RI S	4 ± 10		n.d.	39	2.87	30 [49]
geranic acid	SPME	MS S	7 ± 4		2	23	0.65	-
<i><b>Free C<sub>13</sub>-norisoprenoids</b></i>								
β-damascenone	SPE	MS RI	5 ± 6		n.d.	28	1.27	7 [36]
3-oxo-α-ionol	SPE	MS RI	10 ± 11		n.d.	42	1.11	-
riesling acetal	SPME	MS S	4 ± 2		n.d.	7	0.49	-
vitispirane (isomer 1)	SPME	MS S	4 ± 3		2	14	0.68	800 [38]
1,1,6-trimethyl-1,2-dihydronaphthalene (TDN)	SPME	MS S	5 ± 6		1	28	1.19	20 [38]
vitispirane (isomer 2)	SPME	MS S	14 ± 9		3	37	0.61	800 [38]
<i><b>Bound terpenes</b></i>								
nerol	SPE	MS RI S	3 ± 4		n.d.	20	1.54	

geraniol	SPE	MS RI S	13 ± 13	n.d.	51	1.03
1-hydroxylinalool	SPE	MS RI	2 ± 4	n.d.	15	1.99
geranic acid	SPE	MS RI	8 ± 9	n.d.	36	1.13
<b><u>Bound C<sub>13</sub>-norisoprenoids</u></b>						
3-hydroxy-β-damascone	SPE	MS RI	6 ± 10	n.d.	50	1.83
<i>trans,trans</i> -2,6-dimethyl-2,6-octadiene-1,8-diol (Z8-hydroxygeraniol)	SPE	MS RI	11 ± 13	n.d.	56	1.15
3-oxo-α-ionol	SPE	MS RI	40 ± 32	n.d.	135	0.81
3-oxo-7,8-dihydro-α-ionol (blumenol C)	SPE	MS RI	16 ± 22	n.d.	103	1.38
3-hydroxy-7,8-dihydro-β-ionol	SPE	MS RI	1 ± 2	n.d.	7	2.02

<sup>a</sup> AM: analytical method; <sup>b</sup> IM: identification method (S comparison of mass spectra and retention time with those of standard compounds; RI comparison of order of elution with those reported in literature; MS comparison of mass spectra with those reported in mass spectrum libraries); <sup>c</sup> SD: standard deviation; <sup>d</sup> MIN: minimum value; <sup>e</sup> MAX: maximum value; <sup>f</sup> CV: coefficient of variation; <sup>g</sup> ODT: odor detection threshold; <sup>h</sup> n.d.: not detected

**Table 4. Tryptophan metabolites detected in Ribolla Gialla commercial sparkling wines.**  
**Concentrations are expressed in mg L<sup>-1</sup>. SD: standard deviation; MIN: minimum value; MAX: maximum value; CV: coefficient of variation.**

Compound <sup>a</sup>	Mean	±	SD	MIN	MAX	CV
ILA-SO <sub>3</sub> H	8.53	±	10.31	0.66	54.20	1.21
TOL-SO <sub>3</sub> H	3.73	±	2.55	0.23	12.49	0.68
TYL	2.55	±	0.97	0.93	4.79	0.38
TYR	1.13	±	0.49	0.08	1.98	0.43
ILA-GLU-SO <sub>3</sub> H	0.53	±	0.37	0.12	1.79	0.70
PHE	0.51	±	0.26	0.03	0.99	0.51
TRP-EE	0.44	±	0.21	0.11	0.89	0.48
TOL	0.29	±	0.27	0.01	1.08	0.93
ABA	0.22	±	0.11	0.02	0.46	0.50
IAA-SO <sub>3</sub> H	0.20	±	0.07	0.13	0.39	0.35
ABA-GLU	0.13	±	0.05	0.04	0.26	0.38
TRP	0.13	±	0.10	0.01	0.39	0.77
ILA-GLU	0.10	±	0.05	0.01	0.25	0.50
TYR-EE	0.08	±	0.03	0.02	0.13	0.38
ILA	0.03	±	0.03	n.d. <sup>b</sup>	0.12	1.00
KYNA	0.02	±	0.01	n.d.	0.04	0.50
AA	0.01	±	0.06	n.d.	0.33	6.00
IAA-ASP	4.41	±	2.87	n.d.	12.35	0.65
N-TYR-EE	3.86	±	2.11	n.d.	9.63	0.55
KYN	2.42	±	1.98	n.d.	9.24	0.82
IAA	1.75	±	1.63	n.d.	8.84	0.93
N-TRP-EE	1.08	±	0.81	n.d.	3.47	0.75


<sup>a</sup> ILA-SO<sub>3</sub>H: indole-lactic acid-2-sulfonate; TOL-SO<sub>3</sub>H: tryptophol-2-sulfonate; TYL: tryptophol; TYR: tyrosine; ILA-GLU-SO<sub>3</sub>H: sulfonated indole-3-lactic acid; PHE: phenylalanine; TRP-EE: tryptophan-ethyl ester; TOL: tryptophol; ABA: abscisic acid; IAA-SO<sub>3</sub>H: sulfonated indole 3-acetic acid; ABA-GLU: glucoside of abscisic acid; TRP: tryptophan; ILA-GLU: indole 3-lactic acid glucoside; TYR-EE: tyrosine-ethyl ester; ILA: indole 3-lactic acid; KYNA: kynurenic acid; AA: anthranilic acid; IAA-ASP: indole 3-acetic acid conjugate with aspartic acid; N-TYR-EE: N-acetyl-tyrosine-ethyl ester; KYN: kynurenine; IAA: indole 3-acetic acid; N-TRP-EE: N-acetyl-tryptophan-ethyl ester.

<sup>b</sup> n.d.: not detected

**Table 5. Lipid compounds detected in Ribolla Gialla commercial sparkling wines.**  
**Concentrations are expressed in mg L<sup>-1</sup>. SD: standard deviation; MIN: minimum value;**  
**MAX: maximum value; CV: coefficient of variation.**

Compounds	Mean	±	SD	MIN	MAX	CV
<b><u>Fatty acids</u></b>						
palmitic acid	9.24	±	13.52	5.21	11.49	1.46
stearic acid	6.49	±	9.32	4.43	8.40	1.44
myristic acid	0.31	±	0.84	0.11	0.50	2.68
oleic acid + <i>cis</i> -vaccenic acid	0.19	±	0.32	0.12	0.26	1.72
arachidic acid	0.15	±	0.40	0.08	0.26	2.70
<i>cis</i> -11-eicosanoic acid	0.11	±	0.03	0.10	0.12	0.28
heptadecanoic acid	0.06	±	0.14	0.04	0.10	2.19
myristoleic acid	0.05	±	0.04	0.05	0.06	0.78
linoleic acid	0.04	±	0.08	0.02	0.06	2.29
lignoceric acid	0.03	±	0.16	n.d. <sup>a</sup>	0.09	5.52
behenic acid	0.02	±	0.08	0.01	0.05	4.21
palmitoleic acid	0.02	±	0.36	n.d.	0.14	20.00
linolenic acid	0.01	±	0.04	0.00	0.03	8.00
<b><u>Sterols</u></b>						
ergosterol	0.18	±	0.97	0.03	0.35	5.54
lupeol	0.10	±	1.10	0.01	0.48	10.89
<b><u>Glycerolipids</u></b>						
1-linoleoyl-rac-glycerol	0.02	±	0.02	0.01	0.02	1.18
1-oleoyl-rac-glycerol	0.01	±	0.02	0.01	0.02	1.43
glyceryl tripalmitoleate	0.01	±	0.09	n.d.	0.04	6.43
<b><u>Fatty acid esters</u></b>						
ethyl stearate	0.14	±	0.18	0.10	0.17	1.31
ethyl palmitate	0.13	±	0.22	0.07	0.17	1.64
ethyl oleate	0.03	±	0.05	0.02	0.05	1.47
ethyl linoleate	0.01	±	0.01	0.00	0.01	2.00
methyl palmitate	0.08	±	1.20	n.d.	0.50	15.19
methyl stearate	0.04	±	0.36	n.d.	0.15	9.73
methyl oleate	0.01	±	0.06	n.d.	0.03	6.67
<b><u>Triacylglycerols</u></b>						
tripentadecanoin	0.01	±	0.04	n.d.	0.02	5.00

<sup>a</sup> n.d.: not detected



**Fig. 1.**