



UNIVERSITÀ
DEGLI STUDI
DI UDINE

Università degli studi di Udine

The Zariski-Riemann space of valuation domains associated to pseudo-convergent sequences

Original

Availability:

This version is available <http://hdl.handle.net/11390/1215965> since 2021-12-14T13:59:26Z

Publisher:

Published

DOI:10.1090/TRAN/8185

Terms of use:

The institutional repository of the University of Udine (<http://air.uniud.it>) is provided by ARIC services. The aim is to enable open access to all the world.

Publisher copyright

(Article begins on next page)

The Zariski-Riemann space of valuation domains associated to pseudo-convergent sequences

G. Peruginelli*

D. Spirito[†]

May 6, 2020

Let V be a valuation domain with quotient field K . Given a pseudo-convergent sequence E in K , we study two constructions associating to E a valuation domain of $K(X)$ lying over V , especially when V has rank one. The first one has been introduced by Ostrowski, the second one more recently by Loper and Werner. We describe the main properties of these valuation domains, and we give a notion of equivalence on the set of pseudo-convergent sequences of K characterizing when the associated valuation domains are equal. Then, we analyze the topological properties of the Zariski-Riemann spaces formed by these valuation domains.

Keywords: pseudo-convergent sequence, pseudo-limit, extension of a valuation, residually transcendental extension, Zariski-Riemann space, constructible topology.

MSC Primary 12J20, 13A18, 13F30.

1 Introduction

Let V be a valuation domain with quotient field K . Determining and describing all the extensions of V to the field $K(X)$ of rational functions is an old and well-studied problem, which plays a vital role in several topics in field theory, commutative algebra

*Dipartimento di Matematica "Tullio Levi-Civita", University of Padova, Via Trieste 63, 35121 Padova, Italy. E-mail: gperugin@math.unipd.it

[†]Dipartimento di Matematica e Fisica, University of Roma Tre, Largo San Leonardo Murialdo 1, 00146 Roma. E-mail: spirito@mat.uniroma3.it; *Current address*: Dipartimento di Matematica "Tullio Levi-Civita", University of Padova, Via Trieste 63, 35121 Padova, Italy. E-mail: spirito@math.unipd.it

and beyond (see for example [13] and the references therein). The problem has been approached in a few different ways, with the main ones being through *key polynomials* (starting from the work of MacLane [15] and developed, among many others, by Vaquié [24]), *minimal pairs* (introduced by Alexandru, Popescu and Zaharescu [2, 3]) and *pseudo-convergent sequences*. The latter were introduced by Ostrowski in [17], who used them to describe all rank one extensions of the rank one valuation domain V to $K(X)$; subsequently, Kaplansky used this notion in [11] for valuation domains of any rank to characterize immediate extensions of valuation domains and maximally valued fields. More recently, Chabert in [8] generalized Ostrowski's definition by means of *pseudo-monotone sequences* to characterize the polynomial closure of subsets of valued fields of rank one, an important topic in the study of integer-valued polynomials.

In this paper, we study two constructions of extensions of V to $K(X)$ associated to a pseudo-convergent sequence $E \subset K$. The first one, which we denote by V_E , is the same construction introduced by Loper and Werner [14] for certain kinds of pseudo-convergent sequences on rank one valuation domains; we show that it actually defines a valuation domain for every pseudo-convergent sequence and for valuation domains of any rank (Theorem 3.8). The second one, which we denote by W_E , applies only when V has rank one and E satisfies some conditions, and is defined through its valuation w_E , which was already introduced by Ostrowski [17]; for this reason, we name it the *Ostrowski valuation* associated to E . In Sections 3 and 4, we investigate the structure of these valuation domains, in particular when V has rank one; among other things, we show that $V_E \subseteq W_E$, we characterize when V_E has rank 2 (Theorem 4.9), we find their value group and residue field, and we describe explicitly the valuation v_E associated to V_E as a map from $K(X)$ to \mathbb{R}^2 (Theorem 4.10). Many of these results are based on a general theorem (Theorem 3.3) which expresses the valuation of $\phi(t)$, for a rational function $\phi(X)$, as a linear function of $v(t - s)$, in an annulus of center s which contains neither poles nor zeros of $\phi(X)$.

In Section 5, following Ostrowski, we investigate the notion of equivalence between two pseudo-convergent sequences, analogous to the concept of equivalence between two Cauchy sequences. We show that two pseudo-convergent sequences are equivalent if and only if their associated valuation rings are equal; moreover, if they are of algebraic type then these conditions are also equivalent to the property of having the same set of pseudo-limits (in the algebraic closure of K and with respect to the same extension of v ; see Theorem 5.4). We also give a geometric interpretation of this fact in Section 5.1. Using these results, we show that the extensions of an Ostrowski valuation w_E to $\overline{K}(X)$ is completely determined by its restriction to \overline{K} (Theorem 5.7).

In Section 6 we study the spaces \mathcal{V} and \mathcal{W} formed, respectively, by the rings of the form V_E and by the rings of the form W_E , from a topological point of view; more precisely, we study the Zariski and the constructible topologies they inherit from the Zariski-Riemann space $\text{Zar}(K(X)|V)$. In particular, we first analyze the difference between these two topologies, showing that they coincide on \mathcal{W} (Proposition 6.3), while they coincide on \mathcal{V} if and only if the residue field of V is finite (Proposition 6.11); we also show how V_E can be seen as a limit of valuation domains defined from the members of E , mirroring the fact that (classes of) Cauchy sequences can be associated to their

limit points (Proposition 6.9). In Section 6.1, we show that \mathcal{V} , endowed with the Zariski topology, is a regular space and we deduce a sufficient condition for \mathcal{V} to be metrizable.

2 Background and notation

Throughout the article, V is a valuation domain; we denote by K its quotient field, by M its maximal ideal and by v the valuation associated to V . Its value group is denoted by Γ_v . We denote by \widehat{K} and \widehat{V} the completion of K and V , respectively, with respect to the topology induced by the valuation v . We still denote by v the unique extension of v to \widehat{K} (whose valuation domain is precisely \widehat{V}). We denote by \overline{K} a fixed algebraic closure of K . If u is an extension of v to \overline{K} , then the value group of u is the divisible hull of Γ_v , i.e., $\mathbb{Q}\Gamma_v = \mathbb{Q} \otimes_{\mathbb{Z}} \Gamma_v$.

The basic objects of study of this paper are pseudo-convergent sequences, introduced by Ostrowski in [17] and used by Kaplansky in [11] to describe immediate extensions of valued fields. Related concepts are *pseudo-stationary* and *pseudo-divergent* sequences [8], which we consider in [20].

Definition 2.1. Let $E = \{s_n\}_{n \in \mathbb{N}}$ be a sequence in K . We say that E is a *pseudo-convergent* sequence if $v(s_{n+1} - s_n) < v(s_{n+2} - s_{n+1})$ for all $n \in \mathbb{N}$.

In particular, if $E = \{s_n\}_{n \in \mathbb{N}}$ is a pseudo-convergent sequence and $n \geq 1$, then $v(s_{n+k} - s_n) = v(s_{n+1} - s_n)$ for all $k \geq 1$. We shall usually denote this quantity by δ_n ; following [25, p. 327] we call the sequence $\{\delta_n\}_{n \in \mathbb{N}}$ the *gauge* of E .

We shall make use of the following notation: given a sequence of real numbers $\{r_n\}_{n \in \mathbb{N}}$ and $r \in \mathbb{R} \cup \{\infty\}$, we write $r_n \nearrow r$ if r_n is a strictly increasing sequence with limit r .

Definition 2.2. The *breadth ideal* of E is

$$\text{Br}(E) = \{b \in K \mid v(b) > v(s_{n+1} - s_n), \forall n \in \mathbb{N}\}.$$

In general, $\text{Br}(E)$ is a fractional ideal of V and may not be contained in V .

The following definition has been introduced in [11], even though already in [17, p. 375] an equivalent concept appears (see [17, X, p. 381] for the equivalence).

Definition 2.3. An element $\alpha \in K$ is a *pseudo-limit* of E if $v(\alpha - s_n) < v(\alpha - s_{n+1})$ for all $n \in \mathbb{N}$, or, equivalently, if $v(\alpha - s_n) = \delta_n$ for all $n \in \mathbb{N}$. We denote the set of pseudo-limits of E by \mathcal{L}_E , or \mathcal{L}_E^v if we need to emphasize the valuation.

If $\text{Br}(E)$ is the zero ideal then E is a Cauchy sequence in K and converges to an element of \widehat{K} , which is the unique pseudo-limit of E . In general, Kaplansky proved the following more general result.

Lemma 2.4. [11, Lemma 3] *Let $E \subset K$ be a pseudo-convergent sequence. If $\alpha \in K$ is a pseudo-limit of E , then the set of pseudo-limits of E in K is equal to $\alpha + \text{Br}(E)$.*

If w is an extension of v to a field L containing K , and E is a sequence in K , then E is pseudo-convergent with respect to w if and only if E is pseudo-convergent with respect to v . Moreover, every pseudo-limit of E under v in K is also a pseudo-limit under w .

Suppose now that V has rank one; then we consider Γ_v and $\mathbb{Q}\Gamma_v$ as totally ordered subgroups of \mathbb{R} . The valuation v induces an ultrametric distance d on K , defined by

$$d(x, y) = e^{-v(x-y)}.$$

In this metric, V is the closed ball of center 0 and radius 1. Given $s \in K$ and $\gamma \in \Gamma_v$, we denote the ball of center s and radius $r = e^{-\gamma}$ by:

$$B(s, r) = \{x \in K \mid d(x, s) \leq r\} = \{x \in K \mid v(x - s) \geq \gamma\}.$$

A ball in \overline{K} with respect to an extension u of v is denoted by $B_u(s, r)$.

If $E \subset K$ is a pseudo-convergent sequence, then the gauge $\{\delta_n\}_{n \in \mathbb{N}}$ of E is a strictly increasing sequence of real numbers, and so the following definition makes sense.

Definition 2.5. The *breadth* of a pseudo-convergent sequence $E = \{s_n\}_{n \in \mathbb{N}}$ is the limit

$$\delta_E = \lim_{n \rightarrow \infty} v(s_{n+1} - s_n) = \lim_{n \rightarrow \infty} \delta_n.$$

The breadth δ is an element of $\mathbb{R} \cup \{\infty\}$, and it may not lie in Γ_v . We can use the breadth to characterize the breadth ideal: indeed, $\text{Br}(E) = \{b \in K \mid v(b) \geq \delta_E\}$, or equivalently $\delta_E = \inf\{v(b) \mid b \in \text{Br}(E)\}$. If $\delta = +\infty$, then $\text{Br}(E)$ is just the zero ideal and E is a Cauchy sequence in K . If V is a discrete valuation ring, then every pseudo-convergent sequence is actually a Cauchy sequence. Lemma 2.4 can also be phrased in a geometric way: if $\alpha \in \mathcal{L}_E$, then \mathcal{L}_E is the closed ball of center α and radius $e^{-\delta_E}$, i.e., $\mathcal{L}_E = B(\alpha, e^{-\delta_E})$.

The following concepts have been given by Kaplansky in [11] in order to study the different kinds of immediate extensions of a valued field K . Recall that if L is a field extension of K , a valuation domain W of L *lies over* V if $W \cap K = V$. In this case, the residue field of W is naturally an extension of the residue field of V and similarly the value group of W is an extension of the value group of V . We say that W is immediate over V if both the residue fields and the value groups are the same.

Definition 2.6. Let E be a pseudo-convergent sequence. We say that E is of *transcendental type* if $v(f(s_n))$ eventually stabilizes for every $f \in K[X]$; on the other hand, if $v(f(s_n))$ is eventually increasing for some $f \in K[X]$, we say that E is of *algebraic type*.

The main difference between these two kind of sequences is the nature of the pseudo-limits: if E is of algebraic type, then E has pseudo-limits in \overline{K} (for some extension u of v), while if E is of transcendental type then E admits a pseudo-limit only in a transcendental extension [11, Theorems 2 and 3].

If $L = K(X)$ and W lies over V , then W is said to be a *residually transcendental extension* of V (or simply residually transcendental if V is understood) if the residue field of W is a transcendental extension of the residue field of V [2].

Definition 2.7. Let Γ be a totally ordered group containing Γ_v , and take $\alpha \in K$ and $\delta \in \Gamma$. The *monomial valuation* $v_{\alpha,\delta}$ is defined in the following way: if $f(X) \in K[X]$ is a polynomial, write $f(X) = a_0 + a_1(X - \alpha) + \dots + a_n(X - \alpha)^n$; then,

$$v_{\alpha,\delta}(f) = \inf\{v(a_i) + i\delta \mid i = 0, \dots, n\}.$$

It is well known that $v_{\alpha,\delta}$ naturally extends to a valuation on $K(X)$ [5, Chapt. VI, §. 10, Lemme 1], and $v_{\alpha,\delta}$ is residually transcendental over v if and only if δ has finite order over Γ_v [18, Lemma 3.5]. Furthermore, every residually transcendental extension of V can be written as $W \cap K(X)$, where W is a valuation domain of $\overline{K}(X)$ associated to a monomial valuation [1, 2].

Let D be an integral domain and L be a field containing D (not necessarily the quotient field of D). The *Zariski space of D in L* , denoted by $\text{Zar}(L|D)$, is the set of valuation domains of L containing D endowed with the so-called *Zariski topology*, i.e., with the topology generated by the subbasic open sets

$$B(\phi) = \{V \in \text{Zar}(L|D) \mid \phi \in V\},$$

where $\phi \in L$. Under this topology, $\text{Zar}(L|D)$ is a compact space [27, Chapter VI, Theorem 40] that is almost never Hausdorff nor T_1 (indeed, $\text{Zar}(L|D)$ is a T_1 space if and only if D is a field and L is an algebraic extension of D).

The *constructible topology* (also called *patch topology*) on $\text{Zar}(L|D)$ is the coarsest topology such that the subsets $B(\phi_1, \dots, \phi_k)$ are both open and closed; we denote this space by $\text{Zar}(L|D)^{\text{cons}}$. Clearly, the constructible topology is finer than the Zariski topology; however, $\text{Zar}(L|D)^{\text{cons}}$ is still compact, and furthermore it is always Hausdorff [10, Theorem 1].

3 A valuation domain associated to a pseudo-convergent sequence

The following valuation domain associated to a pseudo-convergent sequence has been introduced by Loper and Werner in [14] in the case of a valuation domain V of K of rank one. We generalize their construction to valuation domains of arbitrary rank.

Definition 3.1. Let $E = \{s_n\}_{n \in \mathbb{N}} \subset K$ be a pseudo-convergent sequence. Let

$$V_E = \{\phi \in K(X) \mid \phi(s_n) \in V, \text{ for all but finitely many } n \in \mathbb{N}\}. \quad (1)$$

The aim of this section is to prove that V_E is a valuation domain of $K(X)$ for every pseudo-convergent sequence E . When the rank of V is one and E is of transcendental type or has zero breadth ideal, this result was already obtained, respectively, in Proposition 5.5 and Theorem 5.8 of [14]. More generally, for any valuation domain, when E is of transcendental type V_E coincides with the valuation domain of $K(X)$ defined by Kaplansky in [11, Theorem 2], which is an immediate extension of V . Since in this case for each $\phi \in K(X)$ we have that $v(\phi(s_n))$ is eventually constant, the value of ϕ with

respect to the above valuation is equal to that constant value that $\phi(X)$ assumes over E . Following this example, our method is heavily based on understanding the values of $\phi(X)$ along a pseudo-convergent sequence.

For the next result, which is not a priori related to pseudo-convergent sequences, we introduce some notations and definitions.

Definition 3.2. Let $\phi \in K(X)$. The *multiset of critical points* of $\phi(X)$ is the multiset Ω_ϕ of zeroes and poles of ϕ in \overline{K} (each of them counted with multiplicity). Given a sub-multiset $S = \{\alpha_1, \dots, \alpha_k\}$ of Ω_ϕ , by the *weighted sum* of S we mean the sum $\sum_{\alpha_i \in S} \epsilon_i$, where ϵ_i is equal either to 1 or to -1 , according to whether α_i is a zero or a pole of ϕ , respectively. By the *S-part* of ϕ we mean the rational function $\phi_S(X) = \prod_{\alpha_i \in S} (X - \alpha_i)^{\epsilon_i}$, where $\epsilon_i \in \{\pm 1\}$ is as above. Note that $\phi_{\Omega_\phi}(X)$ is equal to $\phi(X)$ up to a constant.

Given a convex subset Δ of Γ_v , $\beta \in \overline{K}$ and an extension u of v to \overline{K} , we set

$$\mathcal{C}_u(\beta, \Delta) = \{s \in \overline{K} \mid u(s - \beta) \in \Delta\} \quad (2)$$

and, if $\gamma \in \mathbb{Q}\Gamma_v$, we write $\gamma < \Delta$ ($\gamma > \Delta$, respectively) if $\gamma < \delta$ ($\gamma > \delta$, respectively) for every $\delta \in \Delta$.

Theorem 3.3. Let $\phi \in K(X)$ and let $s \in K$; let u be an extension of v to \overline{K} . Let Δ be a convex subset of $\mathbb{Q}\Gamma_v$ such that $C = \mathcal{C}_u(s, \Delta)$ does not contain any critical point of ϕ . Let $\lambda \in \mathbb{Z}$ be equal to the weighted sum of the multiset S of critical points α of ϕ which satisfy $u(\alpha - s) > \Delta$ and let $\gamma = u\left(\frac{\phi}{\phi_S}(s)\right)$. Then, for all $t \in C \cap K$, we have

$$v(\phi(t)) = \lambda v(t - s) + \gamma. \quad (3)$$

Proof. Over \overline{K} , we can write $\phi(X)$ as a product $c \prod_{i=1}^n (X - \alpha_i)^{\epsilon_i}$, where the α_i are the critical points of ϕ , $\epsilon_i \in \{-1, 1\}$ and $c \in K$. Let $C = \mathcal{C}_u(s, \Delta) \subset \overline{K}$ and let $t \in K \cap C$. If $u(\alpha_i - s) < \Delta$ then $u(t - \alpha_i) = u(s - \alpha_i)$, while if $u(\alpha_i - s) > \Delta$ then $u(t - \alpha_i) = u(s - t)$ (note that, by assumption, there are no other possibilities for the critical points of $\phi(X)$). Therefore, we have

$$\begin{aligned} v(\phi(t)) &= v(c) + \sum_{i: u(\alpha_i - s) < \Delta} \epsilon_i u(t - \alpha_i) + \sum_{i: u(\alpha_i - s) > \Delta} \epsilon_i u(t - \alpha_i) = \\ &= v(c) + \sum_{i: u(\alpha_i - s) < \Delta} \epsilon_i u(\alpha_i - s) + \sum_{i: u(\alpha_i - s) > \Delta} \epsilon_i u(t - s) = \gamma + \lambda v(t - s) \end{aligned}$$

where $\lambda = \sum_{i: u(\alpha_i - s) > \Delta} \epsilon_i$ and $\gamma = v(c) + \sum_{i: u(\alpha_i - s) < \Delta} \epsilon_i u(s - \alpha_i) = u\left(\frac{\phi}{\phi_S}(s)\right)$, with S being the multiset of critical points α_i of ϕ which satisfy $u(\alpha_i - s) > \Delta$. In particular, $\lambda \in \mathbb{Z}$ and $\gamma \in \mathbb{Q}\Gamma_v$ do not depend on t . The claim is proved. \square

Remark 3.4. Let $\alpha_1, \dots, \alpha_n$ be the zeros and the poles of ϕ , and let $\rho_i = u(s - \alpha_i)$; without loss of generality, suppose $\rho_1 < \dots < \rho_n$. Then, the sets $\Delta_i = (\rho_i, \rho_{i+1})$, for $i = 0, \dots, n$ (with the convention $\rho_0 = -\infty$ and $\rho_{n+1} = +\infty$) are the maximal convex sets on which Theorem 3.3 can be applied: that is, they satisfy (by definition) the hypothesis of the theorem, and if $\Delta_i \subsetneq \Delta$ for some other convex set Δ then the theorem cannot be applied to Δ .

In order to apply Theorem 3.3 to pseudo-convergent sequences, we need the following definition.

Definition 3.5. Let $E = \{s_n\}_{n \in \mathbb{N}}$ be a pseudo-convergent sequence in K , let u be an extension of v to \overline{K} and let $\phi \in K(X)$. The *dominating degree* $\deg_{E,u}(\phi)$ of ϕ with respect to E and u is the weighted sum of the critical points of $\phi(X)$ (according to Definition 3.2) which are pseudo-limits of E with respect to u .

Note that, by definition, if E is a pseudo-convergent sequence of transcendental type, then $\deg_{E,u}(\phi) = 0$ for every $\phi \in K(X)$.

The following result shows that the values of a rational function over a pseudo-convergent sequence $E \subset K$ form a sequence that is eventually monotone, either strictly increasing, strictly decreasing or stationary, according to whether the dominating degree of ϕ with respect to E is positive, negative or equal to zero, respectively.

Proposition 3.6. *Let $E = \{s_n\}_{n \in \mathbb{N}} \subset K$ be a pseudo-convergent sequence with gauge $\{\delta_n\}_{n \in \mathbb{N}}$, and let u be an extension of v to \overline{K} . Let $\phi \in K(X)$.*

(a) *If $\lambda = \deg_{E,u} \phi$, then there is $\gamma \in \Gamma_v$ such that, for all sufficiently large n , we have*

$$v(\phi(s_n)) = \lambda \delta_n + \gamma.$$

(b) *If $\beta \in \overline{K}$ is a pseudo-limit of E with respect to u , then $\gamma = u\left(\frac{\phi}{\phi_S}(\beta)\right)$, where S is the set of critical points of $\phi(X)$ which are pseudo-limits of E .*

(c) *The dominating degree of ϕ does not depend on u ; that is, if u' is another extension of v to \overline{K} , then $\deg_{E,u} \phi = \deg_{E,u'} \phi$.*

Proof. If E is of transcendental type, then $\lambda = 0$ and all claims follow from the definition.

Suppose that E is of algebraic type and $\beta \in \mathcal{L}_E^u$; we will prove (a) and (b) together. Let $\Delta = \Delta_E$ be the least initial segment of $\mathbb{Q}\Gamma_v$ containing the gauge of E . There exists $\tau \in \Gamma_v \cap \Delta$ such that $C = \mathcal{C}_u(\beta, \Delta \cap (\tau, +\infty))$ contains no critical points of ϕ . Let λ be the weighted sum of the subset S of Ω_ϕ of those elements α such that $u(\alpha - \beta) > \Delta \cap (\tau, +\infty)$ (or, equivalently, $u(\alpha - \beta) > \Delta$) and $\gamma = u\left(\frac{\phi}{\phi_S}(\beta)\right)$. For all n sufficiently large $s_n \in C$: by Theorem 3.3, it follows that for each such n we have

$$v(\phi(s_n)) = \lambda u(\beta - s_n) + \gamma = \lambda \delta_n + \gamma.$$

Note that $\gamma \in \Gamma_v$ and, by Lemma 2.4, S is the set of critical points of $\phi(X)$ which are pseudo-limits of E , so λ is the dominating degree of ϕ with respect to E .

For (c), we note that $v(\phi(s_n))$ does not depend on the extension u ; hence, if $\lambda = \deg_{E,u} \phi$, $\lambda' = \deg_{E,u'} \phi$, $\gamma = u\left(\frac{\phi}{\phi_S}(\beta)\right)$, $\gamma' = u'\left(\frac{\phi}{\phi_S}(\beta)\right)$, we have

$$v(\phi(s_n)) = \lambda \delta_n + \gamma = \lambda' \delta_n + \gamma'$$

for all large n . However, this clearly implies $\lambda = \lambda'$, as claimed. \square

In view of point (c) of the previous proposition, we denote the dominating degree of ϕ with respect to E simply as $\deg_{\text{dom}_E} \phi$.

The term *dominating degree* comes from the following property. We remark that, in a different context, a similar argument has been given in the proof of [8, Proposition 4.8].

Proposition 3.7. *Let $E = \{s_n\}_{n \in \mathbb{N}} \subset K$ be a pseudo-convergent sequence with pseudo-limit $\beta \in K$, and let $f(X) = \sum_{i=0, \dots, d} a_i(X - \beta)^i \in K[X]$. Then, $\deg_{\text{dom}_E} f$ is the non-negative integer k such that $v(f(s_n)) = v(a_k(s_n - \beta)^k)$ for all large n .*

Proof. Clearly, if $v(f(s_n)) = v(a_k(s_n - \beta)^k)$ for all large n then $v(f(s_n)) = k\delta_n + v(a_k)$ and so $k = \deg_{\text{dom}_E} f$.

Conversely, suppose $k = \deg_{\text{dom}_E} f$. Then, by definition, $v(f(s_n)) = k\delta_n + \gamma$ for some $\gamma \in \Gamma_v$ (for all large n), where $\{\delta_n\}_{n \in \mathbb{N}}$ is the gauge of E . We consider the following linear functions from Γ_v to Γ_v :

$$\lambda_i(\eta) = i\eta + v(a_i), \quad i \in \{0, \dots, d\}.$$

Let Δ be the least initial segment of Γ_v containing the gauge of E : then, since the λ_i are linear, there is a $\tau \in \Delta$ and an $r \in \{0, \dots, d\}$ such that $\lambda_r(\eta) < \lambda_i(\eta)$ for all $\eta \in \Delta \cap (\tau, +\infty)$ and for all $i \neq r$. Therefore, whenever $\delta_n \in \Delta \cap (\tau, +\infty)$ we must have

$$v(f(s_n)) = v\left(\sum_i a_i(s_n - \beta)^i\right) = \inf_i \{v(a_i(s_n - \beta)^i)\} = r\delta_n + v(a_r).$$

In particular, it must be $r = k$, as claimed. \square

Theorem 3.8. *Let $E = \{s_n\}_{n \in \mathbb{N}}$ be a pseudo-convergent sequence. Then $V_E \subset K(X)$ is a valuation domain lying over V with maximal ideal equal to $M_{V_E} = \{\phi \in K(X) \mid v(\phi(s_n)) \in M, \text{ for all but finitely many } n \in \mathbb{N}\}$. Moreover, X is a pseudo-limit of E with respect to the valuation v_E associated to V_E .*

Proof. Clearly, V_E is a ring and $V_E \cap K = V$.

If E is of transcendental type then V_E is exactly the valuation domain of the immediate extension of the valuation v to $K(X)$ induced by E as in [11, Theorem 2]. We have that X is a pseudo-limit of E by [11, Theorem 2].

Suppose now that E is of algebraic type, and let $\phi \in K(X)$. By Proposition 3.6, $v(\phi(s_n))$ is a linear function of δ_n ; hence, it is either eventually positive, eventually zero or eventually negative. Since $v(\phi^{-1}(s_n)) = -v(\phi(s_n))$ (provided that $\phi(s_n) \neq 0$, which happens only finitely many times), we have that $\phi \in V_E$, in the first and second case, while in the third case $\phi^{-1} \in V_E$. Hence, V_E is a valuation domain, and the claim about the maximal ideal follows easily.

Finally, we show that X is a pseudo-limit of E with respect to v_E . Fix $n \in \mathbb{N}$, and let $\phi(X) = \frac{X - s_{n+1}}{X - s_n}$. Then, for $m > n + 1$ we have $v(\phi(s_m)) = \delta_{n+1} - \delta_n > 0$, and thus $v_E(X - s_{n+1}) > v_E(X - s_n)$. It follows that X is a pseudo-limit of E , as claimed. \square

We want now to link the valuation domain V_E to another class of valuation domains, which for example have been recently considered in [19].

Definition 3.9. Let $\alpha \in \widehat{K}$. We denote by W_α the ring of rational functions over K which are integer-valued over α , namely:

$$W_\alpha = \{\phi \in K(X) \mid v(\phi(\alpha)) \geq 0\}$$

It is straightforward to verify that W_α is a valuation domain of $K(X)$ which lies over V (see also [19, Proposition 2.2]).

Remark 3.10. In case E is a pseudo-convergent sequence with zero breadth ideal, and $\alpha \in \widehat{K}$ is the (unique) limit of E , since rational functions are continuous in the topology induced by v , we have $V_E = W_\alpha$. Moreover, α is algebraic (transcendental, respectively) over K if and only if E is of algebraic (transcendental, respectively) type. These kind of valuations domains have been characterized in [19, Proposition 2.2]. We will deal with the case of non-zero breadth ideal in Theorem 4.9.

We conclude this section by describing the valuation v_E , its residue field and its value group when E is a pseudo-convergent sequence of algebraic type.

Proposition 3.11. *Let $E = \{s_n\}_{n \in \mathbb{N}} \subset K$ be a pseudo-convergent sequence and suppose that $\beta \in K$ is a pseudo-limit of E ; let $\alpha \in K$. Then, the following hold.*

(a) $v_E(X - \alpha) \leq v_E(X - \beta)$ and equality holds if and only if $\alpha \in \mathcal{L}_E$.

Let $\Delta_E = v_E(X - \beta) \in \Gamma_{v_E}$ (which by above does not depend on the choice of the pseudo-limit β of E).

(b) Δ_E is not a torsion element in Γ_{v_E}/Γ_v (i.e., if $k \in \mathbb{N}$ is such that $k \cdot \Delta_E \in \Gamma_v$, then $k = 0$). In particular, $v_E = v_{\beta, \Delta_E}$.

(c) $\Gamma_{v_E} = \mathbb{Z}\Delta_E \oplus \Gamma_v$ (as groups).

(d) $V_E/M_E \cong V/M$.

Proof. The condition $v_E(X - \alpha) \leq v_E(X - \beta)$ is equivalent to $\phi(X) = \frac{X - \beta}{X - \alpha} \in V_E \Leftrightarrow \phi(s_n) \in V$, for almost all $n \in \mathbb{N}$. For each $n \in \mathbb{N}$, we have:

$$v(\phi(s_n)) = v(s_n - \beta) - v(s_n - \alpha)$$

Now, we write $v(s_n - \alpha) = v(s_n - \beta + \beta - \alpha)$. Note that $\beta - \alpha \in \text{Br}(E) \Leftrightarrow \alpha \in \mathcal{L}_E$ (Lemma 2.4). If these conditions hold, then $v(\beta - \alpha) > v(s_{n+1} - s_n) = v(s_n - \beta)$ for each $n \in \mathbb{N}$ and therefore $v(s_n - \alpha) = v(s_n - \beta)$. Note that in this case $\phi \in V_E^*$ and so, in particular, $\Delta_E = v_E(X - \beta)$ does not depend on the pseudo-limit β of E we have chosen (in K). If instead $\alpha \notin \mathcal{L}$ then there exists $N \in \mathbb{N}$ such that $v(\beta - \alpha) < v(s_{n+1} - s_n) = v(\beta - s_n)$ for all $n \geq N$. Hence, $v(s_n - \alpha) = v(\beta - \alpha) < v(\beta - s_n)$ for all $n \geq N$, that is, $\phi \in M_E \subset V_E$.

We prove now the other three claims. Suppose there exists $k \in \mathbb{N}$ such that $k \cdot \Delta_E \in \Gamma_v$, that is, $k \cdot v_E(X - \beta) = v_E((X - \beta)^k) = v(a)$, for some $a \in K$. This implies that $\frac{(X - \beta)^k}{a} \in V_E^*$, which is a contradiction, since $k \cdot v(s_n - \beta) - v(a)$ is strictly increasing.

Since $\Delta_E = v_E(X - \beta) \in \Gamma_{v_E}$ is not torsion over Γ_v , by [5, Chapt. VI, §10, Proposition 1] (see also [4, p. 289]) we have that for each $f \in K[X]$, $f(X) = a_0 + a_1(X - \beta) + \dots + a_n(X - \beta)^n$,

$$v_E(f(X)) = \inf\{v(a_i) + i\Delta_E \mid i = 0, \dots, n\}$$

(where the inf is in Γ_{v_E}). In fact, we have $v_E(a_i(X - \beta)^i) \neq v_E(a_j(X - \beta)^j)$, for all $i \neq j \in \{0, \dots, n\}$, otherwise $(i - j)\Delta_E = v(a_j) - v(a_i)$ and Δ_E would be torsion over Γ_v . This implies that $v_E = v_{\beta, \Delta_E}$. Moreover, by the same reference, $\Gamma_{v_E} = \mathbb{Z}\Delta_E \oplus \Gamma_v$ and the residue field of V_E is isomorphic to the residue field of V . \square

In the general case, where E is algebraic but has no pseudo-limits in K , we only need to pass to an extension of V .

Corollary 3.12. *Let $E \subset K$ be a pseudo-convergent sequence of algebraic type, and let u be an extension of v to \bar{K} . Let $\beta \in \bar{K}$ be a pseudo-limit of E with respect to u , and let $\Delta = u_E(X - \beta)$. Then, v_E is equal to the restriction to $K(X)$ of $u_E = u_{\beta, \Delta}$.*

Proof. Let U_E be the valuation domain of $\bar{K}(X)$ associated to E with respect to u . By Proposition 3.11, $u_E = u_{\beta, \Delta}$, where $\Delta = u_E(X - \beta)$. Since $U_E \cap K(X) = V_E$, the claim follows immediately. \square

4 The rank of V_E and the Ostrowski valuation w_E

We assume for the rest of the article that V has rank one.

If W is an extension of V to $K(X)$, then the rank of W is either 1 or 2 ([5, Chapitre VI, §10, Corollaire 1, p. 162]). In this section, we want to determine when each of the two possibilities occurs for $W = V_E$, where E is a pseudo-convergent sequence. To this end, we need to introduce another kind of valuation on $K(X)$ which lies over V ; also this valuation arise from pseudo-convergent sequences and have been first introduced and studied by Ostrowski in [17, 65. p. 374].

Definition 4.1. Let $E = \{s_n\}_{n \in \mathbb{N}} \subset K$ be a pseudo-convergent sequence. We define w_E as the map

$$\begin{aligned} w_E: K(X) &\longrightarrow \mathbb{R} \cup \{\pm\infty\} \\ \phi &\longmapsto \lim_{n \rightarrow \infty} v(\phi(s_n)). \end{aligned}$$

If w_E happens to define a valuation on $K(X)$, we denote by W_E the associated valuation domain, namely, $W_E = \{\phi \in K(X) \mid w_E(\phi) \geq 0\}$.

Note that, for large n , s_n is neither a zero nor a pole of ϕ , so $v(\phi(s_n))$ is defined for all large n . We are going to show under which cases w_E is a valuation on $K(X)$.

One of the main accomplishments of Ostrowski (and also the motivation for the introduction of the notion of pseudo-convergent sequence) in his work [17] is the *Fundamentalsatz*, which we now recall.

Theorem 4.2. [17, 66. IX, p. 378] Let K be an algebraically closed field and let v be a rank one valuation on K . If w is a rank one valuation of $K(X)$ extending v , then there is a pseudo-convergent sequence $E = \{s_n\}_{n \in \mathbb{N}} \subset \overline{K}$ such that $w = w_E$.

When K is not algebraically closed, this means that the rank one valuations of $K(X)$ extending v can be realized as the contraction to $K(X)$ of the valuations w_E on $\overline{K}(X)$ for some pseudo-convergent sequence $E \subset \overline{K}$ and some extension of v to \overline{K} .

For the sake of completeness, in the next two propositions we prove the basic properties of the function w_E .

Proposition 4.3. Let $E = \{s_n\}_{n \in \mathbb{N}}$ be a pseudo-convergent sequence that is either of transcendental type or of algebraic type and non-zero breadth ideal. Then the map $w_E : K(X) \rightarrow \mathbb{R} \cup \{\infty\}$ extends v and is a valuation of rank one on $K(X)$. Furthermore, the valuation ring W_E relative to w_E contains V_E .

Proof. Suppose first that E is of transcendental type. Then for each $\phi \in K(X)$, $v(\phi(s_n))$ is eventually constant, and furthermore $w_E(\phi) = \infty$ if and only if $\phi = 0$.

Suppose now that E is of algebraic type and the breadth ideal is non-zero. Then also in this case w_E is well-defined, since by Proposition 3.6 for every $\phi \in K(X)$ there is a $k \in \mathbb{Z}$ and $\gamma \in \Gamma_v$ such that $v(\phi(s_n)) = k\delta_n + \gamma$, and $\delta_n \rightarrow \delta$ as $n \rightarrow \infty$. Moreover, since $\delta < \infty$, and since ϕ has only finitely many zeros and points where it is not defined, we have $w_E(\phi) = \infty$ if and only if $\phi = 0$.

In either case, if $\phi = a \in K$ is a constant, then $w_E(\phi) = v(a)$; thus, w_E extends v .

If now $\phi_1, \phi_2 \in K(X)$ then

$$v((\phi_1 + \phi_2)(s_n)) = v(\phi_1(s_n) + \phi_2(s_n)) \geq \min\{v(\phi_1(s_n)), v(\phi_2(s_n))\};$$

hence, $w_E(\phi_1 + \phi_2) \geq \min\{w_E(\phi_1), w_E(\phi_2)\}$. In the same way, $w_E(\phi_1\phi_2) = w_E(\phi_1) + w_E(\phi_2)$. Hence, w_E is a valuation.

If now $\phi \in V_E$, then $\phi(s_n) \in V$ for large n , or equivalently $v(\phi(s_n)) \geq 0$ for large n . In particular, $\lim v(\phi(s_n)) \geq 0$, i.e., $w_E(\phi) \geq 0$. Therefore, $\phi \in W_E$. \square

If E is of algebraic type and its breadth ideal is zero, on the other hand, w_E is not a valuation: this is due to the fact that $w_E(\phi)$ tends to ∞ when the pseudo-limit of E (in \overline{K}) is a zero of ϕ . It is, however, very close to a valuation: recall that a *pseudo-valuation* of a field K is a map v from K to $\Gamma_v \cup \{\infty\}$, where Γ_v is a totally ordered abelian group, which satisfies the same axioms of a valuation except that we are not assuming that $v(x) = \infty \Rightarrow x = 0$. The set $\{x \in K \mid v(x) = \infty\}$ is a prime ideal of the valuation domain V of v , called the *socle* of v . The following proposition is straightforward.

Proposition 4.4. Let $E = \{s_n\}_{n \in \mathbb{N}} \subset K$ be a pseudo-convergent sequence of algebraic type and zero breadth ideal. If $q \in K[X]$ is the minimal polynomial of the limit of E in \overline{K} , then the map $w_E : K[X]_{(q)} \rightarrow \mathbb{R} \cup \{\infty\}$ extends v and is a pseudo-valuation with socle $q(X)K[X]_{(q)}$. Moreover, the valuation ring of w_E , that is, $\{\phi \in K[X]_{(q)} \mid w_E(\phi) \geq 0\}$, is equal to V_E .

Definition 4.5. Let $E \subset K$ be a pseudo-convergent sequence which is either of transcendental type or of algebraic type and non-zero breadth ideal. We call the associated rank one valuation $w_E : K(X) \rightarrow \mathbb{R} \cup \{\infty\}$ the *Ostrowski valuation* associated to E , and the corresponding valuation domain W_E the *Ostrowski valuation domain* associated to E .

The following corollary is an easy consequence of Proposition 3.6. It also follows from Lemma 4.8 below.

Corollary 4.6. Let $E = \{s_n\}_{n \in \mathbb{N}} \subset K$ be a pseudo-convergent sequence of algebraic type with breadth δ , and let $\phi \in K(X)$. If $\lambda = \deg_{\text{dom}_E}(\phi)$ and $\gamma \in \Gamma_v$ is as in Proposition 3.6, then we have

$$w_E(\phi) = \lambda\delta + \gamma \quad (4)$$

In particular, $\Gamma_{w_E} = \mathbb{Z}\delta + \Gamma_v$.

Remark 4.7. (a) Let u be an extension of v to \overline{K} . It follows at once from Corollary 4.6 that if $E = \{s_n\}_{n \in \mathbb{N}} \subset K$ is a pseudo-convergent sequence of algebraic type with breadth δ and $\beta \in \mathcal{L}_E^u$, then, for each $s \in K$ we have:

$$w_E(X - s) = \lim_{n \rightarrow \infty} v(s_n - s) = \begin{cases} \delta, & \text{if } s \in \mathcal{L}_E \\ u(s - \beta) < \delta, & \text{if } s \notin \mathcal{L}_E \end{cases} \quad (5)$$

Note that, in case $s \notin \mathcal{L}_E$ and $\beta' \in \mathcal{L}_E^u$, we have $w_E(X - s) = u(s - \beta) = u(s - \beta')$, thus this value is independent of the chosen pseudo-limit of E . Similarly, if E is of transcendental type, then $w_E(X - s) < \delta$ for any $s \in K$.

(b) Under the assumption of Corollary 4.6, let u be a fixed extension of v to \overline{K} and let w_E be extended to $\overline{K}(X)$ along u (i.e., $w_E(\psi) = \lim u(\psi(s_n))$, for any $\psi \in \overline{K}(X)$). Let S be the multiset of critical points of ϕ which are pseudo-limits of E . Then by (4) and (5) we have

$$w_E(\phi) = w_E(\phi_S) + w_E\left(\frac{\phi}{\phi_S}\right) = \lambda\delta + \gamma$$

where $\lambda\delta = w_E(\phi_S)$ and $w_E\left(\frac{\phi}{\phi_S}\right) = u\left(\frac{\phi}{\phi_S}(\beta)\right) = \gamma$, where $\beta \in \mathcal{L}_E^u$, since $w_E(X - \alpha) = v(\beta - \alpha)$ for every $\alpha \notin S$ by the previous remark. We stress the strong analogy between this expression of w_E and the valuation associated to a valuation domain of the form W_α , for $\alpha \in \widehat{K}$ which is algebraic over K . See [6, p. 126] and [19, Remark 2.3].

The next lemma is taken from [17] and gives an important connection between monomial valuations and Ostrowski valuations; we repeat it here for the convenience of the reader.

Lemma 4.8. [17, VII, p. 377] Let $E = \{s_n\}_{n \in \mathbb{N}}$ be a pseudo-convergent sequence of algebraic type and let $\alpha \in \mathcal{L}_E^u$, for some extension u of v to \overline{K} . Then $w_E = (u_{\alpha, \delta_E})|_{K(X)} = v_{\alpha, \delta_E}$.

Proof. We can reduce to proving the statement when K is algebraically closed, so in particular $u = v$. We have to show that $w_E = v_{\alpha, \delta}$, where $\delta = \delta_E$ and $\alpha \in K$ is a pseudo-limit of E . Let $\beta \in K$. To this end, by [17, IV, p. 366], it is sufficient to show that $w_E(X - \alpha + \beta) = \min\{w_E(X - \alpha), v(\beta)\} = \{\delta, v(\beta)\}$. If $\delta \neq v(\beta)$ then this is clear, so suppose that $\delta = v(\beta)$. We have:

$$w_E(X - \alpha + \beta) = \lim_{n \rightarrow \infty} v(s_n - \alpha + \beta) = \lim_{n \rightarrow \infty} v(s_n - \alpha) = \delta$$

so that also in this case we have the claimed equality. \square

We now show under which cases V_E has rank 1 or 2.

Theorem 4.9. *Let $E \subset K$ be a pseudo-convergent sequence.*

- (a) *If E is of transcendental type, then $V_E = W_E$ has rank 1.*
- (b) *If E is of algebraic type and its breadth is infinite, then V_E has rank 2; furthermore, if q is the minimal polynomial of the pseudo-limit of E , then the one-dimensional overring of V_E is $K[X]_{(q)}$.*
- (c) *Suppose that E is of algebraic type with breadth $\delta \in \mathbb{R}$. The following conditions are equivalent:*
 - (i) *δ is not torsion over Γ_v ;*
 - (ii) *W_E is not residually transcendental over V ;*
 - (iii) *V_E has rank one;*
 - (iv) *$V_E = W_E$;*
 - (v) *$V_E \cap K[X] = W_E \cap K[X]$.*

Proof. (a) follows directly from [11, Theorem 2] and the proof of Theorem 3.8, while (b) is a direct consequence of Remark 3.10 (and Proposition 4.4).

(c) Let $E = \{s_n\}_{n \in \mathbb{N}}$ be of algebraic type with finite breadth, and let $\{\delta_n\}_{n \in \mathbb{N}}$ be the gauge of E . Since $V_E \subseteq W_E$ and W_E has rank 1, conditions (iii) and (iv) are clearly equivalent. Since by Lemma 4.8 $w_E = v_{\alpha, \delta}$, by [18, Lemma 3.5] we have that (i) is equivalent to (ii). Clearly, (iv) implies (v).

(v) \implies (iv) Let $\phi \in W_E$, i.e., $w_E(\phi) \geq 0$. Clearly, if $w_E(\phi) > 0$ then $\phi \in V_E$, so suppose $w_E(\phi) = 0$. By Proposition 3.6 and Corollary 4.6, there exist $\lambda \in \mathbb{Z}$ and $\gamma \in \Gamma_v$ such that $v(\phi(s_n)) = \lambda\delta_n + \gamma$ for all large n ; its limit $\lambda\delta + \gamma$ is equal to $w_E(\phi)$, and thus it is 0. If $\lambda \leq 0$, then $v(\phi(s_n))$ is eventually positive, and $\phi \in V_E$.

Suppose $\lambda > 0$ and let $p \in K[X]$ be the minimal polynomial of some pseudo-limit β of E (with respect to some extension u of v in \overline{K}). By Corollary 4.6, there are $\lambda_p \in \mathbb{Z}$, $\gamma_p \in \Gamma_v$ such that $v(p(s_n)) = \lambda_p\delta_n + \gamma_p$ for all large n ; furthermore, $\lambda_p > 0$ since p is a polynomial and one of the roots of $p(X)$ is a pseudo-limit of E . Let $c \in K$ be an element of value $\lambda_p\gamma - \lambda\gamma_p$ (which exists since $\lambda, \lambda_p \in \mathbb{Z}$ and $\gamma, \gamma_p \in \Gamma_v$) and consider $\psi(X) = cp(X)^\lambda \in K[X]$. Then, for all $n \in \mathbb{N}$ sufficiently large we have

$$v(\psi(s_n)) = \lambda_p\gamma - \lambda\gamma_p + \lambda(\lambda_p\delta_n + \gamma_p) = \lambda_p(\gamma + \lambda\delta_n) = \lambda_p v(\phi(s_n)).$$

This quantity has limit 0 as $n \rightarrow \infty$ and is strictly increasing, because $\lambda\lambda_p > 0$; hence $v(\psi(s_n)) < 0$ for all $n \in \mathbb{N}$ sufficiently large. Therefore, $\psi \in W_E \setminus V_E$. However, this contradicts the hypothesis because $\psi(X)$ is a polynomial; hence, the claim is proved.

We show now that (i) \iff (iv), and we claim that it is sufficient to prove the equivalence under the further assumption that E has a pseudo-limit β in K . Suppose that (i) is equivalent to (iv) under this assumption and let $\beta \in \mathcal{L}_E^{v'}$ where v' is an extension of v to $K(\beta)$. Let $V'_E \subseteq W'_E$ be the valuation domains of $K(\beta)(X)$ associated to E with respect to the valuation v' . If δ is not torsion over Γ_v then $V'_E = W'_E$ and contracting down to $K(X)$ we get $V_E = W_E$. Conversely, if the rank of V_E is one (thus, $V_E = W_E$) then also the rank of V'_E is one (because $K(X) \subseteq K(\beta)(X)$ is an algebraic extension) and so δ is not torsion over Γ_v .

Suppose thus that $\beta \in K$ is a pseudo-limit of E . By (5) we have $w_E(X - \beta) = \delta$. If δ is torsion over Γ_v , then $k\delta \in \Gamma_v$ for some $k \in \mathbb{N}$, i.e., there is $c \in K$ such that $w_E((X - \beta)^k) = v(c)$; let $\phi(X) = \frac{(X - \beta)^k}{c}$. Then, $w_E(\phi) = 0$ and thus $\phi \in W_E$, while

$$v\left(\frac{(s_n - \beta)^k}{c}\right) = k\delta_n - v(c) < 0, \quad (6)$$

and thus $\phi(s_n) \notin V$ for every n , which implies $\phi \notin V_E$. Hence, $V_E \neq W_E$.

Conversely, suppose that δ is not torsion over Γ_v , and let $\phi \in W_E$. If $w_E(\phi) > 0$ then ϕ belongs to the maximal ideal of W_E , which is contained in V_E . Suppose $w_E(\phi) = 0$, and let k be the dominating degree of E . By definition we have $w_E(\phi) = k\delta + \gamma$ for some $\gamma \in \Gamma_v$ (see also Corollary 4.6 and (4)); since this quantity is 0 and δ is not torsion, we must have $k = 0$, and so also $\gamma = 0$. But this means that $v(\phi(s_n)) = 0$ for large n ; in particular, $\phi(s_n) \in V$ for large n . Thus $\phi \in V_E$ and $V_E = W_E$. \square

When the rank of V_E is 1, then its valuation v_E is exactly the Ostrowski valuation w_E ; on the other hand, if E is algebraic with infinite breadth, then V_E has been characterized in Remark 3.10 and its valuation is described in [19, Remark 2.3]. When V_E has rank 2 and E has finite breadth, a description of v_E has been obtained in Proposition 3.11; we now want to embed Γ_{v_E} as a totally ordered subgroup in \mathbb{R}^2 , endowed with the lexicographic order (this can be done by Hahn's theorem [21, Théorème 2, p. 22]).

Theorem 4.10. *Let $E = \{s_n\}_{n \in \mathbb{N}}$ be a pseudo-convergent sequence with non-zero breadth ideal such that V_E has rank 2. Then the map*

$$\begin{aligned} \nu: K(X) \setminus \{0\} &\longrightarrow \mathbb{R}^2 \\ \phi &\longmapsto (w_E(\phi), -\deg_{\text{dom}_E}(\phi)) \end{aligned}$$

is a valuation on $K(X)$ whose associated valuation ring is V_E .

Proof. By Theorem 4.9, E is of algebraic type and its breadth δ is torsion over Γ_v . Let $\{\delta_n\}_{n \in \mathbb{N}}$ be the gauge of E . Since w_E is a valuation, we have $w_E(\phi_1\phi_2) = w_E(\phi_1) + w_E(\phi_2)$ for every $\phi_1, \phi_2 \in K(X)$; the same formula holds for the dominating degree, since the multiset of zeros of $\phi_1\phi_2$ is exactly the union of the multisets of zeros of ϕ_1 and ϕ_2 . Hence, $\nu(\phi_1\phi_2) = \nu(\phi_1) + \nu(\phi_2)$.

We now want to show that $\nu(\phi_1 + \phi_2) \geq \min\{\nu(\phi_1), \nu(\phi_2)\}$. Let $\lambda_1 = \deg_{\text{dom}_E}(\phi_1)$, $\lambda_2 = \deg_{\text{dom}_E}(\phi_2)$, $\lambda = \deg_{\text{dom}_E}(\phi_1 + \phi_2)$. By Proposition 3.6, there are $\gamma_1, \gamma_2, \gamma \in \Gamma_v$ such that $v(\phi_i(s_n)) = \lambda_i \delta_n + \gamma_i$, $i = 1, 2$ and $v((\phi_1 + \phi_2)(s_n)) = \lambda \delta_n + \gamma$ for all large n . Furthermore, by Corollary 4.6, $w_E(\phi_i) = \lambda_i \delta + \gamma_i$, $i = 1, 2$ and $w_E(\phi_1 + \phi_2) = \lambda \delta + \gamma$.

We distinguish four cases.

If $w_E(\phi_1) \neq w_E(\phi_2)$, then without loss of generality $w_E(\phi_1) < w_E(\phi_2)$. Hence, we have $v(\phi_1(s_n)) < v(\phi_2(s_n))$ for all large n . Thus,

$$v((\phi_1 + \phi_2)(s_n)) = v(\phi_1(s_n)) = \lambda_1 \delta_n + \gamma_1.$$

Hence, $\lambda_1 \delta_n + \gamma_1 = \lambda \delta_n + \gamma$ infinitely many times. Thus, it must be $\lambda_1 = \lambda$, and so $\nu(\phi_1 + \phi_2) = \nu(\phi_1) = \min\{\nu(\phi_1), \nu(\phi_2)\}$.

If $w_E(\phi_1) = w_E(\phi_2) < w_E(\phi_1 + \phi_2)$, then $\nu(\phi_1 + \phi_2)$ is bigger than both $\nu(\phi_1)$ and $\nu(\phi_2)$, and we are done.

Suppose that $w_E(\phi_1) = w_E(\phi_2) = w_E(\phi_1 + \phi_2)$ and that $\lambda_1 \neq \lambda_2$; without loss of generality, $\lambda_1 > \lambda_2$ (i.e., $\nu(\phi_1) < \nu(\phi_2)$). Then, $\lambda_1 \delta_n + \gamma_1 < \lambda_2 \delta_n + \gamma_2$ for all large n . Therefore, as in the first case, $\lambda_1 \delta_n + \gamma_1 = \lambda \delta_n + \gamma$ for all large n , and so $\nu(\phi_1 + \phi_2) = \nu(\phi_1)$.

Suppose now that $w_E(\phi_1) = w_E(\phi_2) = w_E(\phi_1 + \phi_2)$ and that $\lambda_1 = \lambda_2 =: \lambda'$. Since the sequences $v(\phi_1(s_n)) = \lambda' \delta_n + \gamma_1$ and $v(\phi_2(s_n)) = \lambda' \delta_n + \gamma_2$ have the same limit, they must be eventually equal, and so $\gamma_1 = \gamma_2 =: \gamma'$. Since v is a valuation, $v((\phi_1 + \phi_2)(s_n)) = \lambda \delta_n + \gamma \geq \lambda' \delta_n + \gamma'$. Since $w_E(\phi_1 + \phi_2) = w_E(\phi_1)$, furthermore, the limits $\lambda \delta + \gamma$ and $\lambda' \delta + \gamma'$ are equal; it follows that $\lambda \leq \lambda'$. Hence,

$$\nu(\phi_1 + \phi_2) = (w_E(\phi_1), -\lambda) \geq (w_E(\phi_1), -\lambda') = \nu(\phi_1) = \nu(\phi_2).$$

Therefore, ν is a valuation.

The fact that ν extends v follows from the fact that w_E extends v .

Let V' be the valuation ring associated to ν . Suppose $\phi \in V_E$. If $w_E(\phi) > 0$ then $\nu(\phi) > 0$. If $w_E(\phi) = 0$ then $v(\phi(s_n)) = \lambda \delta_n + \gamma$ must tend to 0 from above, and thus $\lambda \leq 0$, i.e., $\nu(\phi) \geq 0$. Thus, $V_E \subseteq V'$. Conversely, if $\nu(\phi) \geq 0$ then either $w_E(\phi) > 0$ (and so $\phi \in M_{W_E} \subset V_E$) or $w_E(\phi) = 0$ and $\lambda \leq 0$; in the latter case, $\lambda \delta_n + \gamma \geq 0$, and so $v(\phi(s_n)) \geq 0$. Therefore, $V' \subseteq V_E$, and so $V' = V_E$, as claimed. \square

Remark 4.11. Let E be a pseudo-convergent sequence of algebraic type with breadth δ which is torsion over Γ_v . Since ν is a valuation relative to V_E , without loss of generality we can set $v_E = \nu$.

Let $\Delta' = (\delta, -1)$. Take $\phi \in K(X)$ and let $\lambda = \deg_{\text{dom}_E}(\phi)$. By Theorem 4.10, we have

$$v_E(\phi) = (w_E(\phi), -\lambda);$$

by Corollary 4.6, moreover, $w_E(\phi) = \lambda \delta + \gamma_x$ for some $\gamma_x \in \Gamma_v$. It follows that

$$v_E(\phi) = \lambda \Delta' + \gamma,$$

where $\gamma = (\gamma_x, 0) \in \Gamma_{v_E}$. If, furthermore, E has a pseudo-limit $\beta \in K$, then $\Delta' = \Delta = v_E(X - \beta)$.

5 Equivalence of pseudo-convergent sequences

We recall that V is a rank one valuation domain.

Classically, two Cauchy sequences $E, F \subset K$ are equivalent if the distance induced by the valuation v between their corresponding terms goes to zero. If α and β are the limits in \widehat{K} of E and F , respectively, it is known that E and F are equivalent if and only if the valuation domains $V_E = W_\alpha, V_F = W_\beta$ (see Remark 3.10) are the same; in particular, E and F determine the same extension of the valuation v to $K(X)$. Ostrowski investigated in [17, p. 387] the similar problem for the valuation domains of the form W_E , for E a pseudo-convergent sequence in K , which led him to give the notion of equivalent pseudo-convergent sequences. In this section, we consider a definition of equivalence for pseudo-convergent sequence as it appears in [12, Section 3.2], even though we correct a mistake there.

Definition 5.1. Let $E = \{s_n\}_{n \in \mathbb{N}}$ and $F = \{t_n\}_{n \in \mathbb{N}}$ be two pseudo-convergent sequences in K . We say that E and F are *equivalent* if the breadths δ_E and δ_F are equal and, for every $k \in \mathbb{N}$, there are $i_0, j_0 \in \mathbb{N}$ such that, whenever $i \geq i_0, j \geq j_0$, we have

$$v(s_i - t_j) > v(t_{k+1} - t_k).$$

Note that the previous definition boils down to the classical notion of equivalence if E and F are Cauchy sequences.

Remark 5.2. The previous definition was also considered in [12, Section 3.2] without the hypothesis $\delta_E = \delta_F$. However, without this condition the definition is not symmetric: for example, let $F = \{t_n\}_{n \in \mathbb{N}}$ be a sequence in V with $v(t_n) = \delta_n$, where $\{\delta_n\}_{n \in \mathbb{N}}$ is a positive increasing sequence, and let $E = \{s_n = t_n^2\}_{n \in \mathbb{N}}$. Then, for every k and every $i, j \geq k + 1$ we have

$$v(s_i - t_j) = \delta_j > \delta_k = v(t_{k+1} - t_k);$$

on the other hand, if $\delta_k > \frac{1}{2}\delta$, then there are no i, j such that

$$v(s_i - t_j) > 2\delta_k = v(s_{k+1} - s_k);$$

hence, E and F are equivalent according to [12], but F and E are not.

On the other hand, suppose that E and F are two pseudo-convergent sequence of K which are equivalent according to Definition 5.1. Then, for every k there is a k' such that $v(s_{k+1} - s_k) < v(t_{k'+1} - t_{k'})$. If now i_0 and j_0 are such that $v(s_i - t_j) > v(t_{k'+1} - t_{k'})$ for all $i \geq i_0, j \geq j_0$, then clearly $v(s_i - t_j) > v(s_{k+1} - s_k)$, so F and E are equivalent.

We need first the following preliminary lemma.

Lemma 5.3. *Let $E, F \subset K$ two equivalent pseudo-convergent sequences. Then either E and F are both of transcendental type, or E and F are both of algebraic type. In the latter case, $\mathcal{L}_E^u = \mathcal{L}_F^u$ for every extension u of v to \overline{K} .*

Proof. Let $E = \{s_n\}_{n \in \mathbb{N}}$ and $F = \{t_n\}_{n \in \mathbb{N}}$; let $\{\delta_n\}_{n \in \mathbb{N}}, \{\delta'_n\}_{n \in \mathbb{N}}$ be the gauges of E and F , respectively, and δ the breadth of E and F . It is sufficient to prove that if either one of the two pseudo-convergent sequences, say E , is of algebraic type, then also the other is of algebraic type.

Suppose first that K is algebraically closed and let β be a pseudo-limit of E . Fix $k \in \mathbb{N}$. Then there exist $i_0, j_0 \in \mathbb{N}$ such that, for all $m \geq i_0, n \geq j_0$, $v(s_m - t_n) > \delta_k$. For such n and m , suppose also that $m \geq k$. Then $v(t_n - \beta) = v(t_n - s_m + s_m - \beta) \geq \delta_k$. Therefore, $w_F(X - \beta) = \lim_{n \rightarrow \infty} v(t_n - \beta) \geq \delta$.

If $w_F(X - \beta) > \delta$, then there is a n_0 such that, for all $n \geq n_0$, $v(t_n - \beta) > \delta$; since $v(s_m - \beta) = \delta_m < \delta$, this means that, for every m sufficiently large, $v(t_n - s_m) = v(s_m - \beta)$. This would imply that t_n is a pseudo-limit of E for all $n \geq n_0$, and thus that, in particular, $v(t_{n+1} - t_n) \geq \delta$, which is a contradiction since $v(t_{n+1} - t_n) = \delta'_n \nearrow \delta$. Hence, $\delta_n \leq v(t_n - \beta) < \delta$ for all large n and $w_F(X - \beta) = \delta$; this shows that $v(t_n - \beta)$ is eventually strictly increasing, that is, β is a pseudo-limit of F and thus also F is of algebraic type. Moreover, since $\beta \in \mathcal{L}_E$ was arbitrarily chosen, we also have $\mathcal{L}_E \subseteq \mathcal{L}_F$, which shows that these sets are equal since they are closed balls of the same radius (Lemma 2.4).

If now K is not algebraically closed, let u be any extension of v to \overline{K} . Then, E and F are equivalent with respect to u ; applying the previous part of the proof, we have that F is of algebraic type and $\mathcal{L}_E^u = \mathcal{L}_F^u$, as claimed. \square

Theorem 5.4. *Let $E, F \subset K$ be two pseudo-convergent sequences that are of transcendental type or of algebraic type with nonzero breadth ideal. Then, the following are equivalent:*

- (i) E and F are equivalent;
- (ii) $V_E = V_F$;
- (iii) $W_E = W_F$;
- (iv) $w_E = w_F$.

Furthermore, if E and F are of algebraic type, the previous conditions are equivalent to the following:

- (v) $\mathcal{L}_E^u = \mathcal{L}_F^u$ for all extensions u of v to \overline{K} ;
- (vi) $\mathcal{L}_E^u = \mathcal{L}_F^u$ for an extension u of v to \overline{K} .

Proof. As usual, we set $E = \{s_n\}_{n \in \mathbb{N}}$ and $F = \{t_n\}_{n \in \mathbb{N}}$; let $\{\delta_n\}_{n \in \mathbb{N}}, \{\delta'_n\}_{n \in \mathbb{N}}$ be the gauges of E and F , respectively, and δ, δ' the breadths of E, F , respectively. Recall that, by Proposition 4.3, if E and F are of transcendental type, then $V_E = W_E$ and $V_F = W_F$.

The structure of the proof is as follows:

- we first prove (ii) \implies (iii) \iff (iv) \implies (i) in both the algebraic and the transcendental case;
- then we prove (i) \implies (iv) and (iii) \implies (ii) in the transcendental case;

- finally, we prove (i) \implies (v) \implies (vi) \implies (ii) in the algebraic case.

(ii) \implies (iii) and (iv) \implies (iii) are obvious.

(iii) \implies (iv) Suppose there is a $\phi \in K(X)$ such that $w_E(\phi) \neq w_F(\phi)$; without loss of generality, $w_E(\phi) > w_F(\phi)$. We claim that there is a $c \in K$ such that $w_E(\phi) \geq v(c) > w_F(\phi)$. This is obvious if Γ is dense in \mathbb{R} ; otherwise, Γ must be isomorphic to \mathbb{Z} , and V is a discrete valuation ring. In this case, the breadth of E and F must be infinite, and thus (by hypothesis) E and F must be transcendental. However, by [11, Theorem 2], it follows that $V_E = W_E$ is an immediate extension of V ; in particular, the value group of W_E coincide with Γ , and thus we can take a $c \in K$ such that $v(c) = w_E(\phi)$. The existence of c implies that $\frac{\phi}{c} \in W_E$ while $\frac{\phi}{c} \notin W_F$, contradicting $W_E = W_F$. Hence, (iv) holds.

(iv) \implies (i) By definition, for every k and every $l > 0$,

$$\delta'_k = v(t_{k+l} - t_k) = \lim_{n \rightarrow \infty} v(t_n - t_k) = w_F(X - t_k) = w_E(X - t_k) = \lim_{n \rightarrow \infty} v(s_n - t_k). \quad (7)$$

If $\delta_E < \delta_F$, then $\delta'_k > \delta_E$ for large k ; thus,

$$v(s_n - t_{k+1}) = v(s_n - s_{n+1} + s_{n+1} - t_{k+1}) = \delta_n$$

and thus $\delta_n = \delta'_{k+1}$, a contradiction; hence $\delta_E \geq \delta_F$. By symmetry, we have also $\delta_F \geq \delta_E$, and thus $\delta_E = \delta_F = \delta$.

Fix now k , take n'_0 such that $\delta_n > \delta'_k$ for every $n \geq n'_0$; since $\delta'_m > \delta'_k$ if $m > k$, there are $n_0 > n'_0$ and $m_0 > k$ such that $v(s_{n_0} - t_{m_0}) > \delta'_k$. For all $n \geq n_0$, $m \geq m_0$, we have

$$v(s_n - t_m) = v(s_n - s_{n_0} + s_{n_0} - t_{m_0} + t_{m_0} - t_m).$$

The three quantities $v(s_n - s_{n_0})$, $v(s_{n_0} - t_{m_0})$ and $v(t_{m_0} - t_m)$ are all bigger than δ'_k ; hence, so is $v(s_n - t_m)$. Since k was arbitrary, E and F are equivalent.

Suppose now that E is of transcendental type. If (iii) holds, then by the previous part of the proof also (i) holds; thus, by Lemma 5.3 both E and F are of transcendental type, and (ii) follows from Theorem 4.9(a).

If (i) holds, then again F is of transcendental type, and the fact that (iv) holds is exactly [12, Satz 3.10] (though note the slight difference in the definition – see Remark 5.2); we give here a proof for the sake of the reader. Without loss of generality, suppose that K is algebraically closed. In order to show that $w_E(\phi) = w_F(\phi)$ for all $\phi \in K(X)$, it is sufficient to show that $w_E(X - \alpha) = w_F(X - \alpha)$ for every $\alpha \in K$. We have

$$\begin{aligned} w_E(X - \alpha) &= \lim_{n \rightarrow \infty} v(s_n - \alpha) = v(s_n - \alpha), \quad \forall n \geq n_1 \\ w_F(X - \alpha) &= \lim_{n \rightarrow \infty} v(t_n - \alpha) = v(t_n - \alpha), \quad \forall n \geq m_1 \end{aligned}$$

for some $n_1, m_1 \in \mathbb{N}$, since both quantities are eventually constant. We also have that $w_E(X - \alpha)$ and $w_F(X - \alpha)$ are both strictly less δ , since $\alpha \in K$ cannot be a pseudo-limit of E and F , respectively. Hence, there exists $k_1 \in \mathbb{N}$ such that for all $k > k_1$ we have $\delta_k > w_E(X - \alpha)$ and $\delta'_k > w_F(X - \alpha)$. Let $k > \max\{k_1, n_1, m_1\}$. There exists $k_2 \geq k$

such that $\delta'_k < \delta_{k_2}$. Also, there exist $i_0, j_0 \in \mathbb{N}$ such that for each $i \geq i_0$ and $j \geq j_0$ we have $v(s_i - t_j) > \delta'_k$. We have

$$v(s_{k_2} - \alpha) = v(s_{k_2} - s_m + s_m - t_m + t_m - \alpha)$$

Choose $m > \max\{k_2, i_0, j_0\}$. Then $v(s_{k_2} - s_m)$ and $v(s_m - t_m)$ are both strictly bigger than $\delta'_k > v(t_m - \alpha)$. Hence, $w_E(X - \alpha) = v(s_{k_2} - \alpha) = v(t_m - \alpha) = w_F(X - \alpha)$, and the claim is proved.

Suppose now that E is of algebraic type. If (i) holds, then by Lemma 5.3 also F is of algebraic type, and E and F have the same pseudo-limits with respect to any extension u of v to \bar{K} ; hence, (i) \implies (v). Furthermore, (v) \implies (vi) is obvious.

We now show that (vi) implies (ii). Let $\phi \in V_E$. By Proposition 3.6, we have $v(\phi(s_n)) = \lambda\delta_n + \gamma$, where $\lambda = \deg_{\text{dom}_E} \phi$ and $\gamma = u\left(\frac{\phi}{\phi_S}(\beta)\right) \in \Gamma_v$, where β is a pseudo-limit of E (and ϕ_S is defined as in the proposition). Similarly, $v(\phi(t_n)) = \lambda'\delta'_n + \gamma'$; however, since $\mathcal{L}_E^u = \mathcal{L}_F^u$, it follows that $\lambda = \lambda'$ and $\gamma = \gamma'$. Furthermore, by Lemma 2.4, $\delta_E = \delta_F$, and thus $v(\phi(s_n))$ and $v(\phi(t_n))$ have the same limit L as $n \rightarrow \infty$. Since $\phi \in V_E$, we have $v(\phi(s_n)) \geq 0$ for large n , and so $L \geq 0$. If $L > 0$, then also $v(\phi(t_n)) > 0$ for large n ; this implies that $\phi \in V_F$. If $L = 0$, then $\lambda \leq 0$; in particular, it must be $v(\phi(t_n)) \geq 0$ for large n . Again, it follows that $\phi \in V_F$; therefore, $V_E \subseteq V_F$. Symmetrically, $V_F \subseteq V_E$, and thus $V_E = V_F$, as claimed. \square

5.1 A geometric interpretation of equivalence

In this section, we give a geometric interpretation of Theorem 5.4. Let \mathcal{V}_{alg} be the set of the valuation domains V_E , where $E \subset K$ is a pseudo-convergent sequence of algebraic type. Fix an extension u of v to the algebraic closure \bar{K} of K . Then, to every valuation ring $V_E \in \mathcal{V}_{\text{alg}}$ is uniquely associated its set of pseudo-limits $\mathcal{L}_E^u \subset \bar{K}$; furthermore, since $\mathcal{L}_E^u = \beta_E + \text{Br}_u(E)$, where $\beta_E \in \bar{K}$ is a pseudo-limit of E with respect to u (see Lemma 2.4), there is a well-defined and injective map

$$\begin{aligned} \Sigma: \mathcal{V}_{\text{alg}} &\longrightarrow \text{CBall}_u(\bar{K}) \\ V_E &\longmapsto \mathcal{L}_E^u, \end{aligned} \tag{8}$$

where $\text{CBall}_u(\bar{K})$ is the set of closed balls of the ultrametric space \bar{K} , endowed with the metric induced by u .

In general, Σ is not surjective; to find its range, we introduce the following definition. For any $\beta \in \bar{K}$, we consider the minimum distance of the elements of K from β , namely:

$$d_u(\beta, K) = \inf\{d_u(\beta, x) = e^{-u(\beta-x)} \mid x \in K\}.$$

Note that $d_u(\beta, K)$ may be 0 even if $\beta \notin K$: this happens if and only if β is in the completion of K under v . If V is a DVR, then the only closed balls of center $\beta \in \bar{K}$ which can arise as the set of pseudo-limits of a pseudo-convergent sequence $E \subset K$, are those of radius 0 and with $\beta \in \widehat{K}$. If V is non-discrete, we have the following result.

Proposition 5.5. *Let V be a non-discrete rank one valuation domain. Let $\beta \in \overline{K}$, $r \in \mathbb{R}^+$ and u an extension of v to \overline{K} ; let B be the closed ball of center β and radius r with respect to u . Then, $B = \mathcal{L}_E^u$ for some pseudo-convergent sequence $E \subset K$ if and only if $r \geq d_u(\beta, K)$.*

Proof. Suppose $B = \mathcal{L}_E^u$, and let $E = \{s_n\}_{n \in \mathbb{N}}$. Then, $\{d_u(\beta, s_n)\}_{n \in \mathbb{N}}$ is a decreasing sequence of real numbers with limit $e^{-\delta} = r$, where δ is the breadth of E . By definition,

$$d_u(\beta, K) = \inf\{d_u(\beta, s) \mid s \in K\} \leq d_u(\beta, s_n)$$

for every n , and thus $d_u(\beta, K) \leq r$.

Conversely, suppose $r \geq d_u(\beta, K)$. If $r = d(\beta, x)$ for some $x \in K$, take a sequence $Z = \{z_k\}_{k \in \mathbb{N}} \subseteq K$ such that $v(z_k)$ is increasing and has limit $\delta = -\log(r)$. Then, $x + Z = \{x + z_k\}_{k \in \mathbb{N}}$ is a pseudo-convergent sequence whose set of limits (in (\overline{K}, u)) is B .

If $r \neq d(\beta, x)$ for every $x \in K$, we can take a sequence $E = \{s_n\}_{n \in \mathbb{N}}$ such that $d_u(\beta, s_n) = r_n$ decreases to r . Then, E is a pseudo-convergent sequence, and $\mathcal{L}_E^u = B$. \square

If V is not discrete, the next corollary gives a necessary and sufficient condition in order for the map defined in (8) to be surjective. Since V has rank one, \widehat{K} is algebraically closed if and only if the residue field of V is algebraically closed and the value group of V is divisible (see [20, Remark 6.4, 3]), for example).

Corollary 5.6. *Suppose V is not discrete. Then the map Σ defined in (8) is surjective if and only if \widehat{K} is algebraically closed.*

Proof. By Proposition 5.5, Σ is surjective if and only if \widehat{K} contains an algebraic closure of K . This happens if and only if \widehat{K} is algebraically closed. \square

5.2 Extension of an Ostrowski valuation

Let w_E be the Ostrowski valuation on $K(X)$ associated to a pseudo-convergent sequence $E \subset K$, and let u be an extension of v to \overline{K} . The *extension of w_E to $\overline{K}(X)$ along u* is the valuation \overline{w}_E defined by

$$\overline{w}_E(\psi) = \lim_{n \rightarrow \infty} u(\psi(s_n))$$

for every $\psi \in \overline{K}(X)$. Clearly, \overline{w}_E extends w_E and has rank 1. A consequence of Theorem 5.4 is that, if E and F are two pseudo-convergent sequences in K , the equality $w_E = w_F$ implies $\overline{w}_E = \overline{w}_F$, since these equalities are both equivalent to the fact that E and F are equivalent pseudo-convergent sequences (which does not depend on the field containing E and F).

We show now that any extension of an Ostrowski valuation on $K(X)$ to $\overline{K}(X)$ is of this kind.

Theorem 5.7. *Let $E = \{s_n\}_{n \in \mathbb{N}} \subset K$ be a pseudo-convergent sequence such that the associated map w_E is a valuation. If \bar{w} is an extension of w_E to $\bar{K}(X)$ and u is the restriction of \bar{w} to \bar{K} , then \bar{w} is equivalent to the extension \bar{w}_E of w_E to $\bar{K}(X)$ along u (or, equivalently, the valuation domain of \bar{w} is equal to \bar{W}_E).*

Proof. The valuation \bar{w}_E restricts to w_E on $K(X)$ and to u on \bar{K} . Suppose there is another valuation w' on $\bar{K}(X)$ with these properties: then, by [5, Chapt. VI, §8, 6., Corollaire 1], there is a $K(X)$ -automorphism σ of $\bar{K}(X)$ such that $w' \circ \sigma$ is equivalent to \bar{w}_E , that is, $\rho(\bar{W}_E) = W'$, where $\rho = \sigma^{-1}$ and W' is the valuation ring of w' .

Now

$$\begin{aligned} \rho(\bar{W}_E) &= \{\rho \circ \phi \in \bar{K}(X) \mid \lim_n u(\phi(s_n)) \geq 0\} = \\ &= \{\rho \circ \phi \in \bar{K}(X) \mid \lim_n u \circ \sigma \circ \rho(\phi(s_n)) \geq 0\}. \end{aligned}$$

Since $s_n \in K$ and $\rho|_K$ is the identity, $\rho(\phi(s_n)) = (\rho \circ \phi)(s_n)$; hence,

$$\begin{aligned} \rho(\bar{W}_E) &= \{\rho \circ \phi \in \bar{K}(X) \mid \lim_n (u \circ \sigma)((\rho \circ \phi)(s_n)) \geq 0\} = \\ &= \{\psi \in \bar{K}(X) \mid \lim_n (u \circ \sigma)(\psi(s_n)) \geq 0\}. \end{aligned}$$

Since both \bar{W}_E and $W' = \rho(\bar{W}_E)$ are extensions of U , the valuation domain of u , we have $u(t) = (u \circ \sigma)(t)$ for every $t \in \bar{K}$; in particular, this happens for $t = \psi(s_n)$. It follows that $\rho(\bar{W}_E) = W' = \bar{W}_E$, as claimed. \square

Remark 5.8. We note that it is possible for two valuations w_1, w_2 of $\bar{K}(X)$ to be different even if their restriction to $K(X)$ and \bar{K} are equal. For example, let v be a valuation on K , and let w be an extension of v to $K(X)$. If K is complete under the topology induced by v , then there exists a unique extension of v to \bar{K} ; on the other hand, w can have more than one extension to $\bar{K}(X)$.

For an explicit example, suppose that K is complete under v , let \bar{v} be the unique extension of v to \bar{K} and let \bar{V} be the valuation domain of \bar{K} associated to \bar{v} . Let $\alpha, \beta \in \bar{K}$ be two distinct elements which are conjugate over K , and let w be the valuation associated to the valuation domain

$$W = \{\phi \in K(X) \mid \phi(\alpha) \in \bar{V}\} = \{\phi \in K(X) \mid \phi(\beta) \in \bar{V}\};$$

note that the second equality follows from the fact that α and β are conjugate over K (see also [19, Theorem 3.2], where such valuation domains are studied; note that they belong to the same class of the valuation domains considered in Remark 3.10). Then, W extends to the following valuation rings of $\bar{K}(X)$:

$$\bar{W}_\alpha = \{\psi \in \bar{K}(X) \mid \psi(\alpha) \in \bar{V}\}, \quad \bar{W}_\beta = \{\psi \in \bar{K}(X) \mid \psi(\beta) \in \bar{V}\}.$$

However, $\bar{W}_\alpha \neq \bar{W}_\beta$: for example, if $t \in \bar{K}$ satisfies $\bar{v}(t) > \bar{v}(\beta - \alpha)$, then $f(X) = \frac{1}{t}(X - \alpha)$ belongs to \bar{W}_α but not to \bar{W}_β (again, the same conclusion follows from the aforementioned result [19, Theorem 3.2]).

By means of Theorem 5.7, in the next result without loss of generality we assume that the extension of w_E to $\overline{K}(X)$ is equal to \overline{w}_E (for some extension u of v to \overline{K} ; clearly, $u = (\overline{w}_E)|_{\overline{K}}$).

The following is a variant of Theorem 3.3.

Proposition 5.9. *Let $\phi \in K(X)$ and let $E = \{s_n\}_{n \in \mathbb{N}} \subset K$ be a pseudo-convergent sequence. Let \overline{w}_E be an extension of w_E to $\overline{K}(X)$, and let $\theta_1, \theta_2 \in \mathbb{R}$ be such that $C = \{t \in \overline{K} \mid \theta_1 < \overline{w}_E(X - t) < \theta_2\}$ does not contain any critical point of ϕ . Then, there are $\lambda \in \mathbb{Z}$, $\gamma \in \mathbb{Q}\Gamma_v$ such that*

$$v(\phi(t)) = \lambda w_E(X - t) + \gamma$$

for every $t \in K \cap C$. More precisely, if S is the multiset of critical points α of ϕ such that $\overline{w}_E(X - \alpha) \geq \theta_2$, then λ is the weighted sum of S and $\gamma = \overline{w}_E\left(\frac{\phi}{\phi_S}\right)$.

Proof. Let $\phi(X) = c \prod_{\alpha \in S} (X - \alpha)^{\epsilon_\alpha} \prod_{\beta \in S'} (X - \beta)^{\epsilon_\beta}$, where S' is the multiset of critical points of ϕ with $\overline{w}_E < \theta_2$. Let $t \in K \cap C$ and let u be the restriction of \overline{w}_E to \overline{K} . As in the proof of Theorem 3.3, writing $u(t - \alpha) = \overline{w}_E(t - \alpha) = \overline{w}_E(t - X + X - \alpha)$ we see that $u(t - \alpha) = w_E(X - t)$ if $\overline{w}_E(X - \alpha) \geq \theta_2$, while $u(t - \alpha) = \overline{w}_E(X - \alpha)$ if $\overline{w}_E(X - \alpha) \leq \theta_1$ (note that by assumption on C there is no critical point α of ϕ such that $\theta_1 < \overline{w}_E(X - \alpha) < \theta_2$). Hence,

$$v(\phi(t)) = v(c) + \sum_{\alpha \in S} \epsilon_\alpha w_E(X - t) + \sum_{\beta \in S'} \epsilon_\beta \overline{w}_E(X - \beta) = \lambda w_E(X - t) + \gamma,$$

as claimed. \square

Given a pseudo-convergent sequence $E \subset K$, an extension \overline{w}_E of w_E and a rational function $\phi \in K(X)$, we define

$$\delta_{\phi, E} = \max\{\overline{w}_E(X - \alpha) \mid \alpha \text{ is a critical point of } \phi\}$$

(which we simply write δ_ϕ if E is understood from the context). By Remark 4.7(a), $\delta_\phi \leq \delta$, and $\delta_\phi < \delta$ if no critical point of ϕ is a pseudo-limit of E ; in particular, this happens if E is of transcendental type.

Corollary 5.10. *Let $E \subset K$ be a pseudo-convergent sequence and let $\phi \in K(X)$, and suppose that none of the critical points of ϕ is a pseudo-limit of E (with respect to $u = (\overline{w}_E)|_{\overline{K}}$). Then:*

- (a) if $\delta_\phi < w_E(X - t) \leq \delta_E$, then $v(\phi(t)) = w_E(\phi)$;
- (b) if E is of algebraic type and $\alpha \in \mathcal{L}_E^u$, then $w_E(\phi) = u(\phi(\alpha))$.

Proof. Let $E = \{s_n\}_{n \in \mathbb{N}}$. Since no critical point β of ϕ satisfies $\overline{w}_E(X - \beta) \geq \delta_\phi$, by Proposition 5.9 we have $v(\phi(t)) = \overline{w}_E(\phi) = w_E(\phi)$ for every $t \in C = \{s \in \overline{K} \mid \delta_\phi < \overline{w}_E(X - s) \leq \delta_E\}$, so claim (a) is proved. Claim (b) follows by Proposition 3.6(b). \square

6 Spaces of valuation domains associated to pseudo-convergent sequences

We are now interested in studying, from a topological point of view, the sets formed by the valuation rings V_E and W_E induced by the pseudo-convergent sequences E in K ; again we are still assuming that V is a rank one valuation domain. The topologies we are interested in are the Zariski and the constructible topologies (see Section 2 for the definitions). Since we are mainly interested in the former, unless stated otherwise, all the spaces are endowed with the Zariski topology.

We set:

$$\mathcal{V} = \{V_E \mid E \subset K \text{ is a pseudo-convergent sequence}\}$$

and

$$\mathcal{W} = \{W_E \mid E \subset K \text{ is a pseudo-convergent sequence and } w_E \text{ is a valuation}\}.$$

By the results of Section 4, the elements of \mathcal{W} are the rings W_E , when $E \subset K$ is a pseudo-convergent sequence which is either of transcendental type or of algebraic type and non-zero breadth ideal.

When V is discrete, we have the following result.

Theorem 6.1. *[19, Theorem 3.4] Let V be a DVR. Then, \mathcal{V} is homeomorphic to \widehat{K} .*

The homeomorphism can also be described explicitly: indeed, if V is a DVR then \mathcal{V} contains only the rings of the form W_α (see Remark 3.10) and we just send W_α to α . Furthermore, in this context, \mathcal{W} is a subset of \mathcal{V} , and corresponds to the elements of \widehat{K} that are transcendental over V . In view of these facts, we are mainly interested in the case when V is not discrete.

Remark 6.2. If V is discrete and \widehat{K} is algebraic over K (for example, if K is complete; however, this is not a necessary condition, see [22, §1, p. 394]) then \widehat{K} contains no elements that are transcendental over K , and thus \mathcal{W} is empty. Conversely, if \mathcal{W} is empty then V must be discrete, otherwise we have rings W_E coming from non-Cauchy pseudo-convergent sequences E , and \widehat{K} must be algebraic over K .

From now on, we assume that \mathcal{W} is nonempty.

We start by studying \mathcal{W} : indeed, the fact that every Ostrowski valuation domain W_E has rank one has strong consequences on the topology of \mathcal{W} . Recall that a topological space X is said to be *zero-dimensional* if it is T_1 and each point $x \in X$ has a neighborhood base consisting of open-closed sets, or, equivalently, if, for each $x \in X$ and closed set $C \subset X$, there exists an open-closed set containing x and not meeting C [9, f-6].

Proposition 6.3. *The Zariski and the constructible topologies agree on \mathcal{W} . In particular, \mathcal{W} is a zero-dimensional space.*

Proof. The intersection of the maximal ideals of the elements of \mathcal{W} contains the maximal ideal M of V , and thus it is nonzero. Since every W_E has rank 1, the claims follow by [16, Proposition 2.4(b)] and the definition of zero-dimensional space. \square

Let $S \subseteq K$ be a subset. The ring of *integer-valued rational functions* on S is the ring

$$\text{Int}^R(S, V) = \{\phi \in K(X) \mid \phi(S) \subseteq V\}$$

(see [6, Chapter X] for a general reference). Note that $\text{Int}^R(S, V)$ may be equal only to V (for example if $S = K$ and K is algebraically closed, see [7, Proposition 2.4]).

Proposition 6.4. *The space \mathcal{W} is not compact.*

Proof. We claim that

$$\bigcap_{\substack{E \subseteq K \\ E \text{ pseudo-conv.}}} W_E = \text{Int}^R(K, V),$$

Let $\phi \in \text{Int}^R(K, V)$. Then, clearly $\phi \in V_E \subseteq W_E$ for all pseudo-convergent sequences E , by definition of V_E . Conversely, if $\phi(K) \not\subseteq V$, then there is a $t \in K$ such that $\phi(t) \notin V$; since V is closed in K and a rational function induces a continuous function (from the subset of K on which it is defined to K), there is a ball $B(t, r)$ such that $\phi(s) \notin V$ for all $s \in B(t, r)$. Choose $s \in B(t, r)$ which is not a critical point of ϕ and let $E = \{s_n\}_{n \in \mathbb{N}} \subset K$ be a pseudo-convergent Cauchy sequence with limit s . Then $w_E(\phi) = \lim v(\phi(s_n)) = v(\phi(s)) < 0$, that is, $\phi \notin W_E$.

The intersection of the maximal ideals of the Ostrowski valuation overrings is $M \neq (0)$; hence, if \mathcal{W} is compact then by [16, Theorem 5.3] $\text{Int}^R(K, V)$ is a one-dimensional Prüfer domain with quotient field $K(X)$. Hence all rings between $\text{Int}^R(K, V)$ and $K(X)$ have dimension (at most) 1. However, if E is a pseudo-convergent Cauchy sequence with limit in K , then V_E has dimension 2 (by Theorem 4.9(b)) but $\text{Int}^R(K, V) \subset V_E$, a contradiction. Therefore, \mathcal{W} is not compact, as claimed. \square

Remark 6.5. When V is discrete and countable, the space \mathcal{W} is not even locally compact. Indeed, as V is discrete, \mathcal{W} is a subset of \mathcal{V} , and in the homeomorphism of \mathcal{V} into \widehat{K} (Theorem 6.1) \mathcal{W} corresponds to the subset X of those elements which are transcendental over K .

Furthermore, when V is countable, also the algebraic closure is countable, while the completion \overline{K} and all its open subsets are uncountable. Hence, every open ball contains elements that are transcendental over K , and thus X is dense.

If X were locally compact, then by [26, Theorem 18.4] it should be an intersection of an open set O and a closed set C of \widehat{K} (since \widehat{K} is Hausdorff). Since X is dense, we should have $C = \widehat{K}$, and thus $X = O$ should be open. However, also $\widehat{K} \setminus X$ is dense (since all elements of \widehat{K} are limits of sequences in K) and thus X cannot be open. Therefore, \mathcal{W} is not locally compact.

We conjecture that \mathcal{W} , if nonempty, is never locally compact.

In order to study more closely the Zariski topology, we now want to study convergence of sequences of valuation domains. To do so, we give in the next two lemmas two criteria to establish when V_E belongs to $B(\phi)$. We introduce the following notation: if $\beta \in K$, $\gamma_1 \in \Gamma_v$ and $\gamma_2 \in \Gamma_v \cup \{\infty\}$ with $\gamma_1 < \gamma_2$, the *annulus* of center β and radii γ_1 and γ_2 is

$$\mathcal{C}_v(\beta, \gamma_1, \gamma_2) = \{s \in K \mid \gamma_1 < v(\beta - s) < \gamma_2\}.$$

Note that this definition is a special case of the definition given in (2), when V has rank one. When the valuation v is understood from the context, we shall write simply $\mathcal{C}(\beta, \gamma_1, \gamma_2)$ for $\mathcal{C}_v(\beta, \gamma_1, \gamma_2)$.

Lemma 6.6. *Let $E \subset K$ be a pseudo-convergent sequence of algebraic type with breadth δ , let $\beta \in \mathcal{L}_E^u$ and let $\phi \in K(X)$. The following are equivalent:*

- (i) $\phi \in V_E$;
- (ii) there are $\theta_1 \in \mathbb{Q}\Gamma_v$, $\theta_2 \in \mathbb{Q}\Gamma_v \cup \{\infty\}$ such that $\theta_1 < \delta \leq \theta_2$ and such that $\phi(s) \in V$ for all $s \in \mathcal{C}_u(\beta, \theta_1, \theta_2)$;
- (iii) there is $\tau \in \Gamma_v$, $0 < \tau < \delta$ such that $\phi(s) \in V$ for all $s \in \mathcal{C}_u(\beta, \tau, \delta)$.

Proof. (i) \implies (ii) Let $\zeta_1 < \zeta_2$ be two elements in $\mathbb{Q}\Gamma_v$ such that $\zeta_1 < \delta \leq \zeta_2$ and there is no critical point of ϕ in $C = \mathcal{C}_u(\beta, \zeta_1, \zeta_2)$. By Theorem 3.3, there are $\lambda \in \mathbb{Z}$, $\gamma \in \Gamma_v$ such that $v(\phi(s)) = \lambda u(\beta - s) + \gamma$ for all $s \in C$. Let $I = \{h \in (\zeta_1, \zeta_2) \mid \lambda h + \gamma \geq 0\}$; then, I is an interval with endpoints $\theta_1, \theta_2 \in \mathbb{Q}\Gamma_v$, and $\phi(s) \in V$ for all $s \in C' = \mathcal{C}_u(\beta, \theta_1, \theta_2)$; we need only to show that $\theta_1 < \delta \leq \theta_2$.

Since $\phi \in V_E$, and $s_n \in C'$ for large n , we have $\lambda \delta_n + \gamma \geq 0$ for all n large enough, where $\{\delta_n\}_{n \in \mathbb{N}}$ is the gauge of E ; since $\delta_n \nearrow \delta$ and $\zeta_1 < \delta$, it follows that there is an interval $(\tau, \delta) \subseteq I$, and so $\theta_1 < \delta \leq \theta_2$. The claim is proved.

(ii) \implies (iii) is obvious.

(iii) \implies (i) Suppose that there is an annulus $C = \mathcal{C}_u(\beta, \tau, \delta)$ with this property. Since δ is the breadth of E , for large n we have $s_n \in C$; hence, $\phi(s_n) \in V$ and thus $\phi \in V_E$. \square

Remark 6.7. The exact same proof of the previous proposition can be used to show a converse: $\phi \notin V_E$ if and only if there is an annulus $\mathcal{C} = \mathcal{C}(\beta, \tau, \delta)$ such that $\phi(t) \notin V$ for all $t \in \mathcal{C}$ (and similarly for the version with θ_1 and θ_2).

The following technical lemma is based on the Ostrowski valuation \overline{w}_E . Note that the set C defined below is essentially an annulus with respect to \overline{w}_E .

Lemma 6.8. *Let $E = \{s_n\}_{n \in \mathbb{N}} \subset K$ be a pseudo-convergent sequence, and let $\phi \in K(X)$; let u be an extension of v to \overline{K} and let \overline{w}_E be the extension of w_E to $\overline{K}(X)$ along u . There is a $\delta' < \delta_E$ such that, given $C = \{s \in \overline{K} \mid \delta' < \overline{w}_E(X - s) < \delta_E\}$, whenever F is a pseudo-convergent sequence such that $\delta_F \geq \delta_E$ and $\mathcal{L}_F \cap C \neq \emptyset$, we have $\phi \in V_F$ if and only if $\phi \in V_E$.*

Proof. Let $S \subset \overline{K}$ be the multiset of critical points of ϕ , and let $\delta_1 = \sup\{\overline{w}_E(X - \alpha) \mid \alpha \in S, \overline{w}_E(X - \alpha) < \delta_E\}$. Then, there are no critical points of ϕ in $C_1 = \{s \in \overline{K} \mid \delta_1 < \overline{w}_E(X - s) < \delta_E\}$; by Proposition 5.9, there are $\lambda \in \mathbb{Z}$, $\gamma \in \mathbb{Q}\Gamma_v$ such that $u(\phi(t)) = \lambda \overline{w}_E(X - t) + \gamma$ for every $t \in C_1$. Since $v(\phi(s_n)) \rightarrow w_E(\phi)$ and s_n is eventually in C_1 , we can find $\delta' \in [\delta_1, \delta_E)$ such that the quantity $u(\phi(t))$ is either positive, negative or zero for all $t \in C = \{s \in \overline{K} \mid \delta' < \overline{w}_E(X - s) < \delta_E\}$.

Suppose now $F = \{t_m\}_{m \in \mathbb{N}} \subset K$ is a pseudo-convergent sequence with breadth $\delta_F \geq \delta_E$ and such that $\mathcal{L}_F \cap C \neq \emptyset$: then, if $t \in \mathcal{L}_F \cap C$, we have $\overline{w}_E(X - t_m) = \overline{w}_E(X -$

$t + t - t_m) = \bar{w}_E(X - t)$, for all $m \in \mathbb{N}$ sufficiently large, since $\bar{w}_E(X - t) < \delta_E$ and $v(t_m - t) \nearrow \delta_F$ which is greater than or equal to δ (and so, it is eventually greater than $\bar{w}_E(X - t)$). Hence, t_m is eventually in C and thus $v(\phi(t_m))$ is eventually nonnegative if so is $v(\phi(s_n))$ (in which case $\phi \in V_E \cap V_F$), while it is eventually negative if $v(\phi(s_n))$ is eventually negative (and so $\phi \notin V_E$ and $\phi \notin V_F$). The claim is proved. \square

We note that, when we are in the hypothesis of Corollary 5.10 (that is, if ϕ has no critical point which is a pseudo-limit of E), the value δ' of the previous proposition can be taken to be equal to $\delta_{\phi, E}$.

As a first step in the study of \mathcal{V} , we analyze the convergence of sequences in $\text{Zar}(K(X)|V)^{\text{cons}}$.

Proposition 6.9. *Let $E = \{s_n\}_{n \in \mathbb{N}} \subset K$ be a pseudo-convergent sequence of algebraic type with breadth δ , and, for each $n \in \mathbb{N}$, let $\zeta_n \in [\delta, \infty]$. For each $n \in \mathbb{N}$, let $E_n \subset K$ be a pseudo-convergent sequence with pseudo-limit s_n and breadth ζ_n . Then:*

- (a) V_E is a limit of $\{V_{E_n}\}_{n \in \mathbb{N}}$ in $\text{Zar}(K(X)|V)^{\text{cons}}$;
- (b) if $\zeta_n \neq \infty$ for every n , then V_E is a limit of $\{W_{E_n}\}_{n \in \mathbb{N}}$ in $\text{Zar}(K(X)|V)^{\text{cons}}$.

Note that, if V_E is a limit in the constructible topology, it is a limit also in the Zariski topology.

Proof. Let $\mathcal{X} = \text{Zar}(K(X)|V)^{\text{cons}}$; we need to show that, if $V_E \in \Omega$ for some open set Ω , then $V_{E_n}, W_{E_n} \in \Omega$ for large n ; without loss of generality, we can consider only the cases $\Omega = B(\phi)$ and $\Omega = \mathcal{X} \setminus B(\phi)$, where $\phi \in K(X)$. This amounts to prove that $V_E \in B(\phi)$ if and only if $V_{E_n} \in B(\phi)$ (respectively, $W_{E_n} \in B(\phi)$) for all large n .

Suppose first that E has a pseudo-limit $s \in K$. By Proposition 6.6, there is an annulus $\mathcal{C} = \mathcal{C}(s, \tau, \delta)$ such that $\phi(t) \in V$ for all $t \in \mathcal{C}$. There is a N such that $s_n \in \mathcal{C}$ for $n \geq N$; hence, for these n , $\mathcal{L}_{E_n} \cap \mathcal{C} \neq \emptyset$. For all $t \in \mathcal{C}$, we have $w_E(X - t) = u(s - t)$; hence, $\mathcal{C} = \{t \in K \mid \tau < w_E(X - t) < \delta\}$. Therefore, we can apply Lemma 6.8, and so $V_E \in B(\phi)$ if and only if $V_{E_n} \in B(\phi)$ for $n \geq N$. Thus, the sequence V_{E_n} tends to V_E in the constructible topology.

Since $V_{E_n} \subseteq W_{E_n}$, we also have that if $V_E \in B(\phi)$ then $W_{E_n} \in B(\phi)$ for large n . Furthermore, without loss of generality, \mathcal{C} does not contain any critical point of ϕ and $v(\phi(t)) = \lambda v(t - s) + \gamma$, for each $t \in \mathcal{C}$, for some $\lambda \in \mathbb{Z}$ and $\gamma \in \Gamma_v$ by Theorem 3.3; since $v(t_m - s) = v(t - s)$, for all $m \in \mathbb{N}$ sufficiently large, where $t \in \mathcal{L}_F \cap \mathcal{C}$, then $v(\phi(t_m)) = v(\phi(t))$ for all such m , and so $w_F(\phi) = v(\phi(t))$: hence, if $V_E \notin B(\phi)$ then also $W_{E_n} \notin B(\phi)$. It follows that also the sequence W_{E_n} tends to V_E in $\text{Zar}(K(X)|V)^{\text{cons}}$.

Suppose now that E has a limit $\beta \in \bar{K}$ with respect to some extension u of v to \bar{K} ; let $U \subset \bar{K}$ be the valuation domain of u . By the previous part of the proof, U_E is the limit of the sequence U_{E_n} in $\text{Zar}(\bar{K}(X)|U)^{\text{cons}}$. The restriction map $\pi : \text{Zar}(\bar{K}(X)|U)^{\text{cons}} \rightarrow \text{Zar}(K(X)|V)^{\text{cons}}$, $W \mapsto W \cap K(X)$, is continuous; hence, $\pi(U_{E_n}) \rightarrow \pi(U_E)$. However, $\pi(U_{E_n}) = V_{E_n}$ and $\pi(U_E) = V_E$; the claim is proved. The same reasoning applies to the sequence $\{W_{E_n}\}_{n \in \mathbb{N}}$.

The claim about the Zariski topology follows since the constructible topology is finer than the Zariski topology. \square

Example 6.10. Let $E = \{s_n\}_{n \in \mathbb{N}}$ be a pseudo-convergent sequence of algebraic type and, for each $n \in \mathbb{N}$, let $W_{s_n} = \{\phi \in K(X) \mid \phi(s_n) \in V\}$. Then, by the previous lemma, $\{W_{s_n}\}_{n \in \mathbb{N}}$ converges to V_E in the constructible and in the Zariski topology.

Since we are working with the Zariski topology on \mathcal{V} and \mathcal{W} , for ease of notation we set

$$\begin{aligned} B^{\mathcal{V}}(\phi) &= \{V_E \in \mathcal{V} \mid V_E \ni \phi\} = B(\phi) \cap \mathcal{V}, \\ B^{\mathcal{W}}(\phi) &= \{W_E \in \mathcal{W} \mid W_E \ni \phi\} = B(\phi) \cap \mathcal{W}. \end{aligned}$$

We denote by $\mathcal{V}(\bullet, \delta)$ the set of valuation domains V_E such that E has breadth δ .

Proposition 6.11. *Let V be a valuation domain of rank 1 which is not discrete. The following are equivalent:*

- (i) *the residue field of V is finite;*
- (ii) *the Zariski and the constructible topologies coincide on \mathcal{V} ;*
- (iii) *there is a $\delta \in \mathbb{R} \cup \{+\infty\}$ such that the Zariski and the constructible topologies coincide on $\bigcup_{\delta' \leq \delta} \mathcal{V}(\bullet, \delta')$;*
- (iv) *there is a $\delta \in \mathbb{R} \cup \{+\infty\}$ such that the Zariski and the constructible topologies coincide on $\bigcup_{\delta' < \delta} \mathcal{V}(\bullet, \delta')$.*

When V is discrete, \mathcal{V} reduces to $\mathcal{V}(\bullet, \infty)$.

Proof. (i) \implies (ii) To show that the Zariski and the constructible topologies coincide, it is enough to show that $B(\phi)$ is closed in the Zariski topology for every $\phi \in K(X)$. Let thus $E = \{s_n\}_{n \in \mathbb{N}}$ be a pseudo-convergent sequence with breadth δ such that $V_E \notin B(\phi)$; we want to show that there is an open neighborhood of V_E disjoint from $B(\phi)$.

If E is of transcendental type, then $V_E = W_E$; since the Zariski and the constructible topologies agree on \mathcal{W} (Proposition 6.3), the set $B^{\mathcal{W}}(\phi)$ is closed in \mathcal{W} , and thus there are ψ_1, \dots, ψ_k such that $W_E \in B^{\mathcal{W}}(\psi_1, \dots, \psi_k)$ but $B^{\mathcal{W}}(\psi_1, \dots, \psi_k) \cap B^{\mathcal{W}}(\phi) = \emptyset$. In particular, $V_E \in B^{\mathcal{V}}(\psi_1, \dots, \psi_k)$; on the other hand, if $V_F \in B^{\mathcal{V}}(\psi_1, \dots, \psi_k) \cap B^{\mathcal{V}}(\phi)$, then $\psi_1, \dots, \psi_k, \phi \in V_F \subseteq W_F$, and thus $W_F \in B^{\mathcal{W}}(\psi_1, \dots, \psi_k) \cap B^{\mathcal{W}}(\phi)$, a contradiction. Hence, $B^{\mathcal{V}}(\psi_1, \dots, \psi_k)$ and $B^{\mathcal{V}}(\phi)$ are disjoint, and $B^{\mathcal{V}}(\psi_1, \dots, \psi_k)$ is the required neighborhood.

Suppose E is of algebraic type without pseudo-limits in K ; let $\alpha \in \overline{K} \setminus K$ be a pseudo-limit of E with respect to an extension u of v to \overline{K} . By Proposition 6.6 and Remark 6.7, there is an annulus $\mathcal{C} = \mathcal{C}_u(\alpha, \theta_1, \theta_2)$ with $\theta_1, \theta_2 \in \mathbb{Q}\Gamma_v$, $\theta_1 < \delta \leq \theta_2$, such that $\phi(t) \notin V$ for all $t \in \mathcal{C}$. Let $s \in \mathcal{C}$; then $\theta_1 < u(\alpha - s) < \delta$, because otherwise s would be a pseudo-limit of E . Let $d \in K$ be such that $\theta_1 < v(d) < u(\alpha - s)$. Then, $V_E \in B(\frac{X-s}{d})$, since, for large n , $v(s_n - s) - v(d) = u(s_n - \alpha + \alpha - s) - v(d) = u(\alpha - s) - v(d) \geq 0$. On the other hand, if $t \in K$ is such that $v(\frac{t-s}{d}) \geq 0$, then $v(t - s) \geq v(d) > \theta_1$, so $u(t - \alpha) = u(t - s + s - \alpha) > \theta_1$. Since $u(t - \alpha) < \delta$ because E has no pseudo-limits in

K , it follows that $t \in \mathcal{C}$, so that $\phi(t) \notin V$; in particular, $B\left(\frac{X-s}{d}\right)$ is a neighborhood of V_E disjoint from $B(\phi)$.

Suppose now that E is of algebraic type with a pseudo-limit $s \in K$. If $\delta \notin \mathbb{Q}\Gamma_v$, then $V_E = W_E$ by Theorem 5.4, so the claim follows as in the transcendental case. Suppose $\delta \in \mathbb{Q}\Gamma_v$, and let $k \in \mathbb{N}^+$ be such that $k\delta \in \Gamma_v$. By Proposition 6.6 and Remark 6.7, there is an annulus $\mathcal{C}(s, \tau, \delta)$, with $\tau < \delta$, such that $\phi(t) \notin V$ for all $t \in \mathcal{C}(s, \tau, \delta)$. Let $d \in K$ be an element such that $v(d) \in (\tau, \delta)$; then, $V_E \in B\left(\frac{X-s}{d}\right)$.

Let u_1, \dots, u_r be a complete set of representatives for the residue field of V ; suppose that $u_1 \in M$ and $u_i \in V \setminus M$ for $i = 2, \dots, r$. Let $z \in K$ be an element of valuation $k\delta$. Let

$$\psi(X) = \frac{z^r}{((X-s)^k - zu_1) \cdots ((X-s)^k - zu_r)};$$

we claim that $\psi(t) \in V$ if and only if $v(t-s) < \delta$.

Indeed, if $v(t-s) < \delta$ then $v((t-s)^k) = kv(t-s) < k\delta \leq v(zu_i)$ for $i = 1, \dots, r$, and thus

$$v(\psi(t)) = rk\delta - rk v(t-s) > 0.$$

If $v(t-s) > \delta$, then $v((t-s)^k - zu_i) = k\delta$ for $i = 2, \dots, r$ and $v((t-s)^k - zu_1) > k\delta$, and thus

$$v(\psi(t)) < rk\delta - rk\delta = 0.$$

If $v(t-s) = \delta$, then $v((t-s)^k) = k\delta = v(z)$; since u_1, \dots, u_r are a complete set of representatives, there is a (unique) $i \in \{2, \dots, r\}$ such that $v((t-s)^k - zu_i) > k\delta$, while $v((t-s)^k - zu_j) = k\delta$ for $j \in \{1, \dots, r\} \setminus \{i\}$. Hence,

$$v(\psi(t)) = rk\delta - (r-1)k\delta - v((t-s)^k - zu_i) = k\delta - v((t-s)^k - zu_i) < 0.$$

In particular, $V_E \in B(\psi)$ by Proposition 6.6; furthermore, if $\psi(t), \frac{t-s}{d} \in V$, then $t \in \mathcal{C}$. Hence, $B(\psi) \cap B\left(\frac{X-s}{d}\right) \cap B(\phi) = \emptyset$, and thus $B(\psi) \cap B\left(\frac{X-s}{d}\right)$ is a neighborhood of V_E disjoint from $B(\phi)$. It follows that $B(\phi)$ is closed, as claimed.

(ii) \implies (iii) and (iv) are obvious.

Suppose now that either (iii) or (iv) hold for some δ , and let \mathcal{X} be $\bigcup_{\delta' \leq \delta} \mathcal{V}(\bullet, \delta')$ or $\bigcup_{\delta' < \delta} \mathcal{V}(\bullet, \delta')$, accordingly. Suppose that the residue field of V is infinite. Let $c \in K$ such that $\eta = v(c) < \delta$: we claim that $B(c^{-1}X) \cap \mathcal{X}$ is not closed in \mathcal{X} .

Indeed, let $E = \{s_n\}_{n \in \mathbb{N}}$ be a pseudo-convergent sequence with breadth η and having 0 as a pseudo-limit. Then, $V_E \notin B(c^{-1}X)$. Suppose there is a neighborhood of V_E disjoint from $B(c^{-1}X)$: then, there are ψ_1, \dots, ψ_k such that $V_E \in B(\psi_1, \dots, \psi_k)$ and such that $B(\psi_1, \dots, \psi_k) \cap B(c^{-1}X) = \emptyset$.

Fix an extension u of v to \overline{K} . Let β_1, \dots, β_m be the critical points of ψ_1, \dots, ψ_k having valuation η under u (if there are any). Since the residue field of V is infinite, there is a $t \in K$ such that $v(t) = \eta$ and such that $u(t - \beta_i) = \eta$ for all i . We claim that $v(\psi_i(t)) \geq 0$ for all i .

Indeed, fix i , and let $\alpha_1, \dots, \alpha_r$ be the critical points of $\psi = \psi_i$. By construction, we have

$$u(t - \alpha_j) = \begin{cases} u(\alpha_j) & \text{if } u(\alpha_j) < \eta, \\ v(t) = \eta & \text{if } u(\alpha_j) \geq \eta. \end{cases} \quad (9)$$

In particular, a direct calculation gives $v(\psi(t)) = \lambda\eta + \gamma$, where λ is the weighted sum of the critical points of ψ in the closed ball $B(0, e^{-\eta})$ and $\gamma \in \Gamma_v$. By Corollary 4.6, it follows that $v(\psi(t)) = w_E(\psi)$; in particular, $v(\psi(t)) \geq 0$ since $\psi \in V_E$. Therefore, $v(\psi_i(t)) \geq 0$ for all i . Furthermore, we claim that

$$v(\psi_i(t')) = v(\psi_i(t)) \geq 0, \text{ for all } t' \text{ such that } v(t - t') > \eta \quad (10)$$

In fact, by (9) we have $\eta \geq u(t - \alpha_i)$, so $u(t' - \alpha_i) = u(t - \alpha_i)$ for all $i = 1, \dots, r$ and the claim follows.

Hence, if $\eta' > \eta$ and $F = \{t_n\}_{n \in \mathbb{N}}$ is a pseudo-convergent sequence of breadth η' and pseudo-limit t , then $V_F \in B(\psi_1, \dots, \psi_k)$ by (10), since $v(t - t_n) > \eta$ for large n . In particular, we must have $v(t) = v(t_n)$ for every n , since $v(t) = \eta < \eta'$ and $v(t - t_n) \nearrow \eta'$, so 0 is not a pseudo-limit of F (thus, $v(t_n)$ is eventually constant). Hence, V_F also belongs to $B(c^{-1}X)$; therefore, if we choose $\eta' \in (\eta, \delta)$, we have $V_F \in B(\psi_1, \dots, \psi_k) \cap B(c^{-1}X) \cap \mathcal{X}$, against our choice of ψ_1, \dots, ψ_k . Therefore, $B(c^{-1}X) \cap \mathcal{X}$ is not closed, and the constructible topology does not agree with the Zariski topology. By contradiction, (i) holds. \square

To conclude this section, we study the function from \mathcal{W} to \mathcal{V} which maps each W_E to V_E . We need the following lemma.

Lemma 6.12. *Let $\phi \in K(X)$ and $\delta \in \mathbb{R}$. Let S be the set of valuation domains V_F , with $F = \{t_n\}_{n \in \mathbb{N}}$, such that $v(\phi(t_n)) \nearrow \delta$. Then, S is a finite set.*

Proof. Let $V_F \in S$, $F = \{t_n\}_{n \in \mathbb{N}}$ with breadth δ_F , and fix an extension u of v to \overline{K} . By Proposition 3.6 and Corollary 4.6 there are $\lambda \in \mathbb{Z}$, $\gamma \in \Gamma_v$ depending on F and $\phi(X)$ such that

$$\delta = w_F(\phi) = \lambda\delta_F + \gamma. \quad (11)$$

Since $v(\phi(t_n))$ is eventually strictly increasing, F is of algebraic type and by Proposition 3.6 its dominating degree λ is positive, i.e., some zero of ϕ is a pseudo-limit of F with respect to u . Hence, S is the union of $S_\beta = \{V_F \in S \mid \beta \in \mathcal{L}_F^u\}$, as β ranges among the zeroes of ϕ . Since ϕ has only finitely many zeroes, it is enough to show that each S_β is finite.

Let A_β be the set of breadths of the pseudo-convergent sequences in S_β ; then, the cardinality of A_β is equal to the cardinality of S_β , by Theorem 5.4. Let $\theta_1 < \dots < \theta_a$ be the elements of Γ_v such that there is a critical point β' of ϕ with $v(\beta - \beta') = \theta_i$; let $\theta_0 = -\infty$ and $\theta_{a+1} = +\infty$. We claim that $A_\beta \cap (\theta_i, \theta_{i+1})$ has at most one element, for every $i \in \{0, \dots, a\}$.

Let $V_F \in S_\beta$ be such that $\delta_F \in (\theta_i, \theta_{i+1})$, and let $F = \{t_n\}_{n \in \mathbb{N}}$. Note that for such pseudo-convergent sequences F , the values of λ and γ in (11) do not depend on F (explicitly, λ is the weighted sum of critical points β' of ϕ such that $v(\beta - \beta') \geq \delta_F$, which is equivalent to $v(\beta - \beta') > \theta_{i+1}$, and γ is defined as in Proposition 3.6). In particular, by (11), δ_F is uniquely determined in (θ_i, θ_{i+1}) (recall that if $V_F \in S$ then the dominating degree is nonzero), and since we are dealing with pseudo-convergent sequences F having

β as pseudo-limit, by Theorem 5.4 $|A_\beta \cap (\theta_i, \theta_{i+1})| \leq 1$. Therefore,

$$A_\beta \subseteq \{\theta_1 \dots, \theta_a\} \cup \bigcup_{i=0}^a (A_\beta \cap (\theta_i, \theta_{i+1}))$$

is finite. Hence, S_β is finite and the claim is proved. \square

If V is a DVR, then we have already remarked at the beginning of Section 6 that \mathcal{W} is a subset of \mathcal{V} ; in particular, it is a topological embedding. If V is non-discrete, we still have an inclusion, which however is not an embedding.

Proposition 6.13. *Let V be a rank one non-discrete valuation domain. Let Ψ be the map*

$$\begin{aligned} \Psi: \mathcal{W} &\longrightarrow \mathcal{V} \\ W_E &\longmapsto V_E \end{aligned}$$

Then, Ψ is continuous and injective, but it is not a topological embedding.

Proof. By Theorem 5.4, Ψ is injective. To show that Ψ is continuous, it is enough to show that every $\Psi^{-1}(B^\mathcal{V}(\phi))$ is open.

Since $V_E \subseteq W_E$, we have $\Psi^{-1}(B^\mathcal{V}(\phi)) = \{W_E \in \mathcal{W} \mid V_E \ni \phi\} \subseteq B^\mathcal{W}(\phi)$, and the inclusion can be strict; more precisely,

$$C = B^\mathcal{W}(\phi) \setminus \Psi^{-1}(B^\mathcal{V}(\phi)) = \{W_E \mid \phi \in W_E \setminus V_E\} = \{W_E \mid \phi \in W_E^* \setminus V_E\}.$$

If $E = \{s_n\}_{n \in \mathbb{N}}$ is such that $\phi \in W_E^*$, then $w_E(\phi) = 0$; furthermore, if $\phi \notin V_E$ then $v(\phi(s_n))$ is eventually negative. Hence, for every $W_E \in C$ we must have $v(\phi(s_n)) \nearrow 0$, and by Lemma 6.12 the set C is finite (and possibly empty); since \mathcal{W} is T_1 (Proposition 6.3), C is closed. Hence,

$$\Psi^{-1}(B^\mathcal{V}(\phi)) = B^\mathcal{W}(\phi) \cap (\mathcal{W} \setminus C)$$

is open, and so Ψ is continuous.

Let \mathcal{V}_0 be the image of Ψ : to show that Ψ is not a topological embedding, it is enough to show that $\Phi = \Psi^{-1}: \mathcal{V}_0 \longrightarrow \mathcal{W}$ is not continuous. Take a pseudo-convergent sequence E of algebraic type with breadth $\delta \in \Gamma_v$, and let $\zeta > \delta$. By Proposition 6.9, if, for each $n \in \mathbb{N}$, E_n is a pseudo-convergent sequence with limit s_n and breadth ζ , then V_E is the limit of V_{E_n} in the Zariski topology; note that both V_E and the V_{E_n} belong to \mathcal{V}_0 since they have finite breadth.

Hence, if Φ were continuous then $\Phi(V_{E_n}) = W_{E_n}$ would have limit $\Phi(V_E) = W_E$ in \mathcal{W} ; since the Zariski and the constructible topologies agree on \mathcal{W} (by Proposition 6.3), it would follow that W_{E_n} has limit W_E in $\text{Zar}(K(X)|V)^{\text{cons}}$. However, this contradicts Proposition 6.9, since $\text{Zar}(K(X)|V)^{\text{cons}}$ is Hausdorff and $V_E \neq W_E$ by Theorem 4.9 (and the choice of δ). Hence, Φ is not continuous and Ψ is not a topological embedding. \square

6.1 Separation properties of \mathcal{V}

A topological space is *regular* if every point is closed and if, whenever C is a closed set and $x \notin C$ then x and C can be separated by open sets. The space $\text{Zar}(K(X)|V)$ is not regular under the Zariski topology, since it is not even T_1 ; on the other hand, under the constructible topology, \mathcal{V} is regular, since it is a subspace of the regular space $\text{Zar}(K(X)|V)^{\text{cons}}$. In particular, by Proposition 6.11, if the residue field of V is finite then \mathcal{V} is regular even if endowed with the Zariski topology (since in this case the two topologies coincide on \mathcal{V}). In this section, we show that the regularity of \mathcal{V} under the Zariski topology holds without any additional hypothesis.

We say that two subsets C_1, C_2 of a topological space X can be *separated by open-closed sets* (*open sets*, respectively) if there are disjoint open-closed (open, respectively) subsets Ω_1, Ω_2 of X such that $C_i \subseteq \Omega_i$. If $C_1 = \{c_1\}$ is a singleton, we also say that c_1 and C_2 can be separated by open-closed sets (open sets, respectively).

We need a preliminary lemma.

Lemma 6.14. *Let $\gamma \in \mathbb{Q}\Gamma_v$ and $s \in K$. Then, the set*

$$\Omega(s, \gamma) = \{V_E \in \mathcal{V} \mid w_E(X - s) \leq \gamma\}$$

is both open and closed in \mathcal{V} .

Proof. If V is discrete, then by Theorem 6.1 $\Omega(s, \gamma)$ is homeomorphic to the closed ball of \widehat{K} having center s and radius $e^{-\gamma}$, and thus it is both open and closed since \widehat{K} is an ultrametric space.

Suppose V is not discrete, and let $\Omega = \Omega(s, \gamma)$. Let $k > 0$ be an integer such that $k\gamma \in \Gamma_v$, and let $c \in K$ be such that $v(c) = k\gamma$. We claim that

$$\Omega = B\left(\frac{c}{(X-s)^k}\right)$$

and that

$$\mathcal{V} \setminus \Omega = \bigcup_{\substack{d \in K \\ v(d) > \gamma}} B\left(\frac{X-s}{d}\right).$$

Clearly, both right hand sides are open in \mathcal{V} .

Let $E = \{s_n\}_{n \in \mathbb{N}}$ be a pseudo-convergent sequence. Then $v(s_n - s)$ is either eventually increasing or eventually constant, and its limit is $w_E(X - s)$ (see Remark 4.7(a)); hence, $V_E \in \Omega$ if and only if $v(s_n - s) \leq \gamma$ for large n , while $V_E \notin \Omega$ if and only if $v(s_n - s) > \gamma$ for large n .

If $V_E \in \Omega$ then

$$v\left(\frac{c}{(s_n - s)^k}\right) = v(c) - kv(s_n - s) \geq k\gamma - k\gamma = 0$$

and so $V_E \in B\left(\frac{c}{(X-s)^k}\right)$. In the same way, if $V_E \in B\left(\frac{c}{(X-s)^k}\right)$ then $v(s_n - s) \leq \gamma$ and so $V_E \in \Omega$.

Similarly, if $V_E \notin \Omega$ then $v(s_n - s) \geq \gamma' > \gamma$ for some $\gamma' \in \Gamma_v$; if $v(d) = \gamma'$ then $V_E \in B\left(\frac{X-s}{d}\right)$ and so it is in the union. Conversely, if V_E is in the union then $V_E \in B\left(\frac{X-s}{d}\right)$ for some d , and $v(s_n - s) \geq v(d) > \gamma$ for large n , so that $V_E \notin \Omega$. The claim is proved. \square

Theorem 6.15. \mathcal{V} is a regular topological space.

Proof. If V is a DVR, the statement follows from the fact that $\mathcal{V} = \mathcal{V}(\bullet, \infty)$ is an ultrametric space by [19, Theorem 3.4]. Henceforth, we assume that V is not discrete.

We first note that each point of \mathcal{V} is closed: indeed, the closure of a point Z in $\text{Zar}(K(X)|V)$ is equal to the set of valuation domains contained in Z . However, two different domains V_E and V_F are never comparable: if they were, then $W_E = W_F$, and thus $V_E = V_F$ by Theorem 5.4.

Let $E = \{s_n\}_{n \in \mathbb{N}} \subset K$ be a pseudo-convergent sequence of breadth δ , let $\{\delta_n\}_{n \in \mathbb{N}}$ be the gauge of E and let $C \subset \mathcal{V}$ be a closed set which does not contain V_E . Then there are rational functions $\phi_1, \dots, \phi_k \in K(X)$ such that $V_E \in B(\phi_1, \dots, \phi_k)$ while $B(\phi_1, \dots, \phi_k) \cap C = \emptyset$. We let $\Lambda = \{\beta_1, \dots, \beta_m\} \subseteq \overline{K}$ be the set of critical points of ϕ_1, \dots, ϕ_k . Let also u be an extension of v to \overline{K} .

We want to separate V_E and C ; we need to distinguish several cases.

Case 1. E is of transcendental type.

By [25, Theorem 31.18, p. 328], there is an n such that no $\beta \in \Lambda$ satisfies $u(\beta - s_n) \geq \delta_n$. Hence, there is a $\gamma < \delta_n$, $\gamma \in \mathbb{Q}\Gamma_v$ such that each $\beta \in \Lambda$ satisfies $u(\beta - s_n) < \gamma$. Moreover, up to considering a bigger $n \in \mathbb{N}$, we may also suppose that $\phi_i(s_n) \in V$ for all $i = 1, \dots, k$. Let $s = s_n$. By Theorem 3.3, we have $v(\phi_i(t)) = v(\phi_i(s)) \geq 0$ for all t such that $v(t - s) \geq \gamma$ and for all $i = 1, \dots, k$.

We claim that $\Omega(s, \gamma)$ and its complement separate C and V_E ; by Lemma 6.14, this will imply that C and V_E are separated by open-closed sets.

Indeed, clearly $w_E(X - s) = \delta_n > \gamma$ and so $V_E \notin \Omega(s, \gamma)$. On the other hand, if $V_F \in C$ and $F = \{t_n\}_{n \in \mathbb{N}}$, then there is an i such that $v(\phi_i(t_n))$ is eventually negative. By the previous paragraph $v(t_n - s) < \gamma$ for all sufficiently large n ; hence, $w_F(X - s) = \lim_n v(t_n - s) \leq \gamma$ and $V_F \in \Omega(s, \gamma)$. Thus, $C \subseteq \Omega(s, \gamma)$, as claimed.

Case 2. E is of algebraic type without pseudo-limits in K .

Let $\alpha \in \overline{K} \setminus K$ be a pseudo-limit of E with respect to u . By Lemma 2.4, there is no element t of K such that $u(\alpha - t) \geq \delta$. By Proposition 6.6, there is an annulus $\mathcal{C} = \mathcal{C}_u(\alpha, \tau, \delta)$ such that $\phi_i(t) \in V$ for all $i = 1, \dots, k$ and all $t \in \mathcal{C}$; let $s \in \mathcal{C}$ and let $\delta' = u(\alpha - s) \in \mathbb{Q}\Gamma_v$. Note that $\tau < \delta' < \delta$. We claim that $\Omega(s, \tau)$ and its complement separate C and V_E .

Indeed, we have

$$w_E(X - s) = \lim_{n \rightarrow \infty} v(s_n - s) = \lim_{n \rightarrow \infty} u(s_n - \alpha + \alpha - s) = u(\alpha - s) = \delta'$$

since $\delta_n > \delta'$ for large n ; hence, $V_E \notin \Omega(s, \tau)$. On the other hand, if $F = \{t_n\}_{n \in \mathbb{N}}$ is a pseudo-convergent sequence such that $V_F \in C \setminus \Omega(s, \tau)$, then $\tau < v(t_n - s)$ for all $n \in \mathbb{N}$

sufficiently large. Therefore, for each such n we have $\delta > u(t_n - \alpha) = u(t_n - s + s - \alpha) > \tau$. By our assumption this would imply that $\phi_i(t_n) \in V$ for $i = 1, \dots, k$, for all $n \geq N$, which is a contradiction since $C \cap B(\phi_1, \dots, \phi_k) = \emptyset$. Hence, $C \subseteq \Omega(s, \tau)$, and we have proved that C and V_E can be separated by an open-closed set.

Case 3. E is of algebraic type and there exists a pseudo-limit α of E in K .

We partition C into the following three sets:

$$\begin{aligned} C_1 &= \{V_F \in C \mid w_F(X - \alpha) < \delta\}, \\ C_2 &= \{V_F \in C \mid w_F(X - \alpha) > \delta\}, \\ C_3 &= \{V_F \in C \mid w_F(X - \alpha) = \delta\}. \end{aligned}$$

By Theorem 3.3 (and Remark 3.4), we can find $\zeta_1, \zeta_2 \in \mathbb{Q}\Gamma_v$ such that $\zeta_1 < \delta \leq \zeta_2$ and such that $v(\phi_i(t)) = \lambda_i v(t - \alpha) + \gamma_i$ for every $t \in \mathcal{C}(\alpha, \zeta_1, \zeta_2)$, for some $\lambda_i \in \mathbb{Z}$ and $\gamma_i \in \Gamma_v$. Since $V_E \in B(\phi_1, \dots, \phi_k)$, by Proposition 6.6 we can find $\theta_1, \theta_2 \in \mathbb{Q}\Gamma_v$, with $\delta \in (\theta_1, \theta_2] \subseteq (\zeta_1, \zeta_2]$, such that $\phi_i(t) \in V$ for all $t \in \mathcal{C}(\alpha, \theta_1, \theta_2)$ and all $i = 1, \dots, k$.

Consider $\Omega(\alpha, \theta_1)$. We have $w_E(X - \alpha) = \delta > \theta_1$, and so $V_E \notin \Omega(\alpha, \theta_1)$; on the other hand, if $V_F \in C_1$, with $F = \{t_n\}_{n \in \mathbb{N}}$, then $v(t_n - \alpha) < \theta_1$ for all large n (because $C_1 \subseteq C$ has empty intersection with $B(\phi_1, \dots, \phi_k)$) and thus also $w_F(X - \alpha) \leq \theta_1$; hence, $C_1 \subseteq \Omega(\alpha, \theta_1)$. Thus, $\Omega(\alpha, \theta_1)$ and its complement are open-closed subsets separating C_1 and V_E .

Similarly, $w_E(X - \alpha) = \delta \leq \theta_2$ and thus $V_E \in \Omega(\alpha, \theta_2)$; if $V_F \in C_2$, $F = \{t_n\}_{n \in \mathbb{N}}$, then $v(t_n - \alpha) > \theta_2$ for all large n (because $C_2 \subseteq C$ has empty intersection with $B(\phi_1, \dots, \phi_k)$) and, since $v(t_n - \alpha)$ is either eventually strictly increasing or eventually constant, we have $w_F(X - \alpha) > \theta_2$, i.e., $C_2 \cap \Omega(\alpha, \theta_2) = \emptyset$. Hence, $\Omega(\alpha, \theta_2)$ and its complement separate V_E and C_2 . In particular, if $C_3 = \emptyset$ then V_E and C can be separated by open-closed sets.

Suppose $C_3 \neq \emptyset$ and let $V_F \in C_3$, $F = \{t_n\}_{n \in \mathbb{N}}$: then $\delta \in \mathbb{Q}\Gamma_v$, for otherwise $v(t_n - \alpha)$ should increase to δ , and so t_n would enter in any annulus $\mathcal{C}(\alpha, \tau, \delta)$ and by Proposition 3.6 $\phi_i \in V_F$ for $i = 1, \dots, k$, against the fact that $C \cap B(\phi_1, \dots, \phi_k) = \emptyset$. By the same argument, $v(t_n - \alpha)$ is constantly equal to δ (which therefore is in Γ_v). In particular, α is not a pseudo-limit of F so that $\delta_F > v(t_n - \alpha) = \delta = \delta_E$.

Since $C \cap B(\phi_1, \dots, \phi_k) = \emptyset$, for every $V_F \in C$ there is an $i \in \{1, \dots, k\}$ such that $\phi_i(t_n) \notin V$ for all n sufficiently large; for such an i , $w_F(\phi_i) \leq 0$ and if equality holds then $v(\phi_i(t_n)) \nearrow 0$, where $F = \{t_n\}_{n \in \mathbb{N}}$. For each $i = 1, \dots, k$, let

$$\begin{aligned} D_i &= \{V_F \in C_3 \mid w_F(\phi_i) < 0\} \quad \text{and} \\ H_i &= \{V_F \in C_3 \mid w_F(\phi_i) = 0, \phi_i \notin V_F\}, \end{aligned}$$

so that $C_3 = \bigcup_{i=1, \dots, k} (D_i \cup H_i)$.

We claim that every D_i can be separated from V_E by open sets: indeed, let

$$\Omega_i = \bigcup_{\substack{d \in K \\ v(d) < 0}} B\left(\frac{d}{\phi_i(X)}\right).$$

As in the proof of Lemma 6.14, if $V_F \in D_i$ then there is a $\kappa < 0$ such that $v(\phi_i(t_n)) \leq \tau$ for all large n and thus, taking $d \in K$ such that $0 > v(d) \geq \kappa$,

$$v\left(\frac{d}{\phi_i(t_n)}\right) \geq \kappa - v(\phi_i(t_n)) \geq 0$$

and so $V_F \in \Omega_i$. Moreover, $\Omega_i \cap B(\phi_1, \dots, \phi_k) = \emptyset$, since otherwise there should be a $t \in K$ such that

$$\begin{cases} v(\phi_i(t)) \geq 0 \\ v\left(\frac{d}{\phi_i(t)}\right) \geq 0; \end{cases}$$

for some $d \in K$ such that $v(d) < 0$, but the latter condition implies that $v(\phi_i(t)) \leq v(d) < 0$. Hence, $B(\phi_1, \dots, \phi_k)$ and Ω_i separate V_E and D_i .

Since for every $V_F \in H_i$, with $F = \{t_n\}_{n \in \mathbb{N}}$, we have $v(\phi_i(t_n)) \nearrow 0$, every H_i is finite by Lemma 6.12. Furthermore, some zero $\beta \in \overline{K}$ of ϕ_i is a pseudo-limit of F , with respect to some extension u of v to \overline{K} (see the proof of Lemma 6.12). If n is sufficiently large, then $\delta_E < u(t_n - \beta) < \delta_F$. Let $\gamma \in \Gamma_v$ be such that $\delta_E < \gamma < u(t_n - \beta)$. If we let $t = t_n$, then $w_E(X - t) \leq \delta_E < \gamma < w_F(X - t)$ (see Remark 4.7(a)). Then $V_E \in \Omega(t, \gamma)$ and $V_F \notin \Omega(t, \gamma)$. Hence, V_F can be separated from V_E by the open-closed set $\Omega(t, \gamma)$, and since H_i is finite it can also be separated from V_E by open-closed sets.

To summarize, we have

$$C = C_1 \cup C_2 \cup \bigcup_{i=1}^k D_i \cup \bigcup_{i=1}^k H_i,$$

and each of the sets on the right hand side can be separated from V_E by open sets; since the union is finite, C and V_E can be separated and therefore \mathcal{V} is regular. \square

As a consequence, we can show that under some conditions \mathcal{V} is metrizable.

Corollary 6.16. *Let V be a countable valuation domain. Then, \mathcal{V} is metrizable.*

Proof. A basis for \mathcal{V} is $\mathcal{B} = \{B(\phi_1, \dots, \phi_k) \mid \phi_1, \dots, \phi_k \in K(X)\}$. Since V is countable, so are K and $K(X)$; hence, the number of finite subsets of $K(X)$ is countable, and thus also \mathcal{B} is countable. Therefore, \mathcal{V} is second-countable; since it is regular (Theorem 6.15), it follows from Urysohn's metrization theorem [9, e-2] that \mathcal{V} is metrizable. \square

References

- [1] V. Alexandru, N. Popescu, *Sur une classe de prolongements à $K(X)$ d'une valuation sur un corps K* , Rev. Roumaine Math. Pures Appl. 33 (1988), no. 5, 393-400.
- [2] V. Alexandru, N. Popescu, Al. Zaharescu, *A theorem of characterization of residual transcendental extensions of a valuation*. J. Math. Kyoto Univ. 28 (1988), no. 4, 579-592.

- [3] V. Alexandru, N. Popescu, Al. Zaharescu, *Minimal pairs of definition of a residual transcendental extension of a valuation*, J. Math. Kyoto Univ. 30 (1990), no. 2, 207-225.
- [4] V. Alexandru, N. Popescu, Al. Zaharescu, *All valuations of $K(X)$* , J. Math. Kyoto Univ. 30 (1990), no. 2, 281-296.
- [5] N. Bourbaki, *Algèbre commutative*, Hermann, Paris, 1961.
- [6] J.-P. Cahen, J.-L. Chabert, *Integer-valued polynomials*, Amer. Math. Soc. Surveys and Monographs, 48, Providence, 1997.
- [7] P.-J. Cahen, A. Loper, *Rings of integer-valued rational functions*. J. Pure Appl. Algebra 131 (1998), no. 2, 179-193.
- [8] J.-L. Chabert, *On the polynomial closure in a valued field*, J. Number Theory 130 (2010), 458-468.
- [9] K. P. Hart, J. Nagata, and J. E. Vaughan, editors. *Encyclopedia of general topology*. Elsevier Science Publishers, B.V., Amsterdam, 2004.
- [10] M. Hochster, *Prime ideal structure in commutative rings*, Trans. Amer. Math. Soc., 142 (1969), 43-60.
- [11] I. Kaplansky, *Maximal Fields with Valuation*, Duke Math. J. 9 (1942), 303-321.
- [12] C. Karpfinger and H. Wähling, *Äquivalenz und Isomorphie bei Bewertungsfortsetzungen auf einfache-transzendente Erweiterungen*, Math. Z. 238, 461-481 (2001).
- [13] F.-V. Kuhlmann, *Value groups, residue fields, and bad places of rational function fields*. Trans. Amer. Math. Soc. 356 (2004), no. 11, 4559-4600.
- [14] A. Loper, N. Werner, *Pseudo-convergent sequences and Prufer domains of integer-valued polynomials*, J. Commut. Algebra 8 (2016), no. 3, 411-429.
- [15] S. MacLane, *A construction for absolute values in polynomial rings*, Trans. Amer. Math. Soc. 40 (1936), no. 3, 363-395.
- [16] B. Olberding, *A principal ideal theorem for compact sets of rank one valuation rings*, J. Algebra 489 (2017), 399-426.
- [17] A. Ostrowski, *Untersuchungen zur arithmetischen Theorie der Körper*, Math. Z. 39 (1935), 269-404.
- [18] G. Peruginelli, *Prüfer intersection of valuation domains of a field of rational functions*, J. Algebra 509 (2018), 240-262.
- [19] G. Peruginelli, *Transcendental extensions of a valuation domain of rank one*, Proc. Amer. Math. Soc. 145 (2017), no. 10, 4211-4226.

- [20] G. Peruginelli, D. Spirito, *Extending valuations to the field of rational functions using pseudo-monotone sequences*, preprint, arXiv: <https://arxiv.org/abs/1905.02481>.
- [21] P. Ribenboim, *Théorie des valuations*. Séminaire de Mathématiques Supérieures, No. 9 (Été, 1964) Les Presses de l'Université de Montréal, Québec 1968.
- [22] P. Ribenboim, *On the completion of a valuation ring*. Math. Ann. 155 (1964), 392–396.
- [23] L. A. Steen and J. A. Seebach, *Counterexamples in Topology*. Springer-Verlag, New York-Heidelberg, second edition, 1978.
- [24] M. Vaquié, Michel, *Extension d'une valuation*, Trans. Amer. Math. Soc. 359 (2007), no. 7, 3439–3481.
- [25] S. Warner, *Topological Fields*. North-Holland Mathematics Studies, 157.
- [26] S. Willard, *General topology*. Addison-Wesley Publishing Co., Reading, Mass.-London-Don Mills, Ont. 1970.
- [27] O. Zariski, P. Samuel, *Commutative Algebra, vol. II*, Springer-Verlag, New York-Heidelberg-Berlin, 1975.