

Glucose Pulse

A simple method to estimate the amount of glucose oxidized during exercise in type 1 diabetic patients

MARIA PIA FRANCESCATO, MD^{1,2}
 LUIGI CATTIN, MD³
 MARIO GEAT, MD³
 ELENA TOSORATTI, MD^{1,2}

STEFANO LAZZER, PHD^{1,2}
 CLAUDIO NOACCO, MD⁴
 PIETRO ENRICO DI PRAMPERO, MD^{1,2}

In type 1 diabetic patients, exercise contributes to enhance insulin sensitivity (1,2) and may help, together with diet and insulin therapy, to achieve and maintain better metabolic control (3,4). However, lack of insulin regulation during exercise leads to frequent metabolic imbalances (5). Fat and carbohydrates are the main substrates for energy production in skeletal muscle during aerobic exercise in well-fed humans (6), with their relative contribution to total energy production being a function of exercise intensity (7–10). Below the anaerobic threshold, both oxygen consumption and heart rate during exercise increase linearly as a function of exercise intensity (6). On the basis of these relationships, the aim of the present study was to verify the possibility of using heart rate to estimate the amount of glucose oxidized during exercise in type 1 diabetic patients as well as in a control group of healthy subjects.

RESEARCH DESIGN AND METHODS

Fifteen type 1 diabetic patients (9 men, 6 women, HbA_{1c} [A1C] $6 \pm 1.2\%$, reference values 4–6%) (11) aged 18–45 years (average 38 ± 6) and 15 healthy subjects (9 men, 6 women, 34 ± 7 years) gave their voluntary consent to participate after being informed on the nature, purpose, and possible risks.

Duration of diabetes and mean insulin dose were 17 ± 8 years and 0.52 ± 0.1 IU \cdot kg⁻¹ \cdot day⁻¹, respectively. Average body weight and height were 71 ± 9 kg and 175 ± 9 cm in diabetic patients vs. 70 ± 9 kg and 175 ± 8 cm in healthy volunteers (BMI 23 ± 2 vs. 23 ± 2 kg/m²). The study was approved by the local ethical committee and was conducted according to the Declaration of Helsinki.

Inclusion criteria were no clinical evidence of chronic complications (albumin excretion rate, eye examination, deep breathing heart rate variability, Valsalva ratio, tilt test ratio), no medication other than insulin, and self-management to maintain a good metabolic balance.

Patients were asked to maintain their usual diet and insulin dose and to control their blood glucose levels according to the self-management procedures. According to their self-reported physical activity habits, volunteers were subdivided into groups as “aerobically trained” (eight patients, seven healthy), who exercised regularly at least 30 min for 3 days/week, and “sedentary” (seven patients, eight healthy), who exercised occasionally (12). The experimental sessions were performed 3 ± 1 h postprandial.

Thirty minutes before the start, an extra amount of carbohydrates (mainly sugar, sugar drops, or crackers) was given

to the diabetic patients (13). Other carbohydrates (5 g each time) were given during the exercise to prevent falling of glycemia below 6 mmol/l. On average, a total amount of 19.6 ± 11.6 g carbohydrates were given before and/or during the trial. No carbohydrates were given to healthy subjects.

The experiment consisted of four exercises performed on a cycloergometer (Ergomed 839E; Monark, Vansbro, Sweden) lasting 10 min each (total duration of 40 min) of increasing intensity corresponding to 0.7, 1.0, 1.5, and 2.0 W/kg body mass and 0.7, 0.95, 1.25, and 1.75 W/kg body mass in men and women, respectively.

Oxygen consumption (V_{O₂}), carbon dioxide production (V_{CO₂}), and heart rate were measured every 15 s using a metabolic unit (K4; Cosmed, Rome, Italy) at rest and during the last 5 min of each exercise intensity (14) and averaged over the last 2 min. Glycemia was tested before and every 5 min during exercise with reactive strips (Glucotrend; Roche Diagnostics, Basel, Switzerland). Three minutes after the end of the trial, a capillary blood sample was taken to determine lactate concentration using reactive strips (Accusport; Boehringer, Mannheim, Germany).

Respiratory quotient, assumed to be equal to the respiratory exchange ratio (RER), was calculated as the ratio between V_{O₂} and V_{CO₂} (12). Assuming that net protein utilization did not contribute to any significant extent to energy production during exercise (6), glucose oxidation rates (GLU_{ox}) were calculated applying common equations (15). The following regressions were calculated for each subject: heart rate vs. V_{O₂}, V_{O₂} vs. RER, RER vs. GLU_{ox}, and heart rate vs. GLU_{ox}. For the last two regressions, the values corresponding to the highest exercise intensity were discarded when the postexercise blood lactate concentration exceeded 4 mmol/l, which is considered as anaerobic threshold. All of the glucose pulse, i.e., the ratios between GLU_{ox} expressed per unit body weight and heart

From the ¹Department of Biomedical Sciences and Technologies, University of Udine, Udine, Italy; the ²M.A.T.I. Centre of Excellence, University of Udine, Udine, Italy; the ³Department of Clinical, Morphological and Technological Sciences, University of Trieste, Trieste, Italy; and the ⁴Diabetologic Unit, Hospital S. Maria della Misericordia, Udine, Italy.

Address correspondence and reprint requests to Maria Pia Francescato, Department of Biomedical Sciences and Technologies, University of Udine, P.le Kolbe 4, 33100 - UDINE, ITALY. E-mail: mfrancescato@mail.dstb.uniud.it.

Received for publication 10 January 2005 and accepted in revised form 3 May 2005.

Abbreviations: RER, respiratory exchange ratio.

A table elsewhere in this issue shows conventional and Système International (SI) units and conversion factors for many substances.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked “advertisement” in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.




Figure 1—Glucose pulse ($\text{mg} \cdot \text{kg}^{-1} \cdot \text{beat}^{-1}$) as a function of the percentage of maximal heart rate (average of eight measurements during the last 2 min of exercise). The aerobically trained type 1 diabetic patients (●) show lower values compared with the sedentary patients (○). Regression equations are as follows: glucose pulse = $(0.504 \times \% \text{HR}_{\text{max}} - 22.11)/\text{actual heart rate}$; and glucose pulse = $(0.679 \times \% \text{HR}_{\text{max}} - 22.05)/\text{actual heart rate}$ for a 36-year-old trained or sedentary patient, respectively (see text for details).

rate expressed as percentage of maximal theoretical heart rate ($\text{HR}_{\text{max}} = 220 - \text{age}$) (12) were calculated.

ANCOVA was applied to investigate differences between groups. A value of $P < 0.05$ was assumed as statistically significant.

RESULTS— The heart rate vs. VO_2 , VO_2 vs. RER, RER vs. GLU_{ox} , and heart rate vs. GLU_{ox} relationships were statistically significant ($R > 0.90$; $P < 0.05$) for all of the volunteers, type 1 diabetic or healthy subjects.

No significant differences for GLU_{ox} were found between males and females ($F = 0.07$, NS) or between diabetic and healthy subjects ($F = 0.06$, NS), whereas self-reported training habits had a significant effect ($F = 9.3$, $P < 0.005$), with GLU_{ox} higher in sedentary subjects.

The glucose pulse of the sedentary and trained diabetic patients is plotted in Fig. 1 as a function of the percentage of HR_{max} . The two lines represent the glucose pulse that can be estimated from the relationship between GLU_{ox} and percentage HR_{max} in aerobically trained or sedentary patients.

CONCLUSIONS— This study shows that heart rate can be a useful physiological parameter to be used to estimate the amount of glucose oxidized during exercise. The correlation coefficients between

heart rate and GLU_{ox} were greater than 0.95 in all volunteers, both healthy and type 1 diabetic subjects, indicating a highly significant correlation between the two variables. This relationship is the basis of the “glucose pulse” concept, defined as the GLU_{ox} expressed per unit of heart rate. Endurance training leads to a lower GLU_{ox} (16,17), which was also observed in the present study with a lower glucose pulse in aerobically trained patients. The relationship between the glucose pulse and the percentage of maximal theoretical heart rate is not linear. Nevertheless, if only a heart rate below 70% of maximal is considered, the following linear equations apply:

$$\text{trained} = 0.00353 \pm 0.0023 \times \% \text{HR}_{\text{max}} - 0.1428 \pm 0.1328 \quad (n = 23, R = 0.846)$$

$$\text{and} \\ \text{sedentary} = 0.00597 \pm 0.0035 \times \% \text{HR}_{\text{max}} - 0.1801 \pm 0.1891 \quad (n = 24, R = 0.869).$$

In nonobese type 1 diabetic individuals without autonomic neuropathy, the current equations can be used to calculate glucose oxidation during exercise based on easily measurable heart rate. From these concepts, it is possible to use simple tables matching heart rate, glucose oxidation, and dietary equivalents to be ingested, thus providing a potential clinical tool for prevention of exercise-related hypoglycemia.

References

1. Stallknecht B, Larsen JJ, Mikines KJ, Simonsen L, Bulow J, Galbo H: Effect of training on insulin sensitivity of glucose uptake and lipolysis in human adipose tissue. *Am J Physiol* 279:E376–E385, 2000
2. Landt K, Campaigne B, James F, Sperling M: Effect of exercise training on insulin sensitivity in adolescents with type I diabetes. *Diabetes Care* 8:461–465, 1985
3. American Diabetes Association: Diabetes mellitus and exercise. *Diabetes Care* 20: 1908–1912, 1997
4. Rowland T, Swabda LA, Biggs DE, Burke EJ, Reiter EO: Glycemic control with physical training in insulin-dependent diabetes mellitus. *Sports Med* 139:307, 1985
5. MacDonald MJ: Postexercise late-onset hypoglycemia in insulin-dependent diabetic patients. *Diabetes Care* 10:584–588, 1987
6. Astrand PO, Rodahl K, Dahl HA, Stromme SB: *Textbook of Work Physiology: Physiological Bases of Exercise*. McGraw Hill Series in Health Education, Physical Education and Recreation. 4th ed. Windsor, Canada, Human Kinetics, 2003
7. Bergman BC, Brooks GA: Respiratory gas-exchange ratios during graded exercise in fed and fasted trained and untrained men. *J Appl Physiol* 86:479–487, 1999
8. Pirnay F, Crielaard JM, Pallikarakis N, Lacroix M, Mosora F, Krzentowski G, Luyckx AS, Lefebvre PJ: Fate of exogenous glucose during exercise of different intensities in humans. *J Appl Physiol* 53:1620–1624, 1982
9. Romijn JA, Coyle EF, Sidossis LS, Gastaldelli A, Horowitz JF, Ender E, Wolfe RR: Regulation of endogenous fat and carbohydrate metabolism in relation to exercise intensity and duration. *Am J Physiol* 265: E380–E391, 1993
10. Romijn JA, Coyle EF, Sidossis LS, Rosenblatt J, Wolfe RR: Substrate metabolism during different exercise intensities in endurance-trained women. *J Appl Physiol* 88:1707–1714, 2000
11. Sacks DB, Bruns DE, Goldstein DE, MacLaren NK, McDonald JM, Parrott M: Guidelines and recommendations for laboratory analysis in the diagnosis and management of diabetes mellitus. *Clin Chem* 48:436–472, 2002
12. McArdle WD, Katch FI, Katch VL: *Essentials of Exercise Physiology*. 2nd ed. Philadelphia, Lippincott Williams & Wilkins, 2000
13. Francescato MP, Geat M, Fusi S, Stupar G, Noacco C, Cattin L: Carbohydrate requirement and insulin concentration during moderate exercise in type 1 diabetic patients. *Metabolism* 53:1126–1130, 2004
14. Hausswirth C, Bigard A, Le Chevalier J: The Cosmed K4 telemetry system as an

- accurate device for oxygen uptake measurements during exercise. *Int J Sports Med* 18:449–453, 1997
15. Frayn KN: Calculation of substrate oxidation rates in vivo from gaseous exchange. *J Appl Physiol* 55:628–634, 1983
16. van Loon LJC, Greenhaff PL, Constantin-Teodosiu D, Saris WHM, Wagenmakers AJM: The effects of increasing exercise intensity on muscle fuel utilisation in humans. *J Physiol (Lond)* 536:295–304, 2001
17. Brooks GA, Mercier J: Balance of carbohydrate and lipid utilization during exercise: the “crossover” concept. *J Appl Physiol* 76:2253–2261, 1994