



UNIVERSITÀ
DEGLI STUDI
DI UDINE

Università degli studi di Udine

A global bifurcation result for a second order singular equation

Original

Availability:

This version is available <http://hdl.handle.net/11390/1073656> since 2016-01-29T18:14:55Z

Publisher:

Published

DOI:

Terms of use:

The institutional repository of the University of Udine (<http://air.uniud.it>) is provided by ARIC services. The aim is to enable open access to all the world.

Publisher copyright

(Article begins on next page)

A global bifurcation result for a second order singular equation¹

ANNA CAPIETTO, WALTER DAMBROSIO
AND DUCCIO PAPINI

*Dedicated, with gratefulness and friendship, to Professor Fabio Zanolin
on the occasion of his 60th birthday*

ABSTRACT. *We deal with a boundary value problem associated to a second order singular equation in the open interval $(0, 1]$. We first study the eigenvalue problem in the linear case and discuss the nodal properties of the eigenfunctions. We then give a global bifurcation result for nonlinear problems.*

Keywords: self-adjoint singular operator, spectrum, nodal properties, global bifurcation
MS Classification 2010: 34C23, 34B09, 35P05

1. Introduction

We are concerned with a second order ODE of the form

$$-u'' + q(x)u = \lambda u + g(x, u)u, \quad \lambda \in \mathbb{R}, \quad x \in (0, 1], \quad (1)$$

where $q \in C((0, 1])$ satisfies

$$\lim_{x \rightarrow 0^+} \frac{q(x)}{l/x^\alpha} = 1, \quad (2)$$

for some $l > 0$ and $\alpha \in (0, 5/4)$, and $g \in C([0, 1] \times \mathbb{R})$ is such that

$$\lim_{u \rightarrow 0} g(x, u) = 0, \quad \text{uniformly in } x \in (0, 1]. \quad (3)$$

The constant $5/4$ arises in a rather straightforward manner in the study of the differential operator in the left-hand side of (1) (cf. [17, p. 287-288]); details are given in Remark 2.3 below.

¹Under the auspices of GNAMPA-I.N.d.A.M., Italy. The work has been performed in the frame of the M.I.U.R. Projects ‘Topological and Variational Methods in the Study of Nonlinear Phenomena’ and ‘Nonlinear Control: Geometrical Methods and Applications’.

We will look for solutions u of (1) such that $u \in H_0^2(0, 1)$.

When the x -variable belongs to a compact interval, problems of the form (1) have been very widely studied. A more limited number of contributions is available in the literature when the x -variable belongs to a (semi)-open interval, as it is the case in the present paper, or to an unbounded interval [7, 8].

We treat (1) in the framework of bifurcation theory. For this reason, we first discuss in Section 2 the eigenvalue problem

$$-u'' + q(x)u = \lambda u, \quad x \in (0, 1], \quad \lambda \in \mathbb{R}. \quad (4)$$

For such singular problems, the well-known embedding of (4) (by an elementary application of the integration by parts rule, together with the boundary condition $u(0) = 0 = u(1)$) in the setting of eigenvalue problems for compact self-adjoint operators cannot be performed. Thus, the questions of the existence of eigenvalues and of the nodal properties of the associated eigenfunctions have various delicate features. For a comprehensive account on the spectral properties of the Schrödinger operator we refer to the books [12] and [10]; for more specific results on singular problems in $(0, 1)$ we refer, among many others, to [5, 14].

However, the linear spectral theory for singular problems is well-established and can be found, among others, in the classical book by Coddington and Levinson [4] and in the (relatively) more recent text by Weidmann [17]. The former monograph focuses on a generalization of the so-called “expansion theorem” valid for functions in $L^2([0, 1])$ and, by doing this, a sort of “generalized shooting method” is performed. On the other hand, in [17] the singular problem is tackled from an abstract point of view; more precisely, it is considered the general question of the existence of a self-adjoint realization of the formal differential expression $\tau u = -u'' + q(x)u$ and the important Weyl alternative theorem [17, Theorem 5.6] is used. It is interesting to observe that the approach in [4] (based on more elementary ODE techniques) and the abstract one in [17] lead in different ways to the important concepts of “limit point case” and “limit circle case”. The knowledge of one (or the other) case is ensured by suitable assumptions on q and leads to information on the boundary conditions to be added to (4) in order to have a self-adjoint realization of τ .

In the setting of the present paper, the operator τ is regular at $x = 1$; this implies that it is in the limit circle case. Moreover, under assumption (2), from [17, Theorem 6.4] it follows that τ is in the limit circle case also in $x = 0$. Thus, the differential operator $A : u \mapsto \tau u$ with

$$D(A) = \{u \in L^2(0, 1) : u, u' \in AC(0, 1), \tau u \in L^2(0, 1), \lim_{x \rightarrow 0^+} (xu'(x) - u(x)) = 0 = u(1)\}$$

is a self-adjoint realization of τ ([17, p. 287-288]). We prove in Proposition 2.2 that in fact $D(A) = H_0^2(0, 1)$; to do this, we need some knowledge of the behaviour of the solutions of (4) near zero. These estimates are developed in Proposition 2.1 by means of the classical Levinson theorem [6, Theorem 1.8.1]. Finally, at the end of Section 2 we focus on the nodal properties of a solution to (4); more precisely, in Proposition 2.4 we prove that (4) is non-oscillatory and conclude in Proposition 2.5 that the spectrum of A is purely discrete and that, for every $n \in \mathbb{N}$, the eigenfunction associated to the eigenvalue λ_n has $(n - 1)$ simple zeros in $(0, 1)$.

Section 3 contains a global bifurcation result (Theorem 3.2) which follows in a rather straightforward manner as an application of the celebrated Rabinowitz theorem in [11].

In order to exclude alternative (2) in Theorem 3.2, we use a technique that we already applied for Hamiltonian systems in \mathbb{R}^{2N} in [2] and for planar Dirac-type systems in [3]. More precisely, we introduce a continuous integer-valued functional defined on the set of solutions to (1). Due to the singularity at $x = 0$, some care is necessary in order to prove its continuity; this is the content of Proposition 3.4. We can then state and prove our main result (Theorem 3.5).

In what follows, for a given function p we write $p(x) \sim \frac{m}{x^a}, x \rightarrow 0^+$, when

$$\lim_{x \rightarrow 0^+} \frac{p(x)}{m/x^a} = 1 \quad (5)$$

for some $m, a \in \mathbb{R}^+$.

Finally, we write

$$H_0^2(0, 1) = \{u \in H^2(0, 1) : u(0) = 0 = u(1)\},$$

equipped with the norm defined by

$$\|u\|^2 = \|u\|_{L^2(0,1)}^2 + \|u''\|_{L^2(0,1)}^2, \quad \forall u \in H_0^2(0, 1).$$

2. The linear equation

In this section we study a linear second order equation of the form

$$-u'' + q(x)u = \lambda u, \quad x \in (0, 1], \quad \lambda \in \mathbb{R}. \quad (6)$$

We will assume that $q \in C((0, 1])$ and that

$$q(x) \sim \frac{l}{x^\alpha}, \quad x \rightarrow 0^+, \quad (7)$$

for some $l > 0$ and $\alpha \in (0, 5/4)$. Without loss of generality we may suppose that

$$q(x) > 0, \quad \forall x \in (0, 1]. \quad (8)$$

For every $u : (0, 1] \rightarrow \mathbb{R}$ we denote by τu the formal expression

$$\tau u = -u'' + q(x)u;$$

First of all, we study the asymptotic behaviour of solutions of (6) when $x \rightarrow 0^+$; to this aim, let us introduce the change of variables $t = -\log x$ and let

$$w(t) = u(e^{-t}), \quad \forall t > 0.$$

From the relations

$$\begin{aligned} w'(t) &= -e^{-t}u'(e^{-t}) \\ w''(t) &= e^{-t}u'(e^{-t}) + e^{-2t}u''(e^{-t}), \end{aligned} \quad (9)$$

we deduce that u is a solution of (6) on $(0, 1)$ if and only if w is a solution of

$$-w'' - w' + e^{-2t}q(e^{-t})w = \lambda e^{-2t}w \quad (10)$$

on $(0, +\infty)$. Equation (10) can be written in the form

$$Y' = (C + R(t))Y, \quad (11)$$

where $Y = (w, z)^T$ and

$$C = \begin{pmatrix} 0 & 1 \\ 0 & -1 \end{pmatrix}, \quad R(t) = \begin{pmatrix} 0 & 0 \\ e^{-2t}q(e^{-t}) - \lambda e^{-2t} & 0 \end{pmatrix}, \quad \forall t > 0. \quad (12)$$

Now, let us observe that C has eigenvalues $\lambda_1 = 0$, $\lambda_2 = -1$ and corresponding eigenvectors $u_1 = (1, 0)$, $u_2 = (1, -1)$ and that $R \in L^1(0, +\infty)$; therefore, an application of [6, Theorem 1.8.1] implies that (11) has two linearly independent solutions Y_1, Y_2 such that

$$\begin{aligned} Y_1(t) &= u_1 + o(1), \quad t \rightarrow +\infty, \\ Y_2(t) &= (u_2 + o(1))e^{-t}, \quad t \rightarrow +\infty. \end{aligned} \quad (13)$$

As a consequence, we obtain the following result:

PROPOSITION 2.1. *For every $\lambda \in \mathbb{R}$ the equation (6) has two linearly independent solutions $u_{1,\lambda}$, $u_{2,\lambda}$ such that*

$$u_{1,\lambda}(x) = 1 + o(1), \quad u'_{1,\lambda}(x) = o\left(\frac{1}{x}\right) \quad x \rightarrow 0^+, \quad (14)$$

$$u_{2,\lambda}(x) = x + o(x), \quad u'_{2,\lambda}(x) = 1 + o(1), \quad x \rightarrow 0^+,$$

and $u_{2,\lambda} \in H^2(0, 1)$.

For every $f \in L^2(0, 1)$ the solutions of $\tau u = f$ are given by

$$u(x) = c_1 u_{1,0}(x) + c_2 u_{2,0}(x) + u_f(x), \quad \forall x \in (0, 1), \quad c_1, c_2 \in \mathbb{R}, \quad (15)$$

where

$$u_f(x) = \int_0^x G(x, t) f(t) dt, \quad \forall x \in (0, 1), \quad (16)$$

$$G(x, t) = u_{1,0}(t) u_{2,0}(x) - u_{2,0}(t) u_{1,0}(x), \quad \forall x \in (0, 1), \quad t \in (0, 1)$$

fulfill $G \in L^\infty((0, 1)^2)$, $u_f(0) = 0 = u'_f(0)$ and $u_f \in H^2(0, 1)$.

Proof. The estimates in (14) follow from (9) and (13), while (16) is the usual variation of constants formula. Moreover, from (14) we obtain that $u_{2,\lambda}, u'_{2,\lambda} \in L^2(0, 1)$. On the other hand we have

$$q(x) u_{2,\lambda}(x) \sim x^{1-\alpha}, \quad x \rightarrow 0^+, \quad (17)$$

which implies that $q u_{2,\lambda} \in L^2(0, 1)$, since $\alpha < 5/4$ (cf. Remark 2.3 for comments on this restriction); using the fact that $\tau u_{2,\lambda} = \lambda u_{2,\lambda}$, we deduce that

$$u''_{2,\lambda} = \lambda u_{2,\lambda} - q u_{2,\lambda} \in L^2(0, 1).$$

From now on, we will indicate $u_i = u_{i,0}$, $i = 1, 2$. The fact that the function G defined in (16) belongs to the space $L^\infty((0, 1)^2)$ is a consequence of the asymptotic estimates (14). Moreover, from (16) we also deduce that $u_f(0) = 0$ and that

$$u'_f(x) = \int_0^x (u_1(t) u'_2(x) - u_2(t) u'_1(x)) f(t) dt, \quad \forall x \in (0, 1), \quad (18)$$

which implies $u'_f(0) = 0$.

Finally, the condition $u_f(0) = 0 = u'_f(0)$ guarantees that $u_f, u'_f \in L^2(0, 1)$; as far as the second derivative of u_f is concerned, let us observe that we have

$$\tau u_f = f$$

and so

$$u''_f = f - q u_f. \quad (19)$$

Using the fact that $u_f(0) = 0 = u'_f(0)$ and (7), it follows that $q u_f \in L^2(0, 1)$; hence $u_f \in H^2(0, 1)$. \square

In what follows, we study the spectral properties of suitable self-adjoint realizations of τ ; to this aim, let us first observe that the differential operator τ is regular at $x = 1$. As a consequence, it is in the limit circle case at $x = 1$; moreover, from (7), according to [17, Theorem 6.4], τ is in the limit circle case also in $x = 0$.

The differential operator A defined by

$$\begin{aligned} D(A) &= \{u \in L^2(0, 1) : u, u' \in AC(0, 1), \tau u \in L^2(0, 1), \\ &\quad \lim_{x \rightarrow 0^+} (xu'(x) - u(x)) = 0 = u(1)\} \\ Au &= \tau u, \quad \forall u \in D(A), \end{aligned}$$

is then a self-adjoint realization of τ ([17, p. 287-288]). We can show the validity of the following Proposition:

PROPOSITION 2.2. *The relation*

$$D(A) = H_0^2(0, 1)$$

holds true. Moreover, A has a bounded inverse $A^{-1} : L^2(0, 1) \rightarrow H_0^2(0, 1)$.

Proof. 1. Let us start proving that $H_0^2(0, 1) \subset D(A)$. It is well known that $H_0^2(0, 1) \subset C^1(0, 1)$; hence, for every $u \in H_0^2(0, 1)$ we have $u, u' \in AC(0, 1)$. Moreover, using the fact that $u(0) = 0$ we deduce that

$$u(x) = u'(0)x + o(x), \quad x \rightarrow 0^+$$

and

$$q(x)u(x) = u'(0)x^{1-\alpha} + o(x^{1-\alpha}), \quad x \rightarrow 0^+;$$

the condition $\alpha < 5/4$ guarantees again that $qu \in L^2(0, 1)$ and therefore $\tau u = -u'' + qu \in L^2(0, 1)$. Finally, the regularity of u and u' imply that

$$\lim_{x \rightarrow 0^+} (xu'(x) - u(x)) = 0$$

and so also the boundary condition in the definition of $D(A)$ is satisfied.

Now, let us prove that $D(A) \subset H_0^2(0, 1)$; for every $u \in D(A)$ let $f = \tau u \in L^2(0, 1)$. From (15) we deduce that u can be written as

$$u = c_1 u_1 + c_2 u_2 + u_f, \tag{20}$$

for some $c_1, c_2 \in \mathbb{R}$; it is easy to see that the function u_1 does not satisfy the boundary condition given in $x = 0$ in the definition of $D(A)$, while u_2 and u_f do. Hence $u \in D(A)$ if and only if $c_1 = 0$; the last statement of Proposition 2.1 implies then that $u \in H^2(0, 1)$. As in the first part of the proof, the regularity

of u allows to conclude that the boundary condition in $x = 0$ given in $D(A)$ reduces to $u(0) = 0$.

2. Let us study the invertibility of A ; the existence of a bounded inverse of A is equivalent to the fact that $0 \in \rho_A$, being ρ_A the resolvent of A . Since A is self-adjoint on $H_0^2(0, 1)$, this follows from the surjectivity of A (cf. [16, Theorem 5.24]); hence, it is sufficient to prove that A is surjective.

To this aim, let us first observe that condition (8) guarantees that 0 cannot be an eigenvalue of A . Now, let us fix $f \in L^2(0, 1)$ and let us prove that there exists $u \in H_0^2(0, 1)$ such that $Au = f$, i.e. $\tau u = f$; by applying Proposition 2.1 we deduce again that (20) holds true and the same argument of the first part of the proof implies that $c_1 = 0$.

Hence we obtain $u = c_2 u_2 + u_f$; from Proposition 2.1 we deduce that this function belongs to $H^2(0, 1)$ and satisfies the boundary condition $u(0) = 0$. In order to prove that the missing condition $u(1) = 0$ is fulfilled for every $f \in L^2(0, 1)$, let us observe that $u_2(1) \neq 0$, otherwise u_2 would be an eigenfunction of A associated to the zero eigenvalue. Therefore, $u(1) = 0$ is satisfied if

$$c_2 = -\frac{u_f(1)}{u_2(1)},$$

for every $f \in L^2(0, 1)$. □

REMARK 2.3. *As for the restriction $\alpha < 5/4$, we observe that for the proofs of Proposition 2.1 and Proposition 2.2 it is sufficient to require the milder condition $\alpha < 3/2$. The fact that $\alpha < 5/4$ is used (cf. [17, p. 287-288]) in order to obtain that $D(A)$ is the one described above. Finally, we observe that in the particular case when $\alpha < 1$ the problem is regular (cf., among others, [9]).*

The spectral properties of A are related to the oscillatory behaviour of solutions of (6). We first recall the following definition:

DEFINITION 2.4. *The differential equation (6) is oscillatory if every solution u has infinitely many zeros in $(0, 1)$. It is non-oscillatory when it is not oscillatory.*

We observe that the regularity assumptions on q imply that solutions of (6) have a finite number of zeros in any interval of the form $[a, 1)$, for every $0 < a < 1$. Moreover, from (7) we infer that for every $\lambda \in \mathbb{R}$ there exists $c(\lambda) \in (0, 1]$ such that

$$\lambda - q(x) < 0, \quad \forall x \in (0, c(\lambda)).$$

An application of the Sturm comparison theorem proves that every solution of (6) has at most one zero in $(0, c(\lambda))$; as a consequence, we obtain the following result:

PROPOSITION 2.5. *For every $\lambda \in \mathbb{R}$ the differential equation (6) is non-oscillatory.*

Once Proposition 2.5 is obtained, we can provide in a straightforward way some useful information on the spectral properties of A ; more precisely, denoting by σ_{ess} the essential spectrum of a given operator, we have:

PROPOSITION 2.6. *([17, Theorem 14.3, Theorem 14.6 and Theorem 14.9], [12, Theorem XIII.1]) The differential operator A is bounded-below and satisfies*

$$\sigma_{ess}(A) = \emptyset.$$

Moreover, there exists a sequence $\{\lambda_n\}_{n \in \mathbb{N}}$ of simple eigenvalues of A such that

$$\lim_{n \rightarrow +\infty} \lambda_n = +\infty$$

and for every $n \in \mathbb{N}$ the eigenfunction u_n of A associated to the eigenvalue λ_n has $(n-1)$ simple zeros in $(0, 1)$.

REMARK 2.7. *According to [17], operators of the form τ (defined on functions whose domain is $(0, +\infty)$) arise when the time independent Schrödinger equation with spherically symmetric potential*

$$-\Delta u(x) + V(|x|)u(x) = \lambda u(x), \quad u \in L^2(\mathbb{R}^m) \quad (21)$$

is reduced to an infinite system of eigenvalue problems associated to the ordinary differential operators in $L^2(0, +\infty)$

$$\tau_i = -\frac{d^2}{dr^2} + \frac{1}{r^2} \left[i(i+m-2) + \frac{1}{4}(m-1)(m-3) \right] + V(r)$$

($i \in \mathbb{N}$). In Appendix 17.F of [17] it is treated the case of a potential V satisfying assumptions (which enable to consider Coulomb potentials) that lead to (7). More precisely, it is shown that for $m=3, i=0$ the operator is in the limit circle case at zero and self-adjoint extensions of τ_0 are described.

3. The main result

In this section we are interested in proving a global bifurcation result for a nonlinear eigenvalue problem of the form

$$-u'' + q(x)u = \lambda u + g(x, u)u, \quad \lambda \in \mathbb{R}, \quad x \in (0, 1], \quad (22)$$

where $q \in C((0, 1])$ satisfies (7) and $g \in C([0, 1] \times \mathbb{R})$ is such that

$$\lim_{u \rightarrow 0} g(x, u) = 0, \quad \text{uniformly in } x \in [0, 1]. \quad (23)$$

We will look for solutions u of (22) such that $u \in H_0^2(0,1)$. To this aim, let Σ denote the set of nontrivial solutions of (22) in $H_0^2(0,1) \times \mathbb{R}$ and let $\Sigma' = \Sigma \cup \{(0, \lambda) \in H_0^2(0,1) \times \mathbb{R} : \lambda \text{ is an eigenvalue of } A\}$, where A is as in Section 2.

Let M denote the Nemitskii operator associated to g , given by

$$M(u)(x) = g(x, u(x))u(x), \quad \forall x \in [0, 1],$$

for every $u \in H_0^2(0,1)$. We can show the validity of the following:

PROPOSITION 3.1. *Assume $g \in C([0, 1] \times \mathbb{R})$ and (23). Then $M : H_0^2(0,1) \longrightarrow L^2(0,1)$ is a continuous map and satisfies*

$$M(u) = o(\|u\|), \quad u \rightarrow 0. \quad (24)$$

Proof. 1. We first show that $Mu \in L^2(0,1)$ when $u \in H_0^2(0,1)$. When this condition holds, $u \in L^\infty(0,1)$ and the continuity of g implies that there exists $C_u > 0$ such that

$$|g(x, u(x))u(x)| \leq C_u, \quad \forall x \in [0, 1].$$

As a consequence we obtain $Mu \in L^\infty(0,1) \subset L^2(0,1)$.

2. Let us prove that M is continuous. Let us fix $u_0 \in X$ and let $u_n \in X$ such that $u_n \rightarrow u_0$ when $n \rightarrow +\infty$; the continuous embedding

$$H_0^2(0,1) \subset L^\infty(0,1)$$

and the uniform continuity of g on compact subsets of $[0, 1] \times \mathbb{R}$ ensure that

$$g(x, u_n(x)) \rightarrow g(x, u_0(x)) \quad \text{in } L^\infty(0,1). \quad (25)$$

This is sufficient to conclude that $Mu_n \rightarrow Mu_0$ in $L^\infty(0,1)$ and hence $Mu_n \rightarrow Mu_0$ in $L^2(0,1)$.

3. Finally, let us prove (24): using again the fact that $H_0^2(0,1) \subset L^\infty(0,1)$, we have

$$\|Mu\|_{L^2(0,1)} \leq \|g(x, u(x))\|_{L^\infty(0,1)} \|u\|_{L^2(0,1)} \leq \|g(x, u(x))\|_{L^\infty(0,1)} \|u\|,$$

for all $u \in H_0^2(0,1)$; hence, we deduce that

$$\frac{\|Mu\|_{L^2(0,1)}}{\|u\|} \leq \|g(x, u(x))\|_{L^\infty(0,1)}, \quad \forall u \in H_0^2(0,1), \quad u \neq 0.$$

Therefore the result follows from (23) and (25). \square

Now, let us observe that the search of solutions $u \in H_0^2(0, 1)$ of (22) is equivalent to the search of solutions of the abstract equation

$$Au = \lambda u + M(u), \quad (u, \lambda) \in H_0^2(0, 1) \times \mathbb{R}; \quad (26)$$

on the other hand, (26) can be written in the form

$$w = \lambda R w + M(Rw), \quad (w, \lambda) \in L^2(0, 1) \times \mathbb{R}, \quad (27)$$

where $R : L^2(0, 1) \rightarrow H_0^2(0, 1)$ is the inverse of A (cf. Proposition 2.2).

Now, from [17, Theorem 7.10] we deduce that R is compact; this fact and the continuity of M guarantee that the operator $MR : L^2(0, 1) \rightarrow H_0^2(0, 1)$ is compact. Moreover, the condition

$$M(Rw) = o(\|w\|_{L^2(0,1)}), \quad w \rightarrow 0, \quad (28)$$

is a consequence of (24). From an application of the global bifurcation result of Rabinowitz (cfr. [11]) to (27) we then obtain the following result:

THEOREM 3.2. *Assume (7) and (23). Then, for every eigenvalue λ_n of A there exists a continuum C_n of nontrivial solutions of (22) in $H_0^2(0, 1) \times \mathbb{R}$ bifurcating from $(0, \lambda_n)$ and such that one of the following conditions holds true:*

- (1) C_n is unbounded in $H_0^2(0, 1) \times \mathbb{R}$;
- (2) C_n contains $(0, \lambda_{n'}) \in \Sigma'$, with $n' \neq n$.

Now, let us observe that a more precise description of the bifurcating branch, eventually leading to exclude condition (2), can be obtained when there exists a continuous functional $j : \Sigma' \rightarrow \mathbb{N}$ (cf. [2, Pr. 2.1]). In order to define such a functional, we will use the fact that nontrivial solutions of (22) have a finite number of zeros in $(0, 1)$; this will be a consequence of our next result.

For every $\lambda \in \mathbb{R}$ and for every nontrivial solution $u \in H_0^2(0, 1)$ of (22) let us define $q_{u,\lambda} : (0, 1] \rightarrow \mathbb{R}$ by $q_{u,\lambda}(x) = q(x) - \lambda - g(x, u(x))$, for every $x \in (0, 1]$. The following Lemma holds true:

LEMMA 3.3. *For every $\lambda \in \mathbb{R}$ and for every nontrivial solution $u \in H_0^2(0, 1)$ of (22) there exists a neighborhood $U \subset H_0^2(0, 1) \times \mathbb{R}$ of (u, λ) and $x_{u,\lambda} \in (0, 1)$ such that*

$$q_{v,\mu}(x) > 0, \quad \forall (v, \mu) \in U, \quad x \in (0, x_{u,\lambda}]. \quad (29)$$

Proof. Let $(u, \lambda) \in H_0^2(0, 1) \times \mathbb{R}$, $u \neq 0$, be fixed and let U be the neighborhood of radius 1 of (u, λ) in $H_0^2(0, 1) \times \mathbb{R}$; from the continuous embedding $L^\infty(0, 1) \subset H_0^2(0, 1)$ we deduce that if $(w, \mu) \in \Sigma \cap U_1$ then

$$\|w\|_{L^\infty(0,1)} \leq 1 + \|u\|_{L^\infty(0,1)}, \quad |\mu| \leq 1 + |\lambda|$$

and

$$q(x) - \mu - g(x, w(x)) \geq q(x) - |\lambda| - 1 - \max_{\substack{x \in [0,1], \\ |s| \leq 1 + \|u\|_{L^\infty(0,1)}}} |g(x, s)|, \quad \forall x \in (0, 1).$$

From (7) we then deduce that there exists $x_{(u,\lambda)} \in (0, 1)$, depending only on (u, λ) , such that

$$q(x) - \mu - g(x, w(x)) > 0, \quad \forall x \in (0, x_{(u,\lambda)}].$$

□

Now, let us observe that for every $\lambda \in \mathbb{R}$ and for every nontrivial solution $u \in H_0^2(0, 1)$ of (22) the function u is a nontrivial solution of the linear equation

$$-w'' + (q(x) - g(x, u(x)) - \lambda)w = 0. \quad (30)$$

From Lemma 3.3, with an argument similar to the one which led to Proposition 2.5, we deduce that all the nontrivial solutions of (30) (in particular u) have a finite number of zeros in $(0, 1)$. We denote by $n(u)$ this number.

We are then allowed to define the functional j by setting

$$j(u, \lambda) = \begin{cases} n(u) & \text{if } u \not\equiv 0 \\ n - 1 & \text{if } u \equiv 0 \text{ and } \lambda = \lambda_n, \end{cases} \quad (31)$$

for every $(u, \lambda) \in \Sigma'$. Let us observe that the definition $j(0, \lambda_n) = n - 1$ is suggested by Proposition 2.6.

PROPOSITION 3.4. *The function $j : \Sigma' \rightarrow \mathbb{N}$ is continuous.*

Proof. 1. As for the continuity of j in every point of the form $(0, \lambda_n)$, $n \in \mathbb{N}$, we refer to [15, Lemma 2.5].

2. Let us now fix $(u_0, \lambda_0) \in \Sigma$ and let $(u, \lambda) \in U$, with U as in Lemma 3.3; this Lemma guarantees that both u and u_0 have no zeros in $(0, x_{u_0, \lambda_0})$.

On the other hand, in the interval $[x_{u_0, \lambda_0}, 1]$ a standard continuous dependence argument (cf. also [11]) ensures that u and u_0 have the same numbers of zeros if (u, λ) is in a sufficiently small neighborhood of (u_0, λ_0) . As a consequence, we obtain that there exists a neighborhood U_0 of (u_0, λ_0) such that

$$j(u, \lambda) = j(u_0, \lambda_0), \quad \forall (u, \lambda) \in U_0.$$

□

As a consequence, from Theorem 3.2 and Proposition 3.4 we deduce the final result:

THEOREM 3.5. Assume (7) and (23). Then, for every eigenvalue λ_n of A there exists a continuum C_n of nontrivial solutions of (22) in $H_0^2(0, 1) \times \mathbb{R}$ bifurcating from $(0, \lambda_n)$ and such that condition (1) of Theorem 3.2 holds true and

$$j(u, \lambda) = n - 1, \quad \forall (u, \lambda) \in C_n. \quad (32)$$

REMARK 3.6. Theorem 3.2 can be proved as an application of Stuart's result [15, Theorem 1.2] as well. However, since in the situation considered in this paper the singularity at zero does not affect the compactness of the operator R defined after (27), we chose to apply Rabinowitz theorem [11]. We finally mention the interesting paper [1], where global branches of solutions, with prescribed nodal properties, are obtained for a second order degenerate problem in $(0, 1)$.

REFERENCES

- [1] H. BERESTYCKI AND M.J. ESTEBAN, *Existence and bifurcation of solutions for an elliptic degenerate problem*, J. Differential Equations **134** (1997), 1–25.
- [2] A. CAPIETTO AND W. DAMBROSIO, *Preservation of the Maslov index along bifurcating branches of solutions of first order systems in \mathbb{R}^n* , J. Differential Equations **227** (2006), 692–713.
- [3] A. CAPIETTO AND W. DAMBROSIO, *Planar Dirac-type systems: the eigenvalue problem and a global bifurcation result*, J. London Math. Soc. **81** (2010), 477–498.
- [4] E.A. CODDINGTON AND N. LEVINSON, *Theory of Ordinary Differential Equations*, McGraw-Hill, 1955.
- [5] B. ČURGUS AND T.T. READ, *Discreteness of the spectrum of second-order differential operators and associated embedding theorems*, J. Differential Equations **184** (2002), 526–548.
- [6] M.S.P. EASTHAM, *The Asymptotic Solution of Linear Differential Systems*, London Math. Society Monographs New Series, 1989.
- [7] P. FELMER AND J.J. TORRES, *A nonlinear eigenvalue problem in \mathbb{R} and multiple solutions of nonlinear Schrödinger equation*, Adv. Differential Equations **7** (2002), 1215–1234.
- [8] F. HADJ SELEM, *Radial solutions with prescribed numbers of zeros for the nonlinear Schrödinger equation with harmonic potential*, Nonlinearity **24** (2011), 1795–1819.
- [9] R. LEMMERT AND W. WALTER, *Singular nonlinear boundary value problems*, Appl. Anal. **72** (1999), 191–203.
- [10] D.B. PEARSON, *Quantum Scattering and Spectral Theory*, Academic Press, London, 1988.
- [11] P. RABINOWITZ, *Some global results for non-linear eigenvalue problems*, J. Funct. Anal. **7** (1971), 487–513.
- [12] M. REED AND B. SIMON, *Methods of Modern Mathematical Physics. Vol. 4: Analysis of Operators*, Academic Press, London, 1978.
- [13] H. SCHMID AND C. TRETTER, *Eigenvalue accumulation for Dirac operators with spherically symmetric potential*, J. Differential Equations **181** (2002), 511–542.

- [14] I. SIM, R. KAJIKIYA, AND Y.-H. LEE, *On a criterion for discrete or continuous spectrum of p -Laplace eigenvalue problems with singular sign-changing weights*, Nonlinear Anal. **72** (2010), 3515–3534.
- [15] C. STUART, *Global properties of components of solutions of non-linear second order differential equations on the half-line*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) **2** (1975), 265–286.
- [16] J. WEIDMANN, *Linear Operators in Hilbert Spaces*, Graduate Texts in Mathematics, no. 68, Springer, Berlin, 1980.
- [17] J. WEIDMANN, *Spectral Theory of Ordinary Differential Equations*, Lectures Notes in Mathematics, no. 1258, Springer, Berlin, 1987.

Authors' addresses:

Anna Capietto
Dipartimento di Matematica
Università di Torino
Via Carlo Alberto 10, 10123 Torino, Italy
E-mail: anna.capietto@unito.it

Walter Dambrosio
Dipartimento di Matematica
Università di Torino
Via Carlo Alberto 10, 10123 Torino, Italy
E-mail: walter.dambrosio@unito.it

Duccio Papini
Dipartimento di Ingegneria dell'Informazione e Scienze Matematiche,
Università di Siena
Via Roma 56, 53100 Siena, Italy
E-mail: papini@dii.unisi.it

Received May 28, 2012
Revised September 3, 2012