



UNIVERSITÀ
DEGLI STUDI
DI UDINE

Università degli studi di Udine

Influence of lipoproteins at dry-off on metabolism of dairy cows during transition period and on postpartum reproductive outcomes

Original

Availability:

This version is available <http://hdl.handle.net/11390/1105062> since 2017-04-21T08:21:13Z

Publisher:

Published

DOI:10.1016/j.theriogenology.2017.02.007

Terms of use:

The institutional repository of the University of Udine (<http://air.uniud.it>) is provided by ARIC services. The aim is to enable open access to all the world.

Publisher copyright

(Article begins on next page)

1 **Influence of lipoproteins at dry-off on metabolism of dairy cows during transition period and**
2 **on postpartum reproductive outcomes**

3

4 Martina Crociati^a, Lakamy Sylla^a, Claudia Floridi^a, Antonella Comin^b, Gabriele Fruganti^a, Maurizio
5 Monaci^a, Giuseppe Stradaoli^{b*}

6

7 ^aDepartment of Veterinary Medicine, Via San Costanzo, 4 - 06126 – Perugia, Italy;

8 ^bDepartment of Agricultural, Food, Environmental and Animal Sciences (DI4A), Via Delle Scienze,
9 206 - 33100 - Udine, Italy.

10

11

12

13 *Corresponding author: Giuseppe Stradaoli, Department of Agricultural, Food, Environmental and
14 Animal Sciences (DI4A), Via Delle Scienze, 206 - 33100 Udine, Italy, Tel. +39 0432 558580, Fax.
15 +39 0432 558503.

16 Email: giuseppe.stradaoli@uniud.it.

17

18

19 Author contributions: M.C., L.S., M.M., and G.S. designed the study. M.C. conducted most of the
20 fields experimental procedures used in the study, i.e., the samples and data collection, the clinical
21 classification of animals, and data analysis, and drafted the manuscript. C.F. performed lipoprotein
22 and blood's biochemical analysis. A.C. performed P4 determinations. M.C. and L.S. performed data
23 analysis and statistics. G.F., M.M., and G.S. supervised experiments and critically revised the
24 manuscript. All authors have read and approved the manuscript.

25

26

27 **Abstract**

28 High-yielding dairy cows are metabolically challenged during transition, when intense mobilization
29 and hepatic oxidation of lipids is achieved, thus leading to fatty infiltration, ketosis and generalized
30 inflammation. The condition is associated to periparturient diseases and poor fertility. The aim of
31 this study was to assess whether serum lipoprotein concentrations in the dry period could influence
32 the occurrence of postpartum diseases and reproductive performance in dairy cows. The study was
33 carried out on 30 multiparous Holstein Friesian cows. Blood samples were collected at dry-off (-60
34 days), 30 days after dry-off and within 12 hours after parturition for biochemical and serum
35 lipoprotein assays. From 10 to 60 days after parturition milk was collected twice weekly after
36 feeding, for milk whey progesterone assay. The Optimal Cutpoint package identified a threshold of
37 89% for serum High Density Lipoprotein (HDL) concentration at the beginning of the dry period
38 with 95% of confidence interval. Cows with serum HDL greater than 89% (High group, n = 10)
39 showed better reproductive performance when compared to those with low values (Low group, n =
40 18). The odds ratio for reproductive disorders in High group was 0.6875, however, differences were
41 not significant probably due to both the reduced number of animals per group and overall low
42 incidence of postpartum reproductive disease. First postpartum luteal activity occurred around day
43 23, while the second one between days 40 and 48. The average calving to first AI interval was 64.00
44 \pm 3.95 days and 94.50 \pm 12.32 days in High and Low group, respectively ($P < 0.05$). The calving-
45 conception interval was 129.86 \pm 24.42 days and 199.18 \pm 24.73 days in High and Low groups,
46 respectively ($P < 0.05$). Low group displayed an increase in liver markers, that is total bilirubin,
47 with 0.46 \pm 0.09 mg/dL and 0.23 \pm 0.08 mg/dL, in Low and High group respectively ($P < 0.05$), and
48 NEFA/cholesterol ratio, with 0.30 \pm 0.06 and 0.14 \pm 0.03, in Low and High groups, respectively (P
49 < 0.05), at parturition. Concentrations of HDL $> 89\%$ at dry-off could be suggestive of improved
50 liver adaptation to the transition, and probably of enhanced fertility in High group.

51

52 **Key words:** dairy cow; dry period; lipoprotein; fertility; reproductive disorders

53 **1 Introduction**

54 In the last weeks of gestation, cow's metabolism has to sustain fetal growth, mammary gland
55 development and after calving the energy demand further increases due to lactation. Usually, during
56 this period the dry matter intake is not sufficient to meet the requirements. In order to cope with this
57 imbalance, dairy cows mobilize great amounts of body reserves, especially lipids [1]. Lipids are
58 released from periphery into the circulation in the form of non-esterified fatty acids (NEFA) and
59 then they are used in liver for gluconeogenesis and ketogenesis. If negative energy balance is
60 excessive, the intake of NEFA overcomes the possibility of complete oxidation in the liver. In this
61 case, NEFA are re-esterified, turned into triglycerides and stored into the cytoplasm of hepatocytes
62 as lipid droplets. This condition, known as fatty liver, is usually associated to impairment of liver
63 function, subclinical or clinical ketosis, periparturient metabolic diseases and poor fertility, which
64 deeply influence the herd profitability [2-5]. Human and bovine hepatocytes are able to synthesize
65 very low-density lipoprotein (VLDL), in order to export triglycerides towards peripheral tissues [6].
66 However, bovine liver is not able to adjust VLDL synthesis based on NEFA absorption and re-
67 esterification into cytoplasmic droplets [2]; therefore, the prevalence of lipidosis among transition
68 dairy cows can reach 50% [7-8]. Very-low density lipoprotein represents only 3 to 5% of all
69 circulating lipoproteins in cattle, while other classes, such as high-density (HDL) and low-density
70 lipoprotein (LDL), account for 80-85% and 10-15%, respectively. As liver can export triglycerides
71 only through VLDL, it is generally assumed that low levels of circulating VLDL are representative
72 of increased fatty acids infiltration [6, 9].

73 It has been reported a decrease of pregnancy at first artificial insemination when more than 50% of
74 cows in a herd have serum NEFA ≥ 0.5 mEq/L one week before calving, as indicative of intense
75 lipids mobilization [3]. An increased odds ratio for retention of fetal membranes and metritis in
76 cows with prepartum NEFA ≥ 0.3 mEq/L was reported [4]. It has also been highlighted that the
77 peculiar metabolic status of the dairy cows during periods of lipids mobilization could cause lipid
78 accumulation in oocytes and the regenerating endometrium, which impairs fertility via reduction in

79 embryo survival and increased inflammatory changes, respectively [10]. The majority of studies
80 concerning the association between biochemistry profiles of dry cows in late gestation and
81 postpartum performance are focused on the last two or three weeks before calving [3-4, 9].
82 However, Dervishi et al. [11] reported alterations in inflammatory and metabolic profiles as early as
83 eight weeks before calving, in dairy cows that will develop metritis. Brscic et al. [12], recently
84 reported some reference limits for metabolic profiles in Holstein late-pregnant heifers and dry cows,
85 but postpartum performance was not assessed.

86 Moved by the need of finding new predictive indexes of reduced reproductive efficiency, we
87 hypothesized that the level of circulating lipoproteins during the last period of pregnancy could be
88 indicative of the degree of adaptation to transition of dairy cows and of their susceptibility to both
89 postpartum reproductive diseases and reduced fertility. Moreover, we evaluated biochemistry
90 profiles from dry-off until calving as a tool to verify the overall health status in our experimental
91 animals and to exclude alterations in lipoprotein metabolism due to subclinical pathological
92 conditions.

93

94 **2 Materials and Methods**

95 *2.1 Animals and husbandry*

96 In the present study, 30 healthy multiparous Holstein Friesian cows were selected with parity
97 ranging from 2 to 5; they belonged to a commercial dairy farm located in Central Italy (42°95' N,
98 12°39' E), with a herd size ranging between 100 and 120 lactating cows. Each cow was randomly
99 selected through the herd-management software before entering into the dry period. Mean heard
100 intercalving period was 410.27 ± 15.87 days, the mean dry period length was 60.25 ± 10.14 days
101 and the voluntary waiting period averaged 50.16 ± 7.08 days. Average milk production was above
102 10,900 kg/lactation/cow. All lactating cows were housed in free stalls with cubicles and milked with
103 two Automatic milking systems (DeLaval S.p.A., VMS, Milano, Italy); dry cows were kept in a free
104 stall barn with straw. The dry cows had free access to a total mixed ration (TMR) offered *ad*

105 *libitum*, composed of 4.5 kg wheat straw (4.60% CP, 78.90% NDF, 48.40% ADF), 4.5 kg oat hay
106 (8.70% CP, 61.30% NDF, 38.20% ADF), 3.0 kg concentrate (28.50% CP, 20.20% NDF, 9.70%
107 ADF) per head. During the close-up period this ration was supplemented with 10 kg of fresh cow
108 TMR/head. The overall TMR composition was: 4 kg alfalfa and lolium mixed hay (7.67% CP,
109 37.86% NDF, 20.84% ADF), 3.5 kg alfalfa hay (14.91% CP, 42.70% NDF, 33.10% ADF), 11.5 kg
110 concentrate (28.50% CP, 20.20% NDF, 9.70% ADF), 22 kg corn silage (9.20% CP, 45.90% NDF,
111 27.00% ADF) and 0.8 kg molasses (4.30 % CP, 0% NDF, 0% ADF). Mycotoxins content in feed
112 was within the legislative established limits (Italian Law 149/2004). Contents of organic acid in
113 silage and total mixed ration from dry cow feed-bunk were evaluated by HPLC analysis. Only
114 butyric acid was fairly present in both silage and total mixed ration, with mean values of $0.02 \pm$
115 0.01 and 0.03 ± 0.001 g/100g, respectively.

116

117 *2.2 Experimental procedures and samples collection*

118 The study was conducted from January to December 2015; all experimental cows calved before the
119 end of April 2015. Two cows were excluded from the experiment due to abortion and premature
120 calving. Body condition score (BCS), assessed through a five-point scale [13], and blood sampling
121 were performed at 60 (T1) and 30 days (T2) before the expected calving and within 12 hours
122 postpartum (T3). Samples were obtained by coccygeal vein puncture into plain and EDTA vacuum
123 tubes (BD Vacutainer Systems, Plymouth, UK) between 5.30 and 6.00 AM, half an hour before
124 feeding. Samples were stored at 4°C and delivered within 1 hour to the Laboratory; serum was
125 obtained through centrifugation at 1,300 g for 10 minutes and immediately processed. An aliquot of
126 serum from each sample was stored at 4°C until electrophoresis analysis. From 10 to 60 days after
127 parturition milk was collected twice weekly after feeding, between 6.00 and 7.00 AM, in empty
128 eppendorf tubes and immediately frozen at -20°C, until analysis.

129 To evaluate healthy condition of the newborn calf, the APGAR score described by Vannucchi et al.

130 [14] was employed. Briefly, the following symptoms were considered and scored: mucous color (0
131 = cyanotic; 1 = pale; 2 = normal), heart rate (0 = absent; 1 = bradycardia, < 80 beats per minute or
132 irregular; 2 = regular, > 100 beats per minute), muscle tone (0 = flaccid; 1 = slight flexion; 2 =
133 flexion), activity (0 = absent; 1 = some movement; 2 = active calf), and respiration (0 = absent; 1 =
134 irregular < 24 respiration per minute; 2 = regular > 36 respiration per minute).

135 Postpartum uterine diseases were diagnosed according to Sheldon et al. [15]. Briefly, animals were
136 monitored once weekly after calving until complete uterine involution was achieved. The fetal
137 membranes were considered retained when not released within 12 hours after parturition. Cows that
138 presented an enlarged uterus with watery red-brown to viscous off-white purulent uterine discharge,
139 from 0 to 21 days after calving were considered suffering from metritis. Clinical endometritis was
140 defined by the presence of pathological uterine discharge 21 days or more postpartum.

141 Reproductive parameters such as calving to first insemination interval, calving to conception
142 interval and number of inseminations per pregnancy were retrieved from the herd management
143 software, at least until 200 days after parturition.

144 The experimental activity was carried out in accordance to the guidelines of animals experiments as
145 set by the Italian Law 26/2014 (national application of EU Directive 2010/63/EU) and has been
146 approved by the Ethical Committee of the University of Perugia.

147 148 *2.3 Serum lipoprotein's electrophoresis, biochemical profiles and CBC*

149 Electrophoresis of serum lipoproteins was obtained with the Hydrasys – LC Sebia automatic system
150 and Hydragel 7 LIPO + Lp(a) Kit (Sebia® Electrophoresis, Sebia Inc., Norcross, GA USA). Briefly,
151 lipoprotein classes were separated by electrophoresis on agarose gel buffered plates (pH 8.5);
152 different bands were then read with densitometry scanner at 570 nm (Epson Perfection V700
153 PHOTO, Seiko Epson Corporation, Japan) and lipoproteins were expressed in term of relative
154 percentage. Three main bands were identified, namely HDL (or α -lipoproteins), VLDL (or pre- β -
155 lipoproteins) and LDL (or β -lipoproteins).

156 The serum concentrations of albumin (ALB), β -hydroxybutyrate (BHB), total proteins (TP),
157 triglycerides (TG), total bilirubin, direct bilirubin, indirect bilirubin, glucose, urea, creatinine, lactic
158 dehydrogenase (LDH), aspartate transaminase (AST), alanine transaminase (ALT), non-esterified
159 fatty acids (NEFA), gamma-glutamyl transpeptidase (GGT), alkaline phosphatase (ALP), creatine
160 kinase (CK), cholesterol (CHOL), magnesium (Mg), calcium (Ca) and phosphorus (P) were
161 determined with an Hitachi 904 chemistry analyzer (Boehringer Mannheim, Germany).

162 All diagnostic kits were purchased from DiaSys – Diagnostic Systems (GmbH, Alte Strasse 9, D –
163 65558 Holzheim, Germany). Count of blood cells (CBC) was obtained in EDTA samples with a
164 Heco Vet S (SEAC, Florence, Italy) electric impedance system.

165

166 *2.4 Milk whey progesterone assay*

167 Milk progesterone was quantified into the whey fraction through direct enzyme immunoassay as
168 described by Comin et al. [16]. Briefly, whey aliquots were added to micro plates, with 25 μ L/well
169 of anti-progesterone antibody with 50 μ L of progesterone-peroxidase conjugate in ELISA buffer.

170 Plates were incubated one night at 4°C and then washed five times with buffer solution.

171 Chromogenic substrate was added and plates were incubated for 30 minutes at room temperature.

172 Finally, absorbance was read at 450 nm using a plate reader. Resumption of ovarian activity was
173 considered when progesterone concentrations was above 300 pg/mL [16], and presence of a corpus
174 luteum was confirmed by trans rectal ultrasound examination.

175

176 *2.5 Statistical analysis*

177 Data were analyzed through SPSS software [17]. Normal distribution was assessed for each
178 biochemical and reproductive outcome through Kolmogorov-Smirnov test with 95% of confidence
179 interval. However, due to the limited number of animals involved and in order to be conservative,
180 non-parametric statistics were used. All tests were set with $\alpha < 0.05$. Data were expressed through
181 mean and standard error of the mean (SEM). Differences between groups were analyzed with

182 Kruskal-Wallis test. R software v3.2.4 [18] was used to run Optimal Cutpoints package [19] in
183 order to estimate threshold in lipoproteins concentration at T1, T2 and T3 in relation to postpartum
184 diseases. The optimal cutpoint for the diagnostic test (HDL, VLDL and LDL in each of the three
185 periods) was computed by means of a Receiver Operating Characteristic (ROC), using the disease
186 status (e.g. sick or healthy) as dependent categorical variable and optimizing the Younden's Index,
187 so that the sum of Specificity and Sensitivity was maximized. Cows were then assigned to Low and
188 High HDL groups; differences in reproductive outcomes and mean progesterone concentrations
189 were investigated with Kruskal Wallis test in SPSS. If any significant difference was found between
190 High and Low HDL cows, data were discussed as belonging to a single group based on sampling
191 interval.

192

193 **3 Results**

194 *3.1 Clinical outcomes and resumption of ovarian activity*

195 Cows entered the dry period with a mean BCS of 3.23 ± 0.07 that decreased to 3.04 ± 0.05 at
196 calving ($P < 0.05$). Nineteen out of 28 animals had normal postpartum while 3 cows showed
197 retained fetal membranes and 6 metritis. All calving were eutocic and all calves were born healthy
198 and with a mean APGAR score of 8.21 ± 1.15 .

199 Left abomasal displacement was observed in 3 animals without any reproductive disorder.

200 Milk whey progesterone in High ($n = 10$) and Low ($n = 18$) group of cows ranged between 150 and
201 500 pg/mL, but no significant difference was identified. First postpartum luteal activity occurred
202 around day 23, while the second one from day 40 to 48. One animal in High and one in the Low
203 group revealed an increase in P4 concentrations only after 50 days post calving. One cow per group
204 also showed a prolonged period of elevated milk P4.

205

206

207 *3.2 Biochemistry profiles and CBC*

208 3.2.1 Serum lipoprotein electrophoresis

209 Relative proportion of lipoproteins in experimental dairy cows is reported in Table 1. VLDL
210 lipoproteins ranged between 3.27 ± 0.29 % and 2.69 ± 1.43 %, without any difference between
211 groups; HDL decreased ($P < 0.001$) from dry-off (87.52 ± 0.87 %) towards T2 (82.41 ± 1.21 %),
212 then increased at calving (87.39 ± 1.07 %). LDL lipoproteins presented a reverse trend; their
213 concentration was greater at T2, with mean value of 14.32 ± 1.18 % ($P < 0.01$).

214 The Optimal Cutpoint package identified a threshold of 89% for serum HDL concentration at the
215 beginning of the dry period with 95% of confidence interval. As shown in Table 2, cows with serum
216 HDL greater than 89% ($n = 10$) tended to show better reproductive performances when compared to
217 others ($n = 18$). More deeply, the average calving to first AI interval was 64.00 ± 3.95 days and
218 94.50 ± 12.32 days in High and Low group, respectively ($P < 0.05$). The calving - conception
219 intervals were 129.86 ± 24.42 days and 199.18 ± 24.73 days in High and Low groups, respectively
220 ($P < 0.05$). The odds ratio for reproductive disorders in High HDL cows was 0.6875, even if the P
221 was not significant.

222

223 3.2.2 Energy metabolism analytes

224 Beta-hydroxybutyrate, glucose, NEFA and triglyceride concentration are reported in Table 3. Beta-
225 hydroxybutyrate was constantly below reference limit of 1.40 mmol/L and did not show any
226 significant variation throughout the period of study. Mean serum glucose level decreased from
227 66.11 ± 1.85 mg/dL at T1 to 55.75 ± 2.24 mg/dL at parturition ($P < 0.001$) while NEFA increased
228 from 0.18 ± 0.02 mmol/L at dry-off to 0.58 ± 0.08 mmol/L at calving ($P < 0.001$). Triglycerides
229 increased both at T1 and T2, while they declined at calving time ($P < 0.001$).

230

231 3.2.3 Markers of liver function

232 Low group displayed significantly higher concentration of both total and indirect bilirubin, and
233 NEFA/CHOL ratio than High group, only at calving (Table 4).

234 In all cows, urea and LDH did not show any significant variation during the dry period. Aspartate
235 transaminase activity was higher at calving, with a mean value of 106.57 ± 6.98 UI/L, compared to
236 what found during the dry period ($P < 0.001$). Concerning GGT, it decreased at parturition with a
237 mean value of 15.22 ± 1.31 UI/L ($P < 0.05$) and ALT showed a similar trend, averaging $20.21 \pm$
238 0.96 UI/L at calving ($P < 0.001$). Cholesterol declined progressively from 192.79 ± 5.30 mg/dL to
239 112.57 ± 6.96 mg/dL ($P < 0.001$; Table 5).

240

241 *3.2.4 Protein metabolism analytes*

242 Protein metabolism analyses are reported in Table 6. Albumin and CK did not show any significant
243 variation throughout the period of study. Conversely, total serum proteins decreased from T1 (7.56
244 ± 0.16 g/dL) until parturition, when they reached an average value of 6.43 ± 0.11 g/dL ($P < 0.001$);
245 CK remained constant and averaged 265.69 ± 52.59 UI/L throughout the study.

246

247 *3.2.5 Mineral metabolism and other analytes*

248 Mineral and other analytes are reported in Table 7. At dry-off, calcium and phosphorus averaged
249 9.65 ± 0.12 mg/dL and 6.55 ± 0.28 mg/dL, respectively; thereafter they decreased at calving, with
250 mean values of 8.80 ± 0.29 mg/dL ($P < 0.05$) and 4.83 ± 0.28 mg/dL ($P < 0.001$), respectively.

251 Alkaline phosphatase activity was higher at dry-off, with a mean value of 84.57 ± 4.68 UI/L and
252 then decreased to 64.46 ± 5.04 UI/L at calving ($P < 0.001$). Creatinine constantly averaged $1.25 \pm$
253 0.06 mg/dL. Magnesium and Creatinine did not show any significant variation throughout the period
254 of study.

255

256 *3.2.6 Count blood cells*

257 The CBC features were within the referenced limits [20] and there were any significant difference
258 among groups (data not shown).

259

260

261 **4 Discussion**

262 The transition dairy cow has to cope with negative energy balance, lipomobilization and fatty liver,
263 all of them influencing the susceptibility to postpartum diseases and the reproductive outcomes.
264 However, there is lack of data concerning lipoprotein metabolism during the full dry period and
265 their interaction with postpartum health and fertility.

266 High-density lipoproteins, herein reported, were greater at dry-off and at calving, compared to T2,
267 while LDL reached a peak one month before expected delivery. This opposite trend could be due to
268 LDL peripheral catabolism, as mammary gland absorbs triglycerides from LDL for milk fatty acids
269 synthesis; in the middle of dry period, this metabolism is not active and LDL is accumulated, thus
270 determining a relative decline of HDL concentration. We also observed a progressive decline of
271 cholesterol until reaching the lower values at calving, as also previously reported [7, 12, 21-24].

272 Generally, it is accepted that cholesterol is representative of liver synthesis of VLDL and that a
273 decline in VLDL concentration is indicative of poor adaptation of the liver to fat mobilization, even
274 if currently, there is no reference limit for VLDL in dry high-producing dairy cows [9, 21, 25-26].

275 Through Optimal Cutpoint package, we defined a cut-off value of 89% of serum HDL at the
276 beginning of the dry period as a possible threshold to identify cows, which are more prone to
277 experience postpartum reproductive diseases. Cows belonging to High group presented better
278 calving to first AI and calving to conception intervals when compared to Low group ($P < 0.05$).

279 Cows with a greater concentrations of HDL had also lower incidence of retention of fetal
280 membranes and/or metritis, with an odds ratio of 0.6875; however, differences were not significant
281 probably due to both the reduced number of animal per group and the overall low incidence of
282 postpartum reproductive diseases. In the present study, in fact, the prevalence of fetal membrane
283 retention and/or metritis was 32%, which was lower than the rate of 36-50% reported by Sheldon et
284 al. [15]. At parturition, bilirubin and NEFA/CHOL ratio were higher in Low group, as indicative of
285 increased liver metabolic load, even if values remained within reference limits [9, 22].

286 All cows resumed ovarian activity within the first 3 weeks postpartum. Only one cow per group
287 showed a prolonged period of milk P4 due to persistent ovarian luteal cyst. Emergence and growth
288 of first follicular wave after calving is regulated by nutritional status and energy availability which
289 also influence hypothalamic release of gonadotropin releasing hormone [27]. Since resumption of
290 ovarian activity was similar between groups, we hypothesized that serum lipoproteins did not
291 influence the hypothalamic-pituitary-gonadal axis.

292 The reduction in reproductive efficiency observed in the Low group could be related to oocyte-
293 embryo quality. Considering that dairy cows have to cope with generalized inflammatory condition
294 4-5 weeks before calving [23] and that HDL have anti-inflammatory activity [29], we hypothesized
295 that their high concentration in dry cows could protect primordial ovarian follicles against
296 metabolic stress, enhancing oocyte quality, and increasing fertility rate, as observed in High group.

297 In relation to postpartum follicular development and their steroidogenic capacity, Golini et al. [28]
298 showed that there is a down-regulation of genes involved in cell proliferation and steroidogenesis of
299 granulosa cells harvested from preovulatory follicles collected from early lactating cows, until 90
300 days after calving. Moreover, results reported by Wu et al. [30] show that ovarian cells
301 preferentially use HDL-bound cholesterol for steroidogenesis, suggesting their fundamental role in
302 follicle development and maturation. Even if those results are referred to active and growing
303 follicles, it could be speculated that decreased HDL availability in conjunction with lower follicle's
304 steroidogenic capacity and pro-oxidant environment could negatively affect cholesterol absorption
305 storage and metabolism in ovarian follicles during the dry period.

306 To our knowledge, only other two studies reported data about concentration of lipoproteins in dairy
307 cows during dry period [7, 29]. The lack of uniformity in separation and detection methods
308 (electrophoresis *versus* centrifugation) or the investigation of lipoprotein concentration during
309 lactation makes comparison with some studies difficult [25, 31]. Bernabucci et al. [7] used the same
310 analytical procedure as in the present study, but they considered a different period of samples
311 collection (35 days before calving and 3 and 30 days after calving). They reported a lower

312 proportion of serum VLDL and higher HDL after calving, while LDL did not significantly change
313 during the transition period. This difference could be due to the matrix used for the test, since
314 Bernabucci et al. [7] worked on plasma while Sebia Lipo Lp (a) electrophoresis method is
315 standardized for serum. Newman et al. [29] investigated the effect of several diets with different
316 energy density on serum lipoprotein composition, from week 6 before expected calving until week
317 6 of lactation. The study [29] was aimed to verify if a provision of adequate total metabolizable
318 energy would better allow the cow to maintain total cholesterol and an HDL-rich profile throughout
319 the transition period while concurrently limiting the rise in NEFA. However, they did not find
320 influence of energy density in the diet on lipoprotein. This suggests that lipoprotein metabolism is
321 dependent on liver and peripheral secretion and absorption rather than on feeding management.
322 They reported a decline in HDL from week 6 until week 1 before calving and a subsequent increase.
323 In their study, mean concentrations of HDL at the beginning and at the end of the dry period were
324 similar to those found in the present work, for instance 80-85%. Since in their study VLDL was not
325 detectable, LDL relative concentrations were complementary to that of HDL, thus ranging from
326 15% to 25%. This difference could be due to the methods used, such as the duration of
327 electrophoresis and migration, the voltage applied or the staining technique, which may affect the
328 degree of lipoprotein separation and identification.

329 All other analysis carried out to verify animal's overall health status and to exclude alterations in
330 lipoprotein metabolism due to subclinical pathological conditions confirmed the good health
331 condition of the cows. Only during the last month before calving some modifications, indicated a
332 certain degree of lipid mobilization and reactivation of liver function, typical of the dairy cows
333 during the transition period. In particular, BCS was evaluated to assess the overall energy balance
334 and it was fairly constant, while mean NEFA were within reference limit of 0.4 mmol/L, as reported
335 in pending calving cows [4, 9, 32]. Only at calving BCS significantly decreased, as generally
336 described in transition dairy cows [4, 24, 33-34]. Correspondingly, NEFA increased, but remained
337 lower than the threshold value of 0.7 mmol/L established for the postpartum cows [32]. Triglyceride

338 decreased significantly from dry-off to calving, as previously observed by Weber et al. [24]. Herein,
339 cows showed normal glucose concentrations throughout the dry period, similarly to what reported
340 by Weber et al. [24]; however, a significant decrease was found at parturition. We observed BHB
341 was constantly greater than 1.00 mmol/L but below the threshold value of 1.20 - 1.40 mmol/L. Feed
342 analysis excluded the presence of ketogenetic compounds, as only butyric acid was fairly present in
343 both silage and total mixed ration. Subclinical ketosis diagnosis has been defined by BHB
344 concentrations of 1200 or 1400 $\mu\text{mol/L}$ [35-37]. Although BHB above 0.96 mmol/L is associated
345 with increased risk of postpartum disease [38-39], prepartum BHB concentrations are not predictive
346 of disease [4].

347 Concerning the other analytes, mean concentrations of GGT, ALT and TP were in agreement with
348 results reported by Brscic et al. [12], without significant difference between groups. The increase of
349 mean AST levels toward calving in all our experimental groups could be representative of increased
350 muscle labour. As ALP, Ca and P concentrations decreased at parturition but remained within
351 reference limits; we excluded subclinical hypocalcemia or other mineral imbalance.

352

353 **5 Conclusions**

354 The growing knowledge on the role of lipid metabolism in the adaptation of dairy cows to the
355 transition lead to an increased interest in lipoproteins. In our study, animal with HDL levels greater
356 than 89% at dry-off showed better reproductive performance when compared to the other
357 experimental group. We suggest that HDL due to their role of main carrier of cholesterol in ovarian
358 steroidogenic cells and to anti-inflammatory and antioxidant activities can exert some influences
359 on liver adaptation to the transition and on the fertility in cows. However, future studies are needed
360 to improve our knowledge on lipoprotein metabolism and to validate the quantification of HDL in
361 dry and transition phases using larger groups of animals and under different management or
362 nutritional conditions.

363

364 **6 References**

- 365 [1] Bobe G, Young JW, Beitz DC. Invited review: pathology, ethiology, prevention, and treatment of
366 fatty liver in dairy cows. *J Dairy Sci* 2004;87:3105–24.
- 367 [2] Bauchart D. Lipid absorption and transport in ruminants. *J Dairy Sci* 1993;76:3864–81.
- 368 [3] Chapinal N, LeBlanc SJ, Carson ME, Leslie KE, Godden S, Capel M et al. Herd-level
369 association of serum metabolites in the transition period with disease, milk production and early
370 lactation reproductive performance. *J Dairy Sci* 2012;95:5676–82.
- 371 [4] Ospina PA, McArt JA, Overton TR, Stokol T, Nydam DV. Using nonesterified fatty acids and β -
372 hydroxybutyrate concentrations during the transition period for herd-level monitoring of increased
373 risk of disease and decreased reproductive and milking performance. *Vet Clin North Am Food Anim*
374 *Pract* 2013;29:387–412.
- 375 [5] Sordillo LM, Raphael W. Significance of metabolic stress, lipid mobilization, and inflammation
376 on transition cow disorders. *Vet Clin North Am Food Anim Pract* 2013;29:267–78.
- 377 [6] Katoh N. Relevance of apolipoproteins in the development of fatty liver and fatty liver-related
378 peripartum diseases in dairy cows. *J Vet Med Sci* 2002;64: 293–307.
- 379 [7] Bernabucci U, Ronchi B, Basiricò L, Pirazzi D, Rueca F, Lacetera N et al. Abundance of mRNA
380 of apolipoprotein b100, apolipoprotein e, and microsomal triglyceride transfer protein in liver from
381 periparturient dairy cows. *J Dairy Sci* 2004;87:2881–8.
- 382 [8] Roche JR, Kay JK, Friggens NC, Loor JJ, Berry DP. Assessing and managing body condition
383 score for the prevention of metabolic disease in dairy cows. *Vet Clin North Am Food Anim Pract*
384 2013;29:387–412.
- 385 [9] Van Saun RJ. Metabolic profiling of transition dairy cows: can we predict impending problems?
386 *Proceedings Danske Kvægdyrlægers Arsmøde (Danish Bovine Practitioner Seminar),*
387 *Middlefart, Denmark, January 2007;24–5.*
- 388 [10] Wathes DC, Clempson AM, Pollot GE. Association between lipid metabolism and fertility in
389 the dairy cow. *Reproduction, Fertility and Development*, 2013;25, 48-61.

- 390 [11] Dervishi E, Zhang G, Hailemariam D, Goldansaz SA, Deng Q, Dunn SM, Ametaj BN.
391 Alterations in innate immunity reactants and carbohydrate and lipid metabolism precede occurrence
392 of metritis in transition dairy cows. *Res Vet Sci.* 2016 Feb;104:30-9.
- 393 [12] Brscic M, Cozzi G, Lora I, Stefani AL, Contiero B, Ravarotto L et al. Short communication:
394 Reference limits for blood analytes in Holstein late-pregnant heifers and dry cows: Effects of parity,
395 days relative to calving, and season. *J Dairy Sci* 2015;98:7886–92.
- 396 [13] Edmonson AJ, Lean IJ, Weaver LD, Farver T, Webster G. A body condition scoring chart for
397 holstein dairy cows. *J Dairy Sci* 1989;72:68–78.
- 398 [14] Vannucchi CI, Rodrigues JA, Silva LCG, Lúcio CF, Veiga GAI. Effect of dystocia and
399 treatment with oxytocin on neonatal calf vitality and acid-base, electrolyte and haematological
400 status. *The Veterinary Journal* 2015;203:228-232.
- 401 [15] Sheldon IM, Cronin J, Goetze L, Donofrio G, Schuberth HJ. Defining postpartum uterine
402 disease and the mechanisms of infection and immunity in the female reproductive tract in cattle
403 *Biol Reprod* 2009;81:1025–32.
- 404 [16] Comin A, Renaville B, Marchini E, Maiero S, Cairoli F, Prandi A. Technical note: direct
405 enzyme immunoassay of progesterone in bovine milk whey *J Dairy Sci* 2005;88:4239–42.
- 406 [17] SPSS Incorporated: Statistic Package for Social Science (SPSS) Version 16 Chicago, IL USA:
407 SPSS Inc; 2007.
- 408 [18] RStudio, R software v3.2.4, <https://www.rstudio.com/> [accessed 24.03.16].
- 409 [19] Lopez-Raton M, Rodriguez-Alvarez MX Package OptimalCutpoints [https://cran.r-](https://cran.r-project.org/web/packages/OptimalCutpoints/OptimalCutpoints.pdf)
410 [project.org/web/packages/OptimalCutpoints/OptimalCutpoints.pdf](https://cran.r-project.org/web/packages/OptimalCutpoints/OptimalCutpoints.pdf). [accessed 24.03.16].
- 411 [20] Smith BP. Large animal internal medicine. 5th ed. Mosby Elsevier. 2015.
- 412 [21] Stengärde L, Tråvén M, Emanuelson U, Holtenius K, Hultgren J, Niskanen R. Metabolic
413 profiles in five high-producing Swedish dairy herds with a history of abomasal displacement and
414 ketosis. *Acta Veterinaria Scandinavica* 2008;50:31.
- 415 [22] Kalaitzakis E, Panousis N, Roubies N, Giadinis N, Kaldrymidou E, Georgiadis M et al.

- 416 Clinicopathological evaluation of downer dairy cows with fatty liver. *The Canadian Veterinary*
417 *Journal* 2010;51:615–22.
- 418 [23] Trevisi E, Amadori M, Cogrossi S, Razzuoli E, Bertoni G. Metabolic stress and inflammatory
419 response in high-yielding, periparturient dairy cows. *Res Vet Sci* 2012;93:695–704.
- 420 [24] Weber C, Hametner C, Tuchscherer A, Losand B, Kanitz E, Otten W et al. Variation in fat
421 mobilization during early lactation differently affects feed intake, body condition, and lipid and
422 glucose metabolism in high-yielding dairy cows. *J Dairy Sci* 2013;96:165–80.
- 423 [25] Sevinç M, Basoglu A, Guzelbektas H. Lipid and lipoprotein levels in dairy cows with fatty
424 liver *Tusk J Vet Anim Sci* 2003;27:295–9.
- 425 [26] Bertoni G, Trevisi E. Use of the liver activity index and other metabolic variables in the
426 assessment of the metabolic health in dairy herds. *Vet Clin North Am Food Anim Pract*
427 2013;29:413–31.
- 428 [27] Chagas LM, Bass JJ, Blache D, Burke CR, Kay JK, Lindsay DR et al. Invited review: New
429 perspectives on the roles of nutrition and metabolic priorities in the subfertility of high-producing
430 dairy cows. *J Dairy Sci* 2007;90:4022–32.
- 431 [28] Golini VE, Stradaioli G, Sirard MA. Transcriptome analysis of bovine granulosa cells of
432 preovulatory follicles harvested 30, 60, 90, and 120 days postpartum. *Theriogenology* 2014;82:580–
433 91.
- 434 [29] Newman A, Mann S, Nydam DV, Overton TR, Behling-Kelly E. Impact of dietary plane of
435 energy during the dry period on lipoprotein parameters in the transition period in dairy cattle. *J*
436 *Anim Physiol Anim Nutr (Berl)* 2016;100:118–26.
- 437 [30] Wu Q, Sucheta S, Azhar S, Menon KMJ. Lipoprotein enhancement of ovarian theca-interstitial
438 cell steroidogenesis: relative contribution of scavenger receptor class B (Type I) and Adenosine 5'-
439 Triphosphate-Binding Cassette (Type A1) transporter in High-density lipoprotein-cholesterol
440 transport and androgen synthesis. *Endocrinology* 2003;144(6):2437-2446.

- 442 [31] Ileri-Büyükoğlu T, Sahinduran S, Sezer K, Güldür T. Evaluation of changes in serum
443 lipoprotein and apolipoprotein patterns in cows with ketosis. *Am J Vet Res* 2009;70:563–70.
- 444 [32] Mulligan FJ, O’Grady L, Rice DA, Doherty ML. A herd health approach to dairy cow nutrition
445 and production diseases of the transition cow. *Animal Reproduction Science* 2006;96:331–53.
- 446 [33] Ingvarsten, KL and Andersen JB. Integration of metabolism and intake regulation: A review
447 focusing on periparturient animals. *J Dairy Sci* 2000;83:1573–97.
- 448 [34] Drackley JK, Overton TR and Douglas GN. Adaptations of glucose and long-chain fatty acid
449 metabolism in liver of dairy cows during the periparturient period. *J Dairy Sci* 2001;84:E100–E112.
- 450 [35] Duffield TF, Lissemore KD, McBride BW, Leslie KE. Impact of hyperketonemia in early
451 lactation dairy cows on health and production. *J Dairy Sci* 2009;92:571–80.
- 452 [36] LeBlanc S. Monitoring metabolic health of dairy cattle in the transition period. *J Reprod Dev*
453 2010;56:S29–S35.
- 454 [37] Chapinal N, Carson ME, LeBlanc SJ, Leslie KE, Godden S, Capel M et al. The association of
455 serum metabolites in the transition period with milk production and early-lactation reproductive
456 performance. *J Dairy Sci* 2012;95:1301–09.
- 457 [38] Roussel JD, Seybt SH, Toups G. Metabolic profile testing for Jersey cows in Louisiana:
458 reference values. *Am J Vet Res* 1982;43:1075–7.
- 459 [39] Eicher R. Metabolic profile testing in dairy herds: wrong answer or wrong question? *Acta*
460 *Veterinaria Scandinavica* 2003;44(Suppl 1):28.
- 461

462 Table 1. Relative proportion of lipoproteins in the 28 experimental dairy cows.

	%	Time	Mean \pm SEM	P
High-density lipoproteins		T1	87.52 \pm 0.88 ^a	P < 0.001
		T2	82.41 \pm 1.21 ^b	
		T3	87.39 \pm 1.07 ^a	
Low-density lipoproteins		T1	9.65 \pm 0.71 ^a	P < 0.01
		T2	14.32 \pm 1.18 ^b	
		T3	9.92 \pm 1.10 ^a	
Very low-density lipoproteins		T1	2.83 \pm 0.29	NS
		T2	3.27 \pm 0.29	
		T3	2.69 \pm 1.43	

463 T1 = 60 days before expected calving; T2 = 30 days before expected calving; T3 = calving; P =

464 Kruskal-Wallis test significance; NS = not significant. ^{a,b} = Values with different superscripts are

465 significantly different.

466

467

468

469

470 Table 2. Mean (\pm SEM) reproductive outcomes in dairy cows of the Low and High HDL groups.

Cut-off HDL > 89.0 %						
	n	HDL (%)	C - 1AI (days)	C-C (days)	AI (n°)	OR
Low	18	85.07 \pm 0.88	94.50 \pm 12.32	199.18 \pm 24.73	3.36 \pm 0.58	
High	10	91.94 \pm 0.67	64.00 \pm 3.95	129.86 \pm 24.42	3.67 \pm 0.78	0.6875
P		P < 0.001	P < 0.05	P < 0.05	NS	NS

471 n = number of animals; C - 1AI = calving to first artificial insemination interval; C - C = calving to
 472 conception interval; n° AI = number of artificial insemination per pregnancy; OR = odds ratio for
 473 postpartum uterine diseases; P = Kruskal-Wallis test significance; NS = not significant.

474

475

476

477

478 Table 3. Mean (\pm SEM) values of energy metabolism analytes in the 28 experimental dairy cows 60
 479 (T1) and 30 (T2) days before expected calving, and at calving (T3).

	Time	Values	P
BHB (mmol/L)	T1	1.16 \pm 0.08	NS
	T2	1.29 \pm 0.85	
	T3	1.03 \pm 0.53	
NEFA (mmol/L)	T1	0.18 \pm 0.02 ^a	P < 0.001
	T2	0.24 \pm 0.04 ^a	
	T3	0.58 \pm 0.08 ^b	
Glucose (mg/dL)	T1	66.10 \pm 1.85 ^a	P < 0.001
	T2	62.21 \pm 1.73 ^a	
	T3	55.75 \pm 2.24 ^b	
Triglycerides (mg/dL)	T1	16.32 \pm 1.51 ^a	P < 0.001
	T2	18.39 \pm 1.13 ^a	
	T3	11.41 \pm 1.20 ^b	

480 P = Kruskal-Wallis test significance; NS = not significant. ^{a,b} = Values with different superscripts are
 481 significantly different.

482

483

484

485

486 Table 4. Mean (\pm SEM) bilirubin and NEFA/CHOL ratio in the experimental dairy cows of Low and
 487 High HDL groups at calving.

Cut-Off HDL > 89%				
	n	T-Bil	Ind-Bil	NEFA/CHOL
Low	18	0.46 \pm 0.09	0.27 \pm 0.07	0.30 \pm 0.06
High	10	0.23 \pm 0.08	0.12 \pm 0.05	0.14 \pm 0.03
P		P < 0.05	P < 0.05	P < 0.05

488 n = number of animals; T-Bil = total bilirubin; Ind-Bil = indirect bilirubin; NEFA/CHOL = NEFA:
 489 cholesterol ratio; P = Kruskal-Wallis test significance; NS = not significant.

490

491

492

493 Table 5. Mean (\pm SEM) hepatic function analytes values in the 28 experimental dairy cows 60 (T1)
 494 and 30 (T2) days before expected calving, and at calving (T3).

	Time	Values	P
Urea (mg/dL)	T1	29.39 \pm 1.54	NS
	T2	30.54 \pm 1.89	
	T3	30.89 \pm 4.03	
GGT (U/L)	T1	21.00 \pm 2.48 ^a	P < 0.05
	T2	16.71 \pm 1.68 ^b	
	T3	15.22 \pm 1.31 ^b	
LDH (U/L)	T1	883.21 \pm 87.59	NS
	T2	1113.04 \pm 36.80	
	T3	1174.71 \pm 60.14	
AST (U/L)	T1	82.36 \pm 3.95 ^a	P < 0.001
	T2	77.00 \pm 3.93 ^a	
	T3	106.57 \pm 6.98 ^b	
ALT (U/L)	T1	29.21 \pm 0.97 ^a	P < 0.001
	T2	24.54 \pm 1.11 ^b	
	T3	20.21 \pm 0.96 ^b	
Cholesterol (mg/dL)	T1	192.79 \pm 5.30 ^a	P < 0.001
	T2	148.54 \pm 5.72 ^b	
	T3	112.57 \pm 6.97 ^c	

495 P = Kruskal-Wallis test significance; NS = not significant. ^{a,b,c} = Values with different superscripts
 496 are significantly different.

497

498 Table 6. Mean (\pm SEM) protein metabolism analytes values in the 28 experimental dairy cows 60

499 (T1) and 30 (T2) days before expected calving, and at calving (T3).

	Time	Values	P
Albumin (g/L)	T1	4.10 ± 0.08	NS
	T2	4.04 ± 0.08	
	T3	4.05 ± 0.08	
Total Proteins (g/L)	T1	7.56 ± 0.16 ^a	P < 0.001
	T2	7.11 ± 0.13 ^a	
	T3	6.43 ± 0.11 ^b	
CK (U/L)	T1	246.96 ± 59.40	NS
	T2	340.34 ± 144.31	
	T3	209.76 ± 28.52	

500 P = Kruskal-Wallis test significance; NS = not significant. ^{a,b} = Values with different superscripts are
 501 significantly different.

502

503

504

505 Table 7. Mean (\pm SEM) mineral metabolism and non-specific analytes values in the 28 experimental
 506 dairy cows 60 (T1) and 30 (T2) days before expected calving, and at calving (T3).

	Time	Values	P
Creatinine (mg/dL)	T1	1.16 \pm 0.03	
	T2	1.22 \pm 0.05	NS
	T3	1.38 \pm 0.18	
ALP (U/L)	T1	84.57 \pm 4.68 ^a	
	T2	76.07 \pm 5.72 ^a	P < 0.001
	T3	64.46 \pm 5.05 ^b	
Calcium (mg/dL)	T1	9.65 \pm 0.12 ^a	
	T2	9.34 \pm 0.21 ^a	P < 0.05
	T3	8.80 \pm 0.29 ^b	
Magnesium (mg/dL)	T1	2.51 \pm 0.11	
	T2	2.31 \pm 0.08	NS
	T3	2.39 \pm 0.62	
Phosphorous (mg/dL)	T1	6.55 \pm 0.28 ^a	
	T2	6.40 \pm 0.17 ^a	P < 0.001
	T3	4.83 \pm 0.28 ^b	

507 P = Kruskal-Wallis test significance; NS = not significant. ^{a,b} = Values with different superscripts are
 508 significantly different.

509

510

511