



UNIVERSITÀ
DEGLI STUDI
DI UDINE

Università degli studi di Udine

IO and rank-into-rank axioms

Original

Availability:

This version is available <http://hdl.handle.net/11390/1115630> since 2021-03-16T17:09:15Z

Publisher:

Published

DOI:10.1007/s40574-017-0136-y

Terms of use:

The institutional repository of the University of Udine (<http://air.uniud.it>) is provided by ARIC services. The aim is to enable open access to all the world.

Publisher copyright

(Article begins on next page)

I0 and rank-into-rank axioms

Vincenzo Dimonte*

July 7, 2017

Abstract

This is a survey about I0 and rank-into-rank axioms, with some previously unpublished proofs.

Keywords: Large cardinals; Axiom I0; rank-into-rank axioms; elementary embeddings; relative constructibility; embedding lifting; singular cardinals combinatorics.

2010 Mathematics Subject Classifications: 03E55 (03E05 03E35 03E45)

1 Introduction

In 1984 Woodin showed [...] that $\text{AD}^{L(\mathbb{R})}$ follows from a proposition stronger than I1 and straining the limits of consistency [...]. Moreover, Woodin's investigations of I0 was to produce a detailed and coherent structural theory for $L(V_{\delta+1})$.

With attention shifted upward to the strongest large cardinal hypotheses, Woodin boldly formulated I0 just at the edge of the Kunen inconsistency and amplified Martin's embeddings proof to establish the following in early 1984:

Theorem 1.1 (Woodin). *Assume I0. Then $\text{AD}^{L(\mathbb{R})}$.*

This sparse lines in Kanamori's *The Higher Infinite* [?] were probably for years the only published trace of the mysterious I0. This fact is even more surprising as [?] was written in 1994, so ten years after the striking result and even more from its first formulation. What happened? Why all this mystery?

I0 was just one of the many large cardinals that were created in a short span of time: strong cardinals, supercompact cardinals, huge cardinals, Woodin cardinals, rank-into-rank axioms (its smaller siblings), ... were introduced in a rapid fire, shifting the paradigms of set theory. But I0 could not keep up the pace with all of them: its main victory was short-termed, as $\text{AD}^{L(\mathbb{R})}$ was proven to be of much lower consistency strength, there were no other examples of important propositions implied by I0, and it was definitely too large for inner model theory (in fact, the largest cardinal axiom of all times). So, before even a publication, it quietly disappeared, and with it the proof of Theorem ??.

Still, under the surface there was some activity: the bound between I0 and $\text{AD}^{L(\mathbb{R})}$ ran deeper than just Theorem ??, it involved the whole structure of

*Università degli Studi di Udine, via delle Scienze, 206 33100 Udine (UD) *E-mail address:* vincenzo.dimonte@gmail.com

a model of the form $L(V_{\lambda+1})$, that under I_0 had properties very similar than those in $L(\mathbb{R})$ under AD . This peculiarity was explored by Woodin, and the first output was Kafkoulis' lecture notes *An AD-like axiom* [?] (at that time I_0 was called Axiom A), a transcription of Woodin's lectures on the argument, that started to be copied and distributed among mathematicians at the beginning of the Nineties. This underground distribution was, of course, very limited (Internet was at its infancy).

In the meantime the rank-into-rank axioms were enjoying a period of popularity, thanks to their connections with algebra, and Laver, in 2001, tried to push some of his results on rank-into-rank (in particular reflection) towards I_0 . In [?] we can find therefore the first officially published fragments of the theory of I_0 , taken from Kafkoulis notes, and in 2004 Kafkoulis himself published a paper on Coding Lemmata that built on such notes [?]. After that, still silence for many years.

The rhythm of publications changed abruptly in 2011. The reason is Woodin's *Suitable extender models* magnum opus. The aim of such paper (published in two parts) was to push inner model theory up to all large cardinals, so it was necessary to know as much as possible as the top of the hierarchy. Therefore in the sprawling 561 pages there are whole sections dedicated to I_0 , the structure of $L(V_{\lambda+1})$ under it (in a much more developed way respect to Kafkoulis notes) and even stronger axioms. There was a new exciting realm to explore, finally mature, and with connections to other fields of set theory. The people that were Woodin's students in that period (the author included) continued the work, expanding the research on I_0 in different directions.

Such a rapid expansion, after a long silence, carries the risk of excessive fragmentation. It is then time to take stock of the situation and pose solid bases on such a venture, not based on underground notes, folklore, and oral tradition. The objective of this paper is therefore to collect all the results on rank-into-rank axioms and I_0 , with a stress on the latest one, both from the I_0 folklore and from the published papers, so that anybody can have all the tools for reading the latest papers or doing research on her own.

Section ?? glances briefly at some important large cardinals, up until the first inconsistency in the large cardinal hierarchy, i.e., the (non-)existence of $j : V \prec V$.

Starting from this, Section ?? is about rank-into-rank axioms, i.e., axioms of the form $j : V_\lambda \prec V_\lambda$ (I3) and $j : V_{\lambda+1} \prec V_{\lambda+1}$ (I1), with a particular attention to the application operator.

In Section ?? I_0 is introduced, and with it all the structural properties of $L(V_{\lambda+1})$ that we can have for free. Much of this part is taken from [?].

Section ?? is a detailed analysis of many (all?) axioms from I3 to I_0 , with all the proofs of strong implications among them. The techniques of inverse limits and square roots are introduced, that are now a focal centre of the investigations around I_0 (see [?]).

Section ?? uses again [?] as main source, introducing all the results about similarities between I_0 and $AD^{L(\mathbb{R})}$, including the newest developments: measurability of λ^+ , Coding Lemma, large cardinal properties of Θ , perfect set property, an analogous of weakly homogenous Suslin-ness.

In Section ?? all these axioms are treated in relation to forcing, with many results of indestructibility and independence results.

In Section ?? it is noticed how such independence result can actually destroy the similarities between I_0 and $AD^{L(\mathbb{R})}$: the examples treated are the Ultrafilter Axiom, Wadge Lemma, a partition property and Turing Determinacy.

Section ?? describes highly speculatively some possible future lines of inquiry.

On behalf of all authors, the corresponding author states that there is no conflict of interest.

2 Kunen's Theorem

To reach the top of the large cardinal hierarchy, it is worth to have a glance at the lower levels before. In the bottom, large cardinal axioms are very combinatoric in nature: an inaccessible cardinal is a cardinal that is regular and closed under exponentiation, a Ramsey cardinal is a cardinal that satisfies a particular partition property, a Rowbottom cardinal is a cardinal that satisfies another partition property, and so on¹. The pivotal moment was when Scott proved that a combinatoric property on a cardinal is equivalent to a global property of the universe:

Definition 2.1 (Ulam, 1930). *A cardinal κ is measurable iff there exists a κ -complete ultrafilter on κ .*

Definition 2.2. *Let M, N be sets or classes. Then $j : M \rightarrow N$ is an elementary embedding and we write $j : M \prec N$ iff it is injective and for any formula φ and $a_1, \dots, a_n \in M$, $M \models \varphi(a_1, \dots, a_n)$ iff $N \models \varphi(j(a_1), \dots, j(a_n))$.*

In the rest of the paper, if we write $j : M \prec N$ we mean that j is not the identity. If j is the identity, we just write $M \prec N$.

So an elementary embedding is a homeomorphism of the logical structure. In particular, M and N must satisfy the same sentences, so for example $M \models ZF$ iff $N \models ZF$. It is easy to see that for all ordinals α , $j(\alpha) \geq \alpha$. If $M \models AC$ or $N \subseteq M$, then it is possible to prove that there must exist an ordinal α such that $j(\alpha) > \alpha$.

Definition 2.3. *Let M, N be sets or classes such that $M \models AC$ or $N \subseteq M$. Then the critical point of j , $\text{crt}(j)$, is the least α such that $j(\alpha) > \alpha$.*

Theorem 2.4 (Scott, 1961). *A cardinal κ is measurable iff there exist an inner model M and a $j : V \prec M$ such that $\text{crt}(j) = \kappa$.*

This was the start of an intense research on a new breed of large cardinal hypotheses, defined with more requirements for M . For example, one can ask how much of the original universe is in M :

Definition 2.5 (Gaifman, 1974). *A cardinal κ is γ -strong iff there exists $j : V \prec M$, $\text{crt}(j) = \kappa$, $\gamma < j(\kappa)$ and $V_{\kappa+\gamma} \subseteq M$.*

Or one can ask how close M is. The starting point is to notice that if $j : V \prec M$ witnesses that κ is measurable, then $M^\kappa \subseteq M$.

¹Admittedly, this is a very rough classification, as for example indescribable cardinals are difficult to insert in this narrative

Definition 2.6 (Solovay, Reinhardt). *A cardinal κ is γ -supercompact iff there exists $j : V \prec M$, $\text{crt}(j) = \kappa$, $\gamma < j(\kappa)$ and $M^\gamma \subseteq M$.*

It is immediate to see that if κ is γ -supercompact then it is γ -strong. All these definitions fix $j(\kappa)$ as the limit of the closeness of M . Overcoming this means to define stronger large cardinals:

Definition 2.7 (Gaifman, 1974). *A cardinal κ is superstrong iff there exists $j : V \prec M$, $\text{crt}(j) = \kappa$ and $V_{j(\kappa)} \subseteq M$.*

Definition 2.8 (Kunen, 1972). *A cardinal κ is huge iff there exists $j : V \prec M$, $\text{crt}(j) = \kappa$ and $M^{j(\kappa)} \subseteq M$.*

We can go on, defining stronger and stronger large cardinals:

Definition 2.9. *Let M, N be sets or classes such that $M \models \text{AC}$ or $N \subseteq M$. Then the critical sequence of $j : M \prec N$, $\langle \kappa_n : n \in \omega \rangle$, is defined as:*

- $\kappa_0 = \text{crt}(j)$;
- $\kappa_{n+1} = j(\kappa_n)$.

Definition 2.10. *A cardinal κ is n -superstrong iff there exists $j : V \prec M$, $\text{crt}(j) = \kappa$ and $V_{j(\kappa_n)} \subseteq M$.*

Definition 2.11. *A cardinal κ is n -huge iff there exists $j : V \prec M$, $\text{crt}(j) = \kappa$ and $M^{j(\kappa_n)} \subseteq M$.*

This ascent eventually must stop. In 1969 Reinhardt, in his thesis, asked whether it is possible to have directly $j : V \prec V$. Kunen, few months later, proved that it is inconsistent. Such result must be approached cautiously: if we work in ZFC, it is not surprising:

Theorem 2.12 (Naive Kunen's Theorem). (ZFC) *There is no $j : V \prec V$.*

Proof by Suzuki, [?]. As we are working in ZFC, it means that j must be a class that is definable with some first order formula. Let φ be a formula. Let a be a parameter so that $\varphi(a)$ defines an elementary embedding j from V to itself with least critical point κ among all the possible choices of the parameter. Then κ is definable in V , without parameters. So $j(\kappa)$ satisfies the same definition, and it is the least critical point of embeddings definable with φ . But $j(\kappa) > \kappa$, contradiction. \square

Kunen's Theorem is stronger than this, as it involves also classes that are not definable: in the original paper, [?], the proof is in MK, Morse-Kelley theory. It is possible to weaken the theory to NBG, or to ZFC(j), i.e., ZFC with j as a predicate that can be inserted in replacement formulas. We leave the interested reader to [?], where all such problems are solved in great detail, and we state Kunen's Theorem in vague terms, with the only provisos that j is not necessarily definable in V , replacement for j holds and AC holds in V .

Theorem 2.13 (Kunen's Theorem, [?]). (AC) *There is no $j : V \prec V$.*

There are many proofs of this result. Instead of the original proof by Kunen, we show the proof by Woodin ([?], Theorem 23.12, second proof), as it gives information that will be useful in Section ???. It builds on the following classical result by Solovay:

Theorem 2.14 (Solovay). *Let $\kappa > \omega$ be a regular cardinal, and $X \subseteq \kappa$ stationary. Then X can be partitioned in κ disjoint stationary sets, i.e., there is $\langle X_\eta : \eta < \kappa \rangle$ such that $X = \bigcup_{\eta < \kappa} X_\eta$ and all the X_η 's are pairwise disjoint.*

It will be useful this technical remark, that will be used constantly in the paper without comment:

Remark 2.15. *Let $j : M \prec N$, $M \models \text{AC}$ or $N \subseteq M$. Then for any $\gamma < \text{crt}(j)$ and any sequence $\langle a_\alpha : \alpha < \gamma \rangle \in M$, $j(\langle a_\alpha : \alpha < \gamma \rangle) = \langle j(a_\alpha) : \alpha < \gamma \rangle$. So if $A \in M$ is well-orderable (for example $M \models \text{AC}$) and $|A| < \text{crt}(j)$, then $j(A) = \{j(a) : a \in A\}$.*

Proof. Consider the sequence as a function $f \in M$ with domain γ . As $\gamma < \text{crt}(j)$, $j(f)(\alpha) = j(f)(j(\alpha)) = j(f(\alpha))$. \square

Proof of ?? by Woodin, [?]. Let λ be the supremum of the critical sequence of j . Then λ must be a fixed point for j , because

$$j(\lambda) = j(\sup\langle \kappa_n : n \in \omega \rangle) = \sup j(\langle \kappa_n : n \in \omega \rangle) = \sup\langle j(\kappa_n) : n \in \omega \rangle = \lambda.$$

Let $S_\omega^{\lambda^+} = \{\alpha < \lambda^+ : \text{cof}(\alpha) = \omega\}$. Note that $S_\omega^{\lambda^+}$ is stationary: if C is a club in λ^+ , then any ω -limit of elements of C must be in C , as λ^+ is regular, and has cofinality ω^2 . We prove that $S_\omega^{\lambda^+}$ cannot be partitioned in $\text{crt}(j)$ disjoint stationary sets, in contradiction with Solovay's result.

Suppose that $\langle S_\alpha : \alpha < \text{crt}(j) \rangle$ is a partition in stationary subsets of $S_\omega^{\lambda^+}$. Then

$$j(\langle S_\alpha : \alpha < \text{crt}(j) \rangle) = \langle T_\beta : \beta < j(\text{crt}(j)) \rangle$$

must be a partition in stationary subsets of $S_\omega^{\lambda^+}$ (because $j(\lambda^+) = j(\lambda)^+ = \lambda^+$) of length $j(\text{crt}(j))$. Let $C = \{\alpha < \lambda^+ : j(\alpha) = \alpha\}$. Then C is an ω -club in λ^+ : If $\langle \alpha_n : n \in \omega \rangle$ is a sequence of ordinals in C , then

$$\begin{aligned} j(\sup\langle \alpha_n : n \in \omega \rangle) &= \sup j(\langle \alpha_n : n \in \omega \rangle) = \\ &= \sup\langle j(\alpha_n) : n \in \omega \rangle = \sup\langle \alpha_n : n \in \omega \rangle. \end{aligned}$$

To see that it is unbounded, let $\alpha < \lambda^+$, and consider the sequence $\alpha_0 = \alpha$, $\alpha_{n+1} = j(\alpha_n)$. Then $\sup\langle \alpha_n : n \in \omega \rangle \in C$.

As $T_{\text{crt}(j)}$ is stationary, it must intersect the closure of C in some η . But as $\eta \in T_{\text{crt}(j)}$, it has cofinality ω , so it must be in C , therefore $j(\eta) = \eta$. Since $\langle S_\alpha : \alpha < \text{crt}(j) \rangle$ was a partition, there must be some α such that $\eta \in S_\alpha$. But then $\eta = j(\eta) \in j(S_\alpha) = T_{j(\alpha)}$. But $\text{crt}(j)$ cannot be in the image of j (it is not a fixed point, and below it they are all fixed points), so $T_{\text{crt}(j)} \neq T_{j(\alpha)}$, so they should have been disjoint, but they have η in common. Contradiction. \square

We can now analyze such proof, to understand what it really needs of the hypotheses, with the objective of defining new axioms for which the proof does not work, so that there is a hope that they are not inconsistent. The first remark is that Solovay's Theorem uses AC in an essential way, so the proof would not stand without AC.

²In fact, $S_\omega^{\lambda^+}$ is ω -stationary, that is, it intersects all the ω -clubs, i.e., the sets that are cofinal in λ^+ and closed under supremum of ω -sequences

Definition 2.16 (ZF, Reinhardt). *A cardinal κ is Reinhardt iff there exists $j : V \prec V$, with $\text{crt}(j) = \kappa$.*

It is not immediate to see, but a Reinhardt cardinal is consistently stronger than all the hypotheses we will introduce in this paper. It is still open whether a Reinhardt cardinal would imply a contradiction, and the study of Reinhardt and even larger cardinals is still going on.

Another approach would be to weaken the ground theory, for example not permitting replacement with j , so that the critical sequence is not in V and λ is not definable. This is the “Wholeness Axiom”, and it was studied by Paul Corazza ([?]).

A third way would be to consider something smaller than V (but large enough so that the eventual large cardinal would sit at the top of the hierarchy). This will be implemented in the next section.

3 Rank-into-rank embeddings

Analyzing the proof of ??, one realizes that the key object is $\langle S_\alpha : \alpha < \text{crt}(j) \rangle$. Now, if $\beta < \lambda^+$, there is a well-order of λ that has length β , and such well-order can be coded as a subset of λ . Therefore every ordinal less than λ^+ can be coded as an element of $V_{\lambda+1}$, and a subset of λ^+ can be coded as an element of $V_{\lambda+2}$. So the whole proof can be carried on for:

Corollary 3.1. *For any η , there is no $j : V_{\eta+2} \prec V_{\eta+2}$.*

Proof. Suppose that there is $j : V_{\eta+2} \prec V_{\eta+2}$. Let λ be the supremum of its critical sequence. Then $\lambda \leq \eta$, because λ is limit. Then $V_{\lambda+2} \subseteq V_{\eta+2}$, and the proof of ?? gives the contradiction. \square

Therefore the proof of ?? does not exclude the following cases:

I3 There exists $j : V_\lambda \prec V_\lambda$;


I1 There exists $j : V_{\lambda+1} \prec V_{\lambda+1}$.

There is a small ambiguity in the definition of I3. On one hand λ must be at least the supremum of the critical sequence (because $\text{ran}(j) \subseteq V_\lambda$), on the other hand λ must be less than the supremum of the critical sequence +2, because of the Corollary, but, as it is written, it can be the supremum of the critical sequence +1. In general we will assume, without loss of generality, that λ is the supremum of the critical sequence:

I3 There exists $j : V_\lambda \prec V_\lambda$, where λ is the supremum of its critical sequence.

Admittedly, not being excluded by a proof is not the most solid of foundations for new axioms. In fact, the “I” in their name stands for “Inconsistency”, as this was the first thought of the community when they were introduced. Yet, they survived all attempts of proving their inconsistency, and they are now in the center of a rich and complex theory, that has many interesting results, some of which will be shown in this paper.

Let $j : V_\lambda \prec V_\lambda$. The mental picture is the following:



so V_λ is divided in two: under a certain rank, all sets are fixed, above a certain rank, all sets are moved. There are ω “stripes”, and all the sets on one stripe are moved into the next. There is one stripe, then, that contains no image of j .

The members of the critical sequence are large cardinals:

Remark 3.2. *Let $\langle \kappa_n : n \in \omega \rangle$ be the critical sequence of $j : V_\lambda \prec V_\lambda$. Then all the κ_n ’s are measurable.*

Proof. Consider the ultrafilter $U_j = \{X \in \mathcal{P}(\kappa_0) : \kappa_1 \in j(X)\}$. As usual, it is a κ_0 -complete measure, so κ_0 is measurable. Then by elementarity $j(\kappa_0) = \kappa_1$ is measurable, and so on. \square

Therefore λ is strong limit: if $\alpha < \lambda$, then there is a κ_n such that $\alpha < \kappa_n$, and $2^\alpha < \kappa_n < \lambda$. So $|V_\lambda| = \lambda$, and we can code every pair of sets in V_λ (in fact every finite sequence of sets) with a single set of V_λ . Therefore we can code every λ -sequence of elements in $V_{\lambda+1}$ as a single element in $V_{\lambda+1}$ (coding the sequence $\langle a_\eta : \eta < \lambda \rangle$ with $\{(\eta, x) : x \in a_\eta\}$).

The “stripes” structure permits to extend j beyond its scope:

Definition 3.3. *Let $j : V_\lambda \prec V_\lambda$. Define $j^+ : V_{\lambda+1} \rightarrow V_{\lambda+1}$ as*

$$\forall A \subset V_\lambda \quad j^+(A) = \bigcup_{\beta < \lambda} j(A \cap V_\beta).$$

While it is not clear whether j^+ is an elementary embedding, every elementary embedding from $V_{\lambda+1}$ to itself is the ‘plus’ of its restriction to V_λ :

Lemma 3.4. *If $j : V_{\lambda+1} \prec V_{\lambda+1}$, then $(j \upharpoonright V_\lambda)^+ = j$. Thus every $j : V_{\lambda+1} \prec V_{\lambda+1}$ is defined by its behaviour on V_λ , i.e., for every $j, k : V_{\lambda+1} \prec V_{\lambda+1}$,*

$$j = k \quad \text{iff} \quad j \upharpoonright V_\lambda = k \upharpoonright V_\lambda.$$

Proof. The critical sequence $\langle \kappa_n : n \in \omega \rangle$ is a subset of V_λ , so it belongs to

$V_{\lambda+1}$. But then for every $A \subseteq V_\lambda$, $\{A \cap V_{\kappa_n} : n \in \omega\} \in V_{\lambda+1}$, so

$$\begin{aligned} (j \upharpoonright V_\lambda)^+(A) &= \bigcup_{n \in \omega} (j \upharpoonright V_\lambda)(A \cap V_{\kappa_n}) \\ &= \bigcup_{n \in \omega} j(A \cap V_{\kappa_n}) \\ &= j\left(\bigcup_{n \in \omega} (A \cap V_{\kappa_n})\right) \\ &= j(A). \end{aligned}$$

□

It is worth noting that there is a strong connection between first-order formulas in $V_{\lambda+1}$ and second-order formulas in V_λ . In fact, all the elements of $V_{\lambda+1}$ are subsets of V_λ , so they can be replaced with predicate symbols:

Lemma 3.5. *Let $A \in V_{\lambda+1} \setminus V_\lambda$ and $\varphi(v_0, v_1, \dots, v_n)$ be a formula. Fix \hat{A} a predicate symbol, and define $\varphi^*(v_1, \dots, v_n)$ in the language of LST expanded with \hat{A} as following:*

- *for every occurrence of v_0 , substitute \hat{A} ;*
- *for every non-bounded quantified variable x , substitute every occurrence of x with X , a second-order variable.*

Then for every $a_1, \dots, a_n \in V_\lambda$

$$V_{\lambda+1} \models \varphi(A, a_1, \dots, a_n) \quad \text{iff} \quad (V_\lambda, A) \models \varphi^*(a_1, \dots, a_n).$$

Proof. The proof is by induction on the complexity of φ . □

The previous Lemma is key in clarifying the relationship between elementary embeddings in V_λ and $V_{\lambda+1}$, and to finally prove that j^+ is a Σ_0 elementary embedding from $V_{\lambda+1}$ to itself (or, alternatively that j is a Σ_0^1 elementary embedding from V_λ to itself).

Theorem 3.6. *Let $j : V_\lambda \prec V_\lambda$. Then $j^+ : V_{\lambda+1} \rightarrow V_{\lambda+1}$ is a Δ_0 -elementary embedding.*

Proof. Let \hat{A} be a symbol for an 1-ary relation, $\hat{a}_1, \dots, \hat{a}_n$ symbols for constants and let φ be a formula in LST*, the language of LST expanded with \hat{A} and $\hat{a}_1, \dots, \hat{a}_n$. We want to prove that for any $A \subseteq V_\lambda$:

$$(V_\lambda, A, a_1, \dots, a_n) \models \varphi \quad \text{iff} \quad (V_\lambda, j^+(A), j(a_1), \dots, j(a_n)) \models \varphi.$$

Actually we will prove only one direction for every φ , the other one following by considering $\neg\varphi$.

First of all, we skolemize φ , so we find f_1, \dots, f_m functions, $f_i : (V_\lambda)^{m_i} \rightarrow V_\lambda$, such that φ is equivalent to a formula φ^* in LST* expanded with $\hat{f}_1, \dots, \hat{f}_m$, where φ^* is $\forall x_1 \forall x_2 \dots \forall x_m \varphi'$, with φ' a Δ_0 formula.

Let $t_0(x_1, \dots, x_m), \dots, t_p(x_1, \dots, x_m)$ be all the terms that are in φ' . We can express as a logical formula the phrase “ $t_i(x_1, \dots, x_m)$ exists”: we work by induction on the tree of the term, writing a conjunction of formulas in this way

- every time that there is an occurrence of a function, i.e. $\hat{f}_i(t')$, we add to the formula $\exists y f_i(t') = y$;
- every time that there is an occurrence of a constant, i.e. \hat{a}_i , we add to the formula $\hat{a}_i \neq \emptyset$, if $a_i \neq \emptyset$; otherwise we don't write anything.

Then we have that

$$\begin{aligned} (V_\lambda, A, a_1, \dots, a_n, f_1, \dots, f_m) \models \varphi^* \quad \text{iff} \\ \forall \delta < \lambda \ (V_\delta, A \cap V_\delta, a'_1, \dots, a'_n, f_1 \cap V_\delta, \dots, f_m \cap V_\delta) \models \\ \forall x_1, \dots, \forall x_m \left(\bigwedge_{i < p} t_i(x_1, \dots, x_m) \text{ exists} \rightarrow \varphi' \right), \end{aligned}$$

where a'_i is a_i if $a_i \in V_\delta$, otherwise is \emptyset . This is true because, with δ fixed, if only one term does not exist the formula is satisfied, and if all the terms exist, then they are a witness for the satisfaction (or not) of φ' .

So

$$\begin{aligned} (V_\lambda, A, a_1, \dots, a_n) \models \varphi \\ \Leftrightarrow (V_\lambda, A, a_1, \dots, a_n, f_1, \dots, f_m) \models \varphi^* \\ \Leftrightarrow \forall \delta < \lambda \ (V_\delta, A \cap V_\delta, a'_1, \dots, a'_n, f_1 \cap V_\delta, \dots, f_m \cap V_\delta) \models \\ \forall x_1, \dots, \forall x_m \left(\bigwedge_{i < p} t_i(x_1, \dots, x_m) \text{ exists} \rightarrow \varphi' \right) \\ \Leftrightarrow \forall \delta < \lambda \ (V_\delta, j(A \cap V_\delta), j(a'_1), \dots, j(a'_n), j(f_1 \cap V_\delta), \dots, j(f_m \cap V_\delta)) \models \\ \forall x_1, \dots, \forall x_m \left(\bigwedge_{i < p} t_i(x_1, \dots, x_m) \text{ exists} \rightarrow \varphi' \right) \\ \Leftarrow (V_\lambda, j^+(A), j(a_1), \dots, j(a_n), j^+(f_1), \dots, j^+(f_m)) \models \varphi^* \\ \Leftrightarrow (V_\lambda, j^+(A), j(a_1), \dots, j(a_n)) \models \varphi. \end{aligned}$$

The only thing left to prove is the right direction of the fifth line. The bump is in the fact that if a $j^+(f_i)$ is not total, then it would be possible to have the equation in the fourth line satisfied because the relevant terms do not exist, but the fifth line not satisfied on the elements not in the domain of $j^+(f_i)$. We prove that actually the $j^+(f_i)$ are total, in fact even more:

Claim 3.7. *If $j : V_\lambda \prec V_\lambda$ and $f : V_\lambda \rightarrow V_\lambda$, even partial, then $j^+(\text{dom}(f)) = \text{dom}(j^+(f))$.*

Proof of claim. By definition,

$$j^+(\text{dom}(f)) = \bigcup_{m \in \omega} j(\text{dom}(f) \cap V_{\kappa_m}),$$

while

$$\begin{aligned} \text{dom}(j^+(f)) &= \text{dom}\left(\bigcup_{m \in \omega} j(f \cap V_{\kappa_m})\right) = \\ &= \bigcup_{m \in \omega} \text{dom}(j(f \cap V_{\kappa_m})) = \bigcup_{m \in \omega} j(\text{dom}(f \cap V_{\kappa_m})). \end{aligned}$$

It is not immediate to see why the two are the same, because it is possible that $\text{dom}(f) \cap V_{\kappa_m}$ and $\text{dom}(f \cap V_{\kappa_m})$ are different, as on the left we are cutting the dominion, on the right both the dominion and codominion. The solution is to divide everything in even smaller pieces, that would be the same on the left and on the right side.

$$\text{dom}(f) \cap V_{\kappa_m} = \bigcup_{n \in \omega} \text{dom}(f \cap V_{\kappa_n}) \cap V_{\kappa_m},$$

and

$$\text{dom}(f \cap V_{\kappa_m}) = \bigcup_{n \in \omega} \text{dom}(f \cap V_{\kappa_n}) \cap V_{\kappa_n}.$$

Therefore:

$$\begin{aligned} j^+(\text{dom}(f)) &= \bigcup_{m \in \omega} j(\text{dom}(f) \cap V_{\kappa_m}) = \bigcup_{m \in \omega} j\left(\bigcup_{n \in \omega} \text{dom}(f \cap V_{\kappa_n}) \cap V_{\kappa_m}\right) = \\ &= \bigcup_{m \in \omega} \bigcup_{n \in \omega} j(\text{dom}(f \cap V_{\kappa_n}) \cap V_{\kappa_m}) = \bigcup_{n \in \omega} \bigcup_{m \in \omega} j(\text{dom}(f \cap V_{\kappa_n}) \cap V_{\kappa_m}) = \\ &= \bigcup_{n \in \omega} j(\text{dom}(f \cap V_{\kappa_n})) = \text{dom}(j^+(f)). \end{aligned}$$

□

The third and fifth equivalence are a delicate point. In the third equivalence, the ω -sequence $\langle \text{dom}(f \cap V_{\kappa_n}) \cap V_{\kappa_m} : n \in \omega \rangle \in V_\lambda$, even if it is defined with κ_n , therefore j behaves properly. In the fifth equivalence, what seems an infinite union is in fact finite.

□

Therefore every $j : V_\lambda \prec V_\lambda$ can be extended to a unique $j^+ : V_{\lambda+1} \rightarrow V_{\lambda+1}$ that is at least a Δ_0 -elementary embedding. Moreover it is possible to prove

- I3 holds iff for some λ there exists a $j : V_{\lambda+1} \rightarrow V_{\lambda+1}$ that is a Δ_0 -elementary embedding;
- I1 holds iff for some λ there exists a $j : V_{\lambda+1} \prec V_{\lambda+1}$ that is a Σ_n -elementary embedding for every n

Theorem ?? implies that if $j, k : V_\lambda \prec V_\lambda$, then $j^+(k) : V_\lambda \prec V_\lambda$. This operation between elementary embeddings is called *application*, and we write $j^+(k) = j \cdot k$. It creates an interesting algebra.

Theorem 3.8 (Laver, 1992 [?]). *Fix λ and let $\mathcal{E}_\lambda = \{j : V_\lambda \prec V_\lambda\}$. For all $j \in \mathcal{E}_\lambda$ the closure of $\{j\}$ in $(\mathcal{E}_\lambda, \cdot)$ is the free algebra generated by the law*

$$(Left\ Distributive\ Law) \quad i \cdot (j \cdot k) = (i \cdot j) \cdot (i \cdot k).$$

Moreover Laver proved that the free algebra generated by the laws above satisfies the word problem. His proof used extensively I3, but later Dehornoy ([?]) managed to prove the same thing in ZFC. A proof of this can be found in the Handbook of Set Theory [?], Chapter 11.

There are still open problems at the I3 level:

Question 3.9. *Are there $j, k : V_\lambda \prec V_\lambda$ such that the algebra generated by them is free?*

A standard way of using application is to consider $j, j(j), j(j(j)) \dots$, and we write $j^0 = j$ and $j^{n+1} = j(j^n)$. If $\langle \kappa_n : n \in \omega \rangle$ is the critical sequence of j , then by elementarity, $\text{crt}(j(j)) = j(\text{crt}(j)) = \kappa_1$ and $\langle \kappa_n : n \geq 1, n \in \omega \rangle$ is the critical sequence of j^2 . By induction, $\text{crt}(j^n) = \kappa_n$. This is the reason why, when talking about I3 or I1, the critical point is never mentioned, when usually it is the main object of research (see strongness, supercompactness...): if there exists a λ that witnesses I3, then there are infinite critical points tied to the same λ , so it is really λ the relevant cardinal, while the critical points are interchangeable and unbounded in λ . Still, they carry a lot of strength. The following result shows that I3 really sits on top of the large cardinal hierarchy:

Proposition 3.10. *Let $\langle \kappa_i : i < \omega \rangle$ be the critical sequence of j . Then for every i , κ_i is γ -supercompact for any $\gamma < \lambda$ and n -huge for every n .*

We use the characterization via ultrafilter:

Proposition 3.11 (Solovay, Reinhardt). *A cardinal κ is γ -supercompact iff there exists a fine, normal ultrafilter on $\mathcal{P}_\kappa(\lambda)$. It is n -huge iff there is a κ -complete normal ultrafilter U over some $\mathcal{P}(\eta)$ and $\kappa = \eta_0 < \eta_1 < \dots < \eta_n = \eta$ such that for any $i < n$ $\{x \in \mathcal{P}(\eta) : \text{ot}(x \cap \eta_{i+1}) = \eta_i\} \in U$.*

Proof of ??. Let $\kappa_0 \leq \gamma < \lambda$, and suppose $\gamma < \kappa_n$; then it is routine to check that γ -supercompactness of κ_0 is witnessed by

$$U = \{X \subseteq \mathcal{P}_{\kappa_0}(\gamma) : (j^n)''\gamma \in j^n(X)\}$$

and that n -hugeness of κ_0 is witnessed by

$$U = \{X \subseteq \mathcal{P}(\kappa_n) : j''\kappa_n \in j(X)\} \quad \text{and} \quad \kappa_0, \dots, \kappa_n.$$

Then κ_n is γ -supercompact and n -huge by elementarity. □

On the other hand λ (the “true” rank-into-rank cardinal) does not enjoy many large cardinal properties, as it is of cofinality ω , and usually large cardinals are regular. The most one can say is that it is a Rowbottom cardinal.

We end the section with some interesting reflection properties of V_λ :

Lemma 3.12. *Suppose $j : V_\lambda \prec V_\lambda$ and let $\langle \kappa_n : n \in \omega \rangle$ be its critical sequence. Then for every $n \in \omega$, $V_{\kappa_n} \prec V_\lambda$.*

Proof. Since j is the identity on V_{κ_0} , it is easy to see that $V_{\kappa_0} \prec V_{\kappa_1}$. Considering $j(j)$, since $\text{crt}(j(j)) = \kappa_1$ and

$$j(j)(\kappa_1) = j(j)(j(\kappa_0)) = j(j(\kappa_0)) = \kappa_2,$$

we have also that $V_{\kappa_1} \prec V_{\kappa_2}$. We can generalize this to prove that for every $n \in \omega$, $V_{\kappa_n} \prec V_{\kappa_{n+1}}$. But then $\langle (V_{\kappa_n}, \text{id}_{V_{\kappa_n}}), n \in \omega \rangle$ forms a direct system, whose direct limit is V_λ . □

In particular, as κ_0 is inaccessible, $V_\lambda \models \text{ZFC}$.

4 $L(V_{\lambda+1})$ and I0

It would seem that one could not go any further than I1, as already $j : V_{\lambda+2} \prec V_{\lambda+2}$ is inconsistent. But there is a way to enlarge the domain of j without stumbling in Kunen's inconsistency:

Definition 4.1. *Let A be a set. Then the class of sets constructible relative to A is:*

- $L_0(A) =$ the transitive closure of A ;
- $L_{\alpha+1}(A) = \text{Def}(L_\alpha(A))$;
- $L_\gamma(A) = \bigcup_{\alpha < \gamma} L_\alpha(A)$;
- $L(A) = \bigcup_{\alpha \in \text{Ord}} L_\alpha(A)$.

This is the smallest ZF-model that contains A . We can now introduce I0:

Definition 4.2 (Woodin). **I0** *There exists $j : L(V_{\lambda+1}) \prec L(V_{\lambda+1})$ with critical point less than λ .*

As $L(V_{\lambda+1}) \models V = L(V_{\lambda+1})$, it must be that $j(V_{\lambda+1}) = V_{\lambda+1}$. Thanks to the added assumption that the critical point must be less than λ , then, I0 is stronger than I1, as $j \upharpoonright V_{\lambda+1} : V_{\lambda+1} \prec V_{\lambda+1}$ (otherwise it could have been the identity on $V_{\lambda+1}$). This implies that λ must be the supremum of the critical sequence. If one wants to avoid the menace of Kunen's inconsistency, it must be that $\langle S_\alpha : \alpha < \text{crt}(j) \rangle$ is not in $L(V_{\lambda+1})$. This is constructed via the Axiom of Choice on $V_{\lambda+1}$. So, for I0 to hold, it must be that $L(V_{\lambda+1}) \not\models \text{AC}$, and in particular $V_{\lambda+1}$ must not be well-orderable. This is the first affinity with $\text{AD}^{L(\mathbb{R})}$, as in that case we have $L(\mathbb{R}) \not\models \text{AC}$. More affinities will be introduced in section ??.

At a superficial glance I0 seems a Reinhardt cardinal, as it is an embedding in a “universe” of ZF. But the situation is completely different: our ground model is V , we assume that $V \models \text{ZFC}$, and we construct $L(V_{\lambda+1})$ inside V . So j is a definable class of V , and we will see that $L(V_{\lambda+1})$ does not satisfy replacement for j , so such a j is not a witness for Reinhardt, as in that case the theory of the ground model must be richer.

So, every time we want to prove something in $L(V_{\lambda+1})$ under I0, we must be very careful as AC does not hold in $L(V_{\lambda+1})$, but we can actually take advantage of the fact that our base theory (so the theory of V), is actually ZFC. An example of this is Lemma ??.

Before delving in I0, we analyze the structure of $L(V_{\lambda+1})$ itself, without further assumptions, that still carry similarities with $L(\mathbb{R})$.

As $L(V_{\lambda+1})$ is L relativized to $V_{\lambda+1}$, we can relativize in the same way OD and HOD :

Definition 4.3. $OD_{V_{\lambda+1}}$, the class of the sets that are ordinal-definable over $V_{\lambda+1}$, is the Gödel closure of $\{V_\alpha : \alpha \in \text{Ord}\} \cup V_{\lambda+1}$.

$HOD_{V_{\lambda+1}}$, the class of the sets that are hereditarily ordinal-definable over $V_{\lambda+1}$, is the class of sets whose transitive closure is contained in $OD_{V_{\lambda+1}}$.

Lemma 4.4. *There exists a definable surjection $\Phi : \text{Ord} \times V_{\lambda+1} \rightarrow L(V_{\lambda+1})$, therefore $L(V_{\lambda+1}) \models V = \text{HOD}_{V_{\lambda+1}}$*

Proof. This is immediate from the theory of relative constructibility. \square

The first application of this Lemma comes in form of partial Skolem functions. Since $L(V_{\lambda+1}) \not\models \text{AC}$, we possibly could not have Skolem functions. But with the previous Lemma we can define for every formula $\varphi(x, x_1, \dots, x_n)$, $a \in V_{\lambda+1}$, $a_1, \dots, a_n \in L(V_{\lambda+1})$:

$$h_{\varphi, a}(a_1, \dots, a_n) = y \quad \text{where } y \text{ is the minimum in } (\text{OD}_a)^{L(V_{\lambda+1})} \\ \text{such that } L(V_{\lambda+1}) \models \varphi(y, a_1, \dots, a_n).$$

These are partial Skolem functions, and the Skolem Hull of a set is its closure under such Skolem functions.

We indicate with \rightarrow the surjectivity of a function.

Definition 4.5.

$$\Theta_{V_{\lambda+1}}^{L(V_{\lambda+1})} = \sup\{\gamma : \exists f : V_{\lambda+1} \rightarrow \gamma, f \in L(V_{\lambda+1})\}.$$

In the following, we will call $\Theta_{V_{\lambda+1}}^{L(V_{\lambda+1})} = \Theta$, because it is a lighter notation and there is no possibility of misinterpretation.

The role of Θ in $L(V_{\lambda+1})$ is exactly the same of its correspondent in $L(\mathbb{R})$. In the usual setting, under AC , to measure the largeness of a set we fix a bijection from this set to a cardinal or, equivalently, the ordertype of a well-ordering of the set. Since there is no Axiom of Choice in $L(V_{\lambda+1})$, it is not always possible to define cardinality for sets that are not in $V_{\lambda+1}$ in the usual way, so to quantify the “largeness” of a subset of $V_{\lambda+1}$ we will not use bijections, but surjections, or, equivalently, not well-orders, but prewellorderings (pwo for short).

An order is a pwo if it satisfies antireflexivity, transitivity, and every subset has a least element; in other words, it is a well-order without the antisymmetric property. It is easy to see that the counterimage of a surjective function is a pwo. One can think of a pwo as an order whose equivalence classes are well-ordered, or a well-ordered partition. This creates a strong connection between subsets of $V_{\lambda+1}$ and ordinals in Θ :

Lemma 4.6. 1. For every $\alpha < \Theta$, there exists in $L(V_{\lambda+1})$ a pwo in $V_{\lambda+1}$ with ordertype α , that is codeable as a subset of $V_{\lambda+1}$;

2. for every $Z \subseteq V_{\lambda+1}$, $Z \in L(V_{\lambda+1})$ there exists $\alpha < \Theta$ such that $Z \in L_\alpha(V_{\lambda+1})$.

Proof. 1. Let $\rho : V_{\lambda+1} \rightarrow \alpha$. Then

$$R_\alpha = \{(a, b) \in V_{\lambda+1} \times V_{\lambda+1} : \rho(a) \leq \rho(b)\}$$

is a pwo in $V_{\lambda+1}$. Moreover, $V_{\lambda+1} \times V_{\lambda+1}$ can be codified as a subset of $V_{\lambda+1}$, so also R_α can.

2. Let γ be such that $Z \in L_\gamma(V_{\lambda+1})$ and consider $H^{L_\gamma(V_{\lambda+1})}(V_{\lambda+1}, Z)$ the Skolem Hull in $L_\gamma(V_{\lambda+1})$ of $V_{\lambda+1}$ and Z . Then, since $H^{L_\gamma(V_{\lambda+1})}(V_{\lambda+1}, Z) \cong L_\gamma(V_{\lambda+1})$, by condensation its collapse $\mathcal{X} = L_\alpha(V_{\lambda+1})$ for some α . But $H^{L_\gamma(V_{\lambda+1})}(V_{\lambda+1}, Z)$ is the closure under the Skolem functions, and since there is a surjection from $V_{\lambda+1}$ to the Skolem functions, this surjection

transfers to $H^{L_\gamma(V_{\lambda+1})}(V_{\lambda+1}, Z)$ and to $L_\alpha(V_{\lambda+1})$, so $\alpha < \Theta$. Since Z and all its elements are in $H^{L_\gamma(V_{\lambda+1})}(V_{\lambda+1}, Z)$, Z is not collapsed and then $Z \in L_\alpha(V_{\lambda+1})$. \square

We can also have a “local” version of Θ :

Definition 4.7.

$$\Theta_{V_{\lambda+1}}^{L_\alpha(V_{\lambda+1})} = \sup\{\gamma : \exists f : V_{\lambda+1} \twoheadrightarrow \gamma, f \in L(V_{\lambda+1}), \\ \{(a, b) : f(a) < f(b)\} \in L_\alpha(V_{\lambda+1})\}.$$

The reason why we do not just relativize Θ to $L_\alpha(V_{\lambda+1})$ is because there can be pwos in $L(V_{\lambda+1})$ that are not in $L_\alpha(V_{\lambda+1})$, but can be coded in $L_\alpha(V_{\lambda+1})$, so in this way $\Theta_{V_{\lambda+1}}^{L_\alpha(V_{\lambda+1})}$ is more “truthful” to indicate the ordinals that are “hidden” inside $V_{\lambda+1}$.

The notion of “stable” also neatly generalizes. Let δ be the minimum ordinal such that $L_\delta(V_{\lambda+1}) \prec_1 L_\Theta(V_{\lambda+1})$. Then $L_\delta(V_{\lambda+1})$ is the collapse of the closure of $V_\lambda + 1$ under the Skolem functions $h_{\varphi, a}$ such that φ is Σ_1 . The map that associates to any (φ, a, x) the collapse of $h_{\varphi, a}(x)$ is therefore a Σ_1 partial surjection from $V_{\lambda+1}$ to $L_\delta(V_{\lambda+1})$. A study of the fine structure of $L_\delta(V_{\lambda+1})$ has the following results:

- Lemma 4.8.** • *For any $\beta < \delta$, there exists a total surjection from $V_{\lambda+1}$ to $L_\beta(V_{\lambda+1})$ that is in $L_\delta(V_{\lambda+1})$. Therefore $\Theta^{L_\delta(V_{\lambda+1})} = \delta$.*
- *δ is the supremum of the Δ_1^2 pwos of $V_{\lambda+1}$ and $(\Delta_1^2)^{L(V_{\lambda+1})} = L_\delta(V_{\lambda+1}) \cap V_{\lambda+2}$.*
 - *Let $\rho : V_{\lambda+1} \rightarrow L_\delta(V_{\lambda+1})$ be a partial Σ_1 map. Then for any $Z \subseteq \text{dom}(\rho)$, $Z \in L_\delta(V_{\lambda+1})$, $\rho \upharpoonright Z$ is bounded.*

The proofs are the same as in [?].

Definition 4.9.

$$\text{DC}_\lambda : \quad \forall X \forall F : (X)^{<\lambda} \rightarrow \mathcal{P}(X) \setminus \emptyset \exists g : \lambda \rightarrow X \forall \gamma < \lambda g(\gamma) \in F(g \upharpoonright \gamma).$$

Note that this is a generalization of DC, since $\text{DC} = \text{DC}_\omega$: we use directly a function on $< \lambda$ -sequences instead of considering a binary relation because binary relations cannot handle the limit stages.

Lemma 4.10. *In $L(V_{\lambda+1})$ the following hold:*

1. Θ is regular ([?]);
2. DC_λ .

The proof of the first point is as in Exercise 28.19 in [?]. The second point is a consequence of the fact that DC_λ holds in V , and therefore in $V_{\lambda+1}$, and Lemma ??.

Suppose now for the rest of the section that $j : L(V_{\lambda+1}) \prec L(V_{\lambda+1})$, with $\text{crt}(j) < \lambda$. Define

$$U_j = \{X \subseteq V_{\lambda+1} : X \in L(V_{\lambda+1}), j \upharpoonright V_\lambda \in j(X)\}.$$

Then U_j is a normal non-principal $L(V_{\lambda+1})$ -ultrafilter in $V_{\lambda+1}$, and we can construct the ultraproduct $Ult(L(V_{\lambda+1}), U_j)$. Note that for every $f, g : V_{\lambda+1} \rightarrow L(V_{\lambda+1})$

$$\begin{aligned} [f] = [g] &\text{ iff } \{x \in V_{\lambda+1} : f(x) = g(x)\} \in U_j \\ &\text{ iff } j \upharpoonright V_\lambda \in j(\{x \in V_{\lambda+1} : f(x) = g(x)\}) = \\ &\quad = \{x \in V_{\lambda+1} : j(f)(x) = j(g)(x)\} \\ &\text{ iff } j(f)(j \upharpoonright V_\lambda) = j(g)(j \upharpoonright V_\lambda), \end{aligned}$$

and in the same way $[f] \in [g]$ iff $j(f)(j \upharpoonright V_\lambda) \in j(g)(j \upharpoonright V_\lambda)$, so

$$Ult(L(V_{\lambda+1}), U_j) \cong \{j(f)(j \upharpoonright V_\lambda) : \text{dom } f = V_{\lambda+1}\}.$$

Let $i : L(V_{\lambda+1}) \rightarrow Ult(L(V_{\lambda+1}), U_j)$ be the natural embedding of the ultraproduct, then for every $a \in L(V_{\lambda+1})$, $i(a) = [c_a]$ corresponds in the equivalence to $j(c_a)(j \upharpoonright V_\lambda) = j(a)$, so we can suppose $i = j$. Is j an elementary embedding from $L(V_{\lambda+1})$ to $Ult(L(V_{\lambda+1}), U_j)$? Since we do not have AC, the answer is not immediate because we possibly do not have Los' Theorem.

We will prove Los' Theorem for this case, and this will imply that j is an elementary embedding. It is clear that the only real obstacle is to prove that for every formula φ and $f_1, \dots, f_n \in L(V_{\lambda+1})$ such that $\text{dom } f_i = V_{\lambda+1}$

$$\begin{aligned} Ult(L(V_{\lambda+1}), U_j) \models \exists x \varphi([f_1], \dots, [f_n]) \\ \text{ iff } \{x \in V_{\lambda+1} : L(V_{\lambda+1}) \models \exists y \varphi(f_1(x), \dots, f_n(x))\} \in U_j. \end{aligned}$$

The direction from left to right is immediate: if $[g]$ witness the left side, then $g(x)$ witness the right side. For the opposite direction, we need a sort of U_j -choice, i.e., we need to find a function g such that

$$\{x \in V_{\lambda+1} : L(V_{\lambda+1}) \models \varphi(g(x), f_1(x), \dots, f_n(x))\} \in U_j.$$

We re-formulate this considering $f : V_{\lambda+1} \rightarrow L(V_{\lambda+1}) \setminus \emptyset$,

$$f(x) = \{y \in L(V_{\lambda+1}) : L(V_{\lambda+1}) \models \varphi(y, f_1(x), \dots, f_n(x))\}.$$

Lemma 4.11 (Los' Theorem for I0, Woodin, [?]). *Let $j : L(V_{\lambda+1}) \prec L(V_{\lambda+1})$ and U_j as above. Then for every $F : V_{\lambda+1} \rightarrow L(V_{\lambda+1}) \setminus \{\emptyset\}$ there exists $H : V_{\lambda+1} \rightarrow L(V_{\lambda+1}) \setminus \{\emptyset\}$ such that $\{x \in V_{\lambda+1} : H(x) \in F(x)\} \in U_j$.*

Proof. First we consider the case $\forall a \in V_\lambda, F(a) \subseteq V_{\lambda+1}$. We have to define H such that $j(H)(j \upharpoonright V_\lambda) \in j(F)(j \upharpoonright V_\lambda)$. Fix a $b \in j(F)(j \upharpoonright V_\lambda)$, and define

$$H(k) = \begin{cases} c & \text{if } k : V_\lambda \prec V_\lambda, k(k) = j \text{ and } k(c) = b \\ \emptyset & \text{otherwise} \end{cases}$$

Note that

$$K_b := \{k \in V_{\lambda+1} \mid k : V_\lambda \prec V_\lambda, k(k) = j, \exists c \, k(c) = b\} \in U_j,$$

because

$$j(K_b) = \{k \in V_{\lambda+1} \mid k : V_\lambda \prec V_\lambda, k(k) = j(j), \exists c \, k(c) = j(b)\}$$

and $j \upharpoonright V_\lambda \in j(K_b)$ (with $c = b$), so $\{x \in V_{\lambda+1} : H(x) \neq \emptyset\} \in U_j$. Then

$$j(H)(k) = \begin{cases} c & \text{if } k : V_\lambda \prec V_\lambda, k(k) = j(j) \text{ and } k(c) = j(b) \\ 0 & \text{otherwise} \end{cases}$$

so $j(H)(j \upharpoonright V_\lambda) = b \in j(F)(j \upharpoonright V_\lambda)$.

For the more general case $\forall a \in V_\lambda \ F(a) \subseteq L(V_{\lambda+1})$ fix $\Phi : Ord \times V_{\lambda+1} \rightarrow L(V_{\lambda+1})$ definable and define

$$\hat{F}(a) = \{x \in V_{\lambda+1} : \exists \gamma \ \Phi(\gamma, x) \in F(a)\}.$$

Then there exists \hat{H} such that $\{a \in V_{\lambda+1} : \hat{H}(a) \in \hat{F}(a)\} \in U_j$. Let $\gamma_a = \min\{\gamma : \Phi(\gamma, \hat{H}(a)) \in F(a)\}$. Therefore $H(a) = \Phi(\gamma_a, \hat{H}(a))$ is as desired. \square

Therefore, calling

$$\mathcal{Z} = \{j(f)(j \upharpoonright V_\lambda) : f \in L(V_{\lambda+1}), \text{dom}(f) = V_{\lambda+1}\},$$

$j : L(V_{\lambda+1}) \rightarrow \mathcal{Z}$ is an elementary embedding, and $\mathcal{Z} \cong L(V_{\lambda+1})$. Let k_U be the inverse of the collapse of \mathcal{Z} . We've seen in the proof of the previous Lemma that for every $b \in V_{\lambda+1}$ there exists h such that $j(h)(j \upharpoonright V_\lambda)$, so $V_{\lambda+1} \subseteq \mathcal{Z}$ and $k_U : L(V_{\lambda+1}) \prec \mathcal{Z}$. Moreover, if R is a pwo in $V_{\lambda+1}$, then $R = \{a \in V_{\lambda+1} : j(a) \in j(R)\}$, and since $j(a), j(R) \in \mathcal{Z}$ and $V_{\lambda+1} \subseteq \mathcal{Z}$ we have that R is not collapsed, so $\Theta \subseteq \mathcal{Z}$ and $\text{crt}(k_U) > \Theta$.

Theorem 4.12 (Woodin, [?]). *For every $j : L(V_{\lambda+1}) \prec L(V_{\lambda+1})$ there exist a $L(V_{\lambda+1})$ -ultrafilter U in $V_{\lambda+1}$ and $j_U, k_U : L(V_{\lambda+1}) \prec L(V_{\lambda+1})$ such that j_U is the elementary embedding from U , $j = j_U \circ k_U$ and $j \upharpoonright L_\Theta(V_{\lambda+1}) = j_U \upharpoonright L_\Theta(V_{\lambda+1})$.*

Definition 4.13. *Let $j : L(V_{\lambda+1}) \prec L(V_{\lambda+1})$. We say that j is weakly proper if $j = j_U$.*

Note that $U \notin L(V_{\lambda+1})$, otherwise we could run the naive Kunen's Theorem argument, so $L(V_{\lambda+1})$ does not satisfy replacement for j , and this proves that I_0 is not a " $L(V_{\lambda+1})$ -Reinhardt". In fact, one cannot even have $j \upharpoonright L_\Theta(V_{\lambda+1}) \in L(V_{\lambda+1})$, otherwise one could construct U from that, so j is not amenable.

In a certain sense weakly proper embeddings are the "right" embeddings for I_0 , and the most natural extension of I_1 , if one considers that $L_\Theta(V_{\lambda+1})$ is, loosely speaking, the $V_{\lambda+2}$ of $L(V_{\lambda+1})$, i.e., the set of subsets of $V_{\lambda+1}$. Every embedding has therefore a core part, that comes from an ultrapower, it is unique and depends solely on its behaviour on $L_\Theta(V_{\lambda+1})$, and a spurious part (note that k_U is not an I_0 -embedding, as it has critical point $> \lambda$) that moves things up above without really changing the important part of the embedding. For example, if there are indiscernibles, it could be a simple shift of indiscernibles, so nothing that carries real information.

In fact, the behaviour of an embedding that is weakly proper depends only on its V_λ part:

Lemma 4.14 (Woodin, [?]). *For every $j_1, j_2 : L(V_{\lambda+1}) \prec L(V_{\lambda+1})$, if $j_1 \upharpoonright V_\lambda = j_2 \upharpoonright V_\lambda$ then $j_1 \upharpoonright L_\Theta(V_{\lambda+1}) = j_2 \upharpoonright L_\Theta(V_{\lambda+1})$.*

Proof. We can suppose that j_1 and j_2 are weakly proper. By the usual analysis of the ultraprodut, we have that every strong limit cardinal with cofinality bigger than Θ is a fixed point for j_1 and j_2 , so $I = \{\eta : j_1(\eta) = j_2(\eta) = \eta\}$ is a proper class. Let $M = H^{L(V_{\lambda+1})}(I \cup V_{\lambda+1})$. Since $V_{\lambda+1} \subseteq M$, the inverse of the transitive collapse of M , has domain $L(V_{\lambda+1})$. If k^* is not the identity then, as $\Theta \subseteq M$ by Lemma ??, $\text{crt}(k^*) > \Theta$. But in that case $\text{crt}(k^*)$ is a strong limit cardinal with cofinality bigger than Θ , so $\text{crt}(k^*) \in I$, and this is a contradiction, because $I \subseteq \text{ran}(k^*)$ and $\text{crt}(k^*) \notin \text{ran}(k^*)$. So k^* is the identity and $L(V_{\lambda+1}) = H^{L(V_{\lambda+1})}(I \cup V_{\lambda+1})$.

Therefore every element of $L_\Theta(V_{\lambda+1})$ is definable with parameters in $I \cup V_{\lambda+1}$. Let $A \in L(V_{\lambda+1}) \cap V_{\lambda+2}$, $A = \{x \in V_{\lambda+1} : L(V_{\lambda+1}) \models \varphi(\eta, a)\}$ with $\eta \in I$ and $a \in V_{\lambda+1}$. Since $j_1 \upharpoonright V_\lambda = j_2 \upharpoonright V_\lambda$, we have that $j_1(a) = j_2(a)$. Then

$$\begin{aligned} j_1(A) &= \{x \in V_{\lambda+1} : L(V_{\lambda+1}) \models \varphi(\eta, j_1(a))\} = \\ &= \{x \in V_{\lambda+1} : L(V_{\lambda+1}) \models \varphi(\eta, j_2(a))\} = j_2(A). \end{aligned}$$

But every element of $L_\Theta(V_{\lambda+1})$ is definable from an ordinal $\alpha < \Theta$ and an element of $V_{\lambda+1}$, α is definable from some pwo in $L(V_{\lambda+1}) \cap V_{\lambda+2}$, so $j_1 \upharpoonright L_\Theta(V_{\lambda+1}) = j_2 \upharpoonright L_\Theta(V_{\lambda+1})$. \square

Corollary 4.15 (Woodin, [?]). *Let $j : L(V_{\lambda+1}) \prec L(V_{\lambda+1})$ with $\text{crt}(j)$. Then $C = \{\alpha < \Theta : j(\alpha) = \alpha\}$ is unbounded in Θ .*

Proof. We can suppose that j is weakly proper. Then every ordinal $\beta < \Theta$ is definable from some elements of $I \cup V_{\lambda+1}$, say $I_\beta = \{i_0, \dots, i_n\} \in I$.

Let $\alpha_0 < \Theta$ and consider the transitive collapse of the Skolem closure Z of $\bigcup_{\beta < \alpha_0} I_\beta \cup V_{\lambda+1}$. Then by condensation it must be some $L_\gamma(V_{\lambda+1})$. As $\alpha_0 < \Theta$, there exists $\pi : V_{\lambda+1} \rightarrow Z$, therefore $\gamma < \Theta$. As $\alpha_0 \subseteq Z$, we have $\alpha_0 < \gamma$. As $j(Z) = Z$, we have $j(\gamma) = \gamma$. Therefore for any $\alpha_0 < \Theta$ there exists a $\gamma \in C$ such that $\alpha_0 < \gamma < \Theta$. \square

5 Strong implications and inverse limits

When introducing new axioms, it is good practice to prove that they really are adding something, i.e., that their consistency strength is strictly larger (or smaller) than the consistency of the already known ones.

Definition 5.1. *Let $A(x)$ and $B(x)$ two large cardinal properties. We say that $A(x)$ strongly implies $B(x)$ iff for any κ , $A(\kappa)$ implies $B(\kappa)$ and the smallest κ such that $A(\kappa)$ holds is bigger than the smallest η such that $B(\eta)$ holds.*

This is even stronger than just the strict implications, cfr the note after Lemma ??.

Proposition 5.2. *If $j : V_\lambda \prec V_\lambda$ with $\text{crt}(j) = \kappa$, then there is a normal ultrafilter U on κ such that $\{\alpha < \kappa : \alpha \text{ is } n\text{-huge for every } n\} \in U$. Therefore $I3$ strongly implies being n -huge for every n .*

Proof. By ?? κ is n -huge for every $n \in \omega$. Let $U = \{X \subseteq \kappa : \kappa \in j(X)\}$ and let

$$X_0 = \{\alpha < \kappa : \alpha \text{ is } n\text{-huge for every } n\}.$$

Then

$$j(X_0) = \{\alpha < j(\kappa) : \alpha \text{ is } n\text{-huge for every } n\},$$


and $\kappa \in j(X_0)$, so $X_0 \in U$. \square

Note that “ ω -huge”, i.e., $j : V \prec M$ with $M^\lambda \subseteq M^3$, is inconsistent by Kunen’s original proof of Kunen’s Theorem, so this is the best we can do.

Definition 5.3. Let $j : M_0 \prec M_1$. Then we say that (M_α, j^α) is the α -th iterate if:

- $j^0 = j$;
- $j^{\alpha+1} = j^\alpha(j^\alpha)$;
- $M_{\alpha+1} = j^\alpha(M_\alpha)$ is well-founded;
- $j_{\gamma, \alpha+1} = j^\alpha \circ j^{\alpha-1} \circ \dots \circ j^\gamma$. So $j_{\alpha, \alpha+1} = j^\alpha$;
- if α is limit, then $(M_\alpha, j^\alpha, j_{n, \alpha})$ is the direct limit of the system $(M_\beta, j_{\beta, \gamma})$ with $\beta, \gamma < \alpha$, if it is well-founded.

We say that j is α -iterable if (M_α, j^α) exists. We say that j is iterable if it is α -iterable for any α ordinal.



It is possible for the iterate to not exist, whether because $j^\alpha(j^\alpha)$ or $j^\alpha(M_\alpha)$ are not defined, or because the model M_α is not well-founded. The direct limit of the system $(M_\beta, j_{\beta, \gamma})$ is defined as an equivalence relation on the set of (β, a) such that $a \in M_\beta$: if $\beta < \gamma$ then (β, a) is equivalent to (γ, b) iff $j_{\beta, \gamma}(a) = b$, and $[(\beta, a)]$ is in $[(\gamma, b)]$ iff $j_{\beta, \gamma}(a) \in b$. If $\gamma < \beta$, viceversa. Then $j_{\beta, \alpha}(a)$ is defined as $[(\beta, a)]$ and $j^\alpha = j_{0, \alpha}(j)$. Ambiguously, we call M_α both the domain of the model defined via the equivalence class and the \in -model equivalent to it. So if $M_1 \subseteq M_0$, then for all α such that M_α is defined we have $M_\alpha \subseteq M_0$, but it is possible that $M_0 \subseteq M_\alpha$ if seen as \in -models. It will be clear from context which

³This is not a standard definition. For different authors, “ ω -huge” can mean different things.

of the two will be used. Note that if $a \in M_\alpha$, with α limit, then it must be in the range of some $j_{\beta,\alpha}$, with $\beta < \alpha$.

So let $j : V_\lambda \prec V_\lambda$. We have already defined j^n , but

$$j^n(j^n) = j(j^{n-1})(j(j^{n-1})) = j(j^{n-1}(j^{n-1})) = j(j^n) = j^{n+1},$$

by left distributive law and induction, therefore the j^n 's are iterates of j . Moreover, $j(V_\lambda) = V_\lambda$, so the first ω -iterates of j are (V_λ, j^n) and j is finitely iterable. Could we push it any further? As all the $M_\alpha \subseteq V_\lambda$, technically there is no problem in defining it, the only decisive factor being whether M_α is well-founded.

Proposition 5.4. “There exists $j : V_\lambda \prec V_\lambda$ ω -iterable” strongly implies $I3(\lambda)^4$.

Proof. Suppose $j : V_\lambda \prec V_\lambda$ is ω -iterable, let $\langle \kappa_n : n \in \omega \rangle$ be its critical sequence. Then M_ω is the set of equivalence classes of (n, a) such that $a \in V_\lambda$, where if $n < m$ then (n, a) is equivalent to (m, b) iff $j_{n,m}(a) = b$. Also, $j_{n,\omega}(a)$ is the class of $[(n, a)]$.

We prove by induction that $j_{n,\omega}(\alpha) = \alpha$ for $\alpha \in \kappa_n$. Note that if $\beta < \kappa_n$, then for every $m > n$, $j_{n,m}(\beta) = \beta$, therefore $[(n, \beta)] = [(m, \beta)]$. So for any $m \in \omega$, $\beta < \lambda$, $[(m, \beta)] < [(n, \alpha)]$ iff $[(n, \beta)] < [(n, \alpha)]$: If $m \leq n$, then $j_{m,n}(\beta) < \alpha < \kappa_n = j_{m,n}(\kappa_m)$ therefore by elementarity $\beta < \kappa_m$ and $[(m, \beta)] = [(n, \beta)]$. If $m > n$, then $\beta < j_{n,m}(\alpha) = \alpha$, therefore also $[(n, \beta)] < [(n, \alpha)]$. So $j_{n,\omega}(\alpha) = [(n, \alpha)] = \{[(n, \beta)] : \beta < \alpha\}$, that is by induction α .

We want to calculate now $j_{0,\omega}(\kappa_0)$. If $\alpha < j_{0,\omega}(\kappa_n)$, then there exists $n \in \omega$ such that $\alpha = j_{n,\omega}(\beta)$ for some $\beta < \lambda$. But then $j_{n,\omega}(\beta) < j_{0,\omega}(\kappa_0) = j_{n,\omega}(j_{0,n}(\kappa_0)) = \kappa_n$, therefore $\alpha = j_{n,\omega}(\beta) = \beta < \lambda$. On the other hand, if $\alpha < \lambda$ then there exists $n \in \omega$ such that $\alpha < \kappa_n$, so $j_{n,\omega}(\alpha) = \alpha < j_{n,\omega}(\kappa_n) = j_{0,\omega}(\kappa_0)$. Therefore $j_{0,\omega}(\kappa_0) = \lambda$.

If $a \in V_\lambda$, then there exists $n \in \omega$ such that $a \in V_{\kappa_n}$. So $j_{n,\omega}(a) = a \in M_\omega$. Therefore $V_\lambda \subseteq M_\omega$. Let

$$T = \{ \langle e_0, \dots, e_n \rangle : \exists \langle \alpha_i : i \leq n+1 \rangle, \forall i \leq n+1 \alpha_i < \lambda, \\ \forall i \leq n \exists e_i : V_{\alpha_i} \prec V_{\alpha_{i+1}}, \forall i < n e_{i+1} \upharpoonright V_{\alpha_i} = e_i \},$$

the tree of approximations of an I3 elementary embedding. Note that $T = T^{M_\omega}$. As j is a branch of T , T is ill-founded, and by absoluteness of well-foundedness it is ill-founded also in M_ω , with the difference that a branch cannot go up to λ , as λ is regular in M_ω by elementarity. So $M_\omega \models \exists \gamma < \lambda \exists k : V_\gamma \prec V_\gamma$. So by elementarity $\exists \gamma < \kappa_0 \exists k : V_\gamma \prec V_\gamma$. \square

Proposition 5.5. If $j : V_\lambda \prec V_\lambda$ is ω_1 -iterable, then it is iterable.

Proof. This is a standard result in the theory of the iterations of elementary embeddings. Suppose that j is not iterable, and let ν be the least such that M_ν is ill-founded. Let N be a countable substructure of V_λ , let $h : N \prec V_\lambda$ indicate the embedding resulting from the collapse. Let $h(\bar{M}_0) = V_\lambda$, $h(\bar{j}) = j$, and build the iterate sequence in N . Then $h(\bar{M}_\beta) = M_{h(\beta)}$ and let $\bar{\nu} = h^{-1}(\nu)$. In particular $\bar{\nu}$ is countable. For any $\beta \leq \nu$ define h'_β so that the diagram commutes:

⁴This Proposition and Proposition ?? are in collaboration with Alessandro Andretta

$$\begin{array}{ccccccc}
V_\lambda & \cdots & M_\alpha & \xrightarrow{j_{\alpha,\beta}} & M_\beta & \xrightarrow{j_{\beta,\bar{\nu}}} & M_{\bar{\nu}} & \cdots & M_\nu \\
\uparrow h & & \uparrow h'_\alpha & & \uparrow h'_\beta & & \uparrow h'_{\bar{\nu}} & & \nearrow h \\
\bar{M}_0 & \cdots & \bar{M}_\alpha & \xrightarrow{\bar{j}_{\alpha,\beta}} & \bar{M}_\beta & \xrightarrow{\bar{j}_{\beta,\bar{\nu}}} & \bar{M}_{\bar{\nu}} & &
\end{array}$$

Let $h'_0 = h \upharpoonright \bar{M}_0$. If β is a limit ordinal, then every $x \in \bar{M}_\beta$ is some $\bar{j}_{\alpha,\beta}(y)$, with $\alpha < \beta$ and $y \in \bar{M}_\alpha$. Then $h'_\beta(x) = j_{\alpha,\beta}(h'_\alpha(y))$. If $\beta = \gamma + 1$, then by elementarity, as $j : V_\lambda \prec V_\lambda$, $\bar{M}_\beta = \bar{M}_\gamma$ and $M_\beta = M_\gamma$, so $h'_\beta = h'_\gamma$.

But then $h'_{\bar{\nu}} : \bar{M}_{\bar{\nu}} \prec M_{\bar{\nu}}$. As M_ν is illfounded, then by elementarity also $M_{\bar{\nu}}$ is illfounded, and since ν was the least and $\bar{\nu}$ is countable, $\nu = \bar{\nu}$ is countable and j is not ω_1 -iterable. \square

While the “ ω -hugeness” is inconsistent, it is not immediately clear what happens with “ ω -superstrongness”:

Definition 5.6. I2 *There exists $j : V \prec M$, with $M \subseteq V$, such that $V_\lambda \subseteq M$ for some $\lambda = j(\lambda) > \text{crt}(j)$.*

Proposition 5.7. *Suppose j witnesses I2. Then λ is the supremum of its critical sequence, and $j \upharpoonright V_\lambda : V_\lambda \prec V_\lambda$.*

Proof. Let η be the supremum of the critical sequence of j . Then it must be $\eta \leq \lambda$, as η is the first fixed point of j above $\text{crt}(j)$. If $\eta + 2 \leq \lambda$ then there is a contradiction with Kunen’s Theorem, so we only have to check that $\eta + 1 \neq \lambda$, i.e., not all subsets of V_η are in M .

Suppose that $j''\eta \in M$, then define $U = \{X \subseteq \mathcal{P}(\eta) : j''\eta \in j(X)\}$ and let $i : V \prec N$ be the ultrapower via U . Then $i(\eta) = \eta$, and

$$i''\mathcal{P}(\eta) = \left\{ \bigcup_{\alpha < \eta} i(X \cap \alpha) : X \in \mathcal{P}(\eta) \right\} \in N.$$

Now, coding λ^+ as sets in $\mathcal{P}(\lambda)$, $i''\lambda^+ \in N$, and Woodin’s proof of Kunen’s theorem goes smoothly (because C is definable from $i''\lambda^+$). Contradiction.

So $j''\eta \notin M$, and η must be λ . In particular $j \upharpoonright V_\lambda : V_\lambda \prec V_\lambda$. \square

The previous Proposition says that I2 implies I3. We can do more:

Proposition 5.8. *If $j : V \prec M$ witnesses I2(λ) then $j \upharpoonright V_\lambda$ is iterable.*

Proof. Consider an iteration of $j : V \prec M$ with $V_\lambda \subseteq M$. Suppose that $(V_\lambda)^{M_\nu}$ is ill-founded, with ν the least. Let α_0 be the minimum such that $j_{0,\nu}(\alpha_0)$ is ill-founded. In other words, $V_\lambda \models \text{“}\alpha_0 \text{ is the least } \alpha \text{ such that } \exists \mu \text{ such that } j_{0,\mu}(\alpha) \text{ is ill-founded”}$. Let $\beta < j_{0,\nu}(\alpha_0)$ ill-founded. Let $\eta < \nu$ and α_1 such that $j_{\eta,\nu}(\alpha_1) = \beta$. Consider $j_{0,\eta}(\alpha_0)$. As $j_{\eta,\nu}(\alpha_1) < j_{0,\nu}(\alpha_0)$, we must have $\alpha_1 < j_{0,\eta}(\alpha_0)$. But by elementarity

$$\begin{aligned}
(V_\lambda)^{M_\eta} \models \text{“}j_{0,\eta}(\alpha_0) \text{ is the least } \alpha \text{ such that} \\
\exists \mu \text{ such that } [j_{0,\eta}(j)]_{0,\mu}(\alpha) \text{ is ill-founded”}.
\end{aligned}$$

Note that $[j_{0,\eta}(j)]_{0,\mu} = [(j_{\eta,\eta+1})]_{0,\mu} = j_{\eta,\eta+\mu}$. But then α_1 is such that, contradiction because $j_{0,\eta}(\alpha_0)$ should have been the minimum. \square

Another way to extend I3 comes from the way we characterized I3 and I1, respectively as Σ_0 - and Σ_ω -elementary embeddings from $V_{\lambda+1}$ to itself. We can consider all the intermediate steps:

Definition 5.9 (Σ_n^1 Elementary Embedding). *Let $j : V_\lambda \prec V_\lambda$. Then j is Σ_n^1 iff j^+ is a Σ_n -elementary embedding, i.e., iff for every Σ_n^1 -formula $\varphi(X)$, for every $A \subseteq V_\lambda$,*

$$V_\lambda \models \varphi(A) \leftrightarrow \varphi(j^+(A)).$$

It is not immediate, however, to see if these hypotheses are really different, and what is their relationship with I2. In fact, they are not always different: if n is odd, then j is Σ_n^1 iff it is Σ_{n+1}^1 . Moreover, I2 is the same as j is Σ_1^1 .

Theorem 5.10 (Martin). *Let $j : V_\lambda \prec V_\lambda$. Then the following are equivalent:*

- j is Σ_1^1 ;
- j^+ preserves well-founded relations (this is true for $j \upharpoonright V_\lambda$ if j witnesses I2);
- j is Σ_2^1 .

The proof is explained in great detail in [?].

Lemma 5.11 (Laver, [?]). *Let n be odd. Then “ j is Σ_n^1 ” is a Π_{n+1}^1 formula in V_λ , with j as a parameter.*

Lemma 5.12 (Square root Lemma for Σ_n^1 , Laver, [?]). *Let n be odd, $n > 1$. Let j be Σ_n^1 , $\text{crt}(j) = \kappa$. Let $\beta < \kappa$, $A, B \subseteq V_\lambda$. Then there exists a $k : V_\lambda \prec V_\lambda$ that is Σ_{n-1}^1 such that $k(k) = j$, $k(B) = j(B)$, $k(A') = A$ for some $A' \subseteq V_\lambda$ and $\beta < \text{crt}(k) < \kappa$.*

Proof. The formula “ $\exists k, Y, k$ is Σ_{n-2}^1 , $k(k) = j$, $k(B) = j(B)$, $k(Y) = A$, $\beta < \text{crt}(k) < \kappa$ ” is Σ_n^1 by Lemma ???. The following formula is clearly true:

$$\begin{aligned} j \text{ is } \Sigma_{n-2}^1, \quad j(j) = j(j), \quad j(j(B)) = j(j(B)), \\ j(A) = j(A), \quad j(\beta) = \beta < \text{crt}(j) < j(\kappa) \end{aligned}$$

but then, with a smart quantification of some of the parameters of the formula above, we have

$$\begin{aligned} \exists k, Y \quad k \text{ is } \Sigma_{n-2}^1, \quad k(k) = j(j), \quad k(j(B)) = j(j(B)), \\ k(Y) = j(A), \quad j(\beta) < \text{crt}(k) < j(\kappa) \end{aligned}$$

By elementarity the lemma is proved. \square

The lemma therefore provides a “square root” for j , with certain desirable characteristics, but with the catch that for a square root of a Σ_n^1 -elementary embedding we can only assure that it is Σ_{n-2}^1 . It is not possible to do better: suppose that κ is the minimum critical point of $j : V_\lambda \prec V_\lambda$ that is Σ_n^1 . Then the critical point of the square root k of j is less than κ , therefore k cannot be Σ_n^1 . Also note that the square root of an iterable embedding is iterable, as a square root of j is, in a certain sense, j^{-1} .

The possibility of finding a “square root” for an embedding is a central property, that will come up in different parts of the paper. The proof is always the same, it changes only the “strength” of the embedding.

Note that this would be enough to prove strong implication if we considered the existence of the embeddings as a property of the critical point, and we would have also that $I2(\kappa)$ strongly implies iterable $I3(\kappa)$. But we already argued that this seems not the right way to look at this properties. Moreover, it is possible to prove the strong implications of such axioms respect to λ :

Corollary 5.13. *$I2$ strongly implies iterable $I3$.*

Proof. Note that “ j is ω_1 -iterable”, and therefore “ j is iterable”, is Σ_1^1 (there exist $\alpha < \omega_1$, M_0, \dots, M_α , j_0, \dots, j_α such that...). Therefore one can put “ j is iterable” in ?? and the proof works. \square

Theorem 5.14 (Laver, [?]). *Let n be odd. Then if j is Σ_n^1 , it is also Σ_{n+1}^1 .*

Theorem 5.15 (Laver, [?]). *Let h and k be Σ_n^1 embeddings. Then $h \cdot k$ is a Σ_n^1 -embedding.*

Note that the converse is not true: minimizing the critical point, we can find a k that is Σ_n^1 such that any h so that $h(h) = k$ cannot be Σ_n^1 (see comment after Lemma ??).

However, if we switch application with composition, then also the converse is true.

Theorem 5.16 (Laver, [?]). *Let $h, k \in \mathcal{E}_\lambda$. Then h, k are Σ_n^1 iff $h \circ k$ is Σ_n^1 .*

Theorem ?? is promising for our objective, that is proving that being Σ_{n+2}^1 is strictly stronger than being Σ_n^1 for an elementary embedding. The most natural idea for doing this is using some sort of reflection, to prove that if there is a $j \in \mathcal{E}_\lambda$ Σ_{n+2}^1 , then there is a $k \in \mathcal{E}_{\lambda'}$ that is Σ_n^1 for some $\lambda' < \lambda$. It would be tempting to use, as in the case of iterability, direct limit of elementary embeddings, but unfortunately in this setting there are counterexamples:

Theorem 5.17 (Laver, [?]). *There exists a j that is Σ_n^1 that has a stabilizing direct limit of members of \mathcal{A}_j that is not Σ_1^1 ,*

So we will consider inverse limits instead.

Let $\langle j_0, j_1, \dots \rangle$ be a sequence of elements of \mathcal{E}_λ , and let $J = j_0 \circ j_1 \circ \dots$ be the inverse limit of the sequence. By definition the dominion of J is $\{x \in V_\lambda : \exists n_x \forall i > n_x j_i(x) = x\}$. But we know that $j_i(x) = x$ iff $x \in V_{\text{crt}(j_i)}$, so this is $\{x \in V_\lambda : \exists n_x \forall i > n_x x \in V_{\text{crt}(j_i)}\}$. That is, $x \in \text{dom } J$ depends only on the rank of x , and this implies that $\text{dom } J = V_\alpha$ for some α . It is also possible to calculate α , since $\beta < \alpha$ iff $\exists n \forall m \geq n \beta < \text{crt}(j_m)$:


$$\alpha = \sup_{n \geq 0} \inf_{m \geq n} \text{crt}(j_m) = \liminf_{n \in \omega} \text{crt}(j_n).$$

With some cosmetic change, we can also suppose that α as the supremum of the critical points, not only the limit inferior. This will also simplify the following proofs and notations.

So let $\lambda_n = \inf_{m \geq n} \text{crt}(j_m)$. Then $\alpha = \sup_{n \in \omega} \lambda_n$, and λ_n is increasing in n . If the supremum is also a maximum, we incur in the trivial case, where J is

in fact just a finite composition of elementary embeddings: if n is the first one such that $\lambda_n = \alpha$, then all $\text{crt}(j_m)$ with $m > n$ are bigger than α , so they are constant in the domain of J and they don't change anything.

Suppose then that α is a proper supremum of the λ_n sequence. We can suppose $\text{crt}(j_n) = \lambda_n$ by aggregating multiple elementary embeddings in just one: consider the largest n such that $\text{crt}(j_n) = \lambda_0$ (there will be a largest one because α is a proper supremum of the λ_n sequence), define the new k_0 as the old $j_0 \circ \dots \circ j_n$, and repeat this for every λ_n . The following is a graphical example:



The columns represent the behaviours of each j_n on λ , where the column on the left represent j_0 , and the horizontal lines indicate the critical point.

Definition 5.18. Let $J = j_0 \circ j_1 \circ \dots$. Then we define $J_n = j_n \circ j_{n+1} \circ \dots$ and $J_{0(n-1)} = j_0 \circ j_1 \circ \dots \circ j_{n-1}$.

Lemma 5.19. Let $j_m \in \mathcal{E}_\lambda$ for every $m \in \omega$, define $\alpha_m = \text{crt}(j_m)$ and suppose that for every $m \in \omega$, $\alpha_m < \alpha_{m+1}$. Let $\alpha = \sup_{m \in \omega} \alpha_m$ and $J = j_0 \circ j_1 \circ \dots$. Then

- $J''\alpha$ is unbounded in λ ;
- $J : V_\alpha \prec V_\lambda$ is elementary.

Like in the V_λ case, we can extend J to $V_{\alpha+1}$ in the expected way: when $A \subseteq V_\alpha$, $J(A) = \bigcup_{\beta < \alpha} J(A \cap V_\beta)$. Now we want to prove an equivalent of Lemma ??, but for inverse limits.

Lemma 5.20 (Square root for inverse limits of Σ_n^1). Let $J : V_\alpha \prec V_\lambda$ an inverse limit of Σ_{n+1}^1 elementary embeddings. Then for all $A, B \subseteq V_\alpha$ there exist $K : V_\alpha \prec V_\lambda$ inverse limit of Σ_n^1 elementary embeddings and $A' \subseteq V_\alpha$ such that $k(A') = A$ and $k(B) = J(B)$.

We use Lemma ?? to calculate the strength of an inverse limit:

Theorem 5.21 (Laver, [?]). If $J : V_\alpha \prec V_\lambda$ is an inverse limit of Σ_n^1 elementary embeddings, then J is Σ_n^1 .

Proof. The case $n = 0$ is Lemma ?? combined with an obvious generalization of Theorem ??, so we proceed by induction on n .

Suppose that J is Σ_{n-1}^1 , we need to prove that for every φ Π_{n-1}^1 -formula, and any $B \subseteq V_\lambda$,

$$V_\lambda \models \exists X \varphi(X, J(B)) \rightarrow V_\alpha \models \exists X \varphi(X, B).$$

Suppose $V_\lambda \models \exists X \varphi(X, J(B))$, and fix A a witness. Using Lemma ??, we find K inverse limit of Σ_{n-2}^1 elementary embeddings such that $K(A') = A$ and $K(B) = J(B)$ for some $A' \subseteq V_\alpha$. So $V_\lambda \models \varphi(K(A'), K(B))$, and by elementarity $V_\alpha \models \varphi(A', B)$, that is $V_\alpha \models \exists X \varphi(X, B)$. \square

Finally, we can prove that the existence of a Σ_{n+2}^1 elementary embedding strongly implies the existence of a Σ_n^1 elementary embedding.

Theorem 5.22 (Laver, [?]). *Let $j : V_\lambda \prec V_\lambda$ be Σ_{n+2}^1 . Then*

- *for every $B \subseteq V_\lambda$, there exist an $\alpha < \lambda$ and a $k_\alpha : V_\alpha \prec V_\lambda$ such that $k_\alpha(B_\alpha) = B$ for some $B_\alpha \subseteq V_\alpha$. In fact, we can find an ω -club $C \subseteq \lambda$ of such α 's.*
- *there exist an $\alpha < \lambda$ and a $j_\alpha : V_\alpha \prec V_\alpha$ that is Σ_n^1 . Moreover, we can find an ω -club $C \subseteq \lambda$ of such α 's. Therefore “there exists $j : V_\lambda \prec V_\lambda$ Σ_{n+2}^1 ” strongly implies “ $j : V_\lambda \prec V_\lambda$ Σ_{n+2}^1 ”. In particular I2 strongly implies I3.*

The situation for I1 has some peculiarities. First of all, if $j : V_{\lambda+1} \prec V_{\lambda+1}$, then j is definable from $j \upharpoonright V_\lambda$, so it is actually a definable class of $V_{\lambda+1}$. This is not possible in the I3 case, as $V_\lambda \models \text{ZFC}$, so it would contradict the naive Kunen's Theorem. We can reformulate Lemma ?? in this way:

Remark 5.23. *Suppose that $j : V_{\lambda+1} \rightarrow V_{\lambda+1}$. Then there are Π_{n+1} formulas $\varphi_n(x)$ such that for any $n \in \omega$, $V_{\lambda+1} \models \varphi_n(j \upharpoonright V_\lambda)$ iff j is an elementary embedding.*

So let $j : V_{\lambda+1} \prec V_{\lambda+1}$. Consider $j^- = j \upharpoonright V_\lambda$. We can define $j^-(j^-) : V_\lambda \prec V_\lambda$. By the remark above $[j^-(j^-)]^+$ is an elementary embedding, and we write it simply as $j(j)$. So we can iterate also the I1 embeddings.

As $j : V_{\lambda+1} \prec V_{\lambda+1}$ is Σ_n^1 for every n , by Lemma ?? for any n it is possible to find a square root of j that is Σ_n^1 . Therefore for any n it is possible to find $\alpha < \lambda$ and $K : V_\alpha \prec V_\lambda$ inverse limit that is Σ_n^1 , so I1 strongly implies any “There exists $j : V_\lambda \prec V_\lambda$ that is Σ_n^1 ”. We can do more:

Corollary 5.24. *I1(λ) strongly implies $\forall n \exists j : V_\lambda \prec V_\lambda$ that is Σ_n^1*

Proof. Run the proof of Theorem ?? n times, each time considering that j is Σ_n^1 . So there are C_n ω -clubs in λ such that for any $\alpha \in C_n$ there exists $k : V_\alpha \prec V_\alpha$ that is Σ_n^1 . The intersection of ω ω -clubs is an ω -club, so $\alpha \in \bigcap_{n \in \omega} C_n$ is as desired. \square

It is possible to push this kind of proof even more:

Lemma 5.25 (Square root Lemma for $L_1(V_{\lambda+1})$, Laver, [?]). *Let $j : L_1(V_{\lambda+1}) \prec L_1(V_{\lambda+1})$. For every $A, B \subseteq V_\lambda$, $\beta < \text{crt}(j)$ there exists $k : V_{\lambda+1} \prec V_{\lambda+1}$, with $\beta < \text{crt}(k) < \text{crt}(j)$, such that $k(B) = j(B)$ and there exists $A' \subseteq V_\lambda$ such that $k(A') = A$.*

Proof. The only detail we should care of is the fact that “ $j : V_{\lambda+1} \prec V_{\lambda+1}$ ” must be definable in $L_1(V_{\lambda+1})$. But this is true, because the satisfaction relation in $V_{\lambda+1}$ is definable in $L_1(V_{\lambda+1})$, so the proof is, *mutatis mutandis*, the same as Lemma ??. \square

Theorem 5.26 (Laver, [?]). “There exists $j : L_1(V_{\lambda+1}) \prec L_1(V_{\lambda+1})$ ” strongly implies “there exists $j : V_{\lambda+1} \prec V_{\lambda+1}$ ”.

Proof. The proof follows the same method of Theorem ??.

□

Note that in this case there is a fundamental difference between being Σ_n^1 for every n and being an elementary embedding from $V_{\lambda+1}$ to itself. Suppose that we have $j_i : V_{\lambda+1} \prec V_{\lambda+1}$. Then the inverse limit J of $\langle j_i : i \in \omega \rangle$ is *not* an elementary embedding from $V_{\lambda'+1}$ to $V_{\lambda+1}$, because the domain of J is just not $V_{\lambda'+1}$. Consider again the definition of the domain of J :

$$H = \{x \in V_{\lambda+1} : \exists n \in \omega \forall m \geq n \ j_m(x) = x\}.$$

Following this definition, for example, $\lambda' \notin H$, because it is moved by all j_i , while on the other hand $\lambda \in H$, because it is never moved. What we can prove with the methods provided, is that the unique extension of the inverse limits of the $j_i \upharpoonright V_\lambda$ to $V_{\lambda'+1}$ is an elementary embedding.

As I0 is equivalent to $j : L_\Theta(V_{\lambda+1}) \prec L_\Theta(V_{\lambda+1})$, we consider now hypotheses like “there exists $j : L_\alpha(V_{\lambda+1}) \prec L_\alpha(V_{\lambda+1})$ ”, $\alpha < \Theta$, with the aim of finding strong implications among them.

Note that with such aim not any α will be appropriate. We defined strong implications only between properties of λ , while “there exists $j : L_\alpha(V_{\lambda+1}) \prec L_\alpha(V_{\lambda+1})$ ” is a property of α and λ . We should consider therefore only α ’s that depend on λ , for example α ’s that are definable from λ (e.g., λ^+ , $\lambda^+ + \omega$, ...) or just absolutely definable (e.g., ω , ω^ω , ...). This kind of ordinals have also the characteristic of being a fixed point for any large enough embedding, so we already have that I0 implies “there exists $j : L_\alpha(V_{\lambda+1}) \prec L_\alpha(V_{\lambda+1})$ ”.

Our objective is now, given $j : L_\beta(V_{\lambda+1}) \prec L_\beta(V_{\lambda+1})$ to find a $\alpha < \beta$ and $\bar{\alpha}$, $\bar{\lambda} < \lambda$ such that $L_{\bar{\alpha}}(V_{\bar{\lambda}+1}) \equiv L_\alpha(V_{\lambda+1})$ and there exists $k : L_{\bar{\alpha}}(V_{\bar{\lambda}+1}) \prec L_{\bar{\alpha}}(V_{\bar{\lambda}+1})$. With β and α as above, this will imply that “exists $j : L_\beta(V_{\lambda+1}) \prec L_\beta(V_{\lambda+1})$ ” strongly implies “exists $j : L_\alpha(V_{\lambda+1}) \prec L_\alpha(V_{\lambda+1})$ ”.

The strategy is to follow Laver’s proof for strong implications as much as possible. The first step is therefore to prove some square root principle like Lemma ??. The best possible outcome would be to find for any $j : L_{\alpha+1}(V_{\lambda+1}) \prec L_{\alpha+1}(V_{\lambda+1})$ a square root k that is $k : L_\alpha(V_{\lambda+1}) \prec L_\alpha(V_{\lambda+1})$. For this, it is needed that

$$L_{\alpha+1}(V_{\lambda+1}) \models \exists k : L_\alpha(V_{\lambda+1}) \prec L_\alpha(V_{\lambda+1}),$$

i.e., that $j \upharpoonright L_\alpha(V_{\lambda+1}) \in L_{\alpha+1}(V_{\lambda+1})$. Can we assume that?

The next step would be to define an inverse limit of embeddings at the $L_\alpha(V_{\lambda+1})$ level. Here another problem arises following the remarks after Theorem ??: the domain inverse limit of the $k_i : L_\alpha(V_{\lambda+1}) \prec L_\alpha(V_{\lambda+1})$ is not some $L_{\bar{\alpha}}(V_{\bar{\lambda}+1})$, in fact it does not even contain $V_{\bar{\lambda}+1}$, therefore we should do the inverse limit of $k_i \upharpoonright V_\lambda$, and then extend it to some $L_{\bar{\alpha}}(V_{\bar{\lambda}+1})$. How to do it?

Both problems were solved with the study of the structure of the sets $L_\alpha(V_{\lambda+1})$ for $\alpha < \Theta$ made by Laver in [?].

Definition 5.27 (Laver, [?]). Let λ be a cardinal and let $\alpha < \Theta$. Then α is good iff every element of $L_\alpha(V_{\lambda+1})$ is definable in $L_\alpha(V_{\lambda+1})$ from an element in $V_{\lambda+1}$.

The successor of a good ordinal is a good ordinal: let α be good; the largest ordinal in $L_{\alpha+1}(V_{\lambda+1})$ is $\lambda + 1 + \alpha$, therefore α is definable in $L_{\alpha+1}(V_{\lambda+1})$ with λ as a parameter, so $L_\alpha(V_{\lambda+1})$ is definable in $L_{\alpha+1}(V_{\lambda+1})$ with λ as a parameter. But every element in $L_{\alpha+1}(V_{\lambda+1})$ is definable using $L_\alpha(V_{\lambda+1})$ and elements of $L_\alpha(V_{\lambda+1})$ (which in turn are definable with parameters from $V_{\lambda+1}$), and therefore $\alpha + 1$ is good. This proves that the natural numbers are good. But also ω is good: every element of $L_\omega(V_{\lambda+1})$ is in some $L_n(V_{\lambda+1})$, $n \in V_{\lambda+1}$ and n is good for every n . Following the same line of reasoning, every ordinal up to λ is good, and considering that all ordinals less than λ^+ are coded as subsets of λ , and therefore in $V_{\lambda+1}$, every ordinal up to λ^+ is good.

On the other side, non-good ordinals exist. The definition of good ordinal is restricted to ordinals strictly less than Θ because larger ones are trivially not good: if x is definable with a parameter, then it is uniquely determined by its definition, therefore for any α there exists in $L(V_{\lambda+1})$,

$$\pi : V_{\lambda+1} \rightarrow G_\alpha = \{x \in L_\alpha(V_{\lambda+1}) : L_\alpha(V_{\lambda+1}) \models x \text{ is definable from an element in } V_{\lambda+1}\};$$

then if $\Theta \subseteq L_\alpha(V_{\lambda+1})$, G_α must be strictly contained in $L_\alpha(V_{\lambda+1})$, by definition of Θ . But non-good ordinals exist also below Θ : define $L_\gamma(V_{\lambda+1})$ as the collapse of the Skolem closure of $V_{\lambda+1}$ in $L_\Theta(V_{\lambda+1})$; as $L_\Theta(V_{\lambda+1}) \models \exists x x$ not definable from an element in $V_{\lambda+1}$, by elementarity the same must be true in $L_\gamma(V_{\lambda+1})$, as the collapse does not collapse $V_{\lambda+1}$, therefore γ is not good.

One can ask how many good ordinals there are.

Lemma 5.28 (Laver, [?]). *Let λ be a strong limit cardinal. Then the good ordinals are unbounded in Θ .*

Therefore assuming α good is in most cases a reasonable choice.

By the usual condensation argument, if $\alpha < \Theta$ and $i : L_\alpha(V_{\lambda+1}) \rightarrow L_\alpha(V_{\lambda+1})$ then $i \in L_\Theta(V_{\lambda+1})$. If α is good, however, it is possible to be much more precise:

Lemma 5.29 (Laver, Woodin, [?],[?]). *Let λ and α be such that α is good and there exists $i : L_\alpha(V_{\lambda+1}) \prec L_\alpha(V_{\lambda+1})$ with $\text{crt}(i) < \lambda$. Then i is induced by $i \upharpoonright V_\lambda$, and therefore $i \in L_{\alpha+1}(V_{\lambda+1})$.*

Finally:

Lemma 5.30 (Square root Lemma for $L_\alpha(V_{\lambda+1})$). *Let α be good, and let $j : L_{\alpha+1}(V_{\lambda+1}) \prec L_{\alpha+1}(V_{\lambda+1})$. Then for any $A, B \in L_\alpha(V_{\lambda+1})$, for any $\beta < \text{crt}(j)$ there exists $k : L_\alpha(V_{\lambda+1}) \prec L_\alpha(V_{\lambda+1})$ such that $\beta < \text{crt}(k) < \text{crt}(j)$, $k(k \upharpoonright V_\lambda) = j \upharpoonright V_\lambda$, $k(B) = j(B)$ and there exists $A \in \text{ran}(k)$.*

Proof. Exactly as Lemma ???. The key point is that as α is good, “ $\exists k k : L_\alpha(V_{\lambda+1}) \prec L_\alpha(V_{\lambda+1})$ ” is expressible in $L_{\alpha+1}(V_{\lambda+1})$ and it is satisfied by $j \upharpoonright L_\alpha(V_{\lambda+1})$, so it can be put inside the formula instead of “ j is Σ_{n-2}^1 ”. \square

Note that if α is not good, then we can have a similar result: for $j : L_\beta(V_{\lambda+1}) \prec L_\beta(V_{\lambda+1})$, let α be maximum such that $j \upharpoonright L_\alpha(V_{\lambda+1}) \in L_\beta(V_{\lambda+1})$. Then we can find a square root that extend to $L_\alpha(V_{\lambda+1})$. The difference is that in this case the gap can be large, and it is somewhat arbitrary, while for the good case is just one.

Moreover, I0 implies “ $\exists j : L_\alpha(V_{\lambda+1}) \prec L_\alpha(V_{\lambda+1})$ ”:

Lemma 5.31 (Woodin, [?]). *Let λ be a cardinal. If there exists $j : L_\Theta(V_{\lambda+1}) \prec L_\Theta(V_{\lambda+1})$, then for any $\alpha < \Theta$ there exists an $i : L_\alpha(V_{\lambda+1}) \prec L_\alpha(V_{\lambda+1})$.*

Proof. Suppose it is false. Then there is a counterexample α such that every $i : L_\alpha(V_{\lambda+1}) \rightarrow L_\alpha(V_{\lambda+1})$ is not an elementary embedding. All such i 's are in $L_\Theta(V_{\lambda+1})$ (see remark before Lemma ??), therefore $L_\Theta(V_{\lambda+1}) \models \exists \alpha$ (α is a counterexample). Let α_0 be the least counterexample. Then α_0 is definable in $L_\Theta(V_{\lambda+1})$ and $j(\alpha_0) = \alpha_0$. Then $j \upharpoonright L_{\alpha_0}(V_{\lambda+1})$ is as in the lemma, contradiction. \square

Compare this lemma with the fact that $U_j \notin L(V_{\lambda+1})$, and therefore $j \upharpoonright L_\Theta(V_{\lambda+1}) \notin L(V_{\lambda+1})$. Then if $j(\alpha) = \alpha$ and

$$j \upharpoonright L_\alpha(V_{\lambda+1}) : L_\alpha(V_{\lambda+1}) \prec L_\alpha(V_{\lambda+1}),$$

for any $\beta > \alpha$ there must be a j_1 elementary embedding with domain $L_\beta(V_{\lambda+1})$ and such that $j_1 \neq j \upharpoonright L_\beta(V_{\lambda+1})$, otherwise one could build $j \upharpoonright L_\Theta(V_{\lambda+1})$ inside $L(V_{\lambda+1})$. In other words, the tree of embeddings with domain of the type $L_\alpha(V_{\lambda+1})$ has levels of “cardinality” $< \Theta$ (by ??), has height Θ and there is no cofinal branch in $L(V_{\lambda+1})$. In a certain sense, it is a Θ -Aronszajn tree in $L(V_{\lambda+1})$.

Now we can do the inverse limit like in Theorem ??, and so we have $J : V_{\bar{\lambda}} \prec V_{\bar{\lambda}}$, inverse limit of embeddings that can be extended to $L_\alpha(V_{\lambda+1})$. With $\bar{\alpha}$ fixed, we can extend J to $L_{\bar{\alpha}}(V_{\bar{\lambda}+1})$. But how to find a $\bar{\alpha}$ so that J is an elementary embedding?

This point needs more work, and has been solved by Scott Cramer in his PhD Thesis. The idea is always to use ?? repeatedly, to build inverse limits that have in the range witnesses for existential formulas, so that in the end the inverse limit is elementary. The difference is that we need λ different witnesses to be considered every time, so the plan is to build a forcing that at the same time constructs a limit of inverse limits and collapses λ to ω . We follow with a sketch of the ideas behind such a construction, for the details we refer the reader to [?].

The structure of the proof of Theorem ?? is like this:

$$\begin{array}{ccccccc} j \upharpoonright V_{\bar{\lambda}} & j \upharpoonright V_{\bar{\lambda}} & j \upharpoonright V_{\bar{\lambda}} & j \upharpoonright V_{\bar{\lambda}} & \dots \\ K : & k_0 & k_1 & k_2 & k_3 & \dots \end{array}$$

where each embedding below is a square root of the embedding above. We have some freedom for the critical points in the embedding belows, in that for each one we can say that it must be bigger than a certain fixed ordinal, and (optionally) less than $\text{crt}(j)$. In this way, we can push the limit of the critical points of the k_i 's to be λ , so that the inverse limit will be from $V_{\bar{\lambda}}$ to itself, or less than $\text{crt}(j)$, as in Lemma ??, so to have $J : V_{\bar{\lambda}} \prec V_{\bar{\lambda}}$. So, by Lemma ??, the existence of $j : L_{\alpha+1}(V_{\lambda+1}) \prec L_{\alpha+1}(V_{\lambda+1})$ implies the existence of an inverse limit J of square roots of $j \upharpoonright V_{\bar{\lambda}}$ such that $J : V_{\bar{\lambda}} \prec V_{\bar{\lambda}}$ (possibly with $\bar{\lambda} = \lambda$), and all the square roots can be extended to $L_\alpha(V_{\lambda+1})$.

Instead of j , we can start already with an inverse limit:

$$J: \quad j_0 \quad j_1 \quad j_2 \quad j_3 \quad \dots$$

$$K: \quad k_0 \quad k_1 \quad k_2 \quad k_3 \quad \dots$$

We say that K (the inverse limit of the k_i 's) is a *limit root* of J (the inverse limit of the j_i 's) iff there exists an $n \in \omega$ such that $k_i = j_i$ for $i < n$ and k_i is a square root of j_i for $i \geq n$. So for any inverse limit J of embeddings that extend to $L_{\alpha+1}(V_{\lambda+1})$ there exist a limit root of J of embeddings that extend to $L_\alpha(V_{\lambda+1})$. Moreover, by ??, if $J: V_{\bar{\lambda}} \prec V_\lambda$, for any $B \in V_{\bar{\lambda}+1}$ and any $A \in V_{\lambda+1}$ we can have K as before such that $K: V_{\bar{\lambda}} \prec V_\lambda$, $K(B) = J(B)$ and $A \in \text{ran}(K)$.

The next step is to build a limit of inverse limits. Let j be an embedding that extends to $L_{\alpha+\omega}(V_{\lambda+1})$. Then consider something like this:

$$\begin{array}{ccccccc} & j & j & j & j & \dots \\ J: & j_0 & j_1 & j_2 & j_3 & \dots \\ K_1 & j_0 & k_1^1 & k_2^1 & k_3^1 & \dots \\ K_2 & j_0 & k_1^1 & k_2^1 & k_3^2 & \dots \\ & \vdots & \vdots & \vdots & \vdots & \dots \end{array}$$

where J is an inverse limit of square roots of j , and every K_{i+1} is a limit root of K_i . We can choose the k_m^n 's so that they all extend to $L_\alpha(V_{\lambda+1})$: Let n_i be an increasing sequence of integers. Then choose j_i to be a square root of j extendible to $L_{\alpha+n_i}(V_{\lambda+1})$. Once defined k_i^m , if it is extendible to some $L_{\alpha+n}(V_{\lambda+1})$ with $n > 0$, then choose k_i^{m+1} to be a square root of k_i^m that is extendible to $L_{\alpha+n-1}(V_{\lambda+1})$, otherwise $k_i^{m+1} = k_i^m$. As n_i was increasing, the last case can happen only a finite amount of times, so K_{m+1} will always be a limit square root of K_m . Also, at each stage, for any $A_m \subseteq V_{\bar{\lambda}}$ (where $V_{\bar{\lambda}}$) is the domain of J) and any $B_m \subseteq V_\lambda$, we can have $K^{m+1}(A) = K^m(A)$ and $B \in \text{ran}(k_i^{m+1} \circ k_{i+1}^{m+1} \circ \dots)$, where i is the minimum such that $k_i^m \neq k_i^{m+1}$. Note that this is less than a Square root Lemma for inverse limits for $L_\alpha(V_{\lambda+1})$, as we ask only B to be in the range of a tail of the limit, and not of K . But this will be enough.

Theorem 5.32 (Cramer, 2015 [?]). *Let α be good, and let $j: L_{\alpha+\omega}(V_{\lambda+1}) \prec L_{\alpha+\omega}(V_{\lambda+1})$. Let J be as the first step of the construction above, with $V_{\bar{\lambda}}$ its domain. Then there exist $\bar{\alpha}$, $\bar{\lambda} < \lambda$ such that $L_{\bar{\alpha}}(V_{\bar{\lambda}+1}) \equiv L_\alpha(V_{\lambda+1})$.*

Theorem ?? gives us finally the right $\bar{\alpha}$ where to extend J , the inverse limit of embeddings that can be extended to $L_\alpha(V_{\lambda+1})$ in the most natural way: as $\bar{\alpha}$ is $(\bar{\lambda}-)$ good, then every element $b \in L_{\bar{\alpha}}(V_{\bar{\lambda}+1})$ is defined from some parameter $a \in V_{\bar{\lambda}+1}$, and $J(b)$ is the element of $L_\alpha(V_{\lambda+1})$ that is defined with the same formula and parameter $J(a)$. It remains to see that J is actually an elementary embedding. Theorem ?? gives us an elementary embedding, J_* , but it is in $V[G]$. But if we consider the partial order of all possible sequences of K_i 's, that start with J , coupled with the collapse, then J_* will be the common part of the generic of such forcing. Considering J as a sequence of length 1, for any $b \in L_{\bar{\alpha}}(V_{\bar{\lambda}+1})$ we can always extend J so that $J^*(b) = J(b)$, and then elementarity comes from absoluteness. All the details are in [?].

Theorem 5.33 (Cramer, 2015, [?]). *If $j : L_{\alpha+\omega}(V_{\lambda+1}) \prec L_{\alpha+\omega}(V_{\lambda+1})$, then there exist $\bar{\alpha}, \bar{\lambda} < \lambda$ and $k : L_{\bar{\alpha}}(V_{\bar{\lambda}}) \prec L_{\bar{\alpha}}(V_{\bar{\lambda}})$. In particular, if α depends only on λ , then “ $\exists j : L_{\alpha+\omega}(V_{\lambda+1}) \prec L_{\alpha+\omega}(V_{\lambda+1})$ ” strongly implies “ $\exists j : L_{\alpha}(V_{\lambda+1}) \prec L_{\alpha}(V_{\lambda+1})$ ”.*

So all the following are strong implications:

$$\begin{aligned} I0(\lambda) &\Rightarrow \dots \Rightarrow \exists j : L_{\alpha+\omega}(V_{\lambda+1}) \prec L_{\alpha+\omega}(V_{\lambda+1}) \Rightarrow \exists j : L_{\alpha}(V_{\lambda+1}) \prec L_{\alpha}(V_{\lambda+1}) \Rightarrow \\ &\Rightarrow \dots \Rightarrow \exists j : L_{\omega}(V_{\lambda+1}) \prec L_{\omega}(V_{\lambda+1}) \Rightarrow \exists j : L_1(V_{\lambda+1}) \prec L_1(V_{\lambda+1}) \Rightarrow \\ &\Rightarrow I1(\lambda) \equiv \exists j : V_{\lambda} \prec V_{\lambda} \Sigma_{\omega}^1 \Rightarrow \forall n \exists j : V_{\lambda} \prec V_{\lambda} \Sigma_n^1 \Rightarrow \dots \\ &\dots \Rightarrow \exists j : V_{\lambda} \prec V_{\lambda} \Sigma_n^1 \equiv \exists j : V_{\lambda} \prec V_{\lambda} \Sigma_{n+1}^1 \text{ (} n \text{ is odd)} \Rightarrow \dots \\ &\dots \Rightarrow \exists j : V_{\lambda} \prec V_{\lambda} \Sigma_2^1 \equiv j : V_{\lambda} \prec V_{\lambda} \Sigma_1^1 \equiv I2(\lambda) \Rightarrow \text{iterable } I3(\lambda) \Rightarrow I3(\lambda). \end{aligned}$$

6 Similarities with $\text{AD}^{L(\mathbb{R})}$

We already noticed that for $I0(\lambda)$ to hold it is necessary that $L(V_{\lambda+1}) \not\models \text{AC}$, just like for AD to hold in $L(\mathbb{R})$ it is necessary that $L(\mathbb{R}) \not\models \text{AC}$. This is the first sign of something bigger. The research is still ongoing on this respect, but there are many results that indicate a strong similarity between $L(V_{\lambda+1})$ under $I0$ and $L(\mathbb{R})$ under AD . On the other hand, there are also striking differences, so there need to be further work on how much the two models are similar and why (and why they are not too similar).

The first similarity will be the following:

Theorem 6.1 (Solovay). *Suppose $L(\mathbb{R}) \models \text{AD}$. Then $L(\mathbb{R}) \models \omega_1$ is measurable.*

Lemma 6.2 (Woodin, [?]). *Suppose that there exists $j : L(V_{\lambda+1}) \prec L(V_{\lambda+1})$ with $\text{crt}(j) < \lambda$. Then for every $\delta < \lambda$ regular, define*

$$S_{\delta}^{\lambda^+} = \{\eta < \lambda^+ : \text{cof}(\eta) = \delta\},$$

the stationary set of ordinals with the same cofinality and let \mathcal{F} be the club filter on λ^+ . Then there exists $\eta < \lambda$ and $\langle S_{\alpha} : \alpha < \eta \rangle \in L(V_{\lambda+1})$ a partition of $S_{\delta}^{\lambda^+}$ such that for every $\alpha < \eta$ $\mathcal{F} \upharpoonright S_{\alpha}$ is a $L(V_{\lambda+1})$ -ultrafilter in λ^+ . In particular λ^+ is measurable.

It is remarkable that the proof is completely different, yet in both theorems the measure is not just any measure, but it comes from the club filter. Woodin calls the cardinals that are measurable for this reason “ ω -strongly measurable cardinals”, and they have a key role in the development of the inner model descriptive set theory.

The fact that λ^+ is measurable has a lot of $\text{AD}^{\mathbb{R}}$ -like consequences:

Theorem 6.3 (Shi, Trang, 2017 [?]). *Suppose $I0(\lambda)$ holds. Then in $L(V_{\lambda+1})$:*

- *there is no λ^+ -Aronszajn tree;*
- \square_{λ} *fails;*
- *there is no scale at λ in $V_{\lambda+1}$;*

- \diamond_{λ^+} fails;
- there is no λ^+ -sequence of distinct members of $\mathcal{P}(\lambda)$.

In fact Theorem ?? was just the first ω -strongly measurable cardinal found in $L(\mathbb{R})$, when there is a rich structure of them under AD. The same is true in $L(V_{\lambda+1})$ under I0. Moschovakis' Coding Lemma is the way to find them:

Theorem 6.4 (Coding Lemma, Moschovakis). *Suppose $L(\mathbb{R}) \models \text{AD}$. Let $\eta < \Theta$ and $\rho : \mathbb{R} \twoheadrightarrow \eta$, $\rho \in L(\mathbb{R})$. Then there exists $\gamma_\rho < \Theta$ such that for every $A \subseteq \mathbb{R} \times \mathbb{R}$, $A \in L(\mathbb{R})$, there exists $B \subseteq \mathbb{R} \times \mathbb{R}$ such that:*

- $B \subseteq A$;
- $B \in L_{\gamma_\rho}(\mathbb{R})$;
- for every $\alpha < \eta$ if $\exists(a, b) \in A$ $\rho(a) = \alpha$ then $\exists(a, b) \in B$ $\rho(a) = \alpha$.

It is a sort of very weak choice principle for subsets of \mathbb{R} : Consider $A_\xi = \{b : (a, b) \in A, \rho(a) = \xi\}$. Thus A can be seen as an η -sequence of subsets of \mathbb{R} . If we had the Axiom of Choice, we could have a selector for such a sequence. Instead the Coding Lemma gives us a sequence $B_\xi \subseteq A_\xi$ that is in some $L_{\gamma_\rho}(\mathbb{R})$. But however “complex” (i.e., high in the constructive hierarchy) we choose A , γ_ρ is the same, so the “selector” is actually simpler than A .

Theorem 6.5 (Coding Lemma, Woodin, [?]). *Suppose that there exists $j : L(V_{\lambda+1}) \prec L(V_{\lambda+1})$ with $\text{crt}(j) < \lambda$. Let $\eta < \Theta$ and $\rho : V_{\lambda+1} \twoheadrightarrow \eta$, $\rho \in L(V_{\lambda+1})$. Then there exists $\gamma_\rho < \Theta$ such that for every $A \subseteq V_{\lambda+1} \times V_{\lambda+1}$, $A \in L(V_{\lambda+1})$, there exists $B \subseteq V_{\lambda+1} \times V_{\lambda+1}$ such that:*

- $B \subseteq A$;
- $B \in L_{\gamma_\rho}(V_{\lambda+1})$;
- for every $\alpha < \eta$ if $\exists(a, b) \in A$ $\rho(a) = \alpha$ then $\exists(a, b) \in B$ $\rho(a) = \alpha$.

The proof of the Coding Lemma is in [?], with a further generalization. The Coding Lemma is used to prove a sort of inaccessibility of Θ in $L(V_{\lambda+1})$: as $L(V_{\lambda+1}) \not\models \text{AC}$, we cannot say $2^\gamma < \Theta$ for all $\gamma < \Theta$; but we have the following:

Lemma 6.6 (Woodin, [?]). *Suppose that there exists $j : L(V_{\lambda+1}) \prec L(V_{\lambda+1})$ with $\text{crt}(j) < \lambda$. Then in $L(V_{\lambda+1})$ for every $\alpha < \Theta$ there exists a surjection $\pi : V_{\lambda+1} \twoheadrightarrow \mathcal{P}(\alpha)$.*

Proof. Fix $\alpha < \Theta$, $\rho : V_{\lambda+1} \twoheadrightarrow \alpha$, and let γ be the maximum between the least $\beta < \Theta$ such that $\rho \in L_\beta(V_{\lambda+1})$ and the witness for the Coding Lemma for α, ρ . For every $A \subseteq \alpha$ define $A^* = \{(a, 0) \in V_{\lambda+1} : \rho(a) \in A\}$. Then $A^* \in L(V_{\lambda+1})$. So there exists $B^* \subseteq A^*$, $B^* \in L_\gamma(V_{\lambda+1})$ such that $\{\rho(a) : \exists b (a, b) \in B^*\} = A$. But then $A \in L_{\gamma+1}(V_{\lambda+1})$. This means that $\mathcal{P}(\alpha) \subseteq L_{\gamma+1}(V_{\lambda+1})$, and using $\pi : V_{\lambda+1} \twoheadrightarrow \gamma + 1$ we're done. \square

The following theorem is akin to a similar one in $\text{AD}^{L(\mathbb{R})}$ (cfr. with [?]):

Theorem 6.7 (Woodin, [?]). *Suppose that there exists $j : L(V_{\lambda+1}) \prec L(V_{\lambda+1})$ with $\text{crt}(j) = \kappa_0 < \lambda$. Then Θ is a limit of γ such that:*

1. γ is weakly inaccessible in $L(V_{\lambda+1})$;
2. $\gamma = \Theta^{L_\gamma(V_{\lambda+1})}$ and $j(\gamma) = \gamma$;
3. for all $\beta < \gamma$, $\mathcal{P}(\beta) \cap L(V_{\lambda+1}) \in L_\gamma(V_{\lambda+1})$;
4. for cofinally $\kappa < \gamma$, κ is an ω -strongly measurable cardinal in $L(V_{\lambda+1})$;
5. $L_\gamma(V_{\lambda+1}) \prec L_\Theta(V_{\lambda+1})$.

Proof. Fix $\alpha < \Theta$, $j(\alpha) = \alpha$ and let $\gamma < \Theta$ be the minimum such that $L_\gamma(V_{\lambda+1}) \prec_1 L_\Theta(V_{\lambda+1})$. As also $j(\Theta) = \Theta$, we have $j(\gamma) = \gamma$. This will be the γ as in the Theorem. By Corollary ?? we can have γ as large as we want below Θ .

It is tempting to stop here, and say that γ has the same properties of Θ by elementarity, but we do not have this directly, as $\Theta \notin L_\Theta(V_{\lambda+1})$.

Note that, by ?? relativized to $\Sigma_1(\alpha)$ formulas, $\Delta_1^2(\alpha)$ sets, etc. \dots , $\Theta^{L_\gamma(V_{\lambda+1})} = \gamma$, and therefore (2). Also, $\text{cof}(\delta) > \lambda$: Otherwise let $\langle \beta_\xi : \xi < \mu \rangle$ be a sequence cofinal in δ with $\mu < \lambda$. Then any β_ξ is the length of a $\Delta_1^2(\alpha)$ pwo. With DC_λ choose one pwo for any ξ , and then gluing this pwos together it is possible to find a $\Delta_1^2(\alpha)$ pwo of γ , and this contradicts ?. Now the Weak Coding Lemma can be re-proved in $L_\gamma(V_{\lambda+1})$: the key points are that since $\Theta^{L_\gamma(V_{\lambda+1})} = \gamma$, $\gamma_{\xi+1} < \gamma$ and since $\text{cof}(\gamma) > \lambda$ the sequence of the Z_ξ is longer than λ , and we can find the contradiction.

With the Weak Coding Lemma we prove that γ is regular. It is similar to the proof that its cofinality is larger than λ , using the Weak Coding Lemma instead of DC_λ : Let $\beta < \gamma$ and $f : \beta \rightarrow \gamma$ cofinal in γ . Let π be a $\Sigma_1(\alpha)$ partial bounded surjection from $V_{\lambda+1}$ to $L_\gamma(V_{\lambda+1})$, as in Lemma ?. As $\Theta^{L_\gamma(V_{\lambda+1})} = \gamma$, there exists $\rho : V_{\lambda+1} \twoheadrightarrow \beta$ with $\rho \in L_\gamma(V_{\lambda+1})$. Let δ witness the Weak Coding Lemma for ρ in $L_\gamma(V_{\lambda+1})$, and consider

$$A = \{(a, b) : b \in \text{dom}(\pi) \wedge \pi(b) = f(\rho(a))\}.$$

Then there exists $B \subseteq A$, $B \in L_\delta(V_{\lambda+1})$ such that for cofinally $\xi < \beta$ if there exists $(a, b) \in A$ $\rho(a) = \xi$ then there exists $(a, b) \in B$ such that $\rho(a) = \xi$. As for every ξ , $A_\xi \neq \emptyset$, this means that for cofinally ξ , $B_\xi \neq \emptyset$. Let $B^0 = \{a : \exists b (a, b) \in B\}$. Then $\pi'' B^0$ is cofinal in β , so $(f \circ \pi)'' B^0$ is cofinal in γ . Let $Z = \{b : \exists a \in V_{\lambda+1} (a, b) \in B\}$. As $B \subseteq A$, $\rho'' Z = (f \circ \pi)'' B^0$, but that contradicts the boundedness of ρ .

Since γ is regular, then the Coding Lemma can be re-proved relativized to $L_\gamma(V_{\lambda+1})$, so also Lemma ?? holds in $L_\gamma(V_{\lambda+1})$, therefore (3) is proved. So γ is weakly inaccessible because of regularity and (3).

Let $S_\omega^\gamma = \{\alpha < \gamma : \text{cof}(\alpha) = \omega\}$. Let \mathcal{F} be the club filter on S_ω^γ . Instead of proving that \mathcal{F} is an ultrafilter on a stationary set, we define another filter $\mathcal{F}_0 \subseteq \mathcal{F}$, that is an ultrafilter on a stationary set, and that coincides with \mathcal{F} there.

Let $\mathcal{E} = \{k : L_\gamma(V_{\lambda+1}) \prec L_\gamma(V_{\lambda+1})\}$. For each $k \in \mathcal{E}$, let $C_k = \{\eta \in S_\omega^\gamma : k(\eta) = \eta\}$. As γ is regular, C_k is an ω -club. For each $\sigma \subseteq \mathcal{E}$ such that $|\sigma| \leq \lambda$, let $C_\sigma = \cap \{C_k : k \in \sigma\}$. Let \mathcal{F}_0 be the filter on S_ω^γ generated by the sets $C_\sigma \cap S_\omega^\gamma$. It is λ^+ -complete by DC_λ . All its elements are clubs on S_ω^γ , so $\mathcal{F}_0 \subseteq \mathcal{F}$. Also, $j(\mathcal{F}_0) = \mathcal{F}_0$.

As in ?? S cannot be partitioned in κ_0 many \mathcal{F}_0 -positive sets, since $j \restriction L_\gamma(V_{\lambda+1}) \in \mathcal{E}$ and therefore $C_j \in \mathcal{F}_0$. So there must exist a $T \subseteq S$ stationary such that $\mathcal{F}_0 \restriction T$ is an ultrafilter. As it is a maximal filter on T , it must be that $\mathcal{F} \restriction T = \mathcal{F}_0 \restriction T$. It remains to show that \mathcal{F} on T is γ -complete.

Let $\beta < \gamma$ and $\langle A_\xi : \xi < \beta \rangle$ a sequence of subsets of T of \mathcal{F} -measure one. Let $\rho : V_{\lambda+1} \twoheadrightarrow \beta$, with $\rho \in L_\gamma(V_{\lambda+1})$. Let η witness the Coding Lemma in $L_\gamma(V_{\lambda+1})$ for ρ . Ideally, we would like to use the Coding Lemma on

$$A = \{(A_\xi, C_\sigma) : \sigma \subseteq \mathcal{E}, |\sigma| \leq \lambda, \xi < \beta, C_\sigma \subseteq A_\xi\},$$

to “choose” some C_σ inside every A_ξ , so that we can intersect them in the following way: Suppose we can find $B \subseteq A$ “small” enough, so that, if

$$\mathcal{E}_B = \{k \in \mathcal{E} : \exists \xi \exists \sigma \xi < \beta, \sigma \subseteq \mathcal{E}, |\sigma| \leq \lambda, (A_\xi, C_\sigma) \in B \mid k \in \sigma\},$$

for any $\xi < \gamma$, $\sup\{k(\xi) : k \in \mathcal{E}_B\} < \gamma$. Then pick $\xi_0 > \alpha$, $\xi_{n+1} = \sup\{k(\xi_n) : k \in \mathcal{E}_B\}$. Then $\xi_\omega = \sup_{n \in \omega} \xi_n \in T$ and for any $k \in \mathcal{E}_B$ $k(\xi_\omega) = \xi_\omega$. This means that for any A_ξ , $\xi_\omega \in A_\xi$, so we found an intersection.

Therefore we have to code A as a subset of $V_{\lambda+1} \times V_{\lambda+1}$: Note that γ is good, therefore by ?? for any $k_1, k_2 \in \mathcal{E}$, $k_1 = k_2$ iff $k_1 \restriction V_\lambda = k_2 \restriction V_\lambda$, so there is a bijection between \mathcal{E} and $\mathcal{E}_0 = \{k \restriction V_\lambda : k \in \mathcal{E}\} \subseteq V_{\lambda+1}$. Moreover let F be any bijection between $V_{\lambda+1}$ and ${}^\lambda V_{\lambda+1}$: for $a \in V_{\lambda+1}$ we can define $C_a = C_{\sigma(a)}$ with $\sigma(a) = \{k : k \restriction V_\lambda \in F(a)\}$.

Let

$$A = \{(a, b) \in V_{\lambda+1} \times V_{\lambda+1} : F(b) \subseteq \mathcal{E}_0, C_b \subseteq A_{\rho(a)}\}.$$

Let $B \subseteq A$, $B \in L_\eta(V_{\lambda+1})$ given by the Coding Lemma. Let

$$\mathcal{E}_B = \cup\{F(b) : \exists a (a, b) \in B\} \in L_{\eta+1}(V_{\lambda+1}),$$

a set of codes for embeddings k such that $C_k \subseteq A_\xi$ for some ξ . Let $a \in \text{dom}(\pi)$. Note that, as π is $\Sigma_1(\alpha)$ -definable, if $a \in \text{dom}(\pi)$ then $k(a) \in \text{dom}(\pi)$ for any k embedding, so

$$W_a = \{k(a) : k \in \mathcal{E}, k \restriction V_\lambda \in \mathcal{E}_B\} \subseteq \text{dom}(\pi),$$

and $W_a \in L_{\eta+2}(V_{\lambda+1})$. So

$$\pi'' W_a = \{\pi(k(a)) : k \in \mathcal{E}, k \restriction V_\lambda \in \mathcal{E}_B\} = \{k(\pi(a)) : k \in \mathcal{E}, k \restriction V_\lambda \in \mathcal{E}_B\}.$$

Let $\xi < \gamma$, and $a \in V_{\lambda+1}$ such that $\pi(a) = \xi$. So $\pi'' W_a = \{k(\xi) : k \in \mathcal{E}, k \restriction V_\lambda \in \mathcal{E}_B\}$. As W_a is bounded, then $\pi'' W_a$ is bounded, so $\sup\{k(\xi) : k \in \mathcal{E}, k \restriction V_\lambda \in \mathcal{E}_B\} < \gamma$. Now we can find a fixed point for all the extension of embeddings in \mathcal{E}_B , and we are done.

For the point (5) we just sketch the proof. The idea is to consider P , the theory of $L_\Theta(V_{\lambda+1})$, and relativize the previous proof for $L_\gamma(V_{\lambda+1})[P]$. Then

$$(L_\gamma(V_{\lambda+1})[P], P \cap L_\gamma(V_{\lambda+1})) \prec_1 (L_\Theta(V_{\lambda+1}), P)$$

implies full elementarity. \square

Another point in favour of AD is the regularity properties that it entails, and this is partially true also for I0.

We consider on $V_{\lambda+2}$ the topology with basic open sets $O_{(\alpha, a)} = \{x \subseteq V_\lambda : x \cap V_\alpha = a\}$, with $\alpha < \lambda$ and $a \subseteq V_\alpha$. Then a set $X \subseteq V_{\lambda+1}$ is perfect iff $\prod_{n \in \omega} \kappa_n$ can be continuously embedded into X . A set has the perfect set property iff it has cardinality $\leq \lambda$ or it contains a perfect subset.


Theorem 6.8 (Cramer, 2015 [?]). *Suppose there exists $j : L(V_{\lambda+1}) \prec L(V_{\lambda+1})$, with $\text{crt}(j) < \lambda$. Then every set in $L(V_{\lambda+1}) \cap V_{\lambda+2}$ has the perfect set property in $L(V_{\lambda+1})$.*

The proof uses heavily all the inverse limit tools developed in Section ?? . It exploits the fact that there is a multiplicity of possible square roots: Consider $J : L_{\bar{\alpha}+1}(V_{\lambda+1}) \prec L_{\alpha+1}(V_{\lambda+1})$, inverse limit, and let $X \in L_{\alpha+1}(V_{\lambda+1}) \cap V_{\lambda+2}$ and $\bar{X} \in L_{\bar{\alpha}+1}(V_{\lambda+1})$ such that $J(\bar{X}) = X$. Then if $|X| > \lambda$ we can find for any $a \in \bar{X}$ many ($> \lambda$) limit roots of J , all with different images on a . Repeated uses of this kind of pigeonhole principle will be used to construct a perfect set inside X .

Question 6.9. *Is there an equivalent in $L(V_{\lambda+1})$ for Baire property or Lebesgue measurability?*

Some key properties of subsets of \mathbb{R} are proven thanks to the fact that under AD all such sets have a nice tree-structure, namely they are weakly homogeneously Suslin. Woodin introduced a definition that wants to develop a similarity with this concept: $\mathbb{U}(j)$ -representability.

If $x \in V_{\lambda+1}$ and $\langle \lambda_i : i \in \omega \rangle$ is cofinal in λ , then we can see x as the ω -sequence $\langle x \cap V_{\lambda_i} : i \in \omega \rangle$, just like any real can be seen as an ω -sequence of natural numbers. So any set $X \subseteq V_{\lambda+1}$ can be seen as a set of ω -sequences. In \mathbb{R} , homogeneous Suslin-ness consists in labeling any finite sequence of natural numbers with an ultrafilter, so that if a sequence labeled with U_2 extends another labeled with U_1 , then U_2 projects to U_1 . If U_2 projects to U_1 , then there is a related elementary embedding from $\text{Ult}(V, U_1)$, to $\text{Ult}(V, U_2)$, therefore for any infinite sequence one can define the direct limit of all the ultrapowers of the ultrafilters that label its finite initial segments. Then a set Z is homogeneously Suslin iff there is such labeling so that a branch is in Z iff such direct limit is well-founded.



Weakly homogeneous Suslin-ness adds another dimension, considering instead the couples of finite sequences.

So the first step is to decide which ultrafilters are going to label our sequences:

Definition 6.10. *Let $\mathbb{U}(j)$ be the set of $U \in L(V_{\lambda+1})$ such that in $L(V_{\lambda+1})$ the following hold:*

- *U is a λ^+ -complete ultrafilter;*
- *for some $\gamma < \Theta$, $U \subseteq \mathcal{P}(L_\gamma(V_{\lambda+1}))$;*
- *for some $A \in U$ and all sufficiently large $n \in \omega$, $j^n(U) = U$ and $\{a \in A : j^n(a) = a\} \in U$.*

Let $\kappa < \Theta$, $\kappa \leq \Theta^{L_\kappa(V_{\lambda+1})}$. Let $\langle a_i : i < \omega \rangle$ be such that for all $i \in \omega$ there exists $n \in \omega$ such that $j^n(a_i) = a_i$. Then $\mathbb{U}(j, \kappa, \langle a_i : i \in \omega \rangle)$ is the set of $U \in \mathbb{U}(j)$ such that there exists an $n \in \omega$ such that for all $k : L_\kappa(V_{\lambda+1}) \prec L_\kappa(V_{\lambda+1})$ with the property that there are $m, l \in \omega$, $k^m = j^l \restriction L_\kappa(V_{\lambda+1})$, if for all $i \leq n$ $k(a_i) = a_i$, then $\{a \in L_\kappa(V_{\lambda+1}) : k(a) = a\} \in U$.

Now we can introduce $\mathbb{U}(j)$ representability:

Definition 6.11 (Woodin, 2011 [?]). *Let $Z \in L(V_{\lambda+1}) \cap V_{\lambda+2}$. Then Z is $\mathbb{U}(j)$ -representable if there are κ , $\langle a_i : i \in \omega \rangle$, an increasing sequence $\langle \lambda_i : i \in \omega \rangle$, and a function $\pi : \bigcup \{V_{\lambda_{i+1}} \times V_{\lambda_{i+1}} \times \{i\} : i \in \omega\} \rightarrow \mathbb{U}(j, \kappa, \langle a_i : i \in \omega \rangle)$ such that the following hold:*

- *for all $i \in \omega$ and $(a, b, i) \in \text{dom}(\pi)$ there exists $A \subseteq (L(V_{\lambda+1}))^i$ such that $A \in \pi(a, b, i)$;*
- *for all $i \in \omega$ and $(a, b, i) \in \text{dom}(\pi)$, if $m < i$ then $(a \cap V_{\lambda_m}, b \cap V_{\lambda_m}, m) \in \text{dom}(\pi)$ and $\pi(a, b, i)$ projects to $\pi(a \cap V_{\lambda_m}, b \cap V_{\lambda_m}, m) \in \text{dom}(\pi)$, i.e., for every $A \in \pi(a \cap V_{\lambda_m}, b \cap V_{\lambda_m}, m)$*

$$\{x \in (L(V_{\lambda+1}))^i : x \restriction m \in A\} \in \pi(a, b, i);$$

- *for all $x \subseteq V_\lambda$, $x \in Z$ iff there exists $y \subseteq V_\lambda$ such that the tower $\langle \pi(x \cap V_{\lambda_m}, y \cap V_{\lambda_m}, m) : m \in \omega \rangle$ is well-founded.*

The question is now whether there are sets in $L(V_{\lambda+1}) \cap V_{\lambda+2}$ that are $\mathbb{U}(j)$ -representable, and particularly which ones are. The desired result would be, in similarity with AD, that they are all $\mathbb{U}(j)$ -representable. There has been some twists and turns on the still ongoing quest towards this result, and it is worth to see them step by step, with some advice for the reader that wants to know all the details.

The main source for $\mathbb{U}(j)$ -representability is [?]. There the first closure properties are introduced, reminiscent of the closure properties for weakly homogeneously Suslin sets:

Theorem 6.12 (Woodin, Lemma 114, Lemma 115 and Theorem 134 in [?]). *Suppose $j : L(V_{\lambda+1}) \prec L(V_{\lambda+1})$ with $\text{crt}(j) < \lambda$. Then*

- *the λ -union of $\mathbb{U}(j)$ -representable sets is $\mathbb{U}(j)$ -representable*

- the projection of a $\mathbb{U}(j)$ -representable set is $\mathbb{U}(j)$ -representable
- if $Z \in L_\lambda(V_{\lambda+1})$ is $\mathbb{U}(j)$ -representable, then its complement is $\mathbb{U}(j)$ -representable.

Therefore all sets in $L_\lambda(V_{\lambda+1}) \cap V_{\lambda+2}$ are $\mathbb{U}(j)$ -representable.

This was the best result for $\mathbb{U}(j)$ -representability in [?]. Further research was blocked by the Tower Condition. Remember that a tower of ultrafilters $\langle U_n : n \in \omega \rangle$ is well-founded iff there exists an X such that $X \cap \text{dom}(U_n) \in U_n$ for all $n \in \omega$. The Tower Condition holds for $\mathbb{U}(j)$ iff for any $\mathbb{U}(j)$ -representation, it is possible to find such an X for any well-founded tower uniformly:

Definition 6.13 (Woodin, 2011 [?]). *The Tower Condition holds for $\mathbb{U}(j)$ iff for any $A \subseteq \mathbb{U}(j)$, $A \in L(V_{\lambda+1})$, $|A| \leq \lambda$, there is a function $F : A \rightarrow L(V_{\lambda+1})$ such that, for all $\langle U_i : i \in \omega \rangle \in L(V_{\lambda+1})$ that satisfies:*

- for all $i \in \omega$ there exists $Z \in U_i$ such that $Z \subseteq L(V_{\lambda+1})^i$;
- for all $i \in \omega$ $U_i \in A$ and U_{i+1} projects to U_i .

Then the tower is wellfounded iff there exists a function $f : \omega \rightarrow L(V_{\lambda+1})$ such that for all $i \in \omega$, $f \restriction i \in F(U_i)$.

Note that if π is a $\mathbb{U}(j)$ representation and $A = \text{ran}(\pi)$, then all towers in A satisfy the conditions. Woodin immediately noticed that the Tower Condition was central for the study of $\mathbb{U}(j)$ -representable sets:

Lemma 6.14 (Woodin, Lemma 127 in [?]). *If the Tower Condition holds for $\mathbb{U}(j)$, then the complement of a $\mathbb{U}(j)$ -representable set is $\mathbb{U}(j)$ -representable, therefore all the sets in $L_{\lambda^+}(V_{\lambda+1})$ are $\mathbb{U}(j)$ -representable.*

For much of Section 7 of [?], the Tower Condition is a constant and rather cumbersome presence, as at the time of the publication it was still not proven. In [?] Cramer actually managed to prove the Tower Condition, therefore unlocking much of the content of Section 7 of [?]. The suggestion for the completist reader is therefore first to read [?], and reading Section 7 of [?] after, so that the Tower Condition provides the most results.

This approach would also make some proofs in [?] more direct and easier to read. For example:

Lemma 6.15 (Woodin, Lemma 121 of [?]). *Suppose $Z \in L(V_{\lambda+1})$ is $\mathbb{U}(j)$ -representable and $|Z| > \lambda$. Then Z contains a \mathbb{R} -perfect set⁵.*

The proof of such Lemma starts by fixing an elementary embedding $j_0 : V \rightarrow M_0$ such that $(M_0)^{\lambda^+} \subseteq M_0$, $A_0 = \{j_0(U) : U \in \text{ran}(\pi)\}$. All of this is to prove the Tower Condition for $j_0''A_0$. Now that we know that the Tower Condition holds, we can ignore this trick and just consider $M_0 = V$, $j_0 = \text{id}$ and $A_0 = \text{ran}(\pi)$.

It is also possible to improve ??:

Lemma 6.16 (Woodin, Corollary 148 of [?]). *Suppose there is $j : L(V_{\lambda+1}) \prec L(V_{\lambda+1})$ with $\text{crt}(j) < \lambda$. Let $\kappa = \lambda^+$ and $\eta = \sup\{(\kappa^+)^{L[A]} : A \subseteq \lambda\}$. Then every set in $L_\eta(V_{\lambda+1})$ is $\mathbb{U}(j)$ -representable.*

So it is still open:

Question 6.17. *Are all sets in $L(V_{\lambda+1}) \cap V_{\lambda+2}$ $\mathbb{U}(j)$ -representable?*

⁵With \mathbb{R} -perfect set we intend that 2^ω can be continuously embedded in Z , a weaker property than to be a perfect set. Therefore this Lemma is now incorporated in ??

7 Rank-into-rank and forcing

A recurrent image for describing the difference of the extension of the universe of sets via large cardinals or via forcing is that large cardinals extend the universe vertically, while forcing extends the universe horizontally. Such image has definitively its shortcomings, first because stronger large cardinals do not always mean larger large cardinals (we will see an example of this), second because it implies that large cardinals and forcing are orthogonal, while there are cases of interdependence. It is therefore natural, facing new axioms, to ask what is their relationship with forcing. The ideal results are two:

- A forcing always destructs the large cardinal axiom. This would be a strong indication that the propositions forced could be inconsistent with the large cardinal.
- A forcing never destructs the large cardinal axiom. This means that the proposition forced is independent from the large cardinal, and we are actually in a situation of orthogonality.

The test case has always been Cohen forcing: any large cardinal cannot be destructed by Cohen forcing, therefore the Continuum Hypotheses is independent from any large cardinal.

The main lemma for proving the indestructibility of large cardinals via forcing is the following:

Lemma 7.1 (Lifting Lemma). *Let $j : M \prec N$. If $\mathbb{P} \in M$ is a forcing notion, and G is \mathbb{P} -generic, then for any H $j(\mathbb{P})$ -generic such that $j''G \subseteq H$ there exists $j^* : M[G] \prec N[H]$ such that $j^* \upharpoonright M = j$.*

Proof. The key point is that the forcing relation is definable. For any \mathbb{P} -name τ , define $j^*(\tau_G) = (j(\tau))_H$. Then $M[G] \models \varphi(\tau_G)$ iff there exists $p \in G$ such that $p \Vdash \varphi(\tau)$. So by elementarity $j(p) \Vdash \varphi(j(\tau))$. But $j(p) \in H$, i.e., $N[H] \models \varphi((j(\tau))_H)$, i.e., $N[H] \models \varphi(j^*(\tau_G))$. The opposite direction holds considering $\neg\varphi$. \square

This gives immediately a result of lifting: Let $j : V_\lambda \prec V_\lambda$, $\mathbb{P} \in V_{\text{crt}(j)}$. Then $j(\mathbb{P}) = \mathbb{P}$, so $G = H$ in the Lifting Lemma works, as $j''G = G$. Therefore j extends to $j^* : V_\lambda[G] \prec V_\lambda[G]$. But this is still not $V[G] \models I3(\lambda)$, because that is $\exists k : V[G]_\lambda \prec V[G]_\lambda$, where $V[G]_\lambda = (V_\lambda)^{V[G]}$. The difference is subtle, but fundamental: $V_\lambda[G]$ is the set of τ_G such that $\tau \in V_\lambda$, $V[G]_\lambda$ is the set of τ_G with rank of τ_G less than λ in $V[G]$. As the rank of the interpretation of a name is always less or equal than the rank of the name, clearly $V_\lambda[G] \subseteq V[G]_\lambda$, but it is not immediate to see whether the opposite direction is true. This is solved by the nice names lemma:

Lemma 7.2 (Name Rank Lemma). *If τ is a \mathbb{P} -name, $\mathbb{P} \in V_{\gamma+1}$ and $\tau_G \in V[G]_\beta$ for any G \mathbb{P} -generic, then there is a name $\sigma \in V_{\gamma+3 \cdot \beta}$ such that $\sigma_G = \tau_G$ for any G \mathbb{P} -generic.*

In particular, if $\mathbb{P} \in V_{\text{crt}(j)}$, any element of $V[G]_\lambda$ will be in some $V[G]_{\kappa_n}$, and so it will have a name in $V_{3 \cdot \kappa_n} \subseteq V_\lambda$, therefore $V_\lambda[G] = V[G]_\lambda$.

This is peculiar to rank-into-rank axioms: the indestructibility proof must pass two phases: first we have to extend the embedding, then we have to make sure that the domain of the embedding is exactly what we want.

By the discussion before Proposition ??, $\text{crt}(j)$ can be as large as we want below λ , therefore we have the following:

Lemma 7.3. *Suppose $I3(\lambda)$, let $\mathbb{P} \in V_\lambda$ be a forcing notion and G be \mathbb{P} -generic. Then $V[G] \models I3(\lambda)$, So $I3$ is not destructed by “small” forcings.*

The Lifting Lemma holds easily also for $j : V_{\lambda+1} \prec V_{\lambda+1}$ and $\mathbb{P} \in V_{\text{crt}(j)}$, so there exists $j^* : V_{\lambda+1}[G] \prec V_{\lambda+1}[G]$. We have to prove that $V_{\lambda+1}[G] = V[G]_{\lambda+1}$. If $\tau_G \in V[G]_{\lambda+1}$, then $\tau_G \subseteq V[G]_\lambda$, so $\tau_G = \bigcup_{n \in \omega} (\tau_G \cap V[G]_{\kappa_n})$. As $V_\lambda[G] = V[G]_\lambda$, any $\tau_G \cap V[G]_{\kappa_n}$ has a name in V_λ . The name of a union of sets is the union of the names of the sets, therefore there exists a name for τ_G that is the ω -union of sets in V_λ , so it must be in $V_{\lambda+1}$.

Again, $\text{crt}(j)$ can be unbounded below λ , so:

Lemma 7.4. *Suppose $I1(\lambda)$ (or anything slightly weaker, like $I2$ or Σ_n^1), let $\mathbb{P} \in V_\lambda$ be a forcing notion and G be \mathbb{P} -generic. Then $V[G] \models I1(\lambda)$ (or the corresponding weaker axiom). So $I1$ is not destructed by “small” forcings.*

Finally we consider $I0$ and $\mathbb{P} \in V_{\text{crt}(j)}$. Again, the Lifting Lemma holds, so there exists $j^* : L(V_{\lambda+1})[G] \prec L(V_{\lambda+1})[G]$ with $\text{crt}(j^*) < \lambda$. This time we cannot always prove that $L(V_{\lambda+1})[G] = L(V[G]_{\lambda+1})$, but it actually does not matter: as a constructible set will have a constructible name, we have $L(V[G]_{\lambda+1}) \subseteq L(V_{\lambda+1})[G]$. Also, $L(V[G]_{\lambda+1})$ is a definable (with parameter λ) subclass of $L(V_{\lambda+1})[G]$, therefore

$$j^* \upharpoonright L(V[G]_{\lambda+1}) : L(V[G]_{\lambda+1}) \prec L(V[G]_{\lambda+1})$$

and we have $I0(\lambda)$ in the extension.

Can we say that $\text{crt}(j)$ can be unbounded below λ ? For this, we have to prove that a j that witnesses $I0$ is at least finitely iterable. If j is not weakly proper, then it is not clear whether this is possible. But if j is weakly proper, then it is:

Lemma 7.5 (Woodin, Lemma 16 of [?]). *Let $j : L(V_{\lambda+1}) \prec L(V_{\lambda+1})$ with $\text{crt}(j) < \lambda$ and weakly proper. Then j is finitely iterable.*

Proof. Let I be the class of fixed points of ξ (it is a class by the proof of ??). For each $s \in [I]^{<\omega}$, let $Z_s = \text{Hull}^{L(V_{\lambda+1})}(V_{\lambda+1} \cup \{V_{\lambda+1}\} \cup s)$. Then $\langle Z_s : s \in [I]^{<\omega} \rangle$ is a directed system. Let Z be its limit. Then $Z \prec L(V_{\lambda+1})$.

For each $s \in [I]^{<\omega}$, $j \upharpoonright Z_s$ is definable from $\{j \upharpoonright V_\lambda, s\} \in L(V_{\lambda+1})$, therefore $k_s = j(j \upharpoonright Z_s)$ is well-defined. As $j(Z_s) = Z_s$, we have that $k_s : Z_s \prec Z_s$. Then we can define $k^* : Z \prec Z$, its direct limit. Composing with the collapse of Z , we have $k : L(V_{\lambda+1}) \prec L(V_{\lambda+1})$, and $k \upharpoonright V_\lambda = j(j \upharpoonright V_\lambda)$. If k is not weakly proper, then we consider k_U its weakly proper part, and we call it still k .

This proves the existence of j^2 . As j^2 is weakly proper, we can prove by induction that j^n exists and is weakly proper for every $n \in \omega$. \square

In particular $j^n \upharpoonright V_\lambda = (j \upharpoonright V_\lambda)^n$, therefore $\text{crt}(j^n) = \kappa_n$, and so:

Lemma 7.6. *Suppose $I0(\lambda)$, let $\mathbb{P} \in V_\lambda$ be a forcing notion and G be \mathbb{P} -generic. Then $V[G] \models I0(\lambda)$, So $I0$ is not destructed by “small” forcings.*

The reason why we do not know whether $L(V_{\lambda+1})[G] = L(V[G]_{\lambda+1})$ is quite deep:

Theorem 7.7 (Woodin, Theorem 175 in [?]). *Suppose $j : L(V_{\lambda+1}) \prec L(V_{\lambda+1})$ with $\text{crt}(j) < \lambda$. Let $\mathbb{P} \in V_\lambda$, G be \mathbb{P} -generic and $\lambda^\omega \neq (\lambda^\omega)^{V[G]}$. Then $V_{\lambda+1} \notin L(V[G]_{\lambda+1})$, so $L(V_{\lambda+1})[G] \neq L(V[G]_{\lambda+1})$ ⁶.*

Proof. Let $\langle \kappa_i : i \in \omega \rangle$ be the critical sequence of j . As $\mathbb{P} \in V_\lambda$, we can suppose that $\mathbb{P} \in V_{\kappa_0}$, and then there exists $j^* : L(V[G]_{\lambda+1}) \prec L(V[G]_{\lambda+1})$ that extends j . Suppose that $V_{\lambda+1} \in L(V[G]_{\lambda+1})$. Then by Theorem ?? in $L(V[G]_{\lambda+1})$ it is possible to embed continuously $(\lambda^\omega)^{V[G]}$ in $V_{\lambda+1}$.

We do some cosmetic changes so that the proof is easier to read: we fix ρ_n bijections between $V_{\kappa_{n+1}}$ and $|V_{\kappa_{n+1}}|$, rename $|V_{\kappa_{n+1}}|$ as κ_n , and modify the continuous embedding so that its domain is $(\prod_{n \in \omega} \kappa_i)^{V[G]}$. So there exists π that continuously embed $(\prod_{n \in \omega} \kappa_i)^{V[G]}$ to $(\prod_{n \in \omega} \kappa_i)^V$. We prove that $(\prod_{n \in \omega} \kappa_i)^{V[G]} \subseteq V$, and therefore $(\lambda^\omega)^{V[G]} = \lambda^\omega$, contradiction.

Let π be such embedding. We define for every $s \in \bigcup_{n \in \omega} \prod_{i < n} \kappa_n$ a set $S_s \subseteq \kappa_{\text{lh}(s)}$, $S_s \in V$ in such a way:

- $0 \in S_s$;
- Fix $n = \text{lh}(s)$. Let $\alpha \in S_s$. Then let $P_\alpha = \{\beta < \kappa_n : \exists p \in \prod_{i < n} \kappa_i \models \beta = \min\{\dot{\pi}(a)(n) : a \in (\prod_{n \in \omega} \kappa_n)^V, \pi(a) \restriction n = s, \dot{\pi}(a)(n) > \alpha\}\}$. Then $|P_\alpha| < \kappa_0$, therefore P_α is bounded in κ_n . As for any $s \in \bigcup_{n \in \omega} \prod_{i < n} \kappa_n$ $\{\pi(a)(n) : a \in (\prod_{n \in \omega} \kappa_n)^{V[G]}, \pi(a) \restriction n = s\}$ is unbounded in κ_n , $P_\alpha \neq \emptyset$, so let $\alpha' = \max P_\alpha + 1 \in S_s$.

Note that not only $S_s \in V$, but $\{S_s : s \in \bigcup_{n \in \omega} \prod_{i < n} \kappa_n\} \in V$.

By the construction, (*) for every $a \in (\prod_{n \in \omega} \kappa_n)^{V[G]}$ and every $\alpha \in S_{\pi(a) \restriction n}$, there exists $b \in (\prod_{n \in \omega} \kappa_n)^{V[G]}$ such that $\pi(a) \restriction n = \pi(b) \restriction n = s$, and $\alpha < \pi(b)(n) < \min S_s \setminus \alpha$.

We define now another continuous embedding, π' . For any S_s , let $S_s(\alpha)$ be the α -th element of S_s .

Let $a \in (\prod_{n \in \omega} \kappa_n)^{V[G]}$. Then we define by induction b_n :

- Applying (*) with $n = 0$, a anything and $\alpha = S_\emptyset(a(0))$, there exists b_0 such that $S_\emptyset(a(0)) < \pi(b_0)(0) < S_\emptyset(a(0) + 1)$;
- Suppose that b_n is already defined. Applying (*) with $a = b_n$ and $\alpha = S_{\pi(b_n) \restriction n}(a(n + 1))$, there exists b_{n+1} such that $\pi(b_{n+1}) \restriction n = \pi(b_n) \restriction n$ and $S_{\pi(b_n) \restriction n}(a(n + 1)) < \pi(b_{n+1})(n + 1) < S_{\pi(b_n) \restriction n}(a(n + 1) + 1)$.

As π is continuous, the b_n 's are more and more closer. Let b be their limit. Then we define $\pi'(a) = \pi(b)$.

Note that, if α and α' are two consecutive members of S_s , then $|\{\pi(b)(n) : \exists a \in (\prod_{n \in \omega} \kappa_n)^{V[G]}, \pi'(a) = \pi(b), \pi(b) \restriction n = s\} \cap [\alpha, \alpha']| = 1$. Therefore π' is well-defined and injective. But then the inverse of π' is definable in V , as for any $a \in (\prod_{n \in \omega} \kappa_n)^{V[G]}$, $\pi'(a)(n) = |S_{\pi'(a) \restriction n} \cap a(n)|$. \square

On the other hand, if the forcing is ω -closed in V_λ then we have equality:

Theorem 7.8. *Suppose $j : L(V_{\lambda+1}) \prec L(V_{\lambda+1})$ with $\text{crt}(j) < \lambda$. Let $\mathbb{P} \in V_\lambda$ be ω -closed in V_λ , i.e., $(V_\lambda)^\omega = ((V_\lambda)^\omega)^{V[G]}$, G be \mathbb{P} -generic. Then $L(V_{\lambda+1})[G] = L(V[G]_{\lambda+1})$.*

⁶The proof in [?] is more convoluted, as it proves that $V_{\lambda+1}$, if in $L(V[G]_{\lambda+1})$ and $\mathbb{U}(j)$ -representable, is a perfect set, and this brings a contradiction. By ?? we immediately have that $V_{\lambda+1}$ is perfect.

Proof. Fix a bijection in V between V_λ and λ , this extends naturally to a bijection from $V_{\lambda+1}$ and $\mathcal{P}(\lambda)$, so it suffices to define $\mathcal{P}(\lambda)$ inside $L(V[G]_{\lambda+1})$:

$$\mathcal{P}(\lambda) = \{a \in \mathcal{P}^{V[G]}(\lambda) : \forall n \in \omega \ a \cap \kappa_n \in V\}.$$

This is because if $a \cap \kappa_n = a_n \in V$, then $\langle a_n : n \in \omega \rangle \in V$ and $a = \bigcup_{n \in \omega} a_n \in V$. \square

Corollary 7.9. *Suppose $j : L(V_{\lambda+1}) \prec L(V_{\lambda+1})$ with $\text{crt}(j) < \lambda$. Let $\mathbb{P} \in V_\lambda$ be a forcing notion, G be \mathbb{P} -generic. Then $L(V_{\lambda+1})[G] = L(V[G]_{\lambda+1})$ iff $(V_\lambda)^\omega = ((V_\lambda)^\omega)^{V[G]}$.*

So I3, I2, I1 and I0 are not changed by forcing notions bounded in λ . Also, trivially, if a forcing notion is λ -closed, then $V[G]_{\lambda+1} = V_{\lambda+1}$, therefore I3, I2, I1 and I0 are witnessed by the exact same embedding, and are not destructed. In other words, they are not changed by forcings that are above λ .

This case can give some very nice results:

Theorem 7.10 (Shi, Trang, 2017 [?]). *Suppose $I0(\lambda)$ holds. Then all the following statements are true in a generic extension where $I0(\lambda)$ still holds:*

- *special λ^+ -Aronszajn trees exist;*
- *λ^+ -Suslin tree exist;*
- *there is a very good scale at λ^+ ;*
- *Stationary Reflections fails at λ^+ ;*
- \diamond_{λ^+} ;

This is particularly striking if compared with Theorem ??.

It remains to see what happens when the forcing has size λ . The first case is when the forcing is unbounded in λ (the second case will be when the forcing is exactly at λ). The typical case is when the forcing is a λ -iteration of small forcings. In such case, the reverse Easton iteration has been proven in the past to behave very well with large cardinals:

Definition 7.11. *Let \mathbb{P}_α be a forcing iteration of length α , where α is either a strong limit cardinal or is equal to ∞ , the class of all ordinals. Then \mathbb{P}_α is a reverse Easton iteration if nontrivial forcing is done only at infinite cardinal stages, direct limits are taken at all inaccessible cardinal limit stages, and inverse limits are taken at all other limit stages; moreover, \mathbb{P}_α is the direct limit of $\langle \mathbb{P}_\delta : \delta < \alpha \rangle$ if α is regular or ∞ , the inverse limit otherwise.*

Such iteration was introduced by Easton to force different behaviours of the function $\kappa \mapsto 2^\kappa$, the power function. There are two known rules for the power function: if $\kappa < \eta$ then $2^\kappa \leq 2^\eta$, and $\text{cof}(2^\kappa) > \kappa$ for κ regular. Easton proved that any function that satisfies this two rules can be the power function on the regular cardinals:

Definition 7.12. *Let $E : \text{Reg} \rightarrow \text{Card}$ be a class function. Then E is an Easton function iff:*

- $\alpha < \beta \rightarrow E(\alpha) \leq E(\beta)$;

- $\text{cof}(E(\alpha)) > \alpha$ for all $\alpha \in \text{Reg}$.

Then Easton proved:

Theorem 7.13 (Easton, 1970 [?]). *Let E be an Easton function. Then there exists a generic extension $V[G]$ of V such that $V[G] \models \forall \kappa (\kappa \in \text{Reg} \rightarrow 2^\kappa = E(\kappa))$.*

The forcing used was a reverse Easton iteration. The easiest application is to force GCH: in that case for any $\gamma < \alpha$ \mathbb{Q}_γ forces that $2^\gamma = \gamma^+$ if γ is a cardinal, it is trivial otherwise, $\mathbb{P}_{\gamma+1} = \mathbb{P}_\gamma * \dot{\mathbb{Q}}_\gamma$ and G will be the generic filter of the reverse Easton iteration \mathbb{P}_∞ . The problem is when one, for example, starts with a model of GCH and wants to increase everywhere on the regulars the power function: If we force with \mathbb{Q}_ω $2^\omega = \omega_2$ and after that we force $2^{\omega_1} = \omega_3$ via Cohen forcing, then in the final model 2^ω will collapse to ω_1 again. The idea is then that \mathbb{Q}_δ is defined only on δ closed under E , and \mathbb{Q}_δ is the Easton product⁷ of the forcings that make $2^\gamma = E(\gamma)$ for all $\gamma \in [\delta, \delta^*)$, where δ^* is the smallest cardinal closed under E bigger than δ (this is still in [?]).

So let E be an Easton function, and \mathbb{P}_λ that forces $2^\kappa = E(\kappa)$ for any regular κ , with the stages indicated with \mathbb{Q}_δ for $\delta < \lambda$. We want to know for which E 's the forcing \mathbb{P}_λ does not destroy I3, ..., I0.

If for some $\delta < \lambda$ $|\mathbb{Q}_\delta| \geq \lambda$, then there is a $\alpha < \delta^*$ such that in $V[G]$ $2^\alpha \geq \lambda$, therefore λ is not strong limit anymore in $V[G]$ and the embedding is destroyed. Therefore we must have $|\mathbb{Q}_\delta| < \lambda$ for any $\delta < \lambda$.

If for some $\delta < \lambda$ we have that $j(\mathbb{P}_\delta) \neq \mathbb{P}_{j(\delta)}$, then possibly $j(2^\alpha) \neq 2^{j(\alpha)}$ for some $\alpha < \delta$, therefore we should ask that $j(\mathbb{P}_\delta) = \mathbb{P}_{j(\delta)}$. If E is definable then this is trivial.

Also note that \mathbb{Q}_δ is δ -directed closed. This is enough to lift I3, ..., I0 in the generic extension:

Theorem 7.14 ([?]). *Suppose j witnesses I3(λ) (or I1(λ) ...). Let \mathbb{P}_λ be a forcing iteration of length λ that is reverse Easton, directed closed (i.e., for all $\delta < \lambda$, \mathbb{Q}_δ is δ -directed closed), λ -bounded (i.e., for all $\delta < \lambda$, $|\mathbb{Q}_\delta| < \lambda$) and j -coherent (i.e., for all $\delta < \lambda$, $j(\mathbb{P}_\delta) = \mathbb{P}_{j(\delta)}$). Then j lifts to any generic extension via \mathbb{P}_λ .*

Sketch of proof. The proof is in two stages: lifting and right domain.

First notice that if $\gamma < \kappa_n$, then $|\mathbb{P}_\gamma| < \kappa_n$, by elementarity, λ -boundedness and n applications of j . Fixing an $n < \omega$, consider the image of \mathbb{Q}_{κ_n} under j : it is a subset of $\mathbb{Q}_{\kappa_{n+1}}$ of cardinality less than κ_{n+1} , therefore by directed closeness there is an element $\mathbb{Q}_{\kappa_{n+1}}$ that extends all the $j(p)$ with $p \in \mathbb{Q}_{\kappa_n}$. Doing the same for all n and also for $\mathbb{P}_{\kappa_n, \kappa_{n+1}}$, one gets a condition of \mathbb{P}_λ that, if in a generic G , implies that $j''G \subseteq G$. Therefore we can apply the Lifting Lemma.

As for the right domain, since the forcing is directed closed every element in the generic extension is actually in a generic extension for some \mathbb{P}_{κ_n} , therefore the previous proofs still hold. \square

Theorem ?? is just the final step of a series of theorems that goes in that direction: in [?] Hamkins has proved such theorem for I1 and reverse Easton directed closed j -coherent iterations such that $|\mathbb{Q}_\delta| < 2^\delta$; in [?] Corazza has

⁷It is the product with support the Easton ideal, i.e., the sets that are bounded below every inaccessible cardinal.

proved it for I3 and reverse Easton directed closed j -coherent iterations such that $|\mathbb{Q}_\delta| < i(\delta)$, where $i(\delta)$ is the smallest inaccessible larger than δ ; in [?] Friedman has proved it for I2 (called there ω -superstrong) and GCH.

The hypotheses in Theorem ?? are very reasonable, and most of the reverse Easton forcing usually employed satisfy them. So, for example:

Corollary 7.15. *Suppose I3 (or I1, ...). Then in a generic extension can hold:*

- $I3 + \text{GCH}$;
- $I3 + 2^\kappa = \kappa^{+++}$ for any κ regular;
- $I3 + V = \text{HOD}$;
- $I3 + \Diamond$ everywhere;
- $I3 +$ every κ supercompact is Laver indestructible (see [?]);
- ...

Finally the last case is when the forcing is changing λ^+ . This is the hardest case, and at the moment the indestructibility results are very scarce. For example, Gitik short extenders Prikry forcing construction in [?] can be carried on under I3 (it needs $o(\kappa_n) \geq (\kappa_n)^{+m}$ for any $n, m \in \omega$): such construction push 2^λ to $\lambda^{+\delta+1}$ for any desired $\delta < \aleph_1$, but does not add bounded subsets of λ , therefore $V[G]_\lambda = V_\lambda$ and I3 is not destructed. Yet, this construction for I1 does not work, as $V_{\lambda+1} \neq V[G]_{\lambda+1}$.

In [?] Cummings and Foreman, using something more than I2, manage to prove that $\text{I2}(\lambda)$ is consistent with $2^\lambda = \lambda^{++}$.

The most successful strategy has been instead to start with $\text{I0}(\lambda)$, and using some sophisticated tool to prove that, adding a Prikry sequence to the critical point κ_0 , in the extension $\text{I1}(\kappa_0)$ holds.

Definition 7.16. *Let κ be a measurable cardinal, and U a normal measure on κ . Then \mathbb{P} , the Prikry forcing on κ via U , is the set of (s, A) such that $s \in [\kappa]^{<\omega}$, $A \in U$ and $\min A > \max s$.*

We say that $(s, A) < (t, B)$ if $s \sqsupseteq t$, $A \subseteq B$ and for any $n \in \text{lh}(s) \setminus \text{lh}(t)$, $s(n) \in B$.

If (s, A) and (t, B) are in the generic set G , then, s and t must be compatible. Therefore by density $\bigcup \{s : \exists A (s, A) \in G\}$ is an ω -sequence cofinal in κ . So $(\text{cof}(\kappa) = \omega)^{V[G]}$, but it is a very delicate forcing, as it does not add any bounded subset of κ .

There is a convenient condition for an ω -sequence cofinal in κ to be generic:

Theorem 7.17 (Mathias Condition, or geometric condition). *Let κ be a measurable cardinal, \mathbb{P} the Prikry forcing on κ via the normal ultrafilter U and let $\langle \alpha_n : n \in \omega \rangle$ be a cofinal sequence in κ . Then $\langle \alpha_n : n \in \omega \rangle$ is generic for \mathbb{P} iff for any $A \in U$ the set $\langle \alpha_n : n \in \omega \rangle \setminus A$ is finite.*

In Theorem ?? we noticed that if j witnesses $\text{I0}(\lambda)$ is weakly proper, then it is finitely iterable, and its iterates are still weakly proper embeddings from $L(V_{\lambda+1})$ to itself. Woodin proved that iterability can go further:

Theorem 7.18 (Woodin, Lemma 21 in [?]). *Suppose that $j : L(V_{\lambda+1}) \prec L(V_{\lambda+1})$ with $\text{crt}(j) < \lambda$ is weakly proper. Then j is iterable.*

From the proof of Lemma ?? $j_{0,\omega}(\kappa_0) = \lambda$, therefore M_ω is different than $L(V_{\lambda+1})$: in it, by elementarity, λ is regular. Let U be the ultrafilter from j . Then $j_{0,\omega}(U)$ is a normal ultrafilter on λ , therefore λ is measurable in M_ω . If \mathbb{P} is the Prikry forcing on κ_0 via U , then $j_{0,\omega}(\mathbb{P})$ is the Prikry forcing on λ via $j_{0,\omega}(U)$.

Remark 7.19. *If $\langle \kappa_n : n \in \omega \rangle$ is the critical sequence of j , then $\langle \kappa_n : n \in \omega \rangle$ is $j_{0,\omega}(\mathbb{P})$ -generic in M_ω .*

Proof. Note that Mathias' characterization needs the Axiom of Choice, but by elementarity $(V_\lambda)^{M_\omega} \models \text{AC}$, so we can use it. If $A \in j_{0,n}(U)$, then by the definition of U $\kappa_n \in j_{n,\omega}(A)$, as $A \in j_{0,n}(U)$ iff $\kappa_n \in j^n(A)$ and

$$\kappa_n = j_{n+1,\omega}(\kappa_n) \in j_{n+1,\omega}(j^n(A)) = j_{n,\omega}(A).$$

So if $A \in j_{0,\omega}(U)$, there exist $n \in \omega$ and $\bar{A} \in L(V_{\lambda+1})$ such that $A = j_{n,\omega}(\bar{A})$. By elementarity $\bar{A} \in j_{0,n}(U)$ and $j_{n,n+i}(\bar{A}) \in j_{0,n+i}(U)$. So for any $i \in \omega$, $\kappa_{n+i} \in j_{n+i,\omega}(j_{n,n+i}(\bar{A})) = A$. \square

So it make sense to consider $M[\langle \kappa_i : i \in \omega \rangle]$.

Theorem 7.20 (Generic Absoluteness, Woodin, [?]). *Let $j : L(V_{\lambda+1}) \prec L(V_{\lambda+1})$ with $\text{crt}(j)$ be a weakly proper elementary embedding. Let $\langle \kappa_n : n \in \omega \rangle$ be its critical sequence, and M_ω its ω -iterated model.*

Let δ be such that all the sets in $L_\delta(V_{\lambda+1}) \cap V_{\lambda+2}$ are $\mathbb{U}(j)$ -representable. Then there exists an elementary embedding

$$\pi : L_\delta(M_\omega[\langle \kappa_n : n \in \omega \rangle] \cap V_{\lambda+1}) \prec L_\delta(V_{\lambda+1})$$

such that $\pi \upharpoonright V_{\lambda+1}$ is the identity.

Two examples to understand better the theorem:

- As $\langle \kappa_n : n \in \omega \rangle \in V_{\lambda+1}$, clearly

$$M_\omega[\langle \kappa_n : n \in \omega \rangle] \cap V_{\lambda+1} = (V_{\lambda+1})^{M_\omega[\langle \kappa_n : n \in \omega \rangle]} \subseteq V_{\lambda+1}.$$

Since $\pi \upharpoonright V_{\lambda+1} = \text{id}$, the Theorem says that $M_\omega[\langle \kappa_n : n \in \omega \rangle] \cap V_{\lambda+1} \prec V_{\lambda+1}$.

- If A is definable on $M_\omega[\langle \kappa_n : n \in \omega \rangle]$, then $\pi(A)$ will be the set defined in the same way on $V_{\lambda+1}$, and $\pi(A) \cap M_\omega[\langle \kappa_n : n \in \omega \rangle] = A$.

The proof exploits the fact that in the definition of $\mathbb{U}(j)$ -representability we had $j^n(U) = U$ for the ultrafilters used in a representation, therefore given a set in $L_\delta(V_{\lambda+1})$, its representation is also in M_ω , in a certain sense is the set encrypted. When we add the critical sequence to λ , then we can rebuild something similar to the set (if not the set itself) using such representation. The Theorem uses the theory of $L_\delta(V_{\lambda+1})$, as a set, to build an embedding.

This essential tool has an immediate corollary:

Corollary 7.21. *Suppose that there exists $j : L(V_{\lambda+1}) \prec L(V_{\lambda+1})$ with $\text{crt}(j) = \kappa_0 < \lambda$. Let \mathbb{P} be the Prikry forcing on κ_0 . Then there exists G \mathbb{P} -generic such that there exists $k : V[G]_{\kappa_0+1} \prec V[G]_{\kappa_0+1}$.*

Proof. We have seen that $j \upharpoonright V_{\lambda+1}$ is definable from $j \upharpoonright V_\lambda$, therefore it is in $L_1(V_{\lambda+1})$. Then

$$\pi^{-1}(j) = j \cap M_\omega[\langle \kappa_n : n \in \omega \rangle] : (V_{\lambda+1})^{M_\omega[\langle \kappa_n : n \in \omega \rangle]} \prec (V_{\lambda+1})^{M_\omega[\langle \kappa_n : n \in \omega \rangle]}.$$

So in M_ω there is a condition in $j_{0,\omega}(\mathbb{P})$ that forces the existence of a $\text{I1}(\lambda)$ embedding. By elementarity, in V there is a condition on \mathbb{P} that forces the existence of a $\text{I1}(\kappa_0)$ embedding. \square

This corollary, coupled with Theorem ??, can expand the possibilities of consistency results: for example one can force $2^{\kappa_0} > \kappa_0^+$ with the reverse Easton forcing, and then force with Prikry to have $\text{I1}(\kappa_0)$. Prikry will not change $2^{\kappa_0} > \kappa_0^+$, and therefore we have a model for $\text{I1}(\lambda) + 2^\lambda > \lambda^+$.

Theorem 7.22. *Suppose there exists $j : L(V_{\lambda+1}) \prec L(V_{\lambda+1})$ with $\text{crt}(j) < \lambda$. Then for every Easton function E such that $E \upharpoonright \lambda$ is definable over V_λ , there is a generic extension $V[G]$ of V that satisfies $\exists \gamma < \lambda \exists k : V_{\gamma+1} \prec V_{\gamma+1} \wedge 2^\gamma = E(\gamma)$.*

In particular this proves that it is consistent with $\text{I1}(\lambda)$ that there are no strongly compact or supercompact cardinals below λ , since λ is singular and Solovay's Theorem says that above a strongly compact or a supercompact cardinal we have $2^\lambda = \lambda^+$ on singular cardinals. Therefore this is one of the cases where “stronger” does not mean “larger”.

The first way to improve such results is to note that Theorem ?? can express more than just I1 , therefore the same proof works for stronger hypotheses (for example $\exists j : L_\lambda(V_{\lambda+1}) \prec L_\lambda(V_{\lambda+1})$), with the only bound given by which sets are $\text{U}(j)$ -representable, making Question ?? even more relevant.

Another way is to notice that in the proof of Theorem ?? it is not necessary to limit ourselves to Prikry forcing and to $\langle \kappa_n : n \in \omega \rangle$: the important thing is that there is a $G \in V$ $j_{0,\omega}(\mathbb{P})$ -generic and there is an ω -sequence cofinal in λ in $M_\omega[G]$. If $G \notin V$ then there are easy counterexamples: if CH holds in V , then it holds also in M_ω , and a forcing that adds a Cohen real to M_ω would make $V_{\lambda+1}^{M_\omega[G]}$ not elementary in $V_{\lambda+1}$ (not even contained, in fact).

There are numerous variations on Prikry forcing that add ω -sequences to a large cardinal, and as λ is a very large cardinal in M_ω (see ??) they are usually applicable in this context. For example, it is clear in Theorem ?? that the reverse Easton forcing forces $2^\eta = E(\eta)$ for all the regular cardinals $\eta < \gamma$, not only for γ , therefore with the construction of Theorem ??, based on classical Prikry forcing, γ is not the first cardinal on which GCH fails. Being the first one is a stronger property, that is forced usually with Gitik-Magidor long extender Prikry forcing (see [?]). If \mathbb{P} is therefore the Gitik-Magidor long extender Prikry forcing on κ_0 and it could be possible to find a $G \in V$ $j_{0,\omega}(\mathbb{P})$ -generic in M_ω , then by Generic Absoluteness we would have a generic extension in which $\text{I1}(\kappa_0) + 2^{\kappa_0} > \kappa_0^+$ and GCH below κ_0 .

To check if this and other forcing notions work with Generic Absoluteness there is the need of a criterion that, if satisfied, implies that there is a $G \in V$, $j_{0,\omega}(\mathbb{P})$ -generic in M_ω .

Definition 7.23. Let λ be an infinite cardinal. A partially ordered set \mathbb{P} is λ -good if it adds no bounded subsets of λ and for every \mathbb{P} -generic G and every $A \in V[G]$, $A \subset \text{Ord}$, $|A| < \lambda$, there is a non- \subset -decreasing ω -sequence $\langle A_i : i < \omega \rangle \subseteq V$ such that $A = \bigcup_{i \in \omega} A_i$.

Proposition 7.24 (Shi, Proposition 3.20 in [?]). Let $j : L(V_{\lambda+1}) \prec L(V_{\lambda+1})$ with $\text{crt}(j) = \kappa_0 < \lambda$ and let $\mathbb{P} \in V_\lambda$ be a κ_0 -good forcing. Then if $(M_\omega, j_{0,\omega})$ is the ω -iterate of j , there is a $G \in V_{j_{0,\omega}(\mathbb{P})}$ -generic on M_ω .

The original definition of λ -goodness involved families of dense sets, and this is a more convenient way to prove that various Prikry forcings are λ -good. The following is a sufficient condition:

Definition 7.25 ([?]). Let λ be an infinite cardinal. Let \mathbb{P} be a forcing notion equipped with a length measure of the conditions $l : \mathbb{P} \rightarrow \omega$ such that $l(1_\mathbb{P}) = 0$ and for $p, q \in \mathbb{P}$, if $p \leq q$ then $l(p) \geq l(q)$ (for example if \mathbb{P} is the Prikry forcing, then $l(s, A) = \text{lh}(s)$).

We say that \mathbb{P} is λ -geometric if it does not add bounded subsets of λ and for any $\alpha < \lambda$, any $\langle D_\beta : \beta < \alpha \rangle$ collection of open dense sets and any $p \in \mathbb{P}$ there exists a condition $q \leq p$ such that whenever a filter contains q and meets all the dense open sets $E_n = \{p : l(p) > n\}$, it also meets all the D_β 's.

Remark 7.26. Let λ be an infinite cardinal. Let \mathbb{P} be a λ -geometric forcing notion. Then \mathbb{P} is λ -good.

Proof. Let G be \mathbb{P} -generic. Let $A \subseteq \text{Ord}$, $|A| = \mu < \lambda$, $A \in V[G]$. Let $f : \mu \rightarrow \text{Ord}$ such that $\text{ran}(f) = A$, $f \in V[G]$. For any $\alpha < \mu$, let $D_\alpha = \{p \in \mathbb{P} : \exists \beta \ p \Vdash \dot{f}(\alpha) = \beta\}$, the set of the conditions that decide the α -th element of A . By λ -geometricity, the set of q 's such that every filter that contains q and intersects E_n intersects also all D_α 's, is dense in \mathbb{P} , therefore there exists $q_0 \in G$ with such property. Define by induction q_{n+1} that extends q_n such that $q_{n+1} \in G \cap E_n$. Let $B_n = \{\alpha < \mu : q_n \in D_\alpha\} \in V$ be the set of α 's such that q_n decides $f(\alpha)$. Then since $\mathcal{F} = \bigcup_{n \in \omega} \mathcal{F}_{q_n}$, by λ -geometricity, intersects all D_α 's (\mathcal{F}_{q_n} is the filter generated by q_n), for any $\alpha < \mu$ there exists $n \in \omega$ such that $q_n \in D_\alpha$, therefore $\mu = \bigcup_{n \in \omega} B_n$. Define

$$A_n = \{\beta : \exists \alpha \in B_n \ q_n \Vdash \dot{f}(\alpha) = \beta\} \in V.$$

If $\beta \in A$, then there exists $\alpha < \mu$ such that $f(\alpha) = \beta$, and there exists $n \in \omega$ such that $\alpha \in B_n$, i.e., q_n decides $f(\alpha)$. But since $q_n \in G$, it must be that $q_n \Vdash \dot{f}(\alpha) = \beta$, so $\beta \in A_n$, therefore $A = \bigcup_{n \in \omega} A_n$. \square

The key point is that many variations of Prikry forcing satisfy a variation of the Prikry condition: Let $p \leq^* q$ be $p \leq q$ and $l(p) = l(q)$. Then such variation says that for any dense set D and every $p \in \mathbb{P}$ there is a $q \leq^* p$, such that for any $r \leq q$ with $l(r) = l(q) + n$, $r \in D$. In other words every condition can be “changed” so that any extension above a certain length is in the dense set. This condition appears frequently in literature, for example in [?] for Prikry forcing, or [?] for Gitik-Magidor long extender Prikry forcing. In [?] there are other examples, and in both [?] and [?] it is exploited to prove the λ -goodness of Gitik-Magidor long extender Prikry forcing, in different ways. If \leq^* is sufficiently closed, it is easy to see that with repeated uses of this variation

of Prikry condition one can find a q that witnesses λ -geometricity, and therefore λ -goodness.

It is possible also to prove it directly:

Remark 7.27 ([?]). *Let $j : L(V_{\lambda+1}) \prec L(V_{\lambda+1})$ with $\text{crt}(j) = \kappa_0 < \lambda$ and let $\mathbb{P} \in V_\lambda$ be a κ_0 -geometric forcing. Then if $(M_\omega, j_{0,\omega})$ is the ω -iterate of j , there is a $G \in V_{j_{0,\omega}(\mathbb{P})}$ -generic on M_ω .*

Proof. Note that in V there are only λ open dense sets of $j_{0,\omega}(\mathbb{P})$: as $\mathbb{P} \in V_\lambda$ there exists $n \in \omega$ such that $\mathbb{P} \in V_{\kappa_n}$, so $j_{0,\omega}(\mathbb{P}) \in M_\omega \cap V_{j_{0,\omega}(\kappa_n)}$. But $M_\omega \cap V_{j_{0,\omega}(\lambda)}$ is the range of all $j^n \upharpoonright V_\lambda$, therefore $|j_{0,\omega}(\mathbb{P})| < |M_\omega \cap V_{j_{0,\omega}(\lambda)}| = |V_\lambda| = \lambda$.

Let $\langle D_\alpha : \alpha < \lambda \rangle$ be an enumeration of the dense sets of $j_{0,\omega}(\mathbb{P})$ in V . For every $n \in \omega$, fix q_n that witnesses λ -geometricity for $\langle D_\alpha : \alpha < \kappa_n \rangle$. Then for every m there exists a $q'_{n,m} < q_n$ such that $q'_{n,m} \in E_m$. Let H be the filter $\bigcup_{n,m \in \omega} \mathcal{F}_{q'_{n,m}}$, with \mathcal{F}_q the filter generated by q . Then $H \in V$ is generic. \square

Therefore, thanks to the work that in literature has already been done with the structure of Prikry-like forcing, for many variations of Prikry forcing it is possible to prove Generic Absoluteness. Interestingly, for many applications such variations need a forcing preparation that is actually a reverse Easton iteration, therefore Theorem ?? is necessary. In [?] are collected some results:

Theorem 7.28. *Suppose $I0(\lambda)$. Then the following are consistent:*

- $I1(\gamma) + 2^\gamma > \gamma^+ + \forall \kappa < \gamma \ 2^\kappa = \kappa^+$ (see [?], preparation forcing to force GCH, then Gitik-Magidor long extender Prikry forcing);
- $I1(\gamma) + \text{Tree property at } \gamma^+$ (see [?], preparation forcings that make supercompact cardinals Laver indestructible and add many subsets to all regular cardinals, then diagonal supercompact Prikry forcing);
- $I1(\gamma) + \text{Tree property at } \gamma^{++}$ (see [?], preparation forcing is reverse Easton iteration of Sacks forcings, then Prikry forcing);
- ...

8 Dissimilarities with $\text{AD}^{L(\mathbb{R})}$

While there are many similarities between $I0$ and $\text{AD}^{L(\mathbb{R})}$, not everything really matches. The first thing that stands out is that while $L(\mathbb{R}) \models \text{AD}^{L(\mathbb{R})}$, $L(V_{\lambda+1}) \not\models I0(\lambda)$. But the big problem is Theorem ??: the fact that many structural properties below λ are independent from $I0$ in fact implies that $I0$ cannot possibly prove them, and these properties can be the similarities we are interested in. In the $\text{AD}^{L(\mathbb{R})}$ case this situation is avoided as there are no forcings below ω . For example, we have proven that λ^+ is measurable in $L(V_{\lambda+1})$ under $I0$, just like ω_1 was measurable in $L(\mathbb{R})$ under AD . But in the $L(\mathbb{R})$ case we have that the club filter itself is an ultrafilter, while Theorem ?? stops short and proves only that the club filter on a stationary set is measurable. So the question is: how close can we get to the original case?

As $\lambda > \omega$, for any $\alpha < \lambda$ regular $S_\alpha^{\lambda^+}$ is stationary, and for $\alpha \neq \beta$ $S_\alpha^{\lambda^+} \cap S_\beta^{\lambda^+} = \emptyset$, therefore it is not possible to have the club filter to be an ultrafilter. The best outcome would be then to have the club filter to be an ultrafilter on every

$S_\alpha^{\lambda^+}$; this is called Ultrafilter Axiom at λ . But this is consistently false because of Theorem ??:

Proposition 8.1. *Let $j : L(V_{\lambda+1}) \prec L(V_{\lambda+1})$ with $\text{crt}(j) < \lambda$, let $\omega < \gamma < \lambda$ regular. Then in a generic extension the following is true: $I0(\lambda)$ still holds and in $L(V_{\lambda+1})$ there are S_1 and S_2 disjoint stationary sets such that $S_1 \cup S_2 = S_\gamma^{\lambda^+}$.*

Proof. First of all, notice that λ^+ is coded by subsets of λ , so it can be considered a subset of $V_{\lambda+1}$. As $V_{\lambda+1}$ is closed by λ -sequences, the cofinality of a cardinal less than λ^+ is the same in V and in $L(V_{\lambda+1})$ (because is witnessed by an element of $V_{\lambda+1}$), therefore $(S_\eta^{\lambda^+})^V = (S_\eta^{\lambda^+})^{L(V_{\lambda+1})}$ for any η regular. Let $\mathbb{P} \in L(V_{\lambda+1})$ be $\text{Coll}(\gamma^+, \gamma)$. Then by ?? $I0(\lambda)$ holds in $V[G]$. As $\gamma > \omega$, \mathbb{P} is ω -closed. Let G be \mathbb{P} generic. Consider $(S_{\gamma^+}^{\lambda^+})^V$ and $(S_\gamma^{\lambda^+})^V$, both in $L(V_{\lambda+1})$ and therefore in $L(V_{\lambda+1})[G]$. Since \mathbb{P} does not add unbounded sets in λ^+ , $(S_{\gamma^+}^{\lambda^+})^V$ and $(S_\gamma^{\lambda^+})^V$ are still stationary in $L(V_{\lambda+1})[G]$, are both subsets of $(S_\gamma^{\lambda^+})^{V[G]} = (S_\gamma^{\lambda^+})^{L(V_{\lambda+1})[G]}$ and of course are disjoint. Since \mathbb{P} is ω -closed, by ?? $L(V_{\lambda+1})[G] = L(V[G]_{\lambda+1})$, so $(S_{\gamma^+}^{\lambda^+})^V, (S_\gamma^{\lambda^+})^V \in L(V[G]_{\lambda+1})$ are S_1 and S_2 . \square

Note that if $\gamma = \omega$ then the proof breaks down: then $L(V_{\lambda+1})[G] \neq L(V[G]_{\lambda+1})$ and it is not clear why S_1 and S_2 would be in $L(V[G]_{\lambda+1})$. The following problems are still open:

Question 8.2. *Is the Ultrafilter Axiom consistent with $I0$?*

Question 8.3. *Is it consistent with $I0$ to have the club filter an ultrafilter on cofinality ω ? Could it be a consequence of $I0$?*

Towards a solution of the second problem, there are the following results:

Theorem 8.4 (Woodin, Theorem 176 in [?]). *Suppose $I0(\lambda)$. Then there is no partition of $S_\omega^{\lambda^+}$ into two stationary sets such that all the sets definable from them are $\mathbb{U}(j)$ -representable⁸.*

Theorem 8.5 (Cramer, Corollary 4.6 in [?]). *Suppose there exists $j : L_\omega(V_{\lambda+1}^\#, V_{\lambda+1}) \prec L_\omega(V_{\lambda+1}^\#, V_{\lambda+1})$. Then there is no partition of $S_\omega^{\lambda^+}$ into two stationary (in V) sets such that $S_1, S_2 \in L(V_{\lambda+1})$.*

Note that if S is stationary in V , then it is also stationary in $L(V_{\lambda+1})$, but not necessarily viceversa, therefore it is still open whether there can be two disjoint subsets of $S_\omega^{\lambda^+}$ that are in $L(V_{\lambda+1})$, stationary there, but not stationary in V .

Towards a solution for the first problem, we have a weakening of the Ultrafilter Axiom. Theorem ?? says that we can split $S_\gamma^{\lambda^+}$ in $\langle S_\alpha : \alpha < \eta_\gamma \rangle$ disjoint stationary sets for any $\gamma < \lambda^+$ regular. The Ultrafilter Axiom says that $\eta_\gamma = 1$ for any $\gamma < \lambda^+$ regular. The Weak Ultrafilter Axiom states that $\eta_\gamma \leq \gamma^+$ for any $\gamma < \lambda^+$ regular.

⁸This rather clumsy statement comes from the fact that the proof uses the fact that some Z_S defined from the stationary set S is $\mathbb{U}(j)$ -representable. If all the sets in some $L_\delta(V_{\lambda+1})$, with δ limit, are $\mathbb{U}(j)$ -representable, this implies that we cannot find two disjoint stationary sets in $L_\delta(V_{\lambda+1})$. If all the sets are $\mathbb{U}(j)$ -representable, this means that the club filter is an ultrafilter on cofinality ω .

Theorem 8.6 (Woodin, Theorem 16 in [?]). *Suppose $I0(\lambda)$. Then in a generic extension $I0(\lambda) +$ the Weak Ultrafilter Axiom hold.*

Another problematic point is the Wadge hierarchy. Wadge's Lemma states that under AD for any two subsets A, B of \mathbb{R} either there is a continuous function f with $A = f^{-1}[B]$ or there is one with $B = f^{-1}[\mathbb{R} \setminus A]$. Therefore any two subsets of \mathbb{R} are reducible one to another. Under I0 this is consistently false:

Theorem 8.7 (Woodin, Theorem 174 in [?]). *Suppose $I0(\lambda)$. Suppose c is a generic Cohen real of V . Then there exist $X, Y \in V[c]_{\lambda+2}$ such that $X \notin L_\omega(Y, V[c]_{\lambda+1})$ and $Y \notin L_\omega(X, V[c]_{\lambda+1})$.*

As by ?? in $L(V[c]_{\lambda+1})$ $I0(\lambda)$ holds, this proves the consistency of I0 with a negation of Wadge Lemma.

Another aspect one can consider is partition combinatorics. In $L(\mathbb{R})$, under AD, for all $\alpha < \omega_1$, $\omega_1 \rightarrow (\omega_1)_\omega^{\alpha 9}$. A direct generalization always fails:

Remark 8.8 (ZF+DC). *Suppose there exists an ω_1 -sequence of distinct reals. Then there is no κ such that $\kappa \rightarrow (\kappa)_2^{\omega_1}$.*

So the best attempt could only be $\lambda^+ \rightarrow (\lambda^+)_\lambda^\omega$. It is still an open question even whether $\kappa \rightarrow (\kappa)_2^\omega$ is consistent at all. In [?] Lemma 212 proves that a universal strong version of $\lambda^+ \rightarrow (\lambda^+)_\lambda^\omega$ is consistently false with I0.

But there is a partial result:

Proposition 8.9 (Lemma 180 in [?]). *Suppose $I0(\lambda)$. Then for all $\alpha < \lambda$, $L(V_{\lambda+1}) \models \lambda^+ \rightarrow (\lambda^+)_{(\lambda, < \lambda)}^\alpha$, i.e., for all $\pi : \{\sigma \subseteq \lambda^+ : \text{ot}(\sigma) = \alpha\} \rightarrow \lambda$ there exists $H \subseteq \lambda^+$ such that $|\{\pi(\sigma) : \sigma \subseteq H, \text{ot}(\sigma) = \alpha\}| < \lambda$.*

Finally, we consider the measure on Turing degrees. A standard result of AD is that the filter $U = \{A : A \text{ contains a cone}\}$ on sets of Turing degrees is an ultrafilter (Martin), i.e., every set of degrees either contains or is disjoint from a cone. This result is usually called Turing Determinacy, and it is at the base of many results, like Silver Dichotomy for every set in $L(\mathbb{R})$ under AD. In the context of I0 Turing degrees can be substituted with Zermelo degrees: given $a \subseteq \lambda$, let $M(a)$ be the smallest model of Zermelo set theory that contains a . We say that a is equivalent with b if $M(a) = M(b)$, and a is reducible to b if $M(a) \subseteq M(b)$. Then a Zermelo degree is a class of such equivalence relation. Unfortunately Zermelo Degree Determinacy is consistently false under I0:

Theorem 8.10 (Shi, 2015, [?]). *Suppose $I0(\lambda)$ and that all the λ -supercompact cardinals under λ are Laver indestructible. Then there is a set of Zermelo degrees in $L(V_{\lambda+1})$ that neither contains nor is disjoint from a cone.*

This result is significantly more problematic respect the previous results: in ?? and Lemma 212 in [?] we had that small changes would destroy that $\text{AD}^{L(\mathbb{R})}$ -like properties, but in this case, modulo a reasonable indestructibility preparation, it seems that it is intrinsic from I0, and for deep reasons. But there is one ambiguity still to settle: for now, there is no proof that there are λ singular cardinals of cofinality ω such that in $L(V_{\lambda+1})$ Zermelo Degree

⁹Recall that $\kappa \rightarrow (\kappa)_\beta^\alpha$ is that if $\pi : \{\sigma \subseteq \kappa : \text{ot}(\sigma) = \alpha\} \rightarrow \beta$ then there is a set $H \subseteq \kappa$ of cardinality κ homogeneous for π , i.e., $|\{\pi(\sigma) : \sigma \subseteq H, \text{ot}(\sigma) = \alpha\}| = 1$.

Determinacy holds, so maybe its failure is not something peculiar to I_0 , but a general property of such λ 's, just like it is not possible to have a club filter an ultrafilter everywhere or the full partition property.

Question 8.11. *Is there a λ such that Zermelo Degree Determinacy holds in $L(V_{\lambda+1})$?*

9 Further Developments

The main line of research right now is finding an answer for Question ???. The research is ongoing and very promising: if the answer is yes, then not only many already known results will get unlocked (for example the Ultrafilter Axiom at ω or the full power of Generic Absoluteness), but it would give an elegant tree representation to all subsets of $V_{\lambda+1}$. The potential would be huge, just like what happened for $AD^{L(\mathbb{R})}$.

Another approach would be to investigate the inner model theory of I_0 . Here the main result is this:

Theorem 9.1 (Woodin, 2011, [?], [?]). *Let δ be an extendible cardinal. Assume that N is a weak extender model for δ supercompact and $\gamma > \delta$ is a cardinal in N . Let $j : H(\gamma^+)^N \rightarrow H(j(\gamma)^+)^N$ be an elementary embedding with $\delta \leq \text{crt}(j)$ and $j \neq \text{id}$. Then $j \in N$.*

In other words, if there is a (reasonable) canonical inner model for supercompactness and I_0 holds, then I_0 must hold in the inner model. There is no hope therefore to reach results of consistency equivalence to I_0 via inner models, and for this reason Woodin conceived Ultimate- L , a canonical inner model for all large cardinals. If such hypothesis will prove solid, then we would have an actual canonical model, a preferred V , to work with. This would overcome all the problems in the Section ??, and moreover it would be interesting to understand the role of I_0 in this setting.

Question 9.2. *Is it true that $\text{Ultimate-}L \models I_0(\lambda)$ iff $\text{Ultimate-}L \models L(V_{\lambda+1}) \not\models AC$?*

One can ask if there are other axioms above I_0 , still in ZFC. The most natural approach would be to add to $L(V_{\lambda+1})$ the measures that define j , just like for $L[\mu]$, but this is still open:

Question 9.3. *Let $j : L(V_{\lambda+1}) \prec L(V_{\lambda+1})$ weakly proper, and U_j that defines it. Is $V \neq L(V_{\lambda+1})[U_j]$?*

So another approach is to consider $X \subseteq V_{\lambda+1}$, and build from there.

Definition 9.4. *A set $X \subseteq V_{\lambda+1}$ is an Icarus set iff there is an elementary embedding $j : L(X, V_{\lambda+1}) \prec L(X, V_{\lambda+1})$ with $\text{crt}(j) < \lambda$.*

The name comes from the fact that we are trying to go as high as possible before reaching an inconsistency. There are still many open problems in this direction. For example, in the I_0 case we had that a weakly proper embedding was nicely amenable, and for every $a \in L(V_{\lambda+1})$ if we defined $a_0 = a$ and $a_{n+1} = j(a_n)$, then $\langle a_n : n \in \omega \rangle \in L(V_{\lambda+1})$. This is because everything is

definable from $V_{\lambda+1}$ and fixed points, and $V_{\lambda+1}$ is morally closed under ω -sequences. But in the Icarus case it is not clear whether it is necessary that $\langle X_n : n \in \omega \rangle \in L(X, V_{\lambda+1})$.

Definition 9.5 (Woodin, [?]). *Let $X \subseteq V_{\lambda+1}$ be an Icarus set, and let $j : L(X, V_{\lambda+1}) \prec L(X, V_{\lambda+1})$ with $\text{crt}(j) < \lambda$. Then:*

- *j is weakly proper iff j is an ultrapower embedding;*
- *j is proper iff it is weakly proper and $\langle X_n : n \in \omega \rangle \in L(X, V_{\lambda+1})$, with $X_0 = X$ and $X_{n+1} = j(X_n)$.*

Question 9.6. *Is there an Icarus set $X \subseteq V_{\lambda+1}$ and an elementary embedding $j : L(X, V_{\lambda+1}) \prec L(X, V_{\lambda+1})$ that is weakly proper but not proper?*

In [?] and [?] there are partial results in this direction.

It turns out that the sharps build a whole hierarchy of Icarus sets:

Proposition 9.7. *Let λ be a cardinal. Then $V_{\lambda+1}$ is Icarus is strongly implied by $(V_{\lambda+1})^\sharp$ is Icarus, that is strongly implied by $(V_{\lambda+1})^{\sharp\sharp}$ is Icarus, etc. In general, $(V_{\lambda+1})^{(n+1)\sharp}$ is Icarus strongly implies that $(V_{\lambda+1})^{n\sharp}$ is Icarus. Moreover, if $X \subseteq V_{\lambda+1}$ is Icarus and there exists $n \in \omega$ such that $\Theta^{L(X, V_{\lambda+1})} < \Theta^{L((V_{\lambda+1})^{n\sharp}, V_{\lambda+1})}$, then there exists an $m < n$ such that $\Theta^{L(X, V_{\lambda+1})} = \Theta^{L((V_{\lambda+1})^{m\sharp}, V_{\lambda+1})}$ and for any $s \leq m$ $(V_{\lambda+1})^{s\sharp}$ is Icarus.*

Pushing this result above the first ω sharps is more difficult. One can try to define $Y_{\alpha+1} = Y_\alpha^\sharp$ and $Y_\gamma = \bigcup_{\alpha < \gamma} Y_\alpha$ for γ limit, but at a certain point there would be a γ limit such that Y_γ is not a subset of $V_{\lambda+1}$, so then $(Y_\gamma)^\sharp$ is still not a subset of $V_{\lambda+1}$, and so on. The way to do it is to enlarge our analysis to $L(N)$ with $V_{\lambda+1} \subset N \subset V_{\lambda+2}$, and if necessary at the successor of a limit stage adding just less of N^\sharp so that everything is codeable with a subset of $V_{\lambda+1}$. It is a complex endeavour, that takes all Section 4 of [?].

Finally, there is the Reinhardt cardinal approach, i.e., large cardinals in ZF without AC. There is a whole hierarchy above, like super-Reinhardt cardinals and Berkeley cardinals, and it goes frontally against the Ultimate- L approach. Yet, the situation about Reinhardt cardinals is similar to the I0 one: very few published results (for example [?]), and a lot of underground activity, via lecture notes, seminar slides, Wikipedia pages, MathOverflow questions, The possible intersections with inner model theory, though, promise to make this topic a catalyst of set theorists' attention in the near future, and the hope is that this will bring more publications, and maybe even a survey, of this exciting topic.

Acknowledgments. The paper was written under the Italian program “Rita Levi Montalcini 2013”, but it is a collection of notes, thoughts, sketches and ideas that started in the Summer Semester of 2008 at Berkeley, propelled by a course on “Beyond I0” by Woodin. With nine years in the making it is therefore a hard task to thank all the people that helped me writing this paper, as almost all the people I interacted with in my academic career has been an inspiration for this, directly or indirectly. Knowing that this is just a small percentage of the whole picture, I would like to thank first of all Hugh Woodin, the demiurge of the I0 world, that welcomed me in Berkeley when I was still a student and continued to be a creative influence during the years; Alessandro Andretta, for

his steady support in all this years, and because of his (decisive) insistence for this survey to be written. I would also like to thank Sy Friedman and Liuzhen Wu, that helped to push my research in an unexpected direction, and Scott Cramer and Xianghui Shi, for the many, many discussions on I0 and similars. Special thanks also for Luca Motto Ros, that helped me in shaping this paper.

References

- [1] A. W. Apter, G. Sargsyan, *Jonsson-like partition relations and $j : V \rightarrow V$* . Journal of Symbolic Logic **69** (2004), 1267–1281.
- [2] P. Corazza, *The wholeness axiom and Laver sequences*. Annals of Pure and Applied Logic **105** (2000), 157–260.
- [3] P. Corazza, *Lifting elementary embeddings $j : V_\lambda \rightarrow V_\lambda$* . Archive for Mathematical Logic **46** (2007), 61–72.
- [4] S. Cramer, *Inverse limit reflection and the structure of $L(V_{\lambda+1})$* . Journal of Mathematical Logic **15** (2015), 1–38.
- [5] J. Cummings, M. Foreman, *Diagonal Prikry extensions*. Journal of Symbolic Logic **75** (2010), 1383–1402.
- [6] P. Dehornoy, *Braid groups and left distributive operations*. Transactions of the American Mathematical Society **345** (1994), 115–151.
- [7] V. Dimonte, *Totally non-proper ordinals beyond $L(V_{\lambda+1})$* . Archive for Mathematical Logic **50** (2011), 565–584.
- [8] V. Dimonte, *A partially non-proper ordinal beyond $L(V_{\lambda+1})$* . Annals of Pure and Applied Logic **163** (2012), 1309–1321.
- [9] V. Dimonte, S. Friedman, *Rank-into-rank hypotheses and the failure of GCH*. Archive for Mathematical Logic **53** (2014), 351–366.
- [10] V. Dimonte, L. Wu, *A general tool for consistency results related to I1*. European Journal of Mathematics **2** (2016), 474–492.
- [11] V. Dimonte, L. Wu, *The \ast -Prikry condition*. Unpublished manuscript.
- [12] N. Dobrinen, S. Friedman, *The consistency strength of the tree property at the double successor of a measurable cardinal*. Fundamenta mathematicae **208** (2010), 123–153.
- [13] R. Dougherty, T.J. Jech, *Finite left-distributive algebras and embedding algebras*. Advances in Mathematics **130** (1997), 201–241.
- [14] W. B. Easton, *Powers of regular cardinals*. Annals of Mathematical Logic **1** (1970), 139–178.
- [15] M. Foreman, A. Kanamori (eds), *Handbook of Set theory*. Springer, Dordrecht (2010).

- [16] S.-D. Friedman, *Large cardinals and L -like universes*. Set Theory: Recent Trends and Applications (A. Andretta editor). Quaderni di Matematica **17** (2006), 93–110.
- [17] M. Gitik, *Blowing up the power of a singular cardinal - wider gaps*. Annals of Pure and Applied Logic **116** (2002), 1–38.
- [18] M. Gitik, *Prikry-type forcings*. Handbook of set theory, Vols 2, 1351–1447. Springer, Dordrecht 2010.
- [19] G. Kafkoulis, *Coding lemmata in $L(V_{\lambda+1})$* . Archive for Mathematical Logic **43** (2004), 193–213.
- [20] A. Kanamori, *The Higher Infinite*. Springer, Berlin (1994).
- [21] J. Hamkins, *Fragile measurability*. Journal of Symbolic Logic **59** (1994), 262–282.
- [22] J. D. Hamkins, G. Kirmayer, and N. L. Perlmutter, *Generalizations of the Kunen inconsistency*. Annals of Pure and Applied Logic **163** (2012) 1872–1890.
- [23] K. Kunen, *Elementary embeddings and infinite combinatorics*. Journal of Symbolic Logic **36** (1971), 407–413.
- [24] R. Laver, *The left distributive law and the freeness of an algebra of elementary embeddings*. Advances in Mathematics **91** (1992), 209–231.
- [25] R. Laver, *On the algebra of elementary embeddings of a rank into itself*. Advances in Mathematics **110** (1995), 334–346.
- [26] R. Laver, *Implications between strong large cardinal axioms*. Annals of Pure and Applied Logic **90** (1997), 79–90.
- [27] R. Laver, *Reflection of elementary embedding axioms on the $L(V_{\lambda+1})$ hierarchy*. Annals of Pure and Applied Logic **107** (2001), 227–238.
- [28] D.A. Martin, *Infinite games*. Proceedings of the ICM, Helsinki, 1978, Acad. Sci. Fennica, Helsinki, 269–273.
- [29] A. R. D. Mathias, *On sequences generic in the sense of Prikry*. Journal of the Australian Mathematical Society **15** (1973), 409–414.
- [30] C. Merimovich, *Extender Based Radin Forcing*. Transactions of the American Mathematical Society **355** (2003), 1729–1772.
- [31] Y.N. Moschovakis, *Determinacy and prewellorderings of the continuum*. In Y. Bar-Hillel, *Mathematical Logic and Foundations of Set Theory*, Amsterdam, North-Holland, 1970.
- [32] I. Neeman, *Aronszajn trees and failure of the Singular Cardinal Hypothesis*. Journal of Mathematical Logic **9** (2009), 139–157.
- [33] X. Shi, *Axiom I_0 and higher degree theory*. Journal of Symbolic Logic **80** (2015), 970–1021.

- [34] X. Shi, N. Trang, *I_0 and combinatorics at λ^+* . Archive for Mathematical Logic **56** (2017), 131–154.
- [35] J. Steel, *Scales in $L(\mathbb{R})$* . Cabal Seminar 79–81, 107–156, Lecture Notes in Mathematics, 1019, Springer, Berlin, 1983.
- [36] A. Suzuki, *No elementary embedding from V into V is definable from parameters*. Journal of Symbolic Logic, **64** (1999), 1591–1594.
- [37] W. H. Woodin, *An AD-like Axiom*. Lecture notes written by George Kafkoulis, 1989
- [38] W. H. Woodin, *Suitable extender models I*. Journal of Mathematical Logic, **10** (2010) 101–339.
- [39] W. H. Woodin, *Suitable extender models II: beyond ω -huge*. Journal of Mathematical Logic, **11** (2011) 115–436.
- [40] W. H. Woodin, *The Weak Ultrafilter Axiom*. Archive for Mathematical Logic, **55** (2016) 319–351.
- [41] W. H. Woodin, J. Davis, D. Rodríguez, *The HOD Dichotomy*. Unpublished lecture notes.