



UNIVERSITÀ
DEGLI STUDI
DI UDINE

Università degli studi di Udine

CLASSIFICATION OF BLOW-UP LIMITS FOR THE SINH-GORDON EQUATION

Original

Availability:

This version is available <http://hdl.handle.net/11390/1171004> since 2019-12-06T15:04:04Z

Publisher:

Published

DOI:

Terms of use:

The institutional repository of the University of Udine (<http://air.uniud.it>) is provided by ARIC services. The aim is to enable open access to all the world.

Publisher copyright

(Article begins on next page)

CLASSIFICATION OF BLOW-UP LIMITS FOR THE SINH-GORDON EQUATION

ALEKS JEVIKAR, JUNCHENG WEI, WEN YANG

ABSTRACT. The aim of this paper is to use a selection process and a careful study of the interaction of bubbling solutions to show a classification result for the blow-up values of the elliptic sinh-Gordon equation

$$\Delta u + h_1 e^u - h_2 e^{-u} = 0 \quad \text{in } B_1 \subset \mathbb{R}^2.$$

In particular we get that the blow-up values are multiple of 8π . It generalizes the result of Jost, Wang, Ye and Zhou [20] where the extra assumption $h_1 = h_2$ is crucially used.

1. INTRODUCTION

In this paper we mainly focus on the weak limit of the energy sequence for the following equation

$$\Delta u + h_1 e^u - h_2 e^{-u} = 0 \text{ in } B_1 \subset \mathbb{R}^2, \quad (1.1)$$

where h_1, h_2 are smooth positive functions and B_1 is the unit ball in \mathbb{R}^2 .

Equation (1.1) arises in the study of the equilibrium turbulence with arbitrarily signed vortices [11, 31, 29, 32], and it was first proposed by Onsager [35], Joyce and Montgomery [21] from different statistical arguments. When the nonlinear term e^{-u} in (1.1) is replaced by $\tau e^{-\gamma u}$ with $\tau, \gamma > 0$, the equation (1.1) describes a more general type of equation which arises in the context of the statistical mechanics description of 2D-turbulence. For the recent developments of such equation, we refer the readers to [37, 38, 39] and the references therein. Moreover, it also plays a very important role in the study of the construction of constant mean curvature surfaces initiated by Wentz, see [20, 46].

When $h_2 \equiv 0$ the equation (1.1) is reduced to the classic Liouville equation

$$\Delta u + h_1 e^u = 0 \quad \text{in } B_1 \subset \mathbb{R}^2. \quad (1.2)$$

Equation (1.2) is important in the geometry of manifolds as it rules the change of Gaussian curvature under conformal deformation of the metric, see [1, 7, 8, 23, 42]. Another motivation for the study of (1.2) is in mathematical physics as it models the mean field of Euler flows, see [6] and [22]. This equation has attracted a lot of attentions in past decades; now there are many results regarding the existence, compactness of the solutions, the bubbling behavior, etc. We refer the interested reader to the reviews [30] and [45].

2000 *Mathematics Subject Classification.* 35J61, 35R01, 35B44.

Key words and phrases. Geometric PDEs, sinh-Gordon equation, Blow-up analysis.

A.J. is partially supported by the PRIN project *Variational and perturbative aspects of nonlinear differential problems*. J.W. and W.Y. are partially supported by the NSERC of Canada. Part of the paper was written when A.J. was visiting the Mathematics Department at the University of British Columbia. He would like to thank the institute for the hospitality and for the financial support.

Wente's work [46] on the constant mean curvature surfaces and the work of Sacks-Uhlenbeck [40] concerning harmonic maps led to investigate the blow-up phenomena for variational problems that possess a lack of compactness. Later, in a series work by Steffen [43], Struwe [44] and Brezis, Coron [4], the program of understanding the blow-up phenomena for constant mean curvature surfaces was completed.

As many geometric problems, problem (1.1) (and (1.2)) presents loss of compactness phenomena, as its solutions might blow-up. Concerning (1.2) it was proved in [5, 24, 25] that for a sequence of blow-up solutions u_k to (1.2) (relatively to h_1^k) with blow-up point \bar{x} it holds

$$\lim_{\delta \rightarrow 0} \lim_{k \rightarrow \infty} \int_{B_\delta(\bar{x})} h_1^k e^{u_k} = 8\pi. \quad (1.3)$$

Somehow, each blow-up point has a quantized local mass.

On the other hand, the blow-up behavior of the solutions of equation (1.1) is not yet developed in full generality; this analysis was carried out in [33, 34] and [20] under the assumption that $h_1 = h_2$ or h_1, h_2 are constants. In particular, by using the deep connection of the sinh-Gordon equation and differential geometry, Jost, Wang, Ye and Zhou in [20] proved an analogous quantization property as the one in (1.3), namely that the blow-up limits are multiple of 8π , see Theorem 1.1, Corollary 1.2 and Remark 4.5 in [20]. The blow-up phenomena indeed occurs, see [12] and [13]. We point out that the assumption $h_1 = h_2$ (or h_1, h_2 constants) in [20] is crucially used in order to provide a geometric interpretation of equation (1.1) in terms of constant mean curvature surfaces and harmonic maps (see also [46]). In this way they transfer the problem into a blow-up phenomena for harmonic maps. The core of the argument is then to apply a result about no loss of energy during bubbling off for a sequence of harmonic maps, which was proved in [19, 36].

The study of the blow-up limits is interesting by itself. However, it yields also important results: we point out here the compactness property of the following sequence of solutions to a variant of (1.1) which is related to a mean field equation:

$$\Delta u_k + \rho_1^k \left(\frac{H_1 e^{u_k}}{\int_M H_1 e^{u_k}} - \frac{1}{|M|} \right) - \rho_2^k \left(\frac{H_2 e^{-u_k}}{\int_M H_2 e^{-u_k}} - \frac{1}{|M|} \right) = 0 \quad \text{on } M, \quad (1.4)$$

where M is a compact surface without boundary, ρ_1^k, ρ_2^k are non-negative real parameters and H_1, H_2 are two *fixed* positive smooth functions (see [2, 15, 16, 17, 47] and the references therein). In fact, from the local quantization result in [20] and some standard analysis (see [3, 5, 26]) one finds the following global compactness result.

Theorem A. *Let $\rho_1^k, \rho_2^k \notin 8\pi\mathbb{N}$ and suppose that $H_1 = H_2$. Then the solutions to (1.4) are uniformly bounded.*

The latter property is a key ingredient in proving both existence and multiplicity results of (1.4), see for example [2, 15, 16, 17].

Now we return to the topic of this paper. We shall study the same subject of [20] in a more general case (i.e., h_1, h_2 are two different positive C^3 functions) by using pure analytic method. The argument is interesting by itself and it seems the first time that this method is used for this class of equations.

Let u_k be a sequence of blow-up solutions

$$\Delta u_k + h_1^k e^{u_k} - h_2^k e^{-u_k} = 0, \quad (1.5)$$

with 0 being its only blow-up point in B_1 , i.e.:

$$\max_{K \subset \subset B_1 \setminus \{0\}} |u_k| \leq C(K), \quad \max_{x \in B_1} |u_k(x)| \rightarrow \infty. \quad (1.6)$$

Throughout the paper we will call $\int_{B_1} h_1^k e^{u_k}$ the energy of u_k (a corresponding term $\int_{B_1} h_2^k e^{-u_k}$ is defined for the energy of $-u_k$). We assume

$$h_i^k(0) = 1, \quad \frac{1}{C} \leq h_i^k(x) \leq C, \quad \|h_i^k(x)\|_{C^3(B_1)} \leq C, \quad \forall x \in B_1, \quad i = 1, 2 \quad (1.7)$$

for some positive constant C . In addition, we assume u_k has bounded oscillation on ∂B_1 and a uniform bound on its energy:

$$\begin{aligned} |u_k(x) - u_k(y)| &\leq C, \quad \forall x, y \in \partial B_1, \\ \int_{B_1} h_1^k e^{u_k} &\leq C, \quad \int_{B_1} h_2^k e^{-u_k} \leq C, \end{aligned} \quad (1.8)$$

where C is independent of k .

Our main result is concerned with the limit energy of u_k . Let

$$\sigma_1 = \lim_{\delta \rightarrow 0} \lim_{k \rightarrow \infty} \frac{1}{2\pi} \int_{B_\delta} h_1^k e^{u_k}, \quad \sigma_2 = \lim_{\delta \rightarrow 0} \lim_{k \rightarrow \infty} \frac{1}{2\pi} \int_{B_\delta} h_2^k e^{-u_k}, \quad (1.9)$$

be the local blow-up masses and $\Sigma \subset \mathbb{R}^2$ be the following set:

$$\Sigma = \left\{ (\sigma_1, \sigma_2) = (2m(m+1), 2m(m-1)) \text{ or } (2m(m-1), 2m(m+1)), \quad m \in \mathbb{N} \right\}. \quad (1.10)$$

Theorem 1.1. *Let σ_i and Σ be defined as in (1.9) and (1.10) respectively. Suppose u_k satisfies (1.5), (1.6), (1.8) and h_i^k satisfy (1.7). Then $(\sigma_1, \sigma_2) \in \Sigma$. In particular both σ_1 and σ_2 are multiples of 4.*

Remark 1.1. *Theorem 1.1 yields an improvement of the compactness result in Theorem A, which holds now for arbitrary functions H_1, H_2 . As a byproduct we get an improvement of both existence and multiplicity results concerning (1.4) in [2, 16, 17]. Moreover, it is used in [18] where the authors are concerned with the Leray-Schauder topological degree associated to (1.4).*

Remark 1.2. *(1.1) is different from the Liouville equation (1.2) and the system of two equations in [27]. For the latter ones, the blow-up limits (see for example (1.3)) only possess a finite number of possibilities, while for (1.1) we obtain an infinite number of possibilities for (1.9), see (1.10). The reason is that we have a different form of the Pohozaev identity for problem (1.5), see Proposition 3.1.*

The first step in the proof of Theorem 1.1 is to introduce a selection process for describing the situations when blow-up of solutions of (1.5) occurs. This argument has been widely used in the framework of prescribed curvature problems, see for example [9, 23, 41]. It was later modified by Lin, Wei and Zhang in dealing with general systems of n equations to locate the bubbling area which consists of a finite number of disks, see [27]. Roughly speaking, the idea is that in each disk the blow-up solution have the energy of a globally defined system. We shall use the same technique for equation (1.1). Next we prove that in each bubbling disk the local mass of at least one of u_k and $-u_k$ is a perturbation of multiple of 4. Then we combine the areas which are closed to each other and deduce that at least

one component of u_k and $-u_k$ has its local mass equals to a small perturbation of multiples of 4. In this procedure we use the same terminology "group" introduced in [27] to describe bubbling disks which are close to each other and relatively far away from the other disks. Then, Theorem 1.1 is a consequence of an energy quantization for the global Liouville equation with singularities proved in [28] and of the Pohozaev identity Proposition 3.1.

The organization of this paper is as follows. In Section 2 we introduce the selection process for the class of equations as in (1.1), in Section 3 we prove a Pohozaev identity which is the key element in proving Theorem 1.1, in Section 4 we study the asymptotic behavior of the solutions around the blow-up area and in Section 5 we finally prove Theorem 1.1 by a suitable combination of the bubbling areas.

Notation

The symbol $B_r(p)$ stands for the open metric ball of radius r and center p . To simplify the notation we will write B_r for balls which are centered at 0. We will use the notation $a \sim b$ for two comparable quantities a and b .

Throughout the paper the letter C will stand for some constants which are allowed to vary among different formulas or even within the same lines. When we want to stress the dependence of the constants on some parameter (or parameters), we add subscripts to C , as C_δ , etc. We will write $o_\alpha(1)$ to denote quantities that tend to 0 as $\alpha \rightarrow 0$ or $\alpha \rightarrow +\infty$ and use the symbol $O_\alpha(1)$ for bounded quantities.

2. A SELECTION PROCESS FOR THE SINH-GORDON EQUATION

In this section we introduce a selection process for the sinh-Gordon equation (1.1). In particular, we will select a finite number of bubbling areas. This will be the first tool to be used in the proof of the Theorem 1.1.

Proposition 2.1. *Let u_k be a sequence of blow-up solutions to (1.5) that satisfy (1.6) and (1.8), and suppose that h_i^k satisfy (1.7). Then, there exist finite sequences of points $\Sigma_k := \{x_1^k, \dots, x_m^k\}$ (all $x_j^k \rightarrow 0$, $j = 1, \dots, m$) and positive numbers $l_1^k, \dots, l_m^k \rightarrow 0$ such that*

$$(1) |u_k|(x_j^k) = \max_{B_{l_j^k}(x_j^k)} \{|u_k|\} \text{ for } j = 1, \dots, m.$$

$$(2) \exp\left(\frac{1}{2}|u_k|(x_j^k)\right) l_j^k \rightarrow \infty \text{ for } j = 1, \dots, m.$$

$$(3) \text{ Let } \varepsilon_k = e^{-\frac{1}{2}M_k}, \text{ where } M_k = \max_{B_{l_j^k}(x_j^k)} |u_k|. \text{ In each } B_{l_j^k}(x_j^k) \text{ we define the}$$

dilated functions

$$\begin{aligned} v_1^k(y) &= u_k(\varepsilon_k y + x_j^k) + 2 \log \varepsilon_k, \\ v_2^k(y) &= -u_k(\varepsilon_k y + x_j^k) + 2 \log \varepsilon_k. \end{aligned} \tag{2.1}$$

Then it holds that one of the v_1^k, v_2^k converges to a function v in $C_{loc}^2(\mathbb{R}^2)$ which satisfies the Liouville equation (1.2), while the other one tends to minus infinity over all compact subsets of \mathbb{R}^2 .

(4) *There exists a constant $C_1 > 0$ independent of k such that*

$$|u_k|(x) + 2 \log \text{dist}(x, \Sigma_k) \leq C_1, \quad \forall x \in B_1.$$

Proof. Without loss of generality we may assume that

$$u_k(x_1^k) = \max_{x \in B_1} |u_k|(x).$$

By assumption we clearly have $x_1^k \rightarrow 0$. Let (v_1^k, v_2^k) be defined as in (2.1) with x_j^k , M_k replaced by x_1^k and $u_k(x_1^k)$ respectively. By (2.1) and the definition of ε_k , we have $v_i^k \leq 0, i = 1, 2$. Therefore, exploiting the equation (1.5) we can easily see that $|\Delta v_i^k|$ is bounded. By standard elliptic estimate, $|v_i^k(z) - v_i^k(0)|$ is uniformly bounded on any compact subset of \mathbb{R}^2 . By construction, we have $v_1^k(0) = 0$ and v_1^k converges in $C_{loc}^2(\mathbb{R}^2)$ to a function v_1 , while the other component is forced to satisfy $v_2^k \rightarrow -\infty$ over all compact subsets of \mathbb{R}^2 . In addition, v_1 satisfies the following equation

$$\Delta v_1 + e^{v_1} = 0 \quad \text{in } \mathbb{R}^2, \quad (2.2)$$

where we used $\lim_{k \rightarrow +\infty} h_i^k(x_1^k) = 1$. From (1.8), we have

$$\int_{\mathbb{R}^2} h_1 e^{v_1} < C.$$

By the classification result due to Chen and Li [10] it follows that

$$\int_{\mathbb{R}^2} e^{v_1} = 8\pi \quad \text{and} \quad v_1(x) = -4 \log |x| + O(1), \quad |x| > 2.$$

Clearly we can take $R_k \rightarrow \infty$ such that

$$v_1^k(y) + 2 \log |y| \leq C, \quad |y| \leq R_k. \quad (2.3)$$

In other words we can find $l_1^k \rightarrow 0$ such that

$$u_k(x) + 2 \log |x - x_1^k| \leq C, \quad |x - x_1^k| \leq l_1^k,$$

and

$$e^{\frac{1}{2} u_1^k(x_1^k)} l_1^k \rightarrow \infty, \quad \text{as } k \rightarrow \infty.$$

Now we consider the function

$$|u_k(x)| + 2 \log |x - x_1^k|$$

and let q_k be the point where $\max_{|x| \leq 1} (|u_k(x)| + 2 \log |x - x_1^k|)$ is achieved. Suppose that

$$\max_{|x| \leq 1} (|u_k(x)| + 2 \log |x - x_1^k|) \rightarrow \infty. \quad (2.4)$$

Then we define $d_k = \frac{1}{2} |q_k - x_1^k|$ and

$$\begin{aligned} S_1^k(x) &= u_k(x) + 2 \log(d_k - |x - q_k|), \\ S_2^k(x) &= -u_k(x) + 2 \log(d_k - |x - q_k|), \end{aligned} \quad \text{in } B_{d_k}(q_k).$$

We note that $S_i^k(x) \rightarrow -\infty$ as x approaches $\partial B_{d_k}(q_k)$ and

$$\max\{S_1^k(q_k), S_2^k(q_k)\} = |u_k(q_k)| + 2 \log d_k \rightarrow \infty$$

by assumption (2.4). Let p_k be the point where $\max_{x \in \bar{B}_{d_k}(q_k)} \{S_1^k, S_2^k\}$ is attained.

Without loss of generality, we assume that $S_2^k(p_k) = \max_{x \in \bar{B}_{d_k}(q_k)} \{S_1^k, S_2^k\}$. Then

$$-u_k(p_k) + 2 \log(d_k - |p_k - q_k|) \geq \max\{S_1^k(q_k), S_2^k(q_k)\} \rightarrow \infty. \quad (2.5)$$

Let $l_k = \frac{1}{2}(d_k - |p_k - q_k|)$. By the definition of p_k and l_k we observe that, for $y \in B_{l_k}(p_k)$ it holds

$$\begin{aligned} |u_k(y)| + 2 \log(d_k - |y - q_k|) &\leq -u_k(p_k) + 2 \log(2l_k), \\ d_k - |y - q_k| &\geq d_k - |p_k - q_k| - |y - p_k| \geq l_k. \end{aligned}$$

Therefore we get

$$|u_k(y)| \leq -u_k(p_k) + 2 \log 2, \quad \forall y \in B_{l_k}(p_k). \quad (2.6)$$

Let $R_k = e^{-\frac{1}{2}u_k(p_k)}l_k$ and \hat{v}_i^k , $i = 1, 2$ be the following functions:

$$\begin{aligned} \hat{v}_1^k(y) &= u_k(p_k + e^{\frac{1}{2}u_k(p_k)}y) + u_k(p_k), \\ \hat{v}_2^k(y) &= -u_k(p_k + e^{\frac{1}{2}u_k(p_k)}y) + u_k(p_k). \end{aligned}$$

It is not difficult to see that $R_k \rightarrow \infty$ from (2.5). Moreover, $|\Delta \hat{v}_i^k|$ is bounded in $B_{R_k}(0)$. Similarly as before $\hat{v}_2^k(y)$ converges to a function v_2 which satisfies

$$\Delta v_2 + e^{v_2} = 0.$$

On the other hand, $\hat{v}_1^k(y)$ converges uniformly to $-\infty$ over all compact subsets of \mathbb{R}^2 . Next, we study the behavior of $u_k, -u_k$ in $B_{l_k}(p_k)$. Suppose x_2^k is the point where $\max_{B_{l_k}(p_k)} |u_k|$ is obtained: it is not difficult to see that $-u_k(x_2^k) = \max_{B_{l_k}(p_k)} |u_k|$. Moreover, we can find l_2^k such that

$$|u_k(x)| + 2 \log |x - x_2^k| \leq C, \quad \text{for } |x - x_2^k| \leq l_2^k.$$

By (2.6) we have $-u_k(x_2^k) + u_k(p_k) \leq 2 \log 2$ and observe that

$$\hat{v}_2(e^{-\frac{1}{2}u_k(p_k)}(x_2^k - p_k)) - \hat{v}_2(0) = -u_k(x_2^k) + u_k(p_k) \leq 2 \log 2.$$

Therefore we deduce that $e^{-\frac{1}{2}u_k(p_k)}|x_2^k - p_k| = O(1)$. As a consequence, we can choose $l_2^k \leq \frac{1}{2}l_k$ such that $e^{-\frac{1}{2}u_k(p_k)}l_2^k \rightarrow \infty$. Then we re-scale $u_k, -u_k$ around x_2^k and let v_i^k defined in (2.1) and it satisfies (1) and (2) in Proposition 2.1. Moreover, it is easy to see that $B_{l_1^k}(x_1^k) \cap B_{l_2^k}(x_2^k) = \emptyset$. In this way we have defined the selection process. To continue it, we let $\Sigma_{k,2} := \{x_1^k, x_2^k\}$ and consider

$$\max_{x \in B_1} |u_k(x)| + 2 \log \text{dist}(x, \Sigma_{k,2}).$$

If there exists a subsequence such that the quantity above tends to infinity, we can use the same argument to get x_3^k and l_3^k . Since each bubble area contributes a positive energy, the process stops after finite steps due to the bound on the energy (1.8). Finally we get

$$\Sigma_k = \{x_1^k, \dots, x_m^k\}$$

and it holds

$$|u_k(x)| + 2 \log \text{dist}(x, \Sigma_k) \leq C, \quad (2.7)$$

which concludes the proof. \square

Lemma 2.1. *Let $\Sigma_k = \{x_1^k, \dots, x_m^k\}$ be the blow-up set obtained in Proposition 2.1. Then for all $x \in B_1 \setminus \Sigma_k$, there exists a constant C independent of x and k such that*

$$|u_k(x_1) - u_k(x_2)| \leq C, \quad \forall x_1, x_2 \in B(x, d(x, \Sigma_k)/2).$$

Proof. Using the Green's representation formula it is not difficult to prove that the oscillation of u_k on $B_1 \setminus B_{\frac{1}{10}}$ is finite. Hence we can assume $|x_i| \leq \frac{1}{10}, i = 1, 2$. Let

$$G(x, \eta) = -\frac{1}{2\pi} \log |x - \eta| + H(x, \eta)$$

be the Green's function on B_1 with respect to Dirichlet boundary condition. Let $x_0 \in B_1 \setminus \Sigma_k$ and $x_1, x_2 \in B(x_0, d(x_0, \Sigma_k)/2)$. Using the fact u_k has bounded oscillation on ∂B_1 , We have

$$u_k(x_1) - u_k(x_2) = \int_{B_1} \left(G(x_1, \eta) - G(x_2, \eta) \right) \left(h_1^k(\eta) e^{u_k(\eta)} - h_2^k(\eta) e^{-u_k(\eta)} \right) d\eta + O(1).$$

Since $|x_i| \leq \frac{1}{10}, i = 1, 2$ and $\Delta H = 0$ in B_1 , we can use the bound on the energy (1.8) to get

$$\int_{B_1} \left(H(x_1, \eta) - H(x_2, \eta) \right) \left(h_1^k(\eta) e^{u_k(\eta)} - h_2^k(\eta) e^{-u_k(\eta)} \right) d\eta = O(1).$$

It remains to prove

$$\int_{B_1} \log \frac{|x_1 - \eta|}{|x_2 - \eta|} \left(h_1^k(\eta) e^{u_k(\eta)} - h_2^k(\eta) e^{-u_k(\eta)} \right) d\eta = O(1).$$

Let r_k be the distance between x_0 and Σ_k . Then we decompose $B_1 = (B_1 \setminus B_{\frac{3}{4}r_k}(x_0)) \cup B_{\frac{3}{4}r_k}(x_0)$ and compute the integration for each region.

(i) $\eta \in B_1 \setminus B_{\frac{3}{4}r_k}(x_0)$. Then

$$\log \frac{|x_1 - \eta|}{|x_2 - \eta|} = O(1)$$

and the integration in this region is bounded.

(ii) $\eta \in B_{\frac{3}{4}r_k}(x_0)$. Let

$$v_1^k(y) = u_k(x_0 + r_k y) + 2 \log r_k, \quad v_2^k(y) = -u_k(x_0 + r_k y) + 2 \log r_k,$$

for $y \in B_{3/4}$. Let y_1, y_2 be the images of x_1, x_2 after scaling, namely $x_i = x_0 + r_k y_i, i = 1, 2$, we have to prove that

$$\int_{B_{3/4}} \log \frac{|y_1 - \eta|}{|y_2 - \eta|} \left(h_1^k(x_0 + r_k \eta) e^{v_1^k(\eta)} - h_2^k(x_0 + r_k \eta) e^{v_2^k(\eta)} \right) d\eta = O(1).$$

Without loss of generality we may assume that $e_1 = (1, 0)$ is the image after scaling of the blow-up point in Σ_k which is closest to x_0 . From Proposition 2.1 it holds

$$v_i^k(\eta) + 2 \log |\eta - e_1| \leq C.$$

Therefore

$$e^{v_i^k(\eta)} \leq C |\eta - e_1|^{-2}.$$

Moreover, we notice that $|\eta - e_1| \geq C > 0$ for $\eta \in B_{\frac{3}{4}}$. Then for $i, j = 1, 2$, we get

$$\int_{B_{\frac{3}{4}}} \log |y_j - \eta| h_i^k(x_0 + r_k \eta) e^{v_i^k(\eta)} d\eta \leq C \int_{B_{\frac{3}{4}}} \frac{\log |y_j - \eta|}{|\eta - e_1|^2} d\eta \leq C$$

and we finish the proof. \square

3. POHOZAEV IDENTITY AND RELATED ESTIMATES ON THE ENERGY

In this section, we establish a Pohozaev-type identity for (1.1). This identity will be a crucial tool in proving the quantization result of Theorem 1.1.

We start with some observations and terminology. By Lemma 2.1 one can see that the behavior of blowup solutions away from the bubbling area can be described just by its spherical average in a neighborhood of a point in Σ_k . Moreover, the behavior of the solution on a boundary of a ball, say $\partial B_r(x_0)$, will play a crucial role in the forthcoming arguments, see the Remark 3.1. Throughout the paper we will say u_k has fast decay on $\partial B_r(x_0)$ if

$$u_k(x) \leq -2 \log |x - x_0| - N_k, \quad \text{for } x \in \partial B_r(x_0),$$

and some $N_k \rightarrow +\infty$. If instead there exists $C > 0$ independent of k such that

$$u_k(x) \geq -2 \log |x - x_0| - C, \quad \text{for } x \in \partial B_r(x_0),$$

we say u_k has slow decay on $\partial B_r(x_0)$. The same terminology will be used for $-u_k$.

For a sequence of bubbling solutions u_k of (1.5), we have the following result.

Proposition 3.1. *Let u_k satisfy (1.5), (1.6), (1.8) and h_i^k satisfy (1.7). Then we have*

$$4(\sigma_1 + \sigma_2) = (\sigma_1 - \sigma_2)^2.$$

Before we give a proof of Proposition 3.1, we first establish the following auxiliary lemma.

Lemma 3.1. *For all $\varepsilon_k \rightarrow 0$ such that $\Sigma_k \subset B_{\varepsilon_k/2}(0)$, there exists $l_k \rightarrow 0$ such that $l_k \geq 2\varepsilon_k$ and*

$$|\bar{u}_k(l_k)| + 2 \log l_k \rightarrow -\infty, \quad (3.1)$$

where $\bar{u}_k(r) := \frac{1}{2\pi r} \int_{\partial B_r} u_k$.

Proof. Given $\varepsilon_{k,1} \geq \varepsilon_k$ such that $\varepsilon_{k,1} \rightarrow 0$, we claim there exist $r_{k,1}, r_{k,2} \geq \varepsilon_{k,1}$ with the following property:

$$\begin{aligned} u_k(x) + 2 \log r_{k,1} &\rightarrow -\infty, & \forall x \in \partial B_{r_{k,1}}, \\ -u_k(x) + 2 \log r_{k,2} &\rightarrow -\infty, & \forall x \in \partial B_{r_{k,2}}. \end{aligned} \quad (3.2)$$

We prove the claim by contradiction. Suppose it is not true, say for u_k we can find $\varepsilon_{k,1} \rightarrow 0$ with $\varepsilon_{k,1} \geq \varepsilon_k$ such that for all $r \geq \varepsilon_{k,1}$,

$$\sup_{x \in \partial B_r} (u_k(x) + 2 \log |x|) \geq -C,$$

for some $C > 0$. Since $u_k(x)$ has bounded oscillation on each ∂B_r by Lemma 2.1, we get

$$u_k(x) + 2 \log |x| \geq -C$$

for some C and all $x \in \partial B_r$, $r \geq \varepsilon_{k,1}$. Equivalently, we have

$$e^{u_k(x)} \geq C|x|^{-2}, \quad \varepsilon_{k,1} \leq |x| \leq 1.$$

Integrating e^{u_k} on $B_1 \setminus B_{\varepsilon_{k,1}}$ we get $\int_{B_1} h_1 e^{u_k}$ is unbounded and it contradicts to (1.8). Thus (3.2) holds.

Next we choose $\tilde{r}_k \geq \varepsilon_k$ such that

$$\bar{u}_k(\tilde{r}_k) + 2 \log \tilde{r}_k \rightarrow -\infty.$$

Suppose \tilde{r}_k is not tending to 0, then we claim there exists $\hat{r}_k \rightarrow 0$ such that

$$\bar{u}_k(r) + 2 \log r \rightarrow -\infty, \quad \text{for } \hat{r}_k \leq r \leq \tilde{r}_k. \quad (3.3)$$

To prove this we observe that

$$u_k(x) + 2 \log |x| \leq -N_k, \quad |x| = \tilde{r}_k$$

for some $N_k \rightarrow \infty$. Then, for any fixed C , by Lemma 2.1 we obtain

$$u_k(x) + 2 \log |x| \leq -N_k + C_0, \quad \tilde{r}_k/C < |x| < \tilde{r}_k.$$

Based on this fact, it is not difficult to prove that \hat{r}_k can be found such that $\frac{\hat{r}_k}{\tilde{r}_k} \rightarrow 0$ and (3.3) holds. Thus we prove the claim (3.3).

Suppose now $\tilde{r}_k \rightarrow 0$. Similarly as before we can exploit Lemma 2.1 to find $s_k > \tilde{r}_k$ with $s_k \rightarrow 0$ and $\frac{s_k}{\tilde{r}_k} \rightarrow \infty$ such that

$$\bar{u}_k(r) + 2 \log r \rightarrow -\infty, \quad \text{for } \tilde{r}_k \leq r \leq s_k.$$

In both cases we can find r_k with $r_k \in [\hat{r}_k, \tilde{r}_k]$ in the first case, or $r_k \in [\tilde{r}_k, s_k]$ in the second case, such that

$$-\bar{u}_k(r_k) + 2 \log r_k \rightarrow -\infty.$$

Otherwise we would have

$$-\bar{u}_k(r) + 2 \log r \geq -C, \quad \text{for } \hat{r}_k \leq r \leq \tilde{r}_k \quad \text{or} \quad \tilde{r}_k \leq r \leq s_k.$$

Following a similar argument as before, we would get $\int_{B_1} h_2 e^{-u_k}$ is unbounded and contradiction arises. Therefore, we prove the lemma. \square

Proof of Proposition 3.1. We start by observing that there exists $l_k \rightarrow 0$ such that $\Sigma_k \subset B_{l_k/2}(0)$, (3.1) holds and

$$\frac{1}{2\pi} \int_{B_{l_k}} h_1^k e^{u_k} = \sigma_1 + o(1), \quad \frac{1}{2\pi} \int_{B_{l_k}} h_2^k e^{-u_k} = \sigma_2 + o(1). \quad (3.4)$$

In fact, one can first choose l_k so that the property (3.4) is satisfied and then by Lemma 3.1 we can further assume that (3.1) holds true. Let

$$v_1^k(y) = u_k(l_k y) + 2 \log l_k, \quad v_2^k(y) = -u_k(l_k y) + 2 \log l_k.$$

Then v_1^k, v_2^k satisfy

$$\begin{cases} \Delta v_1^k(y) + H_1^k(y) e^{v_1^k} - H_2^k(y) e^{v_2^k} = 0, & |y| \leq 1/l_k, \\ \bar{v}_i^k(1) \rightarrow -\infty, & i = 1, 2, \end{cases} \quad (3.5)$$

where

$$H_i^k(y) = h_i^k(l_k y), \quad i = 1, 2.$$

A modification of the Pohozaev-type identity gives us

$$\begin{aligned} & \sum_{i=1}^2 \int_{B_{\sqrt{R_k}}} (y \cdot \nabla H_i^k) e^{v_i^k} + 2 \sum_{i=1}^2 \int_{B_{\sqrt{R_k}}} H_i^k e^{v_i^k} \\ &= \sqrt{R_k} \int_{\partial B_{\sqrt{R_k}}} \sum_{i=1}^2 H_i^k e^{v_i^k} + \sqrt{R_k} \int_{\partial B_{\sqrt{R_k}}} \left(|\partial_\nu v_1^k|^2 - \frac{1}{2} |\nabla v_1^k|^2 \right), \end{aligned} \quad (3.6)$$

where we used $\nabla v_1^k = -\nabla v_2^k$ and $R_k \rightarrow \infty$ will be chosen later. We rewrite the above formula as

$$\mathcal{L}_1 + \mathcal{L}_2 = \mathcal{R}_1 + \mathcal{R}_2 + \mathcal{R}_3,$$

where the notation is easily understood. First we choose $R_k \rightarrow \infty$ sufficiently smaller than l_k^{-1} such that $\mathcal{L}_1 = o(1)$ by $l_k \rightarrow 0$. Now we consider \mathcal{L}_2 . By Lemma 2.1, $v_i^k(y) \rightarrow -\infty$ over all compact subsets of $\mathbb{R}^2 \setminus B_{1/2}$. Thus we can choose R_k so that

$$\int_{B_{R_k} \setminus B_1} H_i^k e^{v_i^k} = o(1). \quad (3.7)$$

Moreover, by the choice of l_k we have

$$\begin{aligned} \frac{1}{2\pi} \int_{B_1} H_1^k e^{v_1^k} &= \frac{1}{2\pi} \int_{B_{l_k}} h_1^k e^{u_k} = \sigma_1 + o(1), \\ \frac{1}{2\pi} \int_{B_1} H_2^k e^{v_2^k} &= \frac{1}{2\pi} \int_{B_{l_k}} h_2^k e^{-u_k} = \sigma_2 + o(1), \end{aligned} \quad (3.8)$$

which together with (3.7) implies

$$\mathcal{L}_2 = 4\pi \sum_{i=1}^2 \sigma_i + o(1).$$

To estimate \mathcal{R}_1 we notice that by (3.5) and Lemma 2.1

$$v_i^k(y) + 2 \log |y| \rightarrow -\infty, \quad \text{uniformly in } 1 < |y| \leq \sqrt{R_k}. \quad (3.9)$$

As a result, we have $\mathcal{R}_1 = o(1)$.

Next, we shall estimate the terms \mathcal{R}_2 and \mathcal{R}_3 . To do this we have to estimate ∇v_i^k on $\partial B_{\sqrt{R_k}}$. Let

$$G_k(y, \eta) = -\frac{1}{2\pi} \log |y - \eta| + H_k(y, \eta)$$

be the Green's function on $B_{l_k^{-1}}$ with respect to the Dirichlet boundary condition. The regular part is expressed as

$$H_k(y, \eta) = \frac{1}{2\pi} \log \frac{|y|}{l_k^{-1}} \left| \frac{l_k^{-2} y}{|y|^2} - \eta \right|$$

and it holds

$$\nabla_y H_k(y, \eta) = O(l_k), \quad \text{for } y \in \partial B_{\sqrt{R_k}}, \eta \in B_{l_k^{-1}}. \quad (3.10)$$

We start by estimating ∇v_1^k on $\partial B_{\sqrt{R_k}}$. By the Green's representation formula,

$$v_1^k(y) = \int_{B_{l_k^{-1}}} G(y, \eta) \left(H_1^k e^{v_1^k} - H_2^k e^{v_2^k} \right) + F_k,$$

where F_k is a harmonic function satisfying $F_k = v_i^k$ on $\partial B_{l_k^{-1}}$. In particular F_k has bounded oscillation on $\partial B_{l_k^{-1}}$. It follows that $F_k - C_k = O(1)$ for some C_k and

$|\nabla F_k(y)| = O(l_k)$, then we get

$$\begin{aligned} \nabla v_1^k(y) &= \int_{B_{l_k^{-1}}} \nabla_y G(y, \eta) \left(H_1^k e^{v_1^k} - H_2^k e^{v_2^k} \right) d\eta + \nabla F_k(y) \\ &= -\frac{1}{2\pi} \int_{B_{l_k^{-1}}} \frac{y - \eta}{|y - \eta|^2} \left(H_1^k e^{v_1^k} - H_2^k e^{v_2^k} \right) d\eta + O(l_k). \end{aligned} \quad (3.11)$$

In order to estimate the integral of (3.11) we divide the domain into different regions. We first observe that

$$\frac{1}{|y - \eta|} \sim \frac{1}{|\eta|} \leq o\left(R_k^{-\frac{1}{2}}\right), \quad \text{for } y \in \partial B_{\sqrt{R_k}}, \quad \eta \in B_{l_k^{-1}} \setminus B_{R_k^{2/3}}.$$

Hence, using the bound of the energy (1.8), the integral over $B_{l_k^{-1}} \setminus B_{R_k^{2/3}}$ is $o(1)R_k^{-\frac{1}{2}}$.

Consider now the integral over B_1 : we have

$$\frac{y - \eta}{|y - \eta|^2} = \frac{y}{|y|^2} + O\left(\frac{1}{|y|^2}\right), \quad \text{for } y \in \partial B_{\sqrt{R_k}}, \quad \eta \in B_1,$$

which together with (3.8) yields

$$-\frac{1}{2\pi} \int_{B_1} \frac{y - \eta}{|y - \eta|^2} \left(H_1^k e^{v_1^k} - H_2^k e^{v_2^k} \right) = \left(-\frac{y}{|y|^2} + O\left(\frac{1}{|y|^2}\right) \right) (\sigma_1 - \sigma_2 + o(1)).$$

We will see this term is the leading term

For the integral over the region $B_{\sqrt{R_k}/2} \setminus B_1$ we observe that

$$\frac{1}{|y - \eta|} \sim \frac{1}{|y|}, \quad \text{for } y \in \partial B_{\sqrt{R_k}}, \quad \eta \in B_{\sqrt{R_k}/2} \setminus B_1.$$

By the above estimate and (3.7) we obtain

$$\int_{B_{\sqrt{R_k}/2} \setminus B_1} \frac{y - \eta}{|y - \eta|^2} \left(H_1^k e^{v_1^k} - H_2^k e^{v_2^k} \right) = o(1)|y|^{-1}.$$

Similarly it holds that

$$\int_{B_{R_k^{2/3}} \setminus (B_1 \cup B_{\frac{|y|}{2}}(y))} \frac{y - \eta}{|y - \eta|^2} \left(H_1^k e^{v_1^k} - H_2^k e^{v_2^k} \right) = o(1)|y|^{-1}.$$

Moreover, for the integral in $B_{\frac{|y|}{2}}(y)$ we use $e^{v_i^k(\eta)} = o(1)|\eta|^{-2}$ to get

$$\int_{B_{\frac{|y|}{2}}(y)} \frac{y - \eta}{|y - \eta|^2} \left(H_1^k e^{v_1^k} - H_2^k e^{v_2^k} \right) = o(1)|y|^{-1}.$$

Finally, combing all the estimates we deduce

$$\nabla v_1^k(y) = \left(-\frac{y}{|y|^2} \right) (\sigma_1 - \sigma_2 + o(1)) + o(|y|^{-1}), \quad \text{for } y \in \partial B_{\sqrt{R_k}}.$$

Exploiting the latter formula in \mathcal{R}_2 and \mathcal{R}_3 we get

$$\mathcal{R}_2 + \mathcal{R}_3 = \pi(\sigma_1 - \sigma_2)^2 + o(1).$$

Therefore, we end up with

$$4(\sigma_1 + \sigma_2) = (\sigma_1 - \sigma_2)^2 + o(1).$$

Hence, Proposition 3.1 is proved. \square

Remark 3.1. *By the proof of Proposition 3.1 one observes the following fact: the fast decay property is crucial in evaluating the Pohozaev identity. Moreover, let $\Sigma'_k \subseteq \Sigma_k$ and we assume that*

$$\text{dist}(\Sigma'_k, \partial B_{l_k}(p_k)) = o(1) \text{dist}(\Sigma_k \setminus \Sigma'_k, \partial B_{l_k}(p_k)).$$

If both components $u_k, -u_k$ have fast decay on $\partial B_{l_k}(p_k)$, namely

$$|u_k(x)| \leq -2 \log |x - p_k| - N_k, \quad \text{for } x \in \partial B_{l_k}(p_k),$$

and some $N_k \rightarrow +\infty$. Then, we can evaluate a local Pohozaev identity and get

$$(\tilde{\sigma}_1^k(l_k) - \tilde{\sigma}_2^k(l_k))^2 = 4(\tilde{\sigma}_1^k(l_k) + \tilde{\sigma}_2^k(l_k)) + o(1),$$

where

$$\tilde{\sigma}_1^k(l_k) = \frac{1}{2\pi} \int_{B_{l_k}(p_k)} h_1^k e^{u_k}, \quad \tilde{\sigma}_2^k(l_k) = \frac{1}{2\pi} \int_{B_{l_k}(p_k)} h_2^k e^{-u_k}.$$

We note that if $B_{l_k}(p_k) \cap \Sigma_k = \emptyset$, then $\tilde{\sigma}_i^k(l_k) = o(1)$, $i = 1, 2$ and the above formula clearly holds.

This fact will be frequently used in the later arguments.

4. ASYMPTOTIC BEHAVIOR OF SOLUTIONS AROUND EACH BLOW-UP POINT

The goal of this section is to get some energy classification in each blow-up area. We will see in the sequel how the fast decay property of the solutions plays a crucial role in determining the local energy.

By considering suitable translated functions we may assume without loss of generality that $0 \in \Sigma_k$ for any k . Let $\tau_k = \frac{1}{2} \text{dist}(0, \Sigma_k \setminus \{0\})$, we consider the energy limits of $h_1^k e^{u_k}$ and $h_2^k e^{-u_k}$ in B_{τ_k} . Let us define

$$\begin{aligned} v_1^k &= u_k(\delta_k y) + 2 \log \delta_k, & |y| &\leq \tau_k / \delta_k, \\ v_2^k &= -u_k(\delta_k y) + 2 \log \delta_k, \end{aligned} \quad (4.1)$$

where $-2 \log \delta_k = \max_{x \in B(0, \tau_k)} |u_k|$. Thus the equation for v_1^k is

$$\Delta v_1^k(y) + h_1^k(\delta_k y) e^{v_1^k(y)} - h_2^k(\delta_k y) e^{v_2^k(y)} = 0, \quad |y| \leq \tau_k / \delta_k.$$

By the definition of the selection process we have $\tau_k / \delta_k \rightarrow \infty$, see Proposition 2.1.

We note that

$$\begin{aligned} \int_{B_{\tau_k}(0)} h_1^k(x) e^{u_k(x)} dx &= \int_{B_{\tau_k/\delta_k}(0)} h_1^k(\delta_k y) e^{v_1^k(y)} dy, \\ \int_{B_{\tau_k}(0)} h_2^k(x) e^{-u_k(x)} dx &= \int_{B_{\tau_k/\delta_k}(0)} h_2^k(\delta_k y) e^{v_2^k(y)} dy. \end{aligned}$$

As a consequence, we have

$$\begin{aligned} \int_{B_{\tau_k}(0)} h_1^k(x) e^{u_k(x)} dx &= O(1) e^{\bar{v}_1^k(\partial B_{\tau_k/\delta_k}(0))}, \\ \int_{B_{\tau_k}(0)} h_2^k(x) e^{-u_k(x)} dx &= O(1) e^{\bar{v}_2^k(\partial B_{\tau_k/\delta_k}(0))}. \end{aligned} \quad (4.2)$$

Define the following local masses:

$$\sigma_1^k(r) = \frac{1}{2\pi} \int_{B_r} h_1^k e^{u_k}, \quad \sigma_2^k(r) = \frac{1}{2\pi} \int_{B_r} h_2^k e^{-u_k}. \quad (4.3)$$

The main result of this section is the following.

Proposition 4.1. *Suppose (1.5)-(1.8) hold for u_k, h_i^k . Let σ_i^k be defined in (4.3). For any $s_k \in (0, \tau_k)$ such that both $u_k, -u_k$ have fast decay on ∂B_{s_k} , i.e.*

$$|u_k(x)| \leq -2 \log |x| - N_k, \quad \text{for } |x| = s_k \text{ and some } N_k \rightarrow \infty, \quad (4.4)$$

we have that $(\sigma_1^k(s_k), \sigma_2^k(s_k))$ is a small perturbation of one of the following two types:

$$(2m(m+1), 2m(m-1)) \quad \text{or} \quad (2m(m-1), 2m(m+1)),$$

for some $m \in \mathbb{N}$. In particular, they are both a perturbation of multiple of 4.

On ∂B_{τ_k} , either both $u_k, -u_k$ have fast decay as in (4.4) and the conclusion is as before, or one component has fast decay while the other one is not fast decaying. Suppose for example $-u_k$ is not the fast decaying, i.e.,

$$-u_k(x) + 2 \log |x| \geq -C, \quad \text{for } |x| = \tau_k \text{ and some } C > 0,$$

while for u_k it holds that

$$u_k(x) \leq -2 \log |x| - N_k, \quad \text{for } |x| = s_k \text{ and some } N_k \rightarrow \infty.$$

Then $\sigma_1^k(\tau_k)$ is a perturbation of multiple of 4.

In particular, in any case at least one of the two components $u_k, -u_k$ has the local mass in B_{τ_k} equals a perturbation of multiple of 4.

Proof. Let v_i^k be defined in (4.1). Observe that by construction one of the v_i^k 's converges while the other one goes to minus infinity over all compact subsets of \mathbb{R}^2 (see the argument in Proposition 2.1), namely we have just a partially blown-up situation. Without loss of generality we assume that v_1^k converges to v_1 in $C_{loc}^2(\mathbb{R}^2)$ and v_2^k tends to minus infinity over any compact subset of \mathbb{R}^2 . The equation for v_1 is

$$\Delta v_1 + e^{v_1} = 0 \text{ in } \mathbb{R}^2, \quad \int_{\mathbb{R}^2} e^{v_1} < \infty,$$

where we used $\lim_{k \rightarrow \infty} h_1^k(0) = 1$. By the classification result of Chen-Li [10], we have

$$\int_{\mathbb{R}^2} e^{v_1} = 8\pi \quad \text{and} \quad v_1(y) = -4 \log |y| + O(1), \quad |y| > 1.$$

Therefore, we can take $R_k \rightarrow \infty$ (we assume $R_k = o(1)\tau_k/\delta_k$) such that

$$\frac{1}{2\pi} \int_{B_{R_k}} h_1^k(\delta_k y) e^{v_1^k} = 4 + o(1), \quad (4.5)$$

and

$$\frac{1}{2\pi} \int_{B_{R_k}} h_2^k(\delta_k y) e^{v_2^k} = o(1). \quad (4.6)$$

For $r \geq R_k$ we clearly have

$$\sigma_i^k(\delta_k r) = \frac{1}{2\pi} \int_{B_r} h_i^k(\delta_k y) e^{v_i^k}.$$

Up to now we get from (4.5) and (4.6) that

$$\sigma_1^k(\delta_k R_k) = 4 + o(1), \quad \sigma_2^k(\delta_k R_k) = o(1).$$

Let $\bar{v}_i^k(r)$ be the average of v_i^k on ∂B_r , $i = 1, 2$. It will be important to study $\frac{d}{dr} \bar{v}_i^k(r)$, $i = 1, 2$. In fact if

$$\frac{d}{dr} (\bar{v}_i^k(r) + 2 \log r) > 0, \quad \text{for some } i,$$

there is a possibility that for some larger radius s , v_i^k becomes a slow decay component on ∂B_s .

The key fact is to observe that

$$\begin{aligned} \frac{d}{dr} \bar{v}_1^k(r) &= \frac{-\sigma_1^k(\delta_k r) + \sigma_2^k(\delta_k r)}{r}, \\ \frac{d}{dr} \bar{v}_2^k(r) &= \frac{\sigma_1^k(\delta_k r) - \sigma_2^k(\delta_k r)}{r}, \end{aligned} \quad R_k \leq r \leq \tau_k / \delta_k. \quad (4.7)$$

Clearly we have

$$R_k \frac{d}{dr} \bar{v}_1^k(R_k) = -4 + o(1), \quad R_k \frac{d}{dr} \bar{v}_2^k(R_k) = 4 + o(1).$$

To continue the proof of Proposition 4.1 we need the following two lemmas.

Lemma 4.1. *Suppose there exists $L_k \in (R_k, \tau_k / \delta_k)$ such that*

$$v_i^k(y) \leq -2 \log |y| - N_k, \quad \text{for } R_k \leq |y| \leq L_k, \quad i = 1, 2 \quad (4.8)$$

and some $N_k \rightarrow \infty$. Then σ_i^k does not change much from $\delta_k R_k$ to $\delta_k L_k$, i.e.,

$$\sigma_i^k(\delta_k L_k) = \sigma_i^k(\delta_k R_k) + o(1), \quad i = 1, 2.$$

Proof. Suppose the statement is false: then there exists i such that $\sigma_i^k(\delta_k L_k) > \sigma_i^k(\delta_k R_k) + \delta$ for some $\delta > 0$. Let us choose $\tilde{L}_k \in (R_k, L_k)$ such that

$$\max_{i=1,2} \left(\sigma_i^k(\delta_k \tilde{L}_k) - \sigma_i^k(\delta_k R_k) \right) = \varepsilon, \quad (4.9)$$

where $\varepsilon > 0$ is taken sufficiently small. Then by using (4.7) we have

$$\frac{d}{dr} \bar{v}_1^k(r) \leq \frac{-4 + \varepsilon + o(1)}{r} \leq -\frac{2 + \varepsilon}{r}, \quad R_k \leq r \leq \tilde{L}_k. \quad (4.10)$$

By Lemma 2.1 we have that

$$v_i^k(x) = \bar{v}_i^k(|x|) + O(1), \quad x \in B_{\tau_k / \delta_k},$$

where $\bar{v}_i^k(|x|)$ is the average of v_i on $\partial B_{|x|}$. Using the same reason as above and (4.8)-(4.10), we can show that

$$\int_{B_{\tilde{L}_k} \setminus B_{R_k}} e^{v_1^k} = O(1) \int_{B_{\tilde{L}_k} \setminus B_{R_k}} e^{\bar{v}_1^k(\tilde{L}_k)} = o(1),$$

and it implies $\sigma_1^k(\delta_k \tilde{L}_k) = \sigma_1^k(\delta_k R_k) + o(1)$.

It follows that the maximum in (4.9) is attained for $i = 2$, i.e.

$$\sigma_2^k(\delta_k \tilde{L}_k) = \sigma_2^k(\delta_k R_k) + \varepsilon. \quad (4.11)$$

On the other hand, since (4.8) holds, as observed in Remark 3.1 we get

$$(\sigma_1^k(\delta_k \tilde{L}_k) - \sigma_2^k(\delta_k \tilde{L}_k))^2 = 4(\sigma_1^k(\delta_k \tilde{L}_k) + \sigma_2^k(\delta_k \tilde{L}_k)) + o(1).$$

Using

$$\sigma_1^k(\delta_k \tilde{L}_k) = \sigma_1^k(\delta_k R_k) + o(1) = 4 + o(1),$$

we deduce that

$$\sigma_2^k(\delta_k \tilde{L}_k) = o(1) \quad \text{or} \quad \sigma_2^k(\delta_k \tilde{L}_k) = 12 + o(1),$$

which contradicts to (4.9). Thus we prove the lemma. \square

From the argument in Lemma 4.1 we observe the following fact: for $r \geq R_k$ either both v_1, v_2 have fast decay up to $\partial B_{\tau_k/\delta_k}$, namely

$$v_i^k(y) \leq -2 \log |y| - N_k, \quad R_k \leq |y| \leq \tau_k/\delta_k, \quad i = 1, 2, \quad (4.12)$$

for some $N_k \rightarrow +\infty$, or there exists $L_k \in (R_k, \tau_k/\delta_k)$ such that v_2^k has slow decay, i.e.

$$v_2^k(y) \geq -2 \log L_k - C, \quad |y| = L_k, \quad (4.13)$$

for some $C > 0$, while

$$v_1^k(y) \leq -2 \log |y| - N_k, \quad R_k \leq |y| \leq L_k, \quad (4.14)$$

for some $N_k \rightarrow +\infty$. Indeed, we have noticed in Lemma 4.1 that if the local energy changes, σ_2^k has to change first. Moreover, we have seen that L_k can be chosen so that $\sigma_2^k(\delta_k L_k) - \sigma_2^k(\delta_k R_k) = \varepsilon$ for some $\varepsilon > 0$ small. The following lemma treats the latter situation.

Lemma 4.2. *Suppose there exists $L_k \geq R_k$ such that (4.13) and (4.14) hold. For L_k , we assume $L_k = o(1)\tau_k/\delta_k$. Then there exists \tilde{L}_k such that $\tilde{L}_k/L_k \rightarrow \infty$ and $\tilde{L}_k = o(1)\tau_k/\delta_k$ with the following property:*

$$v_i^k(y) \leq -2 \log |y| - N_k, \quad |y| = \tilde{L}_k, \quad i = 1, 2, \quad (4.15)$$

for some $N_k \rightarrow \infty$. Moreover

$$\sigma_1^k(\delta_k \tilde{L}_k) = 4 + o(1), \quad \sigma_2^k(\delta_k \tilde{L}_k) = 12 + o(1). \quad (4.16)$$

Proof. First we observe that from the choice of L_k and the fact $\sigma_2^k(\delta_k R_k) = o(1)$ we can assume $\sigma_2^k(\delta_k L_k) \leq \varepsilon$ for some $\varepsilon > 0$ small, then we get

$$\frac{d}{dr} \bar{v}_1^k(r) \leq \frac{-4 + \varepsilon + o(1)}{r}, \quad R_k \leq r \leq L_k.$$

Now we claim there exists $N > 1$ such that

$$\sigma_2^k(\delta_k(NL_k)) \geq 6 + o(1). \quad (4.17)$$

Suppose this does not hold. Then there exist $\varepsilon_0 > 0$ and $\tilde{R}_k \rightarrow \infty$ such that

$$\sigma_2^k(\delta_k \tilde{R}_k L_k) \leq 6 - \varepsilon_0. \quad (4.18)$$

Moreover, \tilde{R}_k can be chosen to tend to infinity slowly so that by Lemma 2.1 and (4.14) we get

$$v_1^k(y) \leq -2 \log |y| - N_k, \quad L_k \leq |y| \leq \tilde{R}_k L_k. \quad (4.19)$$

which together with Lemma 4.1 implies that $\sigma_1^k(\delta_k L_k) = \sigma_1^k(\delta_k \tilde{R}_k L_k) + o(1)$. Thus by (4.18)

$$\frac{d}{dr} \bar{v}_2^k(r) \geq \frac{-2 + \varepsilon + o(1)}{r}. \quad (4.20)$$

From (4.20) and (4.13) it is not difficult to show

$$\int_{B_{L_k \tilde{R}_k} \setminus B_{L_k}} e^{v_2^k} \rightarrow \infty,$$

and it contradicts to (1.8). Therefore (4.17) holds.

By Lemma 2.1 we have

$$v_i^k(y) + 2 \log NL_k = \bar{v}_i^k(NL_k) + 2 \log(NL_k) + O(1), \quad i = 1, 2, \quad |y| = NL_k.$$

Therefore, we get

$$\begin{aligned} \bar{v}_1^k(NL_k) &\leq -2 \log(NL_k) - N_k, \\ \bar{v}_2^k(NL_k) &\geq -2 \log(NL_k) - C \geq -2 \log(L_k) - C. \end{aligned}$$

Furthermore we can assert that

$$\bar{v}_2^k((N+1)L_k) \geq -2 \log L_k - C,$$

which, jointly with (4.17), yields

$$\int_{B_{(N+1)L_k}} h_2^k(\delta_k y) e^{v_2^k(y)} dy \geq 6 + \varepsilon_0,$$

for some $\varepsilon_0 > 0$. By the above estimate we get

$$\frac{d}{dr} \bar{v}_2^k(r) \leq -\frac{2 + \varepsilon_0}{r}, \quad \text{for } r = (N+1)L_k.$$

Then we can take $\tilde{R}_k \rightarrow \infty$ slowly such that $\tilde{R}_k L_k = o(1)\tau_k/\delta_k$ and

$$\begin{aligned} v_2^k(y) &\leq (-2 - \varepsilon_0) \log |y| - N_k, \quad |y| = \tilde{R}_k L_k, \\ v_1^k(y) &\leq -2 \log |y| - N_k, \quad L_k \leq |y| \leq \tilde{R}_k L_k. \end{aligned}$$

Next, using Lemma 4.1 and (4.5) we get

$$\sigma_1^k(\delta_k \tilde{R}_k L_k) = \sigma_1^k(\delta_k L_k) + o(1) = 4 + o(1).$$

On the other hand, since both components have fast decay on $\partial B_{\tilde{R}_k L_k}$. As explained in Remark 3.1 we can compute the local Pohozaev identity. Combined with (4.17) we get

$$\sigma_2^k(\delta_k \tilde{R}_k L_k) = 12 + o(1).$$

Let $\tilde{L}_k = \tilde{R}_k L_k$ we conclude the proof. \square

Returning to the proof of Proposition 4.1, we are left with the region $\tilde{L}_k \leq |y| \leq \tau_k/\delta_k$. We divide the discussion into two cases.

Suppose first $L_k = O(1)\tau_k/\delta_k$. Then by Lemma 2.1 we directly conclude that one component has fast decay while the other one has slow decay, see for example (4.13) and (4.14). Moreover, we have observed that the local mass in B_{τ_k} of the fast decaying component is a small perturbation of multiple of 4. This is exactly the second alternative of Proposition 4.1 and therefore the proof is concluded.

Suppose now $L_k = o(1)\tau_k/\delta_k$. In this case \tilde{L}_k can be chosen such that $\tilde{L}_k = o(1)\tau_k/\delta_k$. By using the local masses given by Lemma 4.2 we have

$$\frac{d}{dr}\bar{v}_1^k(r) = \frac{8 + o(1)}{r}, \quad \frac{d}{dr}\bar{v}_2^k(r) = -\frac{8 + o(1)}{r}, \quad \text{for } r = \tilde{L}_k.$$

It follows that

$$\frac{d}{dr}\bar{v}_2^k(r) \leq -\frac{2 + \varepsilon}{r}, \quad r = \tilde{L}_k,$$

for some $\varepsilon > 0$. As in Lemma 4.1 we conclude that $\sigma_2^k(r)$ does not change for $r \geq \tilde{L}_k$ unless σ_1^k changes. By the same ideas of Lemmas 4.1, Lemma 4.2 and the argument of the first case $L_k = O(1)\tau_k/\delta_k$, either v_1^k has slow decay up to B_{τ_k/δ_k} or there is $\hat{L}_k = o(1)\tau_k/\delta_k$ such that

$$\sigma_1^k(\delta\hat{L}_k) = 24 + o(1), \quad \sigma_2^k(\delta\hat{L}_k) = 12 + o(1).$$

By the latter local energies we deduce

$$\frac{d}{dr}\bar{v}_1^k(r) = -\frac{12 + o(1)}{r}, \quad \frac{d}{dr}\bar{v}_2^k(r) = \frac{12 + o(1)}{r}, \quad \text{for } r = \hat{L}_k.$$

As before

$$\frac{d}{dr}\bar{v}_1^k(r) \leq -\frac{2 + \varepsilon}{r}, \quad r = \hat{L}_k,$$

for some $\varepsilon > 0$. Now $\sigma_1^k(r)$ does not change for $r \geq \hat{L}_k$ unless σ_2^k changes. Repeating the argument we get either v_2^k has slow decay up to B_{τ_k/δ_k} or there is $\bar{L}_k = o(1)\tau_k/\delta_k$ such that

$$\sigma_1^k(\delta\bar{L}_k) = 24 + o(1), \quad \sigma_2^k(\delta\bar{L}_k) = 40 + o(1).$$

Since after each step one of the local masses changes by a positive number, using the uniform bound on the energy (1.8) the process stops after finite steps. Eventually we get Proposition 4.1. \square

5. COMBINATION OF BUBBLING AREAS AND PROOF OF THEOREM 1.1

In this section we present the argument for combining the blow-up areas. Some parts of this strategy have been already used in several frameworks, see [27]. However, we have to complete the argument with some new ideas developed in [28]. The procedure is the following: we start by considering blow-up points which are close to each other and get a quantization property for each group, see the definition of group below. In particular, in each group the local energy of at least one component is a small perturbation of $4n$, for some $n \in \mathbb{N}$. Similarly, we combine the groups and obtain that the total energy of at least one component is a small perturbation of $4n$, for some $n \in \mathbb{N}$. Then, the conclusion follows by applying a quantization result which we state below and the Pohozaev identity.

As we pointed out, we will exploit the following energy quantization result of global Liouville equations with singularities.

Theorem B. [28] *Let u be a solution of*

$$\begin{cases} \Delta u + 2e^u = \sum_{j=1}^N 4\pi n_j \delta_{p_j} & \text{in } \mathbb{R}^2, \\ \int_{\mathbb{R}^2} e^u < +\infty, \end{cases}$$

where p_j , $j = 1, \dots, N$ are distinct points in \mathbb{R}^2 and $n_j \in \mathbb{N}$, $j = 1, \dots, N$. Then

$$\frac{1}{2\pi} \int_{\mathbb{R}^2} e^u = 2n,$$

for some $n \in \mathbb{N}$.

Before we give the proof of Theorem 1.1, we introduce the following definition which plays a crucial role in our argument.

Definition. Let $G = \{p_1^k, \dots, p_q^k\}$ be a subset of Σ_k with more than one point in it. G is called a group if

- (1) $\text{dist}(p_i^k, p_j^k) \sim \text{dist}(p_s^k, p_t^k)$,
where $p_i^k, p_j^k, p_s^k, p_t^k$ are any points in G such that $p_i^k \neq p_j^k$ and $p_s^k \neq p_t^k$.
- (2) $\frac{\text{dist}(p_i^k, p_j^k)}{\text{dist}(p_i^k, p_k)} \rightarrow 0$,
for any $p_k \in \Sigma_k \setminus G$ and $p_i^k, p_j^k \in G$ with $p_i^k \neq p_j^k$.

Proof of Theorem 1.1: As in Section 4, by considering suitable translation we may assume without loss of generality that $0 \in \Sigma_k$ for any k . Let $2\tau_k$ be the distance between 0 and $\Sigma_k \setminus \{0\}$. To describe the group G_0 that contains 0 we proceed in the following way: if for any $z_k \in \Sigma_k \cap \partial B(0, 2\tau_k)$ we have $\text{dist}(z_k, \Sigma_k \setminus \{z_k\}) \sim \tau_k$, then G_0 contains at least two points. On the other hand, if there exists $z'_k \in \partial B(0, 2\tau_k) \cap \Sigma_k$ such that $\tau_k / \text{dist}(z'_k, \Sigma_k \setminus \{z'_k\}) \rightarrow \infty$ we let G_0 be 0 itself. By the definition of group, all points of G_0 are in $B(0, N\tau_k)$ for some N independent of k . Let

$$\begin{aligned} \tilde{v}_1^k(y) &= u_k(\tau_k y) + 2 \log \tau_k, \\ \tilde{v}_2^k(y) &= -u_k(\tau_k y) + 2 \log \tau_k, \end{aligned} \quad |y| \leq \tau_k^{-1},$$

then \tilde{v}_1^k satisfies

$$\Delta \tilde{v}_1^k(y) + \tilde{h}_1^k(y) e^{\tilde{v}_1^k(y)} - \tilde{h}_2^k(y) e^{\tilde{v}_2^k(y)} = 0, \quad |y| \leq \tau_k^{-1}, \quad (5.1)$$

where $\tilde{h}_i^k(y) = h_i^k(\tau_k y)$, $i = 1, 2$.

Let $0, q_1^k, \dots, q_s^k$ be the images for the members of G_0 after scaling from y to $\tau_k y$. We observe that $q_i^k \in B_N$. By Proposition 4.1 at least one of \tilde{v}_i^k decays fast on ∂B_1 . At first, we study the case that only one component of \tilde{v}_i^k has fast decay. Without loss of generality we assume \tilde{v}_1^k has fast decay:

$$\tilde{v}_1^k \leq -2 \log |y| - N_k, \quad \text{on } \partial B_1,$$

for some $N_k \rightarrow \infty$. Combined with Lemma 2.1 we have \tilde{v}_1^k also has fast decay on $\partial B_1(q_j^k)$, $j = 1, \dots, s$. Using Proposition 4.1, we get

$$\sigma_1^k(\tau_k) = 4\tilde{n} + o(1),$$

for some $\tilde{n} \in \mathbb{N}$ and

$$\sigma_1^k(B_1(q_j^k)) = \frac{1}{2\pi} \int_{B_1(q_j^k)} \tilde{h}_1(y) e^{\tilde{v}_1^k} = 4n_j + o(1), \quad j = 1, \dots, s,$$

for some $n_j \in \mathbb{N}, j = 1, \dots, s$.

By Lemma 2.1, Lemma 4.1 and using the same argument of Lemma 3.1 we can choose L large enough such that $B_L(0)$ contains all the bubbling disks of the group G_0 , the energy of \tilde{v}_1^k in $B_L(0)$ does not change so much and \tilde{v}_1^k still has fast decay on $\partial B_L(0)$: in particular

$$\sigma_1^k(\tau_k L) = 4n + o(1), \quad (5.2)$$

for some $n \in \mathbb{N}$. Now, since \tilde{v}_1^k has fast decay and \tilde{v}_2^k has slow decay, it is not difficult to see that $\tilde{h}_1 e^{\tilde{v}_1^k} \rightarrow \sum_{j=0}^s 8\pi n_{1,j} \delta_{q_j}$, where $n_{1,j} \in \mathbb{N}$ and $q_0 = 0$, while $\tilde{h}_2 e^{\tilde{v}_2^k} \rightarrow \sum_{j=0}^s 8\pi n_{2,j} \delta_{q_j} + F$, where $n_{2,j} \in \mathbb{N}$ and $F \in L^1(\mathbb{R}^2)$. Then we want to prove that the integral of F in \mathbb{R}^2 is multiple of 8π . From the argument in Proposition 4.1, we have $n_{1,j} > n_{2,j}$ for any j . Using equation (5.1), we get that the limit function of \tilde{v}_2^k in $\mathbb{R}^2 \setminus \{q_1, \dots, q_s\}$, which we denote by \tilde{v}_2 , satisfies

$$\Delta \tilde{v}_2 + e^{\tilde{v}_2} = \sum_{j=0}^s 8\pi \tilde{n}_j \delta_{q_j} \quad \text{in } \mathbb{R}^2,$$

where $\tilde{n}_j \in \mathbb{N}, j = 0, \dots, s$. Using a transformation $\tilde{v}_2 = \tilde{w}_2 + \log 2$ and exploiting Theorem B we can conclude

$$\frac{1}{2\pi} \int_{\mathbb{R}^2} e^{\tilde{v}_2} = 4\tilde{n} + o(1),$$

for some $\tilde{n} \in \mathbb{N}$, and it gives the quantization of the function F .

Let $2\tau_k L_k$ be the distance from 0 to the nearest group from G_0 . By the definition of group we have $L_k \rightarrow \infty$. As before we can find $\tilde{L}_k \leq L_k, \tilde{L}_k \rightarrow \infty$ slowly such that

$$\sigma_1^k(\tau_k \tilde{L}_k) = 4n + o(1), \quad \sigma_2^k(\tau_k \tilde{L}_k) = 4\bar{n} + o(1), \quad \text{for some } \bar{n} \in \mathbb{N}$$

and

$$\tilde{v}_i^k(y) \leq -2 \log \tilde{L}_k - N_k, \quad \text{for } |y| = \tilde{L}_k, \quad i = 1, 2,$$

for some $N_k \rightarrow +\infty$.

Since on $\partial B_{\tilde{L}_k}$ both components $\tilde{v}_1^k, \tilde{v}_2^k$ have fast decay, we can apply the argument of Remark 3.1 and compute a local Pohozaev identity. The Pohozaev identity jointly with the fact that both local masses are a small perturbation of multiple of 4 yields that $(\sigma_1^k(\tau_k \tilde{L}_k), \sigma_2^k(\tau_k \tilde{L}_k))$ is a small perturbation of one of the two following types:

$$(2\tilde{m}(\tilde{m} + 1), 2\tilde{m}(\tilde{m} - 1)) \quad \text{or} \quad (2\tilde{m}(\tilde{m} - 1), 2\tilde{m}(\tilde{m} + 1)), \quad (5.3)$$

for some $\tilde{m} \in \mathbb{N}$. In the case where both \tilde{v}_i^k have fast decay one can get the same conclusion without using Theorem B since both local masses are already a small perturbation of multiple of 4.

Without loss of generality we assume the first alternative in (5.3) holds true. As in the proof of Proposition 4.1 we have

$$\begin{aligned} \bar{u}_k(\tau_k \tilde{L}_k) &\leq -2 \log(\tau_k \tilde{L}_k) - N_k, \\ \frac{d}{dr} \bar{u}_k &< -\frac{2 + \varepsilon}{r}, \end{aligned} \quad \text{for } r = \tau_k \tilde{L}_k,$$

for some $\varepsilon > 0$. Now, following the steps in the proof of Proposition 4.1, as r grows from $\tau_k \tilde{L}_k$ to $\tau_k L_k$, the following three cases may happen:

Case 1. Both u_k and $-u_k$ have fast decay up to $|x| = \tau_k L_k$:

$$|u_k(x)| \leq -2 \log |x| - N_k, \quad \tau_k \tilde{L}_k \leq |x| \leq \tau_k L_k,$$

for some $N_k \rightarrow +\infty$. In this case, by Lemma 4.1 we have

$$\sigma_i^k(\tau_k L_k) = \sigma_i^k(\tau_k \tilde{L}_k) + o(1), \quad i = 1, 2.$$

Case 2. There exists $L_{1,k} \in (\tilde{L}_k, L_k)$, $L_{1,k} = o(1)L_k$ such that

$$-u_k(x) \geq -2 \log \tau_k L_{1,k} - C, \quad \text{for } |x| = \tau_k L_{1,k}.$$

By the argument of Lemma 4.2 we can find a suitable $L_{2,k} \geq L_{1,k}$ such that

$$|u_k(x)| \leq -2 \log \tau_k L_{2,k} - N_k, \quad \text{for } |x| = \tau_k L_{2,k},$$

for some $N_k \rightarrow +\infty$ and $(\sigma_1^k(\tau_k L_{2,k}), \sigma_2^k(\tau_k L_{2,k}))$ is a $o(1)$ perturbation of

$$(2\bar{m}(\bar{m} - 1), 2\bar{m}(\bar{m} + 1)),$$

for some $\bar{m} \in \mathbb{N}$.

Case 3. $-u_k$ has slow decay for $|x| = \tau_k L_k$, i.e.

$$-u_k(x) \geq -2 \log \tau_k L_k - C, \quad |x| = \tau_k L_k,$$

for some $C > 0$ and

$$\sigma_1^k(\tau_k L_k) = \sigma_1^k(\tau_k \tilde{L}_k) + o(1) = 4\bar{n} + o(1).$$

Moreover, on $\partial B_{\tau_k L_k}(0)$, u_k is still the fast decaying component.

The only region we have to analyze is $B_{\tau_k L_k}(0) \setminus B_{\tau_k L_{2,k}}(0)$ when the second case above happens. However, the argument in this situation is as same as before. At the end, in any case on $\partial B_{\tau_k L_k}(0)$ at least one of the two components $u_k, -u_k$ has fast decay and its energy is a small perturbation of a multiple of 4.

Finally, we have to combine the groups. The procedure is very similar to the combination of bubbling disks as we have done before. For example, we start by considering groups which are close to each other: take $B_{\varepsilon_k}(0)$ for some $\varepsilon_k \rightarrow 0$ such that all the groups in $B_{2\varepsilon_k}(0)$, say G_0, G_1, \dots, G_t , (namely $(\Sigma_k \setminus (\cup_{i=0}^t G_i)) \cap B(0, 2\varepsilon_k) = \emptyset$) satisfy

$$\begin{aligned} \text{dist}(G_i, G_j) &\sim \text{dist}(G_l, G_q), \quad \forall i \neq j, l \neq q, \\ \text{dist}(G_i, G_j) &= o(1)\varepsilon_k, \quad \forall i, j = 0, \dots, t, i \neq j. \end{aligned}$$

The second property implies that the groups outside $B_{2\varepsilon_k}(0)$ are far away from the groups inside the ball. By the above assumptions, letting $\varepsilon_{1,k} = \text{dist}(G_0, G_j)$, for some $j \in \{1, \dots, t\}$ we have that all G_0, \dots, G_t are in $B_{N\varepsilon_{1,k}}(0)$ for some $N > 0$ which is independent of k . Without loss of generality let u_k be the fast decaying component on $\partial B_{N\varepsilon_{1,k}}(0)$. By Lemma 2.1, we have u_k is also a fast decaying component for G_0, \dots, G_t . As a consequence, we have

$$\sigma_1^k(N\varepsilon_{1,k}) = \sigma_1^k(\tau_k L_k) + 4\hat{m} + o(1),$$

for some $\hat{m} \in \mathbb{N}$. Now, as before we have three possible cases. If $-u_k$ also has fast decay on $\partial B_{N\varepsilon_{1,k}}(0)$, $\sigma_2^k(N\varepsilon_{1,k})$ is also a small perturbation of a multiple of 4 and we can get the quantization as in (5.3).

If instead

$$-u_k(x) \geq -2 \log N\varepsilon_{1,k} - C, \quad |x| = N\varepsilon_{1,k},$$

then as before we can find $\varepsilon_{2,k}$ in $(N\varepsilon_{1,k}, \varepsilon_k)$ such that

$$|u_k(x)| \leq -2 \log \varepsilon_{2,k} - N_k, \quad |x| = \varepsilon_{2,k},$$

for some $N_k \rightarrow \infty$. Moreover

$$\sigma_1^k(\varepsilon_{2,k}) = \sigma_1^k(N\varepsilon_{1,k}) + o(1).$$

Thus, by the usual argument we get the quantization as in (5.3).

The last possibility is

$$\sigma_1^k(\varepsilon_k) = \sigma_1^k(N\varepsilon_{1,k}) + o(1) = \sigma_1^k(\tau_k L_k) + 4\hat{m} + o(1)$$

and

$$-u_k(x) \geq -2 \log \varepsilon_k - C, \quad |x| = \varepsilon_k,$$

for some $C > 0$. In this case u_k is the fast decay component on $\partial B_{\varepsilon_k}(0)$.

It the end, we conclude that in any case on $\partial B_{\varepsilon_k}(0)$ at least one of the two component $u_k, -u_k$ has fast decay and its energy is a small perturbation of a multiple of 4. Then we can conclude as before by applying Theorem B and the local Pohozaev identity.

With this argument we continue to include groups further away from G_0 . Since by construction we have only finite blow-up disks, and this procedure only needs to be applied finite times. Then we can take $s_k \rightarrow 0$ such that $\Sigma_k \subset B_{s_k}(0)$ and both component $u_k, -u_k$ have fast decay on $\partial B_{s_k}(0)$:

$$|u_k(x)| \leq -2 \log s_k - N_k, \quad \text{for } |x| = s_k,$$

for some $N_k \rightarrow \infty$ and both $\sigma_i^k(s_k)$ are a small perturbation of multiple of 4. Combined with the Pohozaev identity, we get that $(\sigma_1^k(s_k), \sigma_2^k(s_k))$ is a small perturbation of one of the following two types:

$$(2m(m+1), 2m(m-1)) \quad \text{or} \quad (2m(m-1), 2m(m+1)),$$

for some $m \in \mathbb{N}$. Finally, by definition (1.9) we have

$$\sigma_i = \lim_{k \rightarrow \infty} \lim_{s_k \rightarrow 0} \sigma_i^k(s_k), \quad i = 1, 2.$$

It follows that σ_1, σ_2 satisfy the quantization property of Theorem 1.1 and the proof is completed. \square

REFERENCES

- [1] A. Bahri and J.M. Coron, The scalar curvature problem on the standard three dimensional sphere. *J. Funct. Anal.* 95 (1991), no. 1, 106-172.
- [2] L. Battaglia, A. Jevnikar, A. Malchiodi and D. Ruiz, A general existence result for the Toda system on compact surfaces. *Adv. Math.* 285 (2015), 937-979.
- [3] L. Battaglia and G. Mancini, A note on compactness properties of singular Toda systems. *Atti Accad. Naz. Lincei Rend. Lincei Mat. Appl.* 26 (2015), no. 3, 299-307.
- [4] H. Brezis and J.M. Coron, Multiple solutions of H-systems and Rellich's conjecture. *Commun. Pure Appl. Math.* 37 (1984), 149-187.
- [5] H. Brezis and F. Merle, Uniform estimates and blow-up behavior for solutions of $-\Delta u = V(x)e^u$ in two dimensions. *Comm. Partial Differential Equation* 16 (1991), 1223-1254.

- [6] E. Caglioti, P.L. Lions, C. Marchioro and M. Pulvirenti, A special class of stationary flows for two-dimensional Euler equations: a statistical mechanics description. *Comm. Math. Phys.* 143 (1992), no. 3, 501-525.
- [7] S.Y.A. Chang, M.J. Gursky and P.C. Yang, The scalar curvature equation on 2- and 3-spheres. *Calc. Var. and Partial Diff. Eq.* 1 (1993), no. 2, 205-229.
- [8] S.Y.A. Chang and P.C. Yang, Prescribing Gaussian curvature on S^2 . *Acta Math.* 159 (1987), no. 3-4, 215-259.
- [9] C.C. Chen and C.S. Lin, Estimate of the conformal scalar curvature equation via the method of moving planes. II. *J. Differential Geom.* 49 (1998), no. 1, 115-178.
- [10] W.X. Chen and C.M. Li, Classification of solutions of some nonlinear elliptic equations. *Duke Math. J.* 63 (1991), no. 3, 615-622.
- [11] A.J. Chorin, Vorticity and Turbulence, Springer, New York, 1994.
- [12] P. Esposito and J. Wei, Non-simple blow-up solutions for the Neumann two-dimensional sinh-Gordon equation. *Calc. Var. PDEs* 34 (2009), no. 3, 341-375.
- [13] M. Grossi and A. Pistoia, Multiple blow-up phenomena for the sinh-Poisson equation. *Arch. Ration. Mech. Anal.* 209 (2013), no. 1, 287-320.
- [14] Z.C. Han, Asymptotic approach to singular solutions for nonlinear elliptic equations involving critical Sobolev exponent. *Annales de l'I.H.P., section C* 8 (1991), no. 2, 159-174.
- [15] A. Jevnikar, An existence result for the mean field equation on compact surfaces in a doubly supercritical regime. *Proc. Royal Soc. Edinb. Sect. A* 143 (2013), no. 5, 1021-1045.
- [16] A. Jevnikar, A note on a multiplicity result for the mean field equation on compact surfaces. *Adv. Nonlinear Stud.* 16 (2016), no. 2, 221-229.
- [17] A. Jevnikar, New existence results for the mean field equation on compact surfaces via degree theory. To appear in *Rend. Semin. Mat. Univ. Padova*.
- [18] A. Jevnikar, J. Wei and W. Yang, On the topological degree of the mean field equation with two parameters. Preprint. Arxiv <http://arxiv.org/pdf/1602.03354v1>.
- [19] J. Jost, Two-dimensional geometric variational problems. Pure and Applied Mathematics, Wiley, Chichester, 1991.
- [20] J. Jost, G. Wang, D. Ye and C. Zhou, The blow-up analysis of solutions of the elliptic sinh-Gordon equation. *Calc. Var. PDEs* 31 (2008), 263-276.
- [21] G. Joyce and D. Montgomery, Negative temperature states for a two dimensional guiding center plasma. *J. Plasma Phys.* 10 (1973), 107-121.
- [22] M.K.H. Kiessling, Statistical mechanics of classical particles with logarithmic interactions. *Comm. Pure Appl. Math.* 46 (1993), no. 1, 27-56.
- [23] Y.Y. Li, Prescribing scalar curvature on S^n and related problems. I. *J. Differential Equations* 120 (1995), no. 2, 319-410.
- [24] Y.Y. Li, Harnack type inequality: the method of moving planes. *Comm. Math. Phys.* 200 (1999), no. 2, 421-444.
- [25] Y.Y. Li and I. Shafrir, Blow-up analysis for solutions of $-\Delta u = V e^u$ in dimension two. *Indiana Univ. Math. J.* 43 (1994), no. 4, 1255-1270.
- [26] C.S. Lin, J.C. Wei and W. Yang, Degree counting and shadow system for $SU(3)$ Toda system: one bubbling. Preprint. ArXiv <http://arxiv.org/pdf/1408.5802v1>.
- [27] C.S. Lin, J.C. Wei and L. Zhang, Classification of blowup limits for $SU(3)$ singular Toda systems. *Anal. PDE* 8 (2015), no. 4, 807-837.
- [28] C.S. Lin, J.C. Wei and L. Zhang, Classification of energy concentration of second order singular Toda systems. In preparation.
- [29] P.L. Lions, On Euler Equations and Statistical Physics, Scuola Normale Superiore, Pisa, 1997.
- [30] A. Malchiodi, Topological methods for an elliptic equation with exponential nonlinearities. *Discrete Contin. Dyn. Syst.* 21 (2008), no. 1, 277-294.
- [31] C. Marchioro and M. Pulvirenti, Mathematical Theory of Incompressible Nonviscous Fluids, Springer, New York, 1994.
- [32] P.K. Newton, The N-Vortex Problem: Analytical Techniques, Springer, New York, 2001.
- [33] H. Ohtsuka and T. Suzuki, Mean field equation for the equilibrium turbulence and a related functional inequality. *Adv. Differ. Equ.* 11 (2006), 281-304.
- [34] H. Ohtsuka and T. Suzuki, A blowup analysis of the mean field equation for arbitrarily signed vortices. *Self-similar solutions of nonlinear PDE* 74 (2006), 185-197.
- [35] L. Onsager, Statistical hydrodynamics. *Nuovo Cimento* 6 (1949), 279-287.

- [36] T.H. Parker, Bubble tree convergence for harmonic maps. *J. Differ. Geom.* 44 (1996), 595-633.
- [37] A. Pistoia and T. Ricciardi, Concentrating solutions for a Liouville type equation with variable intensities in 2D-turbulence, Preprint. ArXiv <http://arxiv.org/pdf/1507.01449v1>.
- [38] T. Ricciardi and R. Takahashi, Blow-up behavior for a degenerate elliptic sinh-Poisson equation with variable intensities, Preprint. ArXiv <http://arxiv.org/pdf/1507.01449>.
- [39] T. Ricciardi, R. Takahashi, G. Zecca and X. Zhang, On the existence and blow-up of solutions for a mean field equation with variable intensities, Preprint. ArXiv <http://arxiv.org/pdf/1509.05204>.
- [40] J. Sacks and K. Uhlenbeck, The existence of minimal immersions of 2-spheres. *Ann. Math.* 113 (1981), 1-24.
- [41] R. Schoen, Stanford Classnotes.
- [42] R. Schoen and D. Zhang, Prescribed scalar curvature on the n-sphere. *Calc. Var. PDEs* 4 (1996), no. 1, 1-25.
- [43] K. Steffen, On the nonuniqueness of surfaces with prescribed constant mean curvature spanning a given contour. *Arch. Rat. Mech. Anal.* 94, (1986) 101-122.
- [44] M. Struwe, Nonuniqueness in the Plateau problem for surfaces of constant mean curvature. *Arch. Rat. Mech. Anal.* 93, (1986) 135-157.
- [45] G. Tarantello, Analytical, geometrical and topological aspects of a class of mean field equations on surfaces. *Discrete Contin. Dyn. Syst.* 28 (2010), no. 3, 931-973.
- [46] H.C. Wente, Counterexample to a conjecture of H. Hopf. *Pacific J. Math.* 121 (1986), no. 1, 193-243.
- [47] C. Zhou, Existence result for mean field equation of the equilibrium turbulence in the super critical case. *Commun. Contemp. Math.* 13 (2011), no 4, 659-673.

ALEKS JEVIKAR, UNIVERSITY OF ROME 'TOR VERGATA', VIA DELLA RICERCA SCIENTIFICA 1, 00133 ROMA, ITALY

E-mail address: jevnikar@mat.uniroma2.it

JUNCHENG WEI, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF BRITISH COLUMBIA, VANCOUVER, BC V6T 1Z2, CANADA

E-mail address: jcwei@math.ubc.ca

WEN YANG, CENTER FOR ADVANCED STUDY IN THEORETICAL SCIENCES (CASTS), NATIONAL TAIWAN UNIVERSITY, TAIPEI 10617, TAIWAN

E-mail address: math.yangwen@gmail.com