



UNIVERSITÀ
DEGLI STUDI
DI UDINE

Università degli studi di Udine

Pseudospectral discretization of delay differential equations in sun-star formulation: Results and conjectures

Original

Availability:

This version is available <http://hdl.handle.net/11390/1171202> since 2020-07-13T17:43:35Z

Publisher:

Published

DOI:10.3934/dcdss.2020196

Terms of use:

The institutional repository of the University of Udine (<http://air.uniud.it>) is provided by ARIC services. The aim is to enable open access to all the world.

Publisher copyright

(Article begins on next page)

1 **PSEUDOSPECTRAL DISCRETIZATION OF DELAY**
2 **DIFFERENTIAL EQUATIONS IN *SUN-STAR* FORMULATION:**
3 **RESULTS AND CONJECTURES**

ODO DIEKMANN

Department of Mathematics, Utrecht University
P.O. Box 80010, 3508 TA Utrecht, The Netherlands

FRANCESCA SCARABEL

Department of Mathematics and Statistics, University of Helsinki
P.O. 68 (Pietari Kalmin katu 5), FI-00014 Helsinki, Finland

ROSSANA VERMIGLIO*

Department of Mathematics, Computer Science and Physics, University of Udine
via delle Scienze 206, I-33100 Udine, Italy

Dedicated to prof. C.T.H. Baker

ABSTRACT. In this paper we study the pseudospectral approximation of delay differential equations formulated as abstract differential equations in the $\odot*$ -space. This formalism also allows us to define rigorously the abstract variation-of-constants formula, where the $\odot*$ -shift operator plays a fundamental role. By applying the pseudospectral discretization technique we derive a system of ordinary differential equations, whose dynamics can be efficiently analyzed by existing bifurcation tools. To better understand to what extent the resulting finite-dimensional system “mimics” the dynamics of the original infinite-dimensional one, we study the pseudospectral approximations of the $\odot*$ -shift operator and of the $\odot*$ -generator in the supremum norm, which is the natural choice for delay differential equations, when the discretization parameter increases. In this context there are still open questions. We collect the most relevant results from the literature and we present some conjectures, supported by various numerical experiments, to illustrate the behavior w.r.t. the discretization parameter and to indicate the direction of ongoing and future research.

2010 *Mathematics Subject Classification.* Primary: 65F35,65L03,65L07,65M70,65P99,65Q20,37M20X,34K07, 34K20,41A10; Secondary: 34K08,34K28, 37K45.

Key words and phrases. Sun-star formalism, delay differential equations, pseudospectral discretization, collocation, Chebyshev differentiation matrix, shift semigroup, differential operators.

The research of the second author was supported by Domast (Doctoral Programme in Mathematics and Statistics, University of Helsinki), and by the Centre of Excellence in Analysis and Dynamics Research, Academy of Finland. F.S. and R.V. are members of the INdAM Research group GNCS, and of CDLab (Computational Dynamics Laboratory), Department of Mathematics, Computer Science and Physics, University of Udine.

* Corresponding author: Rossana Vermiglio.

1 **1. Introduction.** Delay differential equations (DDEs), together with renewal equa-
 2 tions (REs) and systems which couple REs and DDEs (REs/DDEs), are recognized
 3 as a fundamental tool for modelling phenomena in many fields, including, for in-
 4 stance, population dynamics and control theory. For this reason, in the last years
 5 the interest in the study of the dynamics of delay models has been increasing and
 6 important challenges, in particular numerical, have been identified. Indeed, delay
 7 equations describe infinite-dimensional dynamical systems, and theoretical results
 8 should be complemented with efficient numerical methods to approximate solutions
 9 of initial value problems [3, 4, 5, 7, 17, 15, 16, 19, 44, 43], boundary value prob-
 10 lems [48, 49, 50], and to investigate the stability of equilibria and periodic solutions
 11 [10, 11, 12, 13, 14, 47, 52, 53, 66]. In applications the attention is focused not only
 12 on the approximation of the dynamical properties for some given parameter values,
 13 but also on how such properties change when varying some parameters. In particu-
 14 lar, it is interesting to identify the critical thresholds, called bifurcation points, and
 15 to draw stability charts in two or more parameters.

16 While the theory of DDEs is well developed, see for instance [6, 25, 39, 65],
 17 the numerical methods are still unsatisfactory: some software packages, like DDE-
 18 BIFTOOL [27] and Knut [59], are available for the stability and bifurcation analysis
 19 of equations with discrete delays, but cannot be applied, for instance, in the case
 20 of distributed delays. To address this problem, the authors of [8, 56] proposed an
 21 alternative method, which consists in reformulating the original equation as a non-
 22 linear abstract differential equation (ADE) in the state space, and then applying the
 23 pseudospectral discretization (PSD) approach. Then, the dynamics of the resulting
 24 nonlinear systems of ordinary differential equations (ODEs) can be investigated by
 25 one of the continuation packages for the bifurcation analysis of ODEs, like MatCont
 26 [22]. In [8] the approach has been applied also to REs and REs/DDEs. Moreover,
 27 it has been proved that there is a one-to-one correspondence between the equilibria,
 28 and the stability is accurately described. Numerous experiments give evidence that
 29 the PSD approach converges for periodic solutions, too [8, 9], but at the moment
 30 the rigorous proof is not yet completed. In this context, the approximation of the
 31 so-called *shift operator*, which translates a function to the right while extending it
 32 with a constant value, plays a fundamental role. An important question is how
 33 accurately the shift operator is approximated by the resulting discrete operator,
 34 w.r.t. the discretization parameter. The action of the shift operator is described by
 35 the trivial equation $y'(t) = 0$ considered as a DDE with delay $\tau > 0$, and is related
 36 to the partial differential equation (PDE)

$$\begin{cases} \frac{\partial v}{\partial t}(t, \theta) = \frac{\partial v}{\partial \theta}(t, \theta), & \theta \in [-\tau, 0], \\ \frac{\partial v}{\partial \theta}(t, 0) = 0, \end{cases} \quad (1)$$

37 [25, pag. 39]. The PSD approach has been widely applied for the numerical approx-
 38 imation of PDEs [18, 31], and many results are available in the literature. Never-
 39 theless, the estimates of the approximation of the shift operator regard weighted
 40 2-norms in suitable Sobolev spaces [26, 29, 30, 35, 33, 34, 45, 40, 58, 62, 64], whereas,
 41 to our knowledge, no results are available for the supremum norm. The supremum
 42 norm is largely used when studying DDEs with the Banach space of continuous
 43 functions as state space [25, 39]. Nevertheless this choice is not unique and, hav-
 44 ing in mind especially some applications to control theory, L^p -state space has been
 45 considered in [6, 19, 65].

1 In this paper we apply the PSD approach, originally proposed for DDEs in [8],
 2 to the reformulation of the DDE in the $\odot*$ -space. The $\odot*$ -formalism was intro-
 3 duced by [20, 23, 25] for proving the principle of linearized stability for DDEs, REs
 4 and REs/DDEs. This formalism allows to define rigorously both the ADE and the
 5 abstract variation-of-constants formula, where the $\odot*$ -shift operator plays a fun-
 6 damental role. As a further advantage, the $\odot*$ -formulation allows to interpret the
 7 original nonlinear equation as a nonlinear perturbation of a “trivial” linear equation,
 8 separating clearly the action of translation, which is the same for every equation,
 9 from the rule for extension, which is problem-specific and described in terms of
 10 the nonlinear perturbation. From the point of view of the numerical method, the
 11 variation-of-constants formula makes it easier to compare the solution operators of
 12 the continuous and discrete dynamical systems in terms of the $\odot*$ -shift operator
 13 and its discretization. Moreover the readers familiar with spectral approximation
 14 methods as described for instance in [35], will recognize the usual formalism (see
 15 the definitions of *prolongation* and *restriction* operator in Section 3).

16 The PSD approach applied to the $\odot*$ -formulation returns the same approxim-
 17 ating ODE as obtained in [8], but the new interpretation allows one to distinguish
 18 the approximation of the linear part, which relates to the $\odot*$ -shift operator, from
 19 the approximation of the nonlinear perturbation. Then, the comparison of the
 20 variation-of-constants formulas motivates the importance of studying the accuracy
 21 of the approximation of the $\odot*$ -shift operator w.r.t. the discretization parameter.

22 After introducing the PSD and the resulting ODE, we recall the fundamental
 23 results and we present some conjectures and computational experiments about the
 24 asymptotic behavior of the PSD of the $\odot*$ -shift operator and its generator w.r.t.
 25 the discretization parameter. Our aim is to survey what is known and to indi-
 26 cate possible future research directions, to obtain bounds in supremum norm in
 27 order to better understand to which extent the finite-dimensional ODE “mimics”
 28 the dynamics of the infinite-dimensional system described by the DDE. For some
 29 applications of the technique we refer to [2, 8, 9, 32, 37].

30 The paper is organized as follows. In Section 2 we recall the $\odot*$ -formulation of
 31 DDEs, whereas the PSD approach is illustrated in Section 3. In Sections 4 and 5 we
 32 study respectively the discretized version of the infinitesimal generators and of the
 33 shift operators, by using also some numerical experiments. In Section 6 we draw
 34 some conclusions and discuss ongoing and future research.

35 **2. $\odot*$ -formulation of delay differential equations.** In this section we briefly
 36 summarize the basic results on the $\odot*$ -reformulation of DDEs, with the aim to in-
 37 troduce the ADE and to emphasize the role of the $\odot*$ -shift operator in this abstract
 38 framework, in view of the introduction of the PSD approach in Section 3. For a
 39 deeper insight into the theory, we refer to [20, 23, 25].

40 Let d be a positive integer, $\tau > 0$, and consider the Banach space

$$Y := C([-\tau, 0]; \mathbb{R}^d)$$

41 equipped with the norm $\|\psi\|_Y := \max_{\theta \in [-\tau, 0]} |\psi(\theta)|$, for $|\cdot|$ a norm on \mathbb{R}^d .

42 A nonlinear autonomous DDE is

$$y'(t) = G(y_t) \tag{2}$$

43 where $G: Y \rightarrow \mathbb{R}^d$ is a continuous nonlinear function, and the history $y_t \in Y$ is
 44 defined as

$$y_t(\theta) = y(t + \theta), \quad \theta \in [-\tau, 0].$$

1 **Definition 2.1.** [25, p. 231–232] A *solution* of (2) is either a continuously differ-
 2 entiable function $y : \mathbb{R} \rightarrow \mathbb{R}^d$ satisfying (2) everywhere, or a continuous function
 3 $y : [-\tau, t_+) \rightarrow \mathbb{R}^d$, $0 < t_+ \leq \infty$, which is differentiable on $(0, t_+)$ and satisfies (2)
 4 for $0 < t < t_+$.

5 Given the initial condition

$$y(\theta) = \psi(\theta), \quad \theta \in [-\tau, 0], \quad \psi \in Y, \quad (3)$$

6 a solution $y(\cdot; \psi)$ of the initial value problem (2) & (3) is a solution $y : [-\tau, t_+) \rightarrow \mathbb{R}^d$
 7 of (2) which also satisfies (3).

8 Hereafter for ease of formulation we assume that G is globally Lipschitz contin-
 9 uous with Lipschitz constant $Lip(G) > 0$. So the initial value problem (2) & (3)
 10 has a unique solution $y(\cdot; \psi)$ with $t_+ = +\infty$, which depends continuously on ψ
 11 (see [25, 39]).

12 An example of a solution of the first type in Definition 2.1 is given by a periodic
 13 solution \bar{y} of (2), which is differentiable and periodic with period h , i.e. $\bar{y}(t+h) =$
 14 $\bar{y}(t)$, $t \in \mathbb{R}$, and satisfies (2) everywhere.

15 The \odot^* -formulation of DDEs aims to represent the equation (2) in an abstract
 16 framework, where the action of the dynamical system can be described in terms of
 17 two different components: the action of “shift”, which is linear and exactly the same
 18 for every equation, and an action of “extension”, which captures the effect of the
 19 nonlinear function G . In the \odot^* -framework the perturbation theory can be applied,
 20 and the variation-of-constants formula can be rigorously defined [20, 23, 25].

21 The main idea of the \odot^* -formulation is to embed Y into the larger space Y^{\odot^*}
 22 given by

$$Y^{\odot^*} := \mathbb{R}^d \times L^\infty([-\tau, 0], \mathbb{R}^d),$$

which is a Banach space equipped with the norm

$$\|(\alpha, \psi)\|_{Y^{\odot^*}} := \max\{|\alpha|, \|\psi\|_{L^\infty}\}, \quad (\alpha, \psi) \in Y^{\odot^*}.$$

23 The embedding $j : Y \rightarrow Y^{\odot^*}$ is given by $j(\psi) = (\psi(0), \psi)$.

24 As for the “shift” component, it is well known that the generator of translation is
 25 differentiation. So we introduce the operator $\mathcal{A}_0^{\odot^*} : D(\mathcal{A}_0^{\odot^*}) (\subseteq Y^{\odot^*}) \rightarrow Y^{\odot^*}$ given
 26 by

$$\begin{aligned} \mathcal{A}_0^{\odot^*}(\alpha, \psi) &= (0, \psi'), \quad (\alpha, \psi) \in D(\mathcal{A}_0^{\odot^*}), \\ D(\mathcal{A}_0^{\odot^*}) &:= \{(\alpha, \psi) \in Y^{\odot^*} : \psi \text{ is Lipschitz continuous and } \psi(0) = \alpha\}, \end{aligned} \quad (4)$$

27 where “Lipschitz continuous” is a short-hand way to indicate an equivalence class
 28 that contains a Lipschitz continuous function.

29 The operator $\mathcal{A}_0^{\odot^*}$ is the infinitesimal generator in the weak*-sense of the \odot^* -*shift*
 30 *semigroup* $\{\mathcal{T}_0^{\odot^*}(t)\}_{t \geq 0}$, where the \odot^* -*shift operator* is defined by

$$\mathcal{T}_0^{\odot^*}(t)(\alpha, \psi) = (\alpha, \psi_t^\alpha), \quad t \geq 0,$$

31 and

$$\psi_t^\alpha(\theta) := \begin{cases} \psi(t+\theta), & t+\theta \leq 0 \\ \alpha, & t+\theta > 0. \end{cases}$$

But we do not want to lose sight of Y , so we introduce the space

$$Y^{\odot\odot} := \overline{D(\mathcal{A}_0^{\odot^*})} = \{(\alpha, \psi) \in Y^{\odot^*} : \psi \text{ is continuous and } \psi(0) = \alpha\}.$$

1 where, as before, “continuous” indicates an equivalence class that contains a con-
 2 tinuous function. Since the delay is bounded, the space Y is \odot -reflexive in the sense
 3 that j is a bijection between Y and $Y^{\odot\odot}$, with $j^{-1}(\alpha, \psi) = \psi$, see e.g. [20, 25].

4 Finally, by describing the rule for the extension in $Y^{\odot*}$ through the nonlinear
 5 operator $\mathcal{G}: Y^{\odot\odot} \rightarrow Y^{\odot*}$ given by

$$\mathcal{G}(\alpha, \psi) := (G(\psi), 0) \in Y^{\odot*}, \quad (\alpha, \psi) \in Y^{\odot\odot},$$

6 we can construct the operator $\mathcal{A}^{\odot*}: D(\mathcal{A}^{\odot*})(\subseteq Y^{\odot*}) \rightarrow Y^{\odot*}$ defined by

$$\mathcal{A}^{\odot*} = \mathcal{A}_0^{\odot*} + \mathcal{G}, \quad D(\mathcal{A}^{\odot*}) = D(\mathcal{A}_0^{\odot*}), \quad (5)$$

7 whose action describes both the “shift” and “extension”, through $\mathcal{A}_0^{\odot*}$ and \mathcal{G} re-
 8 spectively, while its domain does not depend on the particular equation. Notice that
 9 \mathcal{G} inherits from G the Lipschitz continuity. The operator (5) allows to represent the
 10 DDE (2) as an evolutionary system in the $Y^{\odot*}$ space, namely the semilinear ADE
 11 in $Y^{\odot*}$

$$\frac{djv(t)}{dt} = \mathcal{A}^{\odot*}(jv(t)). \quad (6)$$

12 For $\psi \in Y$, we define *mild solution* of the Cauchy problem

$$\begin{cases} \frac{djv(t)}{dt} = \mathcal{A}^{\odot*}(jv(t)), & t > 0, \\ v(0) = \psi, \end{cases} \quad (7)$$

13 the unique solution of the abstract integral equation (AIE)

$$jv(t) = \mathcal{T}_0^{\odot*}(t)j\psi + \int_0^t \mathcal{T}_0^{\odot*}(t-s)\mathcal{G}(jv(s))ds, \quad t \geq 0, \quad (8)$$

where the integral is a weak*-Riemann integral [20, Theorem 3.1]. From [20, Propo-
 sition 2.1] we have that the integral at the right-hand side of (8) defines an element
 of $Y^{\odot\odot}$, so we can apply j^{-1} to both sides of (8) and write the following AIE in
 the original space Y

$$v(t) = \mathcal{T}_0(t)\psi + j^{-1} \int_0^t \mathcal{T}_0^{\odot*}(t-s)\mathcal{G}(jv(s))ds, \quad t \geq 0,$$

14 where $\mathcal{T}_0(t) := j^{-1}\mathcal{T}_0^{\odot*}(t)j$ is the shift operator on Y . Following [25, Chapter III],
 15 we can conclude that $v(t) = y_t(\cdot; \psi)$, where $y(\cdot; \psi)$ is a solution of the initial value
 16 problem (2) & (3).

17 The part of $\mathcal{A}^{\odot*}$ in $Y^{\odot\odot}$ is the operator $j\mathcal{A}j^{-1}$, with \mathcal{A} defined by

$$\begin{aligned} \mathcal{A}\psi &= \psi', \quad \psi \in D(\mathcal{A}) \\ D(\mathcal{A}) &:= \{\psi \in Y : \psi' \in Y \text{ and } \psi'(0) = G(\psi)\}, \end{aligned} \quad (9)$$

18 whose action is independent of G , whereas now the information about the particular
 19 DDE is incorporated in the domain. For $\psi \in D(\mathcal{A})$, the solution $jv(t)$ of (7) is a
 20 *classical solution*, i.e. a continuously differentiable function $jv(t)$ satisfying (6) for
 21 $t \geq 0$ [20, Theorem 3.6].

22 The key role of the \odot *-shift operator $\mathcal{T}_0^{\odot*}(t)$ and of the generator $\mathcal{A}_0^{\odot*}$ comes to
 23 light clearly in respectively (8) and (6). This will be also useful in the analysis of
 24 their discretization in Section 3. Notice that they describe the \odot *-formulation of
 25 the trivial equation

$$y'(t) = 0, \quad t \geq 0, \quad (10)$$

26 interpreted as a DDE.

1 We remark that the nonlinearity in the domain of (9) makes it hard to handle
 2 the perturbations of (2), since they amount to perturbing the domain of an infin-
 3 itesimal generator. The main advantage of the \odot^* -formulation is indeed to avoid
 4 the dependence of the domains on G , thus treating operators defined on the same
 5 domain. So, although the \odot^* -framework requires some additional technical effort,
 6 the result yields a powerful abstract variation-of-constants formula for a larger class
 7 of perturbations than those bounded from Y into Y . The method also applies to
 8 other classes of delay equations, namely REs of the form

$$x(t) = F(x_t),$$

9 with $x_t \in X := L^1((-\tau, 0), \mathbb{R}^d)$ and $F: X \rightarrow \mathbb{R}^d$ a smooth function, and REs/DDEs

$$\begin{cases} x(t) = F(x_t, y_t) \\ y'(t) = G(x_t, y_t) \end{cases}$$

10 with $(x_t, y_t) \in X \times Y$ and $F, G: X \times Y \rightarrow \mathbb{R}^d$ smooth functions [23].

11 **3. \odot^* -pseudospectral discretization.** In this section we introduce the PSD of
 12 the operator \mathcal{A}^{\odot^*} in (5) with the aim to derive a nonlinear system of ODEs that
 13 “mimics” the dynamics of the original nonlinear DDE (2). The basic idea of the
 14 PSD approach consists in approximating a function with a polynomial, which is
 15 represented as the interpolating polynomial on a suitable set of nodes in the domain
 16 of definition. Then “do to the interpolating polynomial what you would do to the
 17 function” [2].

In order to start from pointwise defined functions and to keep in touch with the original DDE, we will develop the PSD approach in the subspace

$$\widehat{Y} := \mathbb{R}^d \times Y$$

18 of Y^{\odot^*} . Note that $Y^{\odot\odot} \subset \widehat{Y}$.

19 Let M be a positive discretization integer and denote by Π_M the space of all
 20 the \mathbb{R}^d -valued polynomials of degree at most M defined on $[-\tau, 0]$. Consider the
 21 extremal Chebyshev points relative to the delay interval $[-\tau, 0]$, i.e.

$$\theta_{i,M} = \frac{\tau}{2} \left(\cos\left(\frac{i\pi}{M}\right) - 1 \right), \quad i = 0, \dots, M, \quad (11)$$

22 and define

$$\Theta_M = \{\theta_{i,M} : i = 1, \dots, M\}. \quad (12)$$

We first introduce the space

$$Y_M := \mathbb{R}^{dM},$$

with the norm $\|\Psi\|_{Y_M} := \max_{i=1, \dots, M} |\Psi_i|$, where $\Psi \in Y_M$ is a column vector of length dM that consists of M d -dimensional column vectors $\Psi_i, i = 1, \dots, M$ stacked on the top of each other. The space

$$\widehat{Y}_M := \mathbb{R}^d \times Y_M,$$

23 endowed with the norm

$$\|(\alpha, \Psi)\|_{\widehat{Y}_M} := \max\{|\alpha|, \|\Psi\|_{Y_M}\}, \quad (13)$$

24 represents the discretization of index M of \widehat{Y} , in the sense that every (α, ψ) is dis-
 25 cretized by the element $(\alpha, \psi(\Theta_M)) \in \widehat{Y}_M$. The choice of the norm (13) is motivated
 26 by the norm of the state space Y^{\odot^*} .

1 As further ingredients we introduce the *restriction* operator $\widehat{R}_M: \widehat{Y} \rightarrow \widehat{Y}_M$ as

$$\widehat{R}_M(\alpha, \psi) = (\alpha, R_M\psi), \quad (\alpha, \psi) \in \widehat{Y}, \quad (14)$$

2 where $R_M: Y \rightarrow Y_M$ is

$$R_M\psi = (\psi(\Theta_M)), \quad \psi \in Y,$$

3 and the *prolongation* operator $\widehat{P}_M: \widehat{Y}_M \rightarrow D(\mathcal{A}_0^{\circ*}) \subset Y^{\circ\circ}$, which associates to
4 $(\alpha, \Psi) \in \widehat{Y}_M$ the pair (α, ψ_M) , where ψ_M is the M -degree polynomial interpolat-
5 ing (α, Ψ) at the nodes (11). It is important to underline that the construction of
6 ψ_M takes into account the domain condition in (4). By introducing the *Lagrange*

7 *polynomials* $\ell_j(\theta) := \prod_{k=0, k \neq j}^M \frac{\theta - \theta_k}{\theta_j - \theta_k}, j = 0, \dots, M$, and by recalling that they parti-

8 tion unity, i.e. $\sum_{j=0}^M \ell_j = 1$, we obtain the following *Lagrange representation* of the
9 interpolating polynomial ψ_M

$$\begin{aligned} \psi_M(\theta) &= \ell_0(\theta)\alpha + \sum_{j=1}^M \ell_j(\theta)\Psi_j, \\ &= \alpha + \sum_{j=1}^M \ell_j(\theta)(\Psi_j - \alpha), \quad \theta \in [-\tau, 0], \end{aligned} \quad (15)$$

(see e.g. [60, p. 34]). Hereafter, when it will be necessary to emphasize the dependence on (α, Ψ) , we also use the notation

$$\psi_M = P_M(\alpha, \Psi),$$

10 where $P_M := j^{-1}\widehat{P}_M$. Note that $\widehat{R}_M\widehat{P}_M$ is the identity operator on \widehat{Y}_M , while
11 $P_M\widehat{R}_M j$ is the interpolation operator on Y relative to the Chebyshev nodes (11).

12 Now we are ready to define both the PSD of $\mathcal{A}_0^{\circ*}$ as the bounded operator
13 $A_{0,M}: \widehat{Y}_M \rightarrow \widehat{Y}_M$ given by

$$A_{0,M} := \widehat{R}_M \mathcal{A}_0^{\circ*} \widehat{P}_M, \quad (16)$$

14 and the nonlinear map $G_M: \widehat{Y}_M \rightarrow \widehat{Y}_M$ as

$$G_M := \widehat{R}_M \mathcal{G} \circ \widehat{P}_M \quad (17)$$

where \circ denotes the composition of (nonlinear) operators. Their sum furnishes the discretization A_M of $\mathcal{A}^{\circ*}$, i.e.

$$A_M = A_{0,M} + G_M.$$

15 So the PSD approach leads to the following nonlinear ODE in \widehat{Y}_M

$$\frac{d}{dt} v_M(t) = A_M(v_M(t)). \quad (18)$$

16 To write (18) explicitly, we need to find a matrix representation of the operator
17 (16). Let us define the elements $d_{ij} := \ell'_j(\theta_i)$, for $i, j = 0, \dots, M$ and the matrix

$$D_M = (d_{ij})_{i,j=1,\dots,M} \in \mathbb{R}^{M \times M}, \quad (19)$$

18 which will play a fundamental role in the definition and analysis of the discretized
19 equation (18).

1 Since $\ell'_0(\theta_i) = -\sum_{j=1}^M \ell'_j(\theta_i)$, $i = 0, \dots, M$, by denoting $\mathbf{1}$ the vector with all entries
 2 equal to 1 (we do not specify the dimension, since it is clear from the context), from
 3 (15) we get

$$A_{0,M} = \begin{pmatrix} 0 & 0 \\ -D_M \mathbf{1} \otimes I_d & D_M \otimes I_d \end{pmatrix} \quad (20)$$

where \otimes is the tensor product and I_d denotes the identity matrix of dimension d .
 By posing $v_M(t) := (y_M(t), V_M(t))$, and by using (17) and (20), the equation (18)
 can be expressed as

$$\begin{cases} y'_M(t) = G(P_M(y_M(t), V_M(t))), \\ V'_M(t) = D_M \otimes I_d (V_M(t) - \mathbf{1} \otimes y_M(t)). \end{cases}$$

4 **Remark 1.** Since we have to handle ψ such that $j\psi \in D(\mathcal{A}_0^{\odot*})$, the Chebyshev
 5 points (11) are a good choice as interpolation nodes. Let us define the *Lebesgue*
 6 *function*

$$\lambda_M(\theta) = \sum_{j=0}^M |\ell_j(\theta)|, \quad \theta \in [-\tau, 0],$$

7 and the *Lebesgue constant*

$$\Lambda_M = \max_{\theta \in [-\tau, 0]} \sum_{j=0}^M |\ell_j(\theta)|.$$

8 For the choice of nodes (11) the Lebesgue constants satisfy

$$\Lambda_M \leq \frac{2}{\pi} \log(M+1) + 1 \text{ and } \Lambda_M \approx \frac{2}{\pi} \log(M), \quad M \rightarrow \infty,$$

(see e.g. [60, pag.109]). So for the M -degree polynomial ψ_M interpolating a Lips-
 chitz continuous function ψ at the nodes (11), we get the following error bound

$$\|\psi - \psi_M\|_Y \leq \left(1 + \frac{\pi^2}{2}\right) (1 + \Lambda_M) \frac{\tau \text{Lip}(\psi)}{2M},$$

which guarantees the uniform convergence of $\{\psi_M\}_M$ to ψ as $M \rightarrow \infty$ (see e.g. [60,
 pag. 108] and [21, pag. 338]). Moreover, under the same assumptions on ψ , we also
 have

$$|G(\psi) - G(\psi_M)| \leq \text{Lip}(G) \|\psi - \psi_M\|_Y \rightarrow 0 \text{ as } M \rightarrow \infty,$$

and, when also ψ' is a Lipschitz continuous function we have

$$\|\psi' - \psi'_M\|_Y \leq C \log(M+1) \frac{\text{Lip}(\psi')}{M}$$

9 with C a positive constant independent of M and ψ [51, pag. 269], and $\|\psi' -$
 10 $\psi'_M\|_Y \rightarrow 0$ as $M \rightarrow \infty$. Notice that the convergence results mentioned above apply
 11 also to $\psi = \bar{y}_t$, where \bar{y} is a periodic solution of (2).

12 Finally, similarly to the abstract framework in Section 2, we complete the path
 13 by deriving also the integral equation associated to (18). Given the initial condition

$$v_M(0) = (\alpha, \Psi), \quad (\alpha, \Psi) \in \widehat{Y}_M, \quad (21)$$

1 the initial value problem (18) & (21) is equivalent to the following integral equation
 2 in \widehat{Y}_M

$$v_M(t) = T_{0,M}(t)(\alpha, \Psi) + \int_0^t T_{0,M}(t-s)G_M(v_M(s)) ds, \quad t \geq 0, \quad (22)$$

3 where $T_{0,M}(t) : \widehat{Y}_M \rightarrow \widehat{Y}_M$ is given by $T_{0,M}(t) = e^{A_{0,M}t}$ and admits the following
 4 representation

$$T_{0,M}(t) = \begin{pmatrix} I_d & 0 \\ (\mathbf{1} - e^{D_M t} \mathbf{1}) \otimes I_d & e^{D_M t} \otimes I_d \end{pmatrix}, \quad t \geq 0. \quad (23)$$

5 We underline the close analogy in structure between the integral formulations
 6 (22) and (8). We will return to this in Section 5.

7 In the paper [8] the PSD approach has been applied to the operator (9) obtaining
 8 the same ODE (18). So what did we gain by working in the \odot^* framework? It has
 9 furnished a deep insight into the connection between the infinite-dimensional and
 10 the finite-dimensional equations, including not only the differential equations (6)
 11 and (18), but also the integral equations (8) and (22). Moreover, the regularity
 12 condition of the functions in the domain has been relaxed w.r.t. those in $D(\mathcal{A})$ in
 13 (9), and so we can handle Lipschitz continuous functions.

14 In order to use the ODE (18) to gain insight into the dynamics of the original
 15 DDE (2), we need to understand how the dynamical properties of the two equations
 16 are related. In the paper [8] it has been proved that the equilibria of (2) and
 17 (18) are in one-to-one correspondence, and the stability is the same for M large
 18 enough. In order to understand the behavior of more complicated solutions, like
 19 periodic solutions, it is fundamental to study the solution operators generated by
 20 the nonlinear equations, which are described by the integral equations (8) and (22).
 21 The integral formulation is indeed one of the advantages of the \odot^* -formalism and
 22 it is particularly useful for the convergence analysis of solution operators and, in
 23 turn, of periodic solutions. Indeed, thanks to the integral equations, we can relate
 24 the convergence of the nonlinear semigroups to the convergence of the trivial linear
 25 semigroups $T_{0,M}(t)$ to $\mathcal{T}_0^{\odot^*}(t)$. Moreover (4) and (20) suggest that the matrix (16)
 26 "mimics" the infinitesimal generator $\mathcal{A}_0^{\odot^*}$ of the semigroup $\{\mathcal{T}_0^{\odot^*}(t)\}_{t \geq 0}$ associated
 27 to the trivial DDE (10). Therefore, to understand the dynamical behavior of (18), it
 28 is crucial to investigate the properties of the matrix $A_{0,M}$ and of the corresponding
 29 semigroup $\{T_{0,M}(t)\}_{t \geq 0}$ as $M \rightarrow \infty$.

30 **4. The generators $\mathcal{A}_0^{\odot^*}$ and $A_{0,M}$: results.** In this section we focus on the PSD
 31 $A_{0,M}$ of the infinitesimal generator $\mathcal{A}_0^{\odot^*}$ and on their connection. From (20) it is
 32 clear that the matrix D_M plays a major role. We remark that D_M can be viewed
 33 as the Lagrange representation of the PSD w.r.t. the nodes (12) of the derivative
 34 operator $\mathcal{D} : D(\mathcal{D}) \subset L^\infty \rightarrow L^\infty$

$$\mathcal{D}\psi = \psi', \quad (24)$$

35 with domain $D(\mathcal{D})$ the set of equivalence classes containing a Lipschitz continuous
 36 function ψ such that $\psi(0) = 0$ [25, pag. 124]. In the following we collect some of the
 37 results about the matrix D_M available in the literature, which are mainly obtained
 38 in the context of the approximations of PDEs of the form (1).

39 For the sake of presentation hereafter we assume $d = 1$, and we use $\|\cdot\|$ for all
 40 the norms, omitting the subscripts, since the space we are working with will be
 41 clear from the context. Moreover in the numerical experiments we consider $\tau = 1$.

1 Indeed the results for general $\tau > 0$ can be easily obtained by a suitable scaling of
 2 the time variable.

3 All the experiments are made with Matlab 2018b, whose machine precision is
 4 $u = \epsilon/2$ with $\epsilon = 2^{-52} \approx 2.2204 \times 10^{-16}$.

5 Finally, hereafter we consider complex-valued functions and complex Banach
 6 spaces. For a detailed presentation of the complexification procedure, the interested
 7 reader is referred to [25, Sections III.7 and IV.2].

8 Since the derivative operator (24) is unbounded, we expect $\|D_M\|$ to diverge
 9 as $M \rightarrow \infty$. The following proposition confirms this claim and defines the order
 10 of divergence w.r.t. M , which is determined by the asymptotic behavior of the
 11 Chebyshev polynomials.

12 **Proposition 1.** [55, section 2.7.4] $\|D_M\| = O(M^2)$.

13 Indeed the numerical results in Table 1 give evidence that $\|D_M\| = 2M^2 - 1$ for
 14 $\tau = 1$.

M	$\ D_M\ $	order	$\ D_M\ /M^2$
4	31		1.9375
8	127	2.0345	1.9844
16	511	2.0085	1.9961
32	2047	2.0021	1.9990
64	8191	2.0005	1.9998
128	32767	2.0001	1.9999
256	131071	2.0000	2.0000

TABLE 1. $\|D_M\|$ and estimation of the order $\log_2 \frac{\|D_{2M}\|}{\|D_M\|}$ varying M .

14
 15 We now study whether the discrete operator preserves the spectral properties
 16 of the continuous operator. The spectrum of the operator $\mathcal{A}_0^{\circ*}$ contains only the
 17 eigenvalue $\lambda = 0$, and every non-null constant function is an eigenfunction. The
 18 spectrum of $A_{0,M}$ contains, in addition to the eigenvalue $\lambda = 0$, also the eigenvalues
 19 of the matrix D_M : the pseudospectral approximation introduces M eigenvalues
 20 which are “spurious”, i.e., due only to the discretization procedure. Therefore it is
 21 important to study the behavior of the spectrum of D_M when $M \rightarrow \infty$.

22 The eigenvalues of D_M are not explicitly known, but many theoretical and em-
 23 pirical results have been derived by various authors. In fact their spectral properties
 24 are relevant in the convergence analysis of the pseudospectral methods, otherwise
 25 known as spectral collocation, for first order hyperbolic PDEs.

26 So the first fundamental results have been established within this context [26,
 27 35, 33, 34, 40, 58]. Various further properties can be found also in [30, 29, 31,
 28 18, 45, 54, 63, 64]. Both the derivative operator in Hilbert space and the matrix
 29 D_M are non-normal, and we recall the illuminating book [62], where the fundamen-
 30 tal properties of non-normal operators in Hilbert spaces and matrices have been
 31 investigated through the analysis of their pseudospectra.

32 The following proposition allows us to analyze the position in the complex plane
 33 of the eigenvalues of D_M .

34 **Proposition 2.** [58, Appendix B] *For every $M \geq 1$, the eigenvalues of the matrix*
 35 *D_M have negative real part.*

Concerning the asymptotic behavior of the spectral abscissa $\alpha(D_M)$, which is defined as

$$\alpha(D_M) := \max\{\operatorname{Re}(\lambda) \mid \lambda \in \sigma(D_M)\},$$

1 where $\sigma(D_M)$ denotes the spectrum of D_M , the first result goes back to Dubiner
 2 [26], and was revised recently by Wang & Waleffe [64].

3 **Proposition 3.** [26, 64] $\lim_{M \rightarrow \infty} \alpha(D_M) = -\infty$. Moreover $\alpha(D_M) = O(\log(M))$, for
 4 sufficiently large M .

5 Therefore, the eigenvalues of D_M “disappear” as $M \rightarrow \infty$, consistent with the
 6 fact that the operator $\mathcal{A}_0^{\odot*}$ contains the eigenvalue $\lambda = 0$ only. Indeed, for un-
 7 bounded operators the concept of *extended spectrum* can be defined [25, pag. 479].
 8 It consists of the spectrum of the operator and the point ∞ . So, to some extent, we
 9 can say that the spectrum of $A_{0,M}$ “converges” to the extended spectrum of $\mathcal{A}_0^{\odot*}$
 10 as $M \rightarrow \infty$.

11 We refer to Figure 1 for a representation of the spectrum of D_M for different val-
 12 ues of M (left) and for a plot of $-\alpha(D_M)/\log(M)$ versus M (right). As remarked in
 13 [62, 63], the computation of the spectrum of pseudospectral differentiation matrices
 14 is extremely sensitive to rounding errors. In particular, the authors observed that
 15 eigenvalues $\lambda \in \mathbb{C}$ with $\operatorname{Re}\lambda < \log \epsilon$ are affected by rounding errors because the
 16 precision in the corresponding eigenvectors is lost. This phenomenon is observable
 17 in Figure 1 (left): for $M = 64$, a part of the computed spectrum falls to the left of
 18 the line $\operatorname{Re}\lambda = \log \epsilon$ and we expect such results to be affected by rounding errors.
 19 We note that the rounding errors affect mostly the eigenvalues which are large in
 20 modulus, while the computation of the $\alpha(D_M)$ seems to be more stable, (see Figure
 21 1). The right panel of Figure 1 shows the oscillatory convergence of $\frac{-\alpha(D_M)}{\log(M)}$.




FIGURE 1. Left: plot of $\sigma(D_M)$ for different M , together with the line $\operatorname{Re}\lambda = \log \epsilon \approx -36.0437$ connected with the instability phenomenon studied in [63]. Right: plot of $\frac{-\alpha(D_M)}{\log(M)}$, versus M .

22 We now focus our attention on the resolvent operators and their approximations.
 23 For any $\lambda \in \mathbb{C} \setminus \{0\}$, the resolvent of $\mathcal{A}_0^{\odot*}$ exists and it is the bounded operator

1 given by

$$\begin{aligned} (\lambda I - \mathcal{A}_0^{\odot*})^{-1}(\beta, \varphi) &= \left(\frac{\beta}{\lambda}, \frac{\beta}{\lambda} e^{\lambda \cdot} - \int_0^{\cdot} e^{\lambda(\cdot-s)} \varphi(s) ds\right), \\ &= \frac{\beta}{\lambda}(1, e^{\lambda \cdot}) + (0, - \int_0^{\cdot} e^{\lambda(\cdot-s)} \varphi(s) ds), \end{aligned} \quad (25)$$

2 for $(\beta, \varphi) \in Y^{\odot*}$.

Let $(\alpha, \psi) = (\lambda I - \mathcal{A}_0^{\odot*})^{-1}(\beta, \varphi)$. From (25) we obtain that

$$\begin{aligned} |\psi(\theta)| &\leq |\beta| \frac{e^{\theta \operatorname{Re} \lambda}}{|\lambda|} + \|\varphi\| \frac{1 - e^{\theta \operatorname{Re} \lambda}}{\operatorname{Re} \lambda} \\ &\leq |\beta| \frac{e^{\theta \operatorname{Re} \lambda}}{|\operatorname{Re} \lambda|} + \|\varphi\| \frac{1 - e^{\theta \operatorname{Re} \lambda}}{\operatorname{Re} \lambda} \\ &\leq \begin{cases} \frac{1}{\operatorname{Re} \lambda} \|(\beta, \varphi)\|, & \operatorname{Re} \lambda > 0 \\ \frac{e^{-\tau \operatorname{Re} \lambda}}{|\operatorname{Re} \lambda|} \|(\beta, \varphi)\|, & \operatorname{Re} \lambda < 0, \end{cases} \end{aligned}$$

for all $\theta \in [-\tau, 0]$, which leads to the following bound of the resolvent norm

$$\|(\lambda I - \mathcal{A}_0^{\odot*})^{-1}\| \leq \begin{cases} \frac{1}{\operatorname{Re} \lambda}, & \operatorname{Re} \lambda > 0 \\ \frac{e^{-\tau \operatorname{Re} \lambda}}{|\operatorname{Re} \lambda|}, & \operatorname{Re} \lambda < 0 \end{cases}.$$

Note that the upper bound depends on $\operatorname{Re} \lambda$ only, and, though it is finite for every $\lambda \in \mathbb{C} \setminus \{0\}$, it increases exponentially when $\operatorname{Re} \lambda \rightarrow -\infty$. This phenomenon has been already documented w.r.t. L^2 -norm in [62]. When $\beta = 0$, we get

$$(\lambda I - \mathcal{A}_0^{\odot*})^{-1}(0, \varphi) = (0, \int_0^{\cdot} e^{\lambda(\cdot-s)} \varphi(s) ds), \quad \varphi \in L^\infty,$$

3 and the second component is also defined for $\lambda = 0$ through the integral operator

4 $\mathcal{V}: L^\infty \rightarrow L^\infty$ given by

$$(\mathcal{V}\varphi)(\theta) := \int_0^\theta \varphi(s) ds, \quad \theta \in [-\tau, 0], \quad \varphi \in L^\infty. \quad (26)$$

5 What can we say about the resolvent operator of the discrete operator $A_{0,M}$?

6 Given $\lambda \neq 0$, Proposition 3 states that there exists $M_0 := M_0(\lambda)$ such that $\lambda \notin$

7 $\sigma(D_M)$ for all $M \geq M_0$. Therefore for all $M \geq M_0$ the resolvent operator of $A_{0,M}$

8 is defined by the matrix

$$(\lambda I - A_{0,M})^{-1} = \begin{pmatrix} \frac{1}{\lambda} & 0 \\ -\frac{1}{\lambda}(\lambda I - D_M)^{-1} D_M \mathbf{1} & (\lambda I - D_M)^{-1} \end{pmatrix}. \quad (27)$$

9 Let $\lambda \in \mathbb{C} \setminus \{0\}$ and $(\beta, \varphi) \in \widehat{Y}$. The function

$$\psi(\theta) := \psi(\theta; \lambda, \beta, \varphi) = j^{-1}(\lambda I - \mathcal{A}_0^{\odot*})^{-1}(\beta, \varphi), \quad \theta \in [-\tau, 0], \quad (28)$$

10 is the solution of the following problem

$$\begin{cases} \psi'(\theta) = \lambda \psi(\theta) - \varphi(\theta), & \theta \in [-\tau, 0], \\ \psi(0) = \frac{\beta}{\lambda}. \end{cases} \quad (29)$$

1 We denote by $\psi_M(\theta) := \psi_M(\theta; \lambda, \beta, \varphi)$, $\theta \in [-\tau, 0]$, the M -degree collocation poly-
 2 nomial for the problem (29) such that

$$\begin{cases} \psi'_M(\theta) = \lambda\psi_M(\theta) - \varphi(\theta), & \theta \in \Theta_M, \\ \psi_M(0) = \frac{\beta}{\lambda}. \end{cases} \quad (30)$$

3 We state the following theorem.

4 **Theorem 4.1.** *Let $\lambda \in \mathbb{C} \setminus \{0\}$ and $(\beta, \varphi) \in \widehat{Y}$. There exists $M_0 := M_0(\lambda)$ such
 5 that, for all $M \geq M_0$, the collocation equations (30) for the problem (29) uniquely
 6 define the collocation polynomial ψ_M and*

$$\|\psi_M - \psi\| \leq 2 \left(\frac{1 - e^{-\operatorname{Re}\lambda\tau}}{\operatorname{Re}\lambda} \right) \|r_M\|, \quad (31)$$

7 where

$$r_M := \lambda(\mathcal{L}_{M-1}\psi - \psi) + (\varphi - \mathcal{L}_{M-1}\varphi). \quad (32)$$

8 and \mathcal{L}_{M-1} is the interpolation operator relative to the nodes Θ_M in (12). Moreover,
 9 for all $M \geq M_0$, ψ_M admits the following representation

$$\psi_M = P_M(\lambda I - A_{0,M})^{-1} \widehat{R}_M(\beta, \varphi), \quad (33)$$

10 which is the discrete counterpart of (28).

11 *Proof.* The proof is based on the approach of [14, Proposition 5.1]. By using the
 12 integral operator (26), we can rewrite both (29) and (30) as integral equations in
 13 the space Y to obtain respectively

$$\psi = \frac{\beta}{\lambda} + \lambda\mathcal{V}\psi - \mathcal{V}\varphi,$$

14 and

$$\psi_M = \frac{\beta}{\lambda} + \lambda\mathcal{V}\mathcal{L}_{M-1}\psi_M - \mathcal{V}\mathcal{L}_{M-1}\varphi.$$

15 By introducing the error $e_M := \psi_M - \psi$, we obtain the following equation for e_M :

$$(I - \lambda\mathcal{V}\mathcal{L}_{M-1})e_M = \mathcal{V}r_M, \quad (34)$$

16 where r_M is defined in (32). We have that e_M is a solution of (34) if and only if
 17 $e_M = \mathcal{V}z_M$ with $z_M \in Y$ solution of the equation

$$(I - \lambda\mathcal{L}_{M-1}\mathcal{V})z_M = r_M. \quad (35)$$

18 By observing that the operator $I - \lambda\mathcal{V}$ is invertible with inverse

$$\left[(I - \lambda\mathcal{V})^{-1} f \right] (\theta) = f(\theta) + \lambda \int_0^\theta e^{\lambda(\theta-s)} f(s) ds, \quad f \in Y, \theta \in [-\tau, 0], \quad (36)$$

19 and that $\|\lambda(\mathcal{L}_{M-1} - I)\mathcal{V}\| \rightarrow 0$, as $M \rightarrow +\infty$ (see e.g. [51, pag. 269]), we can
 20 apply the Banach Perturbation Lemma (see e.g. [46, Theorem 10.1]) to conclude
 21 that there exists $M_0 := M_0(\lambda)$ such that for all $M \geq M_0$,

$$\|\lambda(\mathcal{L}_{M-1} - I)\mathcal{V}\| \cdot \|(I - \lambda\mathcal{V})^{-1}\| \leq \frac{1}{2}, \quad (37)$$

22 and the operator $I - \lambda\mathcal{L}_{M-1}\mathcal{V}$ is invertible. Therefore, for all $M \geq M_0$, the equation
 23 (35) has a unique solution z_M and, as a consequence, $e_M = \mathcal{V}(I - \lambda\mathcal{L}_{M-1}\mathcal{V})^{-1} r_M$

1 is the unique solution of (34). Moreover, for all $M \geq M_0$, from (36) and (37) we
 2 easily get the bound

$$\|\mathcal{V}(I - \lambda\mathcal{L}_{M-1}\mathcal{V})^{-1}\| \leq 2\|\mathcal{V}(I - \lambda\mathcal{V})^{-1}\| \leq 2\left(\frac{1 - e^{-\operatorname{Re}\lambda\tau}}{\operatorname{Re}\lambda}\right),$$

3 and (31) follows.

4 Finally, by using the Lagrange representation (15) of ψ_M and the matrix (19), we
 5 can easily obtain from the collocation equation (30) that ψ_M is given by (33). \square

6 The bound (31) states that the error depends on the interpolation error of both,
 7 ψ and φ , and so we need to assume more regularity on φ to obtain convergence. It
 8 is sufficient to consider $j\varphi \in D(\mathcal{A}_0^{\circ*})$. Moreover when φ is an analytic function, we
 9 get the so-called *spectral accuracy*, i.e. the error $\|\psi_M - \psi\|$ decays as $O(\rho^{-M})$ for
 10 some $\rho > 1$ [60, Chapter 8]. In Figure 2 we plot the error $\|\psi_M - \psi\|$ when $\beta = 0$
 11 and $\varphi = 1$. In this case $\psi = \frac{1 - e^{-\lambda x}}{\lambda}$ is analytic and $\mathcal{L}_{M-1}\mathbf{1} = \mathbf{1}$. Notice the spectral
 12 accuracy and that the convergence is slower when λ has larger modulus.




FIGURE 2. Error $\|\psi_M - \psi\|$ for the resolvent operators applied to $\beta = 0$, $\varphi = 1$, versus M . Left: $\lambda = 1$; right: $\lambda = 10$. Note the spectral accuracy and that the convergence is slower when λ has larger modulus.

Finally for $\beta = 0$ and $\lambda = 0$, we get that

$$\|\mathcal{V}\varphi - P_M(0, D_M^{-1}\varphi(\Theta_M))\| \leq \tau\|\mathcal{L}_{M-1}\varphi - \varphi\|, \quad \varphi \in Y.$$

13 The integral operator \mathcal{V} has norm equal to τ , and consistently with this result, we
 14 have the analogous property for the matrix D_M^{-1} .

15 **Proposition 4.** [57] *For any $M \geq 1$, D_M is non-singular and $\|D_M^{-1}\| = \tau$.*

16 **5. The solution operators $\mathcal{T}_0^{\circ*}(t)$ and $T_{0,M}(t)$: results and conjectures.**

17 In this section, we focus our attention on the connection between $\mathcal{T}_0^{\circ*}$ and $T_{0,M}$,
 18 when $M \rightarrow \infty$, which is a crucial question for the convergence analysis. Indeed

1 this is motivated by the integral equations (8) and (22), which can be reformulated
 2 respectively as

$$jv(t) = \mathcal{T}_0^{\odot*}(t)j\psi + \int_0^t \mathcal{T}_0^{\odot*}(t-s)(z(s), 0) ds, \quad t \geq 0, \quad (38)$$

3 and

$$v_M(t) = T_{0,M}(t)(\alpha, \Psi) + \int_0^t T_{0,M}(t-s)(z(s), 0) ds, \quad t \geq 0, \quad (39)$$

4 where $z(t)$ is $G(v(t))$ in (38) and $G(P_M v_M(t))$ in (39).

5 We first introduce the operator $\mathcal{H}_0(t) : L^\infty \rightarrow L^\infty$ defined as

$$\mathcal{H}_0(t)\psi = \psi_t^0, \quad t \geq 0, \quad (40)$$

and we reformulate the action of the $\odot*$ -shift operator as follows

$$\begin{aligned} \mathcal{T}_0^{\odot*}(t)(\alpha, \psi) &= (\alpha, (I - \mathcal{H}_0(t))\alpha) + (0, \mathcal{H}_0(t)\psi) \\ &= (\alpha, \alpha) + (0, \mathcal{H}_0(t)(\psi - \alpha)) \end{aligned}, \quad t \geq 0, (\alpha, \psi) \in Y^{\odot*}.$$

As usual α denotes either a scalar or the constant function. Similarly, by using (23) we can rewrite the discrete counterpart as

$$\begin{aligned} T_{0,M}(t)(\alpha, \Psi) &= (\alpha, (I - e^{D_M t})\mathbf{1}\alpha) + (0, e^{D_M t}\Psi) \\ &= (\alpha, \mathbf{1}\alpha) + (0, e^{D_M t}(\Psi - \mathbf{1}\alpha)) \end{aligned}, \quad t \geq 0, (\alpha, \Psi) \in \widehat{Y}_M,$$

6 which highlights that $e^{D_M t}$ plays the role of $\mathcal{H}_0(t)$ for all $t \geq 0$. According to these
 7 reformulations and without loss of generality, we can assume that $\alpha = 0$ and focus
 8 on the behavior of $e^{D_M t}$ w.r.t. M for $t \geq 0$.

9 As already pointed out in Section 4, many results about D_M and $e^{D_M t}$ have been
 10 derived in the applications of the spectral collocation methods to hyperbolic PDEs.
 11 In fact, within this context, the *space stability*, i.e. the norm boundedness of $e^{D_M t}$
 12 w.r.t. M , plays a key role in the convergence analysis [18, 28, 40]. PDEs are usually
 13 studied in suitable Hilbert spaces and so the results mainly regard weighted 2-norms
 14 in Y_M , which correspond to the discretization of L^2 norms (see [31, 33, 36, 45, 58]).
 15 Hyperbolic PDEs are strongly related to DDEs, but the latter ones require to work
 16 in infinite-dimensional Banach spaces endowed with supremum norm and, to our
 17 knowledge, no bounds are available in this case.

18 To address the question and have a first insight, we have performed some nu-
 19 merical simulations. The computation of the exponential matrices, especially non-
 20 normal, can be difficult and strongly affected by rounding errors, and the research
 21 on this subject is still in progress (see for instance [42] and the references therein).
 22 Here we use the built-in Matlab function `expm`, which is based on the algorithms
 23 in [1, 41].

24 The semigroup \mathcal{H}_0 in (40) is contractive, i.e. $\|\mathcal{H}_0(t)\| \leq 1, t \in [0, \tau]$ and nilpotent,
 25 i.e. $\mathcal{H}_0(t) = 0, t \geq \tau$. What can we say about the discrete counterpart $e^{D_M t}$?

Propositions 2 and 3 allow us to derive the *time stability* of $e^{D_M t}$, i.e.

$$\|e^{D_M t}\| \rightarrow 0, \quad \text{as } t \rightarrow \infty,$$

uniformly in M . But since the matrix D_M is non-normal, the eigenvalues do not
 tell us the whole story. In fact the behavior of $\|e^{D_M t}\|$ w.r.t. t is different in the
 initial, transient and asymptotic phases. The spectral abscissa $\alpha(D_M)$ allows one

to describe the asymptotic behavior of $\|e^{D_M t}\|$, whereas the initial growth rate is determined by the *logarithmic norm* of D_M , defined by

$$\mu(D_M) = \lim_{t \rightarrow 0^+} \frac{\|I + tD_M\| - 1}{t}$$

(see [38, 62]). The logarithmic norm admits the useful reformulation

$$\mu(D_M) = \lim_{t \rightarrow 0^+} \frac{\log \|e^{D_M t}\|}{t},$$

- 1 and the explicit expression $\mu(D_M) = \sup_i (\operatorname{Re}(d_{ii}) + \sum_{j:j \neq i} |d_{ij}|)$. In Table 2 we com-
- 2 pute $\mu(D_M)$ varying M to experimentally investigate the order of convergence w.r.t.
- 3 M . The results show that $\mu(D_M) \approx \mu M^2$, as $M \rightarrow \infty$, with $\mu \approx 0.6$.

M	$\mu(D_M)$	order	$\mu(D_M)/M^2$
4	6.6569e+00		4.1605e-01
8	3.8921e+01	2.5476	6.0814e-01
16	1.6698e+02	2.1011	6.5227e-01
32	6.7900e+02	2.0237	6.6308e-01
64	2.7270e+03	2.0058	6.6577e-01
128	1.0919e+04	2.0015	6.6644e-01
256	4.3687e+04	2.0004	6.6661e-01

TABLE 2. $\mu(D_M)$ and estimation of the order $\log_2 \frac{\mu(D_{2M})}{\mu(D_M)}$ varying M .

4 Here our main interest is in the behavior of $\|e^{D_M t}\|$ w.r.t. M in the transient
 5 phase, which has no connection with the eigenvalues and looks quite different from
 6 the asymptotic behavior. Indeed the left panel of Figure 3 shows that, in the first
 7 interval $[0, 1]$, $\|e^{D_M t}\|$ increases as M increases.

8 The results in Figure 3 give also evidence that there exists \hat{t}_M where $\|e^{D_M t}\|$
 9 changes its behavior from not convergent to convergent, and that the convergence
 10 is attained for $t \geq \hat{t}_M$. The relevant fact emerging from the simulations is that
 11 $\hat{t}_M \in (\tau, 1.5\tau)$ for sufficiently large M . In truth the upper bound is quite safe.

12 Since the norm of the interpolation operator for Chebyshev nodes gives $\Lambda_M =$
 13 $O(\log(M))$, the divergence behavior of $\|e^{D_M t}\|$ w.r.t. M in the first interval $[0, \tau]$
 14 is not totally surprising and it is quite natural to wonder if there is a connection
 15 between the two phenomena. Indeed, even if Faber's Theorem states that Cheby-
 16 shev interpolation points could not ensure the convergence for all functions, quoting
 17 Trefethen [61], the "*polynomial interpolation in Chebyshev points is a powerful and*
 18 *reliable method for approximation of functions*", since "*for Lipschitz continuous*
 19 *functions or better, as is easily done in almost any application, Faber's Theorem*
 20 *ceases to be applicable*". By extending this remark to $\|e^{D_M t}\|$, we conjecture that its
 21 divergence w.r.t. M in the transient phase is due to a "non-generic" set of vectors
 22 Ψ , which are constructed in a particularly nasty way. The left panel of Figure 4
 23 makes evident that also random data do not capture the "bad" data, whereas the
 24 right panel of Figure 4 suggests that $\|e^{D_M t}\|$ diverges as $\log(M)$.

25 Having in mind Proposition 3 and the simulations in Figure 3 & Figure 4, we
 26 come to the following conjecture: there exist $\hat{\tau} \in (\tau, 1.5\tau)$ and positive constants




FIGURE 3. Left: plot of $\log \|e^{D_M t}\|$ for $t \in [0, 2]$ with different values of M . Notice divergence in $[0, 1]$ for increasing M . Right: plot of $\max_{t \in I} \frac{\log \|e^{D_M t}\|}{\log(M)}$ versus M , for I specified in the legend. Notice that the convergence gains one order at every time interval.




FIGURE 4. $\|e^{D_M t}\|$ computed with different routines: built-in Matlab function `expm` and `norm(., inf)` versus maximum over $n = 100$ random initial vectors of the solution of the ODE system. Left: $\|e^{D_M t}\|$ versus time. Right: $\max_{t \in [0, 2]} \|e^{D_M t}\|$ versus M . The dotted line is the reference line $\log(M)$. The fact that the effective norm of the exponential matrix diverges, while the norm computed by selecting a random set of vectors is uniformly bounded, suggests that the “bad” behavior of the norm is due to a small set of vectors.

- 1 a, K independent of M such that

$$\begin{aligned} \|e^{D_M t}\| &= O(\log(M)), \quad t \in [0, \hat{\tau}], \\ \|e^{D_M t}\| &\leq KM^{-at}, \quad t > \hat{\tau}. \end{aligned} \tag{41}$$

- 2 for $M \rightarrow \infty$.

1 But in the first term of the right-hand side of (39), the vector Ψ derives from
 2 the discretization of a function ψ , namely $\Psi = \psi(\Theta_M)$, and ψ is at least Lipschitz
 3 continuous with $\psi(0) = 0$. May we expect to attain the space stability or, even
 4 better, its convergence w.r.t. M in the transient phase? In case of a positive answer,
 5 is the asymptotic convergence rate w.r.t. M faster for smoother functions? In other
 6 words, we wonder if $\|e^{D_M t} R_M \psi\|$ retains memory of the interpolants. Indeed we
 7 have that

$$e^{D_M t} R_M \psi = \sum_{k \geq 0} \frac{R_M (D P_M^0 R_M t)^k \psi}{k!}, \quad t \geq 0, \quad (42)$$

where \mathcal{D} is defined in (24) and

$$P_M^0 \Psi := P_M(0, \Psi), \quad \Psi \in Y_M.$$

8 The formula above makes clear the role of interpolation through $P_M^0 R_M$, and that
 9 the action of every term corresponds to interpolation with null value at $\theta = 0$ and
 10 derivation, imposing, to some extent, that the value at $\theta = 0$ of next derivatives is
 11 zero.

12 So let us get another actor on stage to give us some suggestions, namely the
 13 operator $H_{0,M}(t) : Y_M \rightarrow Y_M$ defined as

$$H_{0,M}(t) \Psi = R_M \mathcal{H}_0(t) P_M^0 \Psi, \quad \Psi \in Y_M, t \geq 0. \quad (43)$$

14 From (15), (43) admits the following representation

$$H_{0,M}(t) \Psi = R_M \sum_{i=1}^M \mathcal{H}_0(t) \ell_i \Psi_i, \quad t \geq 0. \quad (44)$$

15 Notice that, contrary to $\{e^{D_M t}\}_{t \geq 0}$, the family $\{H_{0,M}(t)\}_{t \geq 0}$ does not define a
 16 semigroup on Y_M . Moreover $H_{0,M}(t) = 0$ for $t \geq \tau$, while $e^{D_M t}$ is not. From (44)
 17 we easily get the following bound

$$\|H_{0,M}(t)\| \leq C_M(t), \quad t \in [0, \tau],$$

18 where

$$C_M(t) := \max_{j: t+\theta_j \leq 0} \sum_{i=1}^M |\ell_i(t + \theta_j)|, \quad t \in [0, \tau]. \quad (45)$$

19 and $\sup_{t \geq 0} C_M(t) \leq (1 + \frac{2}{\pi} \log(M+1))$.

20 In the right panel of Figure 5, we notice that the behavior of $\|e^{D_M t}\|$ resem-
 21 bles the behavior of the function $C_M(t)$, and that, moreover, $\sup_{t \geq 0} C_M(t)$ "domi-
 22 nates" $\sup_{t \geq 0} \|e^{D_M t}\|$. So, encouraged by this experiment, we also conjecture that
 23 $H_{0,M}(t) \Psi$ can predict, to some extent, the behavior of $\|e^{D_M t} \Psi\|$ in the transient
 24 phase. To support this we first analyze $H_{0,M}(t) \Psi$ for $\Psi = \psi(\Theta_M)$ by using well-
 25 known interpolation results. Second we experimentally explore the behavior of
 26 $\|e^{D_M t} \psi(\Theta_M)\|$ by selecting some test functions ψ with different regularity prop-
 27 erties.

Suppose that ψ and its derivatives through $\psi^{(k-1)}$ are absolutely continuous on $[-\tau, 0]$ and that $\psi^{(k)}$ has bounded variation $V(\psi^{(k)})$, for $k \geq 1$. Then for any $M > k$ we have

$$\begin{aligned} \|H_{0,M}(t) R_M \psi - R_M \mathcal{H}_0(t) \psi\| &\leq \|R_M \mathcal{H}_0(t) (\psi - P_M^0 R_M \psi)\| \\ &\leq \frac{4(\tau-t)V(\psi^{(k)})}{\pi k(M-k)^k}, \quad t \in [0, \tau], \end{aligned}$$




FIGURE 5. Left: plot of the function $\sum_{i=1}^M |\ell_i(\theta)|$ for $M = 8$. Right: plot of $\|e^{D_M t}\|$ (blue) and $C_M(t)$ in (45) (red) versus time for $M = 8$.

- 1 which implies the convergence at algebraic rate $O(M^{-k})$ [60, Chapter 7]. A similar
 2 result holds under the condition that $\psi^{(k)}$ is continuous. When ψ is an analytic
 3 function, the interpolants converge spectrally [60, Chapter 8]. Finally, regarding
 4 the following function

$$H(\theta) = \begin{cases} 1, & -\tau \leq \theta < 0, \\ 0, & \theta = 0. \end{cases} \quad (46)$$

the interpolation process does not converge, but the interpolation error is uniformly bounded in supremum norm, i.e.,

$$\lim_{M \rightarrow \infty} \|P_M^0 \mathbf{1}\| = c_1,$$

for a suitable constant c_1 [60, Chapter 9], and therefore we can conclude that

$$\|P_M^0 H_{0,M}(t) \mathbf{1} - \mathcal{H}_0(t) H\| = O(1), \quad M \rightarrow \infty, \quad t \in [0, \tau].$$

- 5 This effect is due to the well-known Gibbs phenomenon for step functions: the in-
 6 terpolants always “overshoot” their target and, as M increases, the overshooting
 7 does not get lower, although the region of overshooting gets narrower. This phe-
 8 nomenon prevents the norm-convergence of the interpolants and is responsible for
 9 the uniform bound. Although we will not go further into details here, we note that
 10 the Gibbs phenomenon still ensures the pointwise convergence of the interpolants
 11 of H for all $\theta \in [-\tau, 0]$.

- 12 For our simulations we choose some functions ψ with different regularity prop-
 13 erties and we estimate the behavior of the error $\|e^{D_M t} \Psi - R_M \mathcal{H}_0(t) \Psi\|$ with $\Psi =$
 14 $\psi(\Theta_M)$ at some selected points t . All the results in the right panels of Figures 7,
 15 8, 9 & 10 indicate that the convergence occurs in the first interval ($t = 0.5$) with
 16 the rate predicted by the interpolation results for the respective functions (see their
 17 plots in the left panels), whereas the order increases in the next intervals ($t > 1$).
 18 Moreover Figure 10 presents the spectral convergence behavior. Finally Figure 6,

- 1 Figure 11 & Figure 12 show there is no convergence in first interval, in accordance with the interpolation results, but the uniform boundedness holds in all cases.




FIGURE 6. Left: plot of $\psi(\theta) = H(\theta)$ (black) and its interpolating polynomial for $M = 8$ (red). Right: log log plot of the error $\|e^{D_M t} \psi(\Theta_M) - R_M \mathcal{H}_0(t) \psi\|$ for $t = 0.5, 1.5, 2.5$ versus M . Note the uniform bound for $t = 0.5$ and the convergence for $t > 1$. The dotted lines are the reference lines M^{-k} , $k = 1, \dots, 4$.

2




FIGURE 7. Same as Figure 6 for $\psi(\theta) = 0.5 - |\theta + 0.5|$. Note that $\psi \in Y$ and $\psi^{(1)}$ has bounded variation. The convergence rate for $t = 0.25, 0.5$ is $O(M^{-1})$ (right).

- 3 The numerical experiments indicate that, in the transient phase, $\|e^{D_M t} \Psi\| / \|\psi\|$
 4 is uniformly bounded w.r.t. M when applied to vectors $\Psi = \psi(\Theta_M)$ deriving from
 5 the discretization of functions ψ , for which the interpolation error is $O(M^{-k})$ with
 6 $k \geq 0$, and it is converging for sufficiently smooth ψ . This supports the effectiveness
 7 of the PSD in our context and in particular for the analysis of periodic solutions.




FIGURE 8. Same as Figure 6 for $\psi(\theta) = e^{\theta^2} - 1$. Note that the interpolating polynomial is indistinguishable from the function (left) and that $\psi^{(2)}$ has bounded variation. The convergence rate for $t = 0.5$ is $O(M^{-2})$ (right).




FIGURE 9. Same as Figure 6 for $\psi(\theta) = -\theta^3$. Note that the interpolating polynomial is indistinguishable from the function (left) and that $\psi^{(3)}$ has bounded variation. The convergence rate for $t = 0.5$ is $O(M^{-3})$ (right).

- 1 Now we consider the second term of both (38) and (39). Let $z : \mathbb{R}^+ \rightarrow \mathbb{R}$ be a
- 2 continuous function. Then the function $w : \mathbb{R}^+ \rightarrow Y^{\odot\odot}$, defined as

$$w(t) = j^{-1} \int_0^t \mathcal{T}_0^{\odot*}(t-s)(z(s), 0) ds, \quad t \geq 0, \quad (47)$$




FIGURE 10. Same as Figure 6 for $\psi(\theta) = e^{-1/\theta^2}$. Note that the interpolating polynomial is indistinguishable from the function (left) and the spectral convergence for $t = 0.5, 1.5, 2.5$ (right).




FIGURE 11. Same as Figure 6 for $\psi(\theta) = \sin \frac{1}{\theta}$, if $\theta < 0$, $\psi(0) = 0$. Note there is no convergence for $t = 0.5$.

is continuous [25, Lemmas III.2.3] and

$$\begin{cases} w(t)(0) = \int_0^t z(s) ds \\ w(t)(\theta) = \begin{cases} \int_0^{t+\theta} z(s) ds, & t+\theta \geq 0 \\ 0, & t+\theta < 0 \end{cases} \end{cases}$$

- 1 for all $t \geq 0$ [25, III.4.3]. By applying the convergence results to the function (47)
- 2 [60, Thorem 7.2], we have that

$$\|w(t) - P_M(w(t)(0), w(t)(\Theta_M))\| \leq \frac{4 \max_{s \in [0, t]} |z(s)|}{\pi(M-1)}, \quad t \in [0, \tau].$$




FIGURE 12. Same as Figure 6 for $\psi(\theta) = \sin(6\theta) + \text{sign}(\sin(\theta + e^{2\theta}))$ [60, pag.10]. Note there is no convergence for $t = 0.5$.

1 So, to guarantee the convergence of

$$\|w(t) - P_M \left(\int_0^t T_{0,M}(t-s)(z(s), 0) \right)\|, \quad t \in [0, \tau],$$

- 2 we conjecture that $\|\int_0^t e^{(t-s)D_M} \mathbf{1}z(s) ds\|$ inherits the convergence behavior in $[0, \tau]$.
 3 To this aim we experimentally estimate the error $\|\int_0^t e^{D_M(t-s)} \mathbf{1}z(s) ds - w(t)(\Theta_M)\|$
 4 at some selected values of t for two test continuous functions z varying M . Figures
 5 13 & 14 illustrate that the behavior of the errors is as expected.




FIGURE 13. Left: plot of the function z (saw function) Right: error $\|\int_0^t e^{(t-s)D_M} z(s) ds - w(t)(\Theta_M)\|$ for $t = 0.5, 1, 1.5$ versus M . The dotted line is the reference line $\log(M)/M$.

6 **6. Conclusions, ongoing and future research.** In this paper we have focused
 7 on the shift semigroup in the \odot -* formulation of the DDE (2), and its PSD of index




FIGURE 14. Same as Figure 13 for the function z built so that it is continuously differentiable with jumps in the second derivative at $t = 0.5, 1, 1.5$.

1 M . The generator of the discrete semigroup is a finite dimensional operator that
 2 can be described in terms of a matrix D_M , which has been widely studied in the
 3 context of PDEs.

4 In Section 4 we have collected the most relevant results, especially about the
 5 spectrum of the matrix D_M , of the generator $A_{0,M}$, and about their relation with
 6 the continuous counterpart $\mathcal{A}_0^{\odot*}$. We also specified the convergence of the resolvent
 7 operators by adapting the proof in [11, 14].

8 About the convergence of the discrete semigroup $T_{0,M}(t)$ to the continuous semi-
 9 group $\mathcal{T}_0^{\odot*}(t)$, very few results are available in the supremum norm, which is the
 10 norm of the natural state space of DDEs. In the absence of theoretical results, we
 11 have presented a series of numerical tests and compiled some conjectures about the
 12 behavior of the semigroup, both in time and as $M \rightarrow \infty$.

13 To arrive to a rigorous proof, we could start from the representation (42) and
 14 consider the properties of the Chebyshev polynomials [55]. Another possibility for
 15 proving convergence of the discrete semigroup $T_{0,M}(t)$ to the continuous semigroup
 16 $\mathcal{T}_0(t)$ is to use the convergence results of Theorem 4.1 and to recall that the resolvent
 17 operator can be expressed, for $\operatorname{Re}\lambda > 0$, in terms of the solution operator via Laplace
 18 transform as

$$(\lambda I - \mathcal{A}_0^{\odot*})^{-1}(\alpha, \psi) = \int_0^\infty e^{-\lambda s} \mathcal{T}_0^{\odot*}(s)(\alpha, \psi) ds, \quad \lambda \notin \sigma(\mathcal{A}_0^{\odot*}),$$

19 We are currently working in this direction.

20 The study of the discrete and continuous shift semigroups is motivated by the
 21 PSD of DDEs in $\odot*$ -formulation, which is introduced in this manuscript for the
 22 first time. The $\odot*$ -formalism, despite introducing some “complications” due to
 23 the embedding into larger spaces, appears to be useful in the context of numeri-
 24 cal methods for several reasons. The main advantage is to provide us with the
 25 variation-of-constants formula (8), which is particularly useful to relate the nonlin-
 26 ear solution semigroup to the trivial linear semigroup $\mathcal{T}_0^{\odot*}(t)$. This is particularly
 27 useful in the prospect of comparing the discrete and continuous nonlinear semi-
 28 groups. The convergence of the nonlinear semigroups is the next step for the final

1 goal of understanding in which sense the PSD approximates the periodic solutions
 2 of (2) and their stability. This is ongoing research of the authors and collaborators
 3 announced in the paper [8].

4 The \odot^* framework exploits naturally the pairing between dual spaces and there-
 5 fore comes with two topologies, the norm topology and the weak* topology. So even
 6 if approximations do not necessarily converge with respect to the norm, they still
 7 may converge in a weaker sense which, probably, is sufficient for our purposes. We
 8 intend to investigate this idea in the near future.

9 In the case of DDEs, the \odot^* -formulation allowed us to give a direct description
 10 of the PSD of the operators by means of projections into the discretization space
 11 and injections into the bigger space. Moreover, it provides us with a variation-of-
 12 constants formula, which is helpful to relate the nonlinear semigroup to the trivial
 13 shift semigroup. We mention also that the \odot^* -framework is particularly important
 14 also in the PSD of REs and REs/DDEs, where the domain of the infinitesimal
 15 generator is described by means of an algebraic, rather than differential, equation.
 16 In this context, the \odot^* -formulation may allow to overcome the inversion of the
 17 nonlinear algebraic equation proposed in [8], treating the condition more efficiently.
 18 This is currently under investigation by the authors.

19 Finally, the PSD approach has been proposed also for DDEs, REs, and REs/DDEs
 20 with infinite delay in [37]. Moreover, the analogous theory of \odot^* -calculus for infinite
 21 delay has been developed in [24]. Similarly to the case with finite delay, the PSD
 22 of the shift operator in this context is described by a matrix derived by Laguerre
 23 differentiation matrices. Therefore, similar questions to the case of finite delay arise:
 24 does the discrete shift operator converge as $M \rightarrow \infty$? Is it possible to describe the
 25 spectral values, and do they converge to the exact ones? How does the choice of
 26 the nodes influence the convergence of the operators and their spectra?

27 The theoretical aspects are still work-in-progress, and there are still many ques-
 28 tions to address. The results and the numerical experiments are encouraging.

29 **Acknowledgments.** We would like to thank Sebastiaan G. Janssens for his valu-
 30 able comments and suggestions.

31 REFERENCES

- 32 [1] A. H. Al-Mohy and N. J. Higham, A new scaling and squaring algorithm for the matrix
 33 exponential, *SIAM J. Matrix Anal. Appl.*, **31** (2009), 970–989.
 34 [2] A. Andò, D. Breda, L. Davide, S. Maset, F. Scarabel and R. Vermiglio, 15 years or so of
 35 pseudospectral collocation methods for stability and bifurcation of delay equations, in print
 36 on *Advances on Delays and Dynamics*, Springer, New York, 2019.
 37 [3] C. T. H. Baker, *The numerical treatment of integral equations*, Monographs on Numerical
 38 Analysis, Clarendon Press, Oxford, 1977.
 39 [4] C. Baker, Numerical analysis of volterra functional and integral equations. the state of the
 40 art in numerical analysis, in *Inst. Math. Appl. Conf. Ser. New Ser.*, **63**, 1997.
 41 [5] H. T. Banks, J. A. Burns and E. M. Cliff, Spline-based approximation methods for control
 42 and identification of hereditary systems, in *International Symposium on Systems Optimiza-
 43 tion and Analysis (Rocquencourt, 1978)*, Lecture Notes in Control and Information Sci.,**14**,
 44 Springer, Berlin-New York, 1979, 314–320.
 45 [6] A. Batkai and S. Piazzera, *Semigroup for Delay Equations*, Research Notes in Mathematics,
 46 A K Peters, Wellesley Massachusetts, 2005.
 47 [7] A. Bellen and M. Zennaro, *Numerical methods for delay differential equations*, Numerical
 48 Mathematics and Scientific Computation, Oxford university press, 2013.
 49 [8] D. Breda, O. Diekmann, M. Gyldenber, F. Scarabel and R. Vermiglio, Pseudospectral dis-
 50 cretization of nonlinear delay equations: new prospects for numerical bifurcation analysis,
 51 *SIAM J. Appl. Dyn. Syst.*, **15** (2016), 1–23.

- 1 [9] D. Breda, O. Diekmann, D. Liessi and F. Scarabel, Numerical bifurcation analysis of a class
2 of nonlinear renewal equations, *Electron. J. Qual. Theory Differ. Equ.*, **64** (2016), 1–24.
- 3 [10] D. Breda, P. Getto, J. Sánchez Sanz and R. Vermiglio, Computing the eigenvalues of realistic
4 Daphnia models by pseudospectral methods, *SIAM J. Sci. Comput.*, **37** (2015), 2607–2629.
- 5 [11] D. Breda, S. Maset and R. Vermiglio, Pseudospectral differencing methods for characteristic
6 roots of delay differential equations, *SIAM J. Sci. Comput.*, **27** (2005), 482–495.
- 7 [12] D. Breda, S. Maset and R. Vermiglio, Pseudospectral approximation of eigenvalues of deriv-
8 ative operators with non-local boundary conditions, *Appl. Num. Math.*, **56** (2006), 318–331.
- 9 [13] D. Breda, S. Maset and R. Vermiglio, Approximation of eigenvalues of evolution operators for
10 linear retarded functional differential equations, *SIAM J. Num. Anal.*, **50** (2012), 1456–1483.
- 11 [14] D. Breda, S. Maset and R. Vermiglio, *Stability of linear delay differential equations. A numer-
12 ical approach with MATLAB*, SpringerBriefs in Control, Automation and Robotics, Springer,
13 New York, 2015.
- 14 [15] H. Brunner, *Collocation methods for Volterra integral and related functional differential equa-
15 tions*, Cambridge Monographs on Applied and Computational Mathematics, **15**, Cambridge
16 University Press, Cambridge, 2004.
- 17 [16] H. Brunner, *Volterra integral equations. An introduction to theory and applications*, Cam-
18 bridge Monographs on Applied and Computational Mathematics, **30**, Cambridge University
19 Press, Cambridge, 2017.
- 20 [17] H. Brunner and P. van der Houwen, *The numerical solution of Volterra equations*, CWI
21 Monographs, North-Holland Publishing Co., Amsterdam, 1986.
- 22 [18] C. Canuto, Y. M. Hussaini, A. Quarteroni and T. A. J. Zang, *Spectral methods in fluid
23 dynamics*, Springer Series in Computational Physics, Springer-Verlag, New York, 1988.
- 24 [19] M. D. Chekroun, M. Ghil, H. Liu and S. Wang, Low-dimensional galerkin approximations
25 of nonlinear delay differential equations, *Discrete and Continuous Dynamical Systems*, **36**
26 (2016), 4133–4177.
- 27 [20] P. Clément, O. Diekmann, M. Gyllenberg, H. Heijmans and H. Thieme, Perturbation theory
28 for dual semigroups. iii. nonlinear lipschitz continuous perturbations in the sun-reflexive case,
29 in *Proceedings of Volterra Integrodifferential Equations in Banach Spaces and Applications*
30 *1987*, 1989, 67–89.
- 31 [21] P. J. Davis, *Interpolation and Approximation*, Dover Books on Advanced Mathematics, Dover
32 Publication, Inc. New York, 1975.
- 33 [22] A. Dhooge, W. Govaerts, Y. A. Kuznetsov, H. G. E. Meijer and B. Sautois, New features
34 of the software matcont for bifurcation analysis of dynamical systems, *Mathematical and
35 Computer Modelling of Dynamical Systems*, **14** (2008), 147–175, Available from: <https://sourceforge.net/projects/matcont/>.
- 36 [23] O. Diekmann, P. Getto and M. Gyllenberg, Stability and bifurcation analysis of Volterra
37 functional equations in the light of suns and stars, *SIAM J. Math. Anal.*, **39** (2007), 1023–
38 1069.
- 39 [24] O. Diekmann and M. Gyllenberg, Equations with infinite delay: blending the abstract and
40 the concrete, *J. Differential Equations*, **252** (2012), 819–851.
- 41 [25] O. Diekmann, S. A. van Gils, S. M. Verduyn Lunel and H.-O. Walther, *Delay equations.
42 Functional, complex, and nonlinear analysis*, Applied Mathematical Sciences, **110**, Springer-
43 Verlag, New York, 1995.
- 44 [26] M. Dubiner, Asymptotic analysis of spectral methods, *J. Sci. Comput.*, **2** (1987), 3–31.
- 45 [27] K. Engelborghs, T. Luzyanina, G. Samaey, D. Roose and K. Verheyden, DDE-BIFTOOL:
46 a MATLAB package for bifurcation analysis of delay differential equations, Available from:
47 <http://ddebiftool.sourceforge.net>, Accessed: 11.01.2019.
- 48 [28] B. Fornberg, *A Practical Guide to Pseudospectral Methods*, Cambridge Monographs-on Ap-
49 plied and Computational Mathematics, **1**, Cambridge University Press, 1996.
- 50 [29] D. Funaro, A preconditioning matrix for the chebyshev differencing operator, *SIAM J. Num.
51 Anal.*, **24** (1987), 1024–1031.
- 52 [30] D. Funaro, Some results about the spectrum of the chebyshev differencing operator, in *North-
53 Holland Mathematics Studies*, **133**, Elsevier, 1987, 271–284.
- 54 [31] D. Funaro, Polynomial approximation of differential equation, in *Lecture Notes in Physics.
55 New Series m: Monographs*, **8**, Springer-Verlag, Berlin, 1992.
- 56 [32] P. Getto, M. Gyllenberg, Y. Nakata and F. Scarabel, Stability analysis of a state-dependent
57 delay differential equation for cell maturation: analytical and numerical methods, *J. Math.
58 Bio.*, **79** (2019), 281–328.
- 59

- 1 [33] D. Gottlieb, The stability of pseudospectral-Chebyshev methods, *Mathematics of Computa-*
2 *tion*, **36** (1981), 107–118.
- 3 [34] D. Gottlieb and L. Lustman, The spectrum of the Chebyshev collocation operator for the
4 heat equation, *SIAM J. Num. Anal.*, **20** (1983), 909–921.
- 5 [35] D. Gottlieb and S. A. Orszag, *Numerical analysis of spectral methods: theory and applications*,
6 SIAM, 1977.
- 7 [36] D. Gottlieb and E. Turkel, Topics in spectral methods, in *Numerical Methods in Fluid Dy-*
8 *namics*, Lecture Notes in Mathematics, **1127**, Springer, Berlin, 1985, 115–155.
- 9 [37] M. Gyllenberg, F. Scarabel and R. Vermiglio, Equations with infinite delay: numerical bifur-
10 cation analysis via pseudospectral discretization, *Appl. Math. Comput.*, **333** (2018), 490–505.
- 11 [38] E. Hairer, S. P. Nørsett and G. Wanner, *Solving ordinary differential equations I: Nonstiff*
12 *problems*, **8** of Springer Series in Computational Mathematics, Springer-Verlag Berlin, 1993.
- 13 [39] J. K. Hale and S. M. Verduyn Lunel, *Introduction to functional differential equations*, Applied
14 Mathematical Sciences, **99** Springer-Verlag, New York, 1993.
- 15 [40] J. S. Hesthaven, S. Gottlieb and D. Gottlieb, *Spectral methods for time-dependent problems*,
16 **21**, Cambridge University Press, 2007.
- 17 [41] N. J. Higham, The scaling and squaring method for the matrix exponential revisited, *SIAM*
18 *J. Matrix Anal. Appl.*, **26** (2005), 1179–1193.
- 19 [42] N. J. Higham and A. H. Al-Mohy, Computing matrix functions, *Acta Numerica*, **19** (2010),
20 159–208.
- 21 [43] K. Ito and F. Kappel, A uniformly differentiable approximation scheme for delay systems
22 using splines, *Appl. Math. Optim.*, **23** (1991), 217–262.
- 23 [44] K. Ito and F. Kappel, Two families of approximation schemes for delay systems, *Results*
24 *Math.*, **21** (1992), 93–137.
- 25 [45] Z. Jackiewicz and B. D. Welfert, Stability of Gauss–Radau pseudospectral approximations of
26 the one-dimensional wave equation, *J. Sci. Comput.*, **18** (2003), 287–313.
- 27 [46] R. Kress, *Linear Integral Equations*, **82** of Applied Mathematical Sciences, Springer-Verlag,
28 Berlin, 1989.
- 29 [47] D. Liessi, *Pseudospectral methods for stability of periodic solutions of delay models*, PhD
30 thesis, University of Udine, 2018.
- 31 [48] S. Maset, The collocation method in the numerical solution of boundary value problems for
32 neutral functional differential equations. Part I: Convergence results, *SIAM J. Num. Anal.*,
33 **53** (2015), 2771–2793.
- 34 [49] S. Maset, The collocation method in the numerical solution of boundary value problems
35 for neutral functional differential equations. Part II: Differential equations with deviating
36 arguments, *SIAM J. Num. Anal.*, **53** (2015), 2794–2821.
- 37 [50] S. Maset, An abstract framework in the numerical solution of boundary value problems for
38 neutral functional differential equations, *Numerische Mathematik*, **133** (2016), 525–555.
- 39 [51] G. Mastroianni and G. V. Milovanović, *Interpolation processes. Basic theory and applications*,
40 Springer Monographs in Mathematics, Springer-Verlag, Berlin, 2008.
- 41 [52] W. Michiels, I. Boussaada and S.-I. Niculescu, An explicit formula for the splitting of multiple
42 eigenvalues for nonlinear eigenvalue problems and connections with the linearization for the
43 delay eigenvalue problem, *SIAM J. Matrix Anal. Appl.*, **38** (2017), 599–620.
- 44 [53] W. Michiels and S.-I. Niculescu, *Stability, control, and computation for time-delay systems.*
45 *An eigenvalue-based approach, 2nd edn.*, **27** of Advances in Design and Control, SIAM, 2014.
- 46 [54] S. C. Reddy and L. N. Trefethen, Lax-stability of fully discrete spectral methods via stability
47 regions and pseudo-eigenvalues, *Computer Methods in Applied Mechanics and Engineering*,
48 **80** (1990), 147–164.
- 49 [55] T.-J. Rivlin, *Chebyshev Polynomials: From Approximation Theory to Algebra and Number*
50 *Theory*, Pure and Applied Mathematics: A Wiley Series of Texts, Monographs and Tracts,
51 **10**, Wiley, 1990.
- 52 [56] F. Scarabel, *Capture the past to portray the future: Numerical bifurcation analysis of delay*
53 *equations, with a focus on population dynamics*, PhD thesis, University of Helsinki, 2018.
- 54 [57] D. M. Sloan, On the norms of inverses of pseudospectral differentiation matrices, *SIAM J.*
55 *Num. Anal.*, **42** (2004), 30–48.
- 56 [58] A. Solomonoff and E. Turkel, Global properties of pseudospectral methods, *J. Comput. Phys.*,
57 **81** (1989), 239–276.
- 58 [59] R. Szalai, Knut, Available from: <http://rs1909.github.io/knut/>, Accessed: 11.01.2019.
- 59 [60] L. N. Trefethen, *Approximation theory and approximation practice*, SIAM, 2013.

- 1 [61] L. N. Trefethen, Inverse yogiisms, *Notices of the American Mathematical Society*, **63** (2016),
2 1281–1285.
- 3 [62] L. N. Trefethen and M. Embree, *Spectra and pseudospectra: the behavior of nonnormal ma-*
4 *trices and operators*, Princeton University Press, 2005.
- 5 [63] L. N. Trefethen and M. R. Trummer, An instability phenomenon in spectral methods, *SIAM*
6 *J. Num. Anal.*, **24** (1987), 1008–1023.
- 7 [64] J. Wang and F. Waleffe, The asymptotic eigenvalues of first-order spectral differentiation
8 matrices, *J. Appl. Math. Phys.*, **2** (2014), 176–188.
- 9 [65] G. Webb, Functional differential equations and nonlinear semigroup in l_p -spaces, *J. Differen-*
10 *tial Equations*, **20** (1976), 71–89.
- 11 [66] Z. Wu and W. Michiels, Reliably computing all characteristic roots of delay differential equa-
12 tions in a given right half plane using a spectral method, *J. Comput. Appl. Math.*, **236** (2012),
13 2499–2514.

14 Received xxxx 20xx; revised xxxx 20xx.

15 *E-mail address:* o.diekman@uu.nl

16 *E-mail address:* francesca.scarabel@helsinki.fi

17 *E-mail address:* rossana.vermiglio@uniud.it