

# Università degli studi di Udine

# Compositional characterization of commercial sparkling wines from cv. Ribolla Gialla produced in Friuli Venezia Giulia

**Original** 

Availability: This version is available http://hdl.handle.net/11390/1157659 since 2020-03-27T11:15:01Z

Publisher:

Published DOI:10.1007/s00217-019-03334-9

Terms of use:

The institutional repository of the University of Udine (http://air.uniud.it) is provided by ARIC services. The aim is to enable open access to all the world.

Publisher copyright

(Article begins on next page)



Draft Manuscript for Review

## **Characterization of the metabolomic profile of Ribolla Gialla commercial sparkling wines**



Manuscripts

 $\mathbf{1}$  $\overline{2}$  $\overline{3}$  $\overline{4}$  $\overline{7}$ 



#### **Abstract**

From the properties of properties and  $\beta$ -<br>rietal aromas were below the odor thresh<br>e found (e.g., palmitic acid), which could<br>However, high concentrations of tryptopl<br>ing atypical ageing defects.<br>la Gialla; sparkling wi Ribolla Gialla (RG) is a white grape variety used in the production of high-quality sparkling wines. It is widespread in North-East Italy, particularly Friuli Venezia Giulia (FVG), as well as Slovenia. Because of its limited area of cultivation, however, there is little information about the composition and chemical characteristics of the sparkling wines produced. This work used different analytical approaches to characterize thirty-three commercial sparkling RG wines from different areas of FVG. The characteristics included the overall volatile profile and content of terpenes, C13-norisoprenoids, lipids, and tryptophan metabolites. The aroma profile of RG wines was mainly characterized by fermentative esters and β-damascenone, whereas other norisoprenoids and varietal aromas were below the odor threshold. Appreciable amounts of certain fatty acids were found (e.g., palmitic acid), which could be potentially correlated with greater foam stability. However, high concentrations of tryptophan metabolites highlighted a higher risk of developing atypical ageing defects.

**KEYWORDS:** Ribolla Gialla; sparkling wine; volatile compounds; tryptophan metabolites;

lipids

#### **Introduction**

 Ribolla Gialla is a promising white grape variety that has recently been used for the production of premium sparkling wines, which have received good appreciation in international wine markets. Ribolla has been cultivated since ancient times in North-Eastern Italy (in the Friuli Venezia Giulia region), as well as Slovenia and the Ionian Islands (Kefalonia), where it is known as Rebula and Robola, respectively [1]. A certain number of studies have been carried out on the genetic identity and characterization of Ribolla grapes [1–4], but very few studies have dealt with the metabolomic profile of the wines [5, 6]. In addition, none of them have focused on the aromatic characteristics of sparkling wines obtained from Ribolla grapes, including their volatile organic compound (VOC) content, lipids, polyphenols, and tryptophan metabolites. All of these compounds are important because they may be linked with positive and negative aspects of wine quality.

etabolomic profile of the wines [5, 6]. In<br>titic characteristics of sparkling wines ob<br>e organic compound (VOC) content, lipids<br>see compounds are important because they<br>f wine quality.<br>I for determining the sensory charact VOCs are fundamental for determining the sensory characteristics of wine and its profile. These compounds are classified into four groups [7]: i) primary grape VOCs, which are present in the cells of grapes; ii) secondary grape aroma, which are formed during the pre-fermentation phases (crushing, pressing, and skin contact) or by thermal, chemical, and enzymatic reactions in must; iii) fermentation bouquet, which includes aroma compounds that form during alcoholic fermentation; iv) and maturation bouquet, which refers to the aroma compounds that develop when ageing the wine. Primary aroma is particularly important for the production of certain wine typologies, such as white sparkling wines. Furthermore, different aroma compounds may be produced after fermentation and ageing that further affect the aroma profile of the final wine, which are mainly due to pre-fermentative processes and yeast metabolism (e.g., C6-compounds, norisoprenoids, fatty acids, alcohols, esters, carbonyl compounds) [8–11].

 Moreover, yeast autolysis takes place during the ageing of sparkling wine, which leads to the release of substances in the medium. Lipids represent a wide class of compounds among these substances [8]. Fatty acids may have a significant impact on the sensory properties of the wine.  A large part of the fatty acids originates from the firm tissues of the grapes, but the greatest amount is formed during alcoholic fermentation. Therefore, fatty acids may be present in wine in free or bound forms as ethyl esters. Both forms directly contribute to the flavor of the wine, while unsaturated fatty acids such as oleic, linolenic, and linoleic acids act as precursors of C6- aldehydes and alcohols with herbaceous notes, and are critical for yeast growth during fermentation [12]. Another part of the sensory assessment of sparkling wine is the relationship between the foaming properties and lipid compounds. Therefore, it has been reported that medium-chain free fatty acids C8, C10, and C12 are negatively correlated with foamability, while the ethyl esters of hexanoic, octanoic, and decanoic acids have a positive effect on the formation and stability of the foam [13].

of hexanoic, octanoic, and decanoic acids<br>
of the foam [13].<br>
cid tryptophan (TRP) and its metabolites, e<br>
to be potential precursors of 2-aminoacetop<br>
causes an atypical ageing off-flavor (ATA<br>
l with aroma descriptors s The essential amino acid tryptophan (TRP) and its metabolites, especially indole-3-acetic acid (IAA), are considered to be potential precursors of 2-aminoacetophenone (2-AAP), which is an aroma compound that causes an atypical ageing off-flavor (ATA) in *Vitis vinifera* wines. The off-flavor is described with aroma descriptors such as "acacia blossom," "furniture polish," "wet wool," "mothballs," or "fusel alcohol," which are combined with a loss of the typical bouquet of the grape variety [14]. Depending on the wine matrix, the detection threshold of 2- 81 AAP varies from 0.5 to 1.5 µg L<sup>-1</sup>. Strongly aromatic wines are able to integrate more than 1.5  $\mu$ g L<sup>-1</sup> of 2-AAP, while meager wines might be rejected as tainted by ATA with less than 0.5  $\mu$ g L<sup>-1</sup>. It is generally accepted that the ultimate cause of ATA development in white wines is a stress reaction in the vineyard triggered by drought, nutritional deficiency, and other viticultural factors, such as the time of harvest and leaf removal [14, 15].

 Most papers on sparkling wines focus on the characterization of internationally known varieties with a worldwide distribution [16–19], but there are few studies on local or less relevant cultivars. Therefore, the aim of this work was to investigate the compositional profile of different Ribolla Gialla sparkling wines produced in different DOC (*Denominazione di Origine Controllata*) districts in the Friuli Venezia Giulia region in North-East Italy. The wines were

 characterized for their chemical composition (pH, titratable acidity, residual sugars, alcohol, and sulfur dioxide), volatile profile (overall profile of aroma compounds, free and bound terpenes, and norisoprenoids), and content of lipids and tryptophan metabolites. The results are critically discussed with the aim of defining the metabolomic profile of Ribolla Gialla sparkling wines.

**Materials and Methods**

#### *Reagents and materials*

30% w/w), ethanol (96% v/v), ACS gra<br>fate and citric acid were purchased from<br>le (99.5%) was obtained from Honeywel<br>solvents dichloromethane, *n*-pentane, a<br>*i*, 2-propanol, chloroform, formic acid,<br>l, 2-octanol, ethyl he Hydrogen peroxide (30% w/w), ethanol (96% v/v), ACS grade hydrochloric acid (37%), anhydrous sodium sulfate and citric acid were purchased from Carlo Erba Reagents (Milan, Italy). Sodium chloride (99.5%) was obtained from Honeywell Fluka (Morris Plains, New Jersey). HPLC-grade solvents dichloromethane, *n*-pentane, and methanol, LC-MS grade methanol, acetonitrile, 2-propanol, chloroform, formic acid, ammonium formate, ethyl 103 heptanoate, 1-heptanol, 2-octanol, ethyl hexanoate-d<sub>11</sub>, 3-(2-hydroxy ethyl)-indole, kynurenic acid, D-tryptophan methyl ester, L-tyrosine ethyl ester, *N*-acetyl tyrosine ethyl ester, and 3,5- di-*tert*-4-butylhydroxytoluene (BHT) were obtained from Sigma-Aldrich (St. Louis, MO, 106 USA). C<sub>7</sub>-C<sub>30</sub> *n*-alkane solution in *n*-hexane was purchased from Supelco (Bellefonte, PA, USA), while cholesterol-d <sup>7</sup> and octadecanoic acid-d <sup>3</sup> were obtained from CDN Isotopes (Quebec, Canada). Finally, the glycosidase preparation (Rapidase Revelation Aroma) used for the determination of bound monoterpenes was obtained from Oenobrands (Montellier, France).

#### *Wine samples*

 Local wineries located in the North-East Italian region of Friuli Venezia Giulia provided thirthy-three commercial Ribolla Gialla sparkling wines. The locations of the wineries are shown in Figure 1.

 The basic quality control parameters (reducing sugars, alcoholic strength, total acidity, volatile acidity, pH, malic, lactic and tartaric acid) were determined by FTIR spectroscopy with a 117 Winescan<sup>TM</sup> FT-120 instrument (FOSS, Hillerød, Denmark). All of the samples were analyzed two times, and the mean value of the two measurements was considered for the data analysis.

#### *Determination of free and bound terpenes and norisoprenoids by SPE-GC-MS*

Superiorus in rice and boand forms were expected<br>Superiorus in rice and boand forms were expected<br>A GCMS-QP-2010 system (Shimadzu, Ky<br>compounds were separated on a J&W DB<br>um film thickness) provided by Agilent Te<br>perating 120 Terpenes and  $C_{13}$ -norisoprenoids in free and bound forms were extracted on Isolute ® 500-mg, 6-mL, C18 SPE cartridges (Biotage, Uppsala, Sweden) according to the method reported by Comuzzo et al. [20]. A GCMS-QP-2010 system (Shimadzu, Kyoto, Japan) was used for GC- MS analyses. Volatile compounds were separated on a J&W DB-Wax capillary column (30 m  $124 \times 0.25$  mm i.d., 0.25  $\mu$ m film thickness) provided by Agilent Technologies Inc. (Santa Clara, CA, USA) using the operating conditions described by Loira et al. [21].

 Electron impact mass spectra were recorded at 70 eV and volatile compounds were tentatively identified by comparison of their mass spectra and retention times with those of standard compounds or by comparison of the mass spectrum with those reported in the Wiley 6, NIST 21, and NIST 107 mass spectrum libraries (provided by the manufacturer). Linear retention indices were calculated according to the retention times of *n*-alkanes and compared with those reported in literature. The semi-quantitative analysis was based on the internal 132 standard method using 1-heptanol (312 mg L<sup>-1</sup> in 96 % v/v ethanol) as the internal standard while considering the response factor to be equal to 1.00.

*Determination of volatile compounds*

 The volatile compounds in the commercial wines were determined using two analytical approaches: liquid-liquid extraction (LLE) and solid-phase microextraction (SPME). Samples

 were degassed before analysis by placing them in an ultrasonic bath (Falc Labsonic, Treviglio, Italy) for 2 min.

#### *Liquid-liquid extraction (LLE-GC-MS)*

 Aroma compounds were determined as reported by Loira et al. [21] by mixing 5 mL of wine with 5 mL of a 30% (w/v) sodium chloride solution and 200 μL of internal standard (ethyl 142 heptanoate, 422 mg  $L^{-1}$  in 96% v/v ethanol). The mixture was subjected to three extractions using 2.5 mL of pentane:dichloromethane (2:1 v/v) each. The organic phase was collected in a Pyrex tube, dehydrated with anhydrous sodium sulfate, and concentrated under nitrogen flow to a final volume of about 1 mL. The samples obtained were subjected to GC-MS analysis as reported previously[21].

#### *SPME-GC-MS/MS*

d with anhydrous sodium sulfate, and convolut 1 mL. The samples obtained were sul<br>
1. The samples obtained were sul<br>
1. Subsetequency of 2-octanol at 2.<br>
1. Subsetequency of 2-octanol at 2.<br>
1.5 g of sodium chlc<br>
1.5 g of Degassed wine (1 mL) was spiked with 50 µL of 2-octanol at 2.13 mg L-1 (IS) in ethanol and placed in a 20-mL headspace vial containing 1.5 g of sodium chloride. Two technical replicates were prepared for each sample, along with a blank sample containing only sodium chloride. GC analysis was performed using a Trace GC Ultra gas chromatograph coupled with a TSQ Quantum Tandem mass spectrometer, which was upgraded to the XLS configuration and equipped with a Triplus autosampler (Thermo Fisher Scientific, Waltham, MA, USA). The method used was adopted by Carlin et al. [22]. Samples were incubated for 5 min at 35°C,s and VOCs were extracted for 20 min with a 2-cm-long 50/30-µm coated divinylbenzene/carboxen/polydimethylsiloxane (DVB/CAR/PDMS) SPME fiber (Supelco, Sigma Aldrich, Milan, Italy). After solid-phase microextraction, the fiber was desorbed for 3 min at 250°C in the GC system with the injector set in splitless mode. The fiber was reconditioned between each sample at 270°C for 7 min. Helium was used as a carrier gas at a flow rate of 1.2 mL min-1 .

161 The GC oven was equipped with a  $30-m \times 0.25$ -mm VF-WAXms column (Agilent Technologies Inc., Santa Clara, CA, USA) with a film thickness of 0.25 µm. The oven temperature was held for 2 min at 40°C after the injection, ramped at 6°C min-1 up to 250°C, 164 and held for 5 min. The ion source was set at 230°C, and electron impact mass spectra were recorded at 70 eV. Data acquisition and analyses were performed using the Xcalibur Workstation software supplied by the manufacturer.

#### *Determination of tryptophan metabolites by UHPLC-MS/MS*

les were filtered at 0.22  $\mu$ m by a Millemd directly collected in 2-mL HPLC am<br>aity UHPLC system provided with an aut<br>pectrometer equipped with an electrospray<br>Fryptophan metabolites were separated at a<br>nm Waters Acquity Sparkling wine samples were filtered at 0.22 µm by a Millex-GV filtration unit (Merc, Darmstadt, Germany) and directly collected in 2-mL HPLC amber vials. The analyses were performed on an Acquity UHPLC system provided with an autosampler and coupled with a XEVO TQMS mass spectrometer equipped with an electrospray source (Waters Corporation, Milford, MA, USA). Tryptophan metabolites were separated at a flow rate of 0.4 mL min-1 on 173 a 1.8-um 150  $\times$  2.1-mm Waters Acquity HSS T3 column (Waters Corporation), which was conditioned at 40°C. A linear gradient of water containing 0.1% formic acid (solvent A) and 0.1% formic acid in acetonitrile (solvent B) was used according to the conditions reported by Arapitsas et al.[23]. The injection volume was 10 µL, and the wine samples were stored in the 177 autosampler at 6<sup>o</sup>C during analyses.

178 The MS conditions included a capillary voltage of 3.5 kV in positive mode and  $-2.7$  kV in negative mode. The ion source temperature was set at 150°C, while the desolvation temperature 180 was 500 $^{\circ}$ C. Nitrogen was used as the cone gas with a flow rate of 50 L h<sup>-1</sup>. The data were processed using Waters MassLynx (version 4.1) and TargetLynx software (Milford, MA, USA).

#### *Analysis of lipid molecules by UHPLC-MS/MS*

 Lipid analysis was performed as described by Della Corte et al. [24] with slight modifications. An aliquot (3 mL) of degassed wine was introduced in a 50-mL Falcon tube and spiked with 186 30 µL of a methanolic solution containing cholesterol- $d_7$  and octadecanoic acid- $d_3$  (IS) at 187 concentrations of 1.01 mg mL<sup>-1</sup> and 1.002 mg mL<sup>-1</sup>, respectively. Lipids were extracted two 188 times in 21 mL of a chloroform–methanol solution (2:1 v/v) containing BHT (10 mg  $L^{-1}$ ). The total lower lipid-rich layer was collected in 100-mL flasks, and the solvent was evaporated to dryness using a rotary evaporator. The samples were reconstituted in 300 µL of acetonitrile/2- propanol/water (65:30:5 v/v/v) and filtered (0.22 µm) into 2-mL HPLC amber vials for UHPLC-MS/MS analysis.

evaporator. The samples were reconstitute<br>  $(0.22 \mu m)$  into 2<br>  $s$  iss.<br>
s performed on a Dionex 3000 chromatogra<br>
equipped with an autosampler and coup<br>
trometer (Sciex, Concord, Vaughan, ON, on<br>
source (ESI). Lipids wer UHPLC separation was performed on a Dionex 3000 chromatograph (Thermo Fisher Scientific, Waltham, MA, USA) equipped with an autosampler and coupled with an API 5500 triple- quadrupole mass spectrometer (Sciex, Concord, Vaughan, ON, Canada), which was provided 196 with an electrospray ion source (ESI). Lipids were separated on a 2.7-um 150  $\times$  2.1-mm RP Ascentis Express column (Sigma-Aldrich, Milan, Italy) set at 55°C. The injection volume was  $\frac{1}{2}$  µL, and the samples were stored in an autosampler at 10 $\degree$ C during the analyses. The mobile phase and chromatographic conditions are those described by Della Corte et al. [24], and the flow rate was 0.260 mL min-1 .

 The spray voltage of the ESI source was set at 5500 V for positive mode and –4500 V for 202 negative mode, and the source temperature was  $250^{\circ}$ C. The nebulizer (Gas 1) and heater gas (Gas 2) pressures were set at 40 and 20 psi, respectively. Ultra-high-purity nitrogen (99.999%) was used as both a curtain gas and collision gas (CAD) at 20 and 9 psi, respectively. Instrument control and data acquisition were performed by Analyst software (Applera Corporation, Norwalk, CT, USA), and the data were processed using MultiQuant, version 2.1 (Sciex, Concord, Vaughan, ON, Canada).

 Statistica for Windows Version 8 (Statsoft, Tulsa, OK, USA) was used to calculate the means, standard deviations (SDs), coefficients of variation (CVs), minimum values (MIN), and maximum values (MAX) of the different analytical parameters determined for wine samples.

## **Results and Discussion**

#### *Basic quality parameters*

ults related to the basic chemical compos<br>s show that the samples are very homog<br>dity, tartaric acid, pH, and even volatile ac<br>tly higher CV. This relatively homogene<br>at the sparkling wines analyzed are all<br>ment of the st Table 1 shows the results related to the basic chemical composition of the wines. The mean values, SDs, and CVs show that the samples are very homogeneous concerning alcoholic strength, titratable acidity, tartaric acid, pH, and even volatile acidity, which shows quite low values despite a slightly higher CV. This relatively homogeneous composition is probably related to the fact that the sparkling wines analyzed are all DOC productions and were manufactured in fulfillment of the standards reported in the DOC production guidelines.

 However, some differences that are more relevant from a practical point of view were observed 221 for residual sugars (min-max range:  $6-19 \text{ g L}^{-1}$ ), malic acid (0.01-3.79 g L<sup>-1</sup>), and lactic acid  $(0.00-2.75 \text{ g L}^{-1})$ . These differences could be related to the application of different winemaking protocols by the producers, leading to different styles of wine for Ribolla Gialla DOC, even if they are in agreement to the DOC production guidelines. In regard to sugars, most of the wines 225 analyzed showed a residual sugar content of less than 12 g  $L^{-1}$  or between 12 and 17 g  $L^{-1}$ . According to the European Commission Regulation (EC) n. 607/2009 [25], these wines are classified as *brut* and *extra dry*, respectively. Only one sample had higher sugar content (about 228 19 g  $L^{-1}$ ) and was categorized as *sec* (or *dry*).

 The variations of malic and lactic acid highlight the different ways of managing malolactic fermentation (MLF), which are probably connected with the different storage times of the wines on yeast lees after refermentation. Most of the wines did not contain lactic acid in appreciable

 amounts, which was probably due to the lack of malolactic fermentation. MLF generally occurs during ageing on the lees because of the positive effect that they have on promoting the growth of lactic acid bacteria [26]. For this reason, most winemakers would consider Ribolla Gialla sparkling wine as a fresh and young wine according to the data.

concentration. This evidence may be related and with some products being kept in atoclave. Such *sur-lies* aging may be prologe). In addition, certain winemakers also prologe). In addition, certain winemakers also promome Ribolla Gialla is generally refermented via the Martinotti/Charmat short method, which involves refermentation in stainless steel autoclaves without contact with the lees at the end of refermentation. However, six wine samples had a higher content of lactic acid. This accounted for the increase of the CV calculated for lactic acid in Table 1 and was combined with the decrease of malic acid concentration. This evidence may be related to the manufacture of certain Ribolla Gialla wines, with some products being kept in contact with the lees after refermentation in an autoclave. Such *sur-lies* aging may be prolonged for up to several months (2-4 months on average). In addition, certain winemakers also produce Ribolla Gialla by the 244 traditional refermentation method in bottles.

#### *Wine aroma profile*

 Table 2 shows the results of the qualitative and quantitative determination of the aroma compounds detected in wine samples by the different analytical techniques. Concerning non- varietal aromas, a total number of fifty-eight volatiles were tentatively identified in the wines, including acids, alcohols, esters, C6 compounds, diols, and carbonyls. Fatty acids are some of the most representative volatiles in the wines analyzed, and generally, they are described as having cheese, rancid, and fatty notes. If their amounts are higher than the odor detection threshold (ODT), they may negatively affect the organoleptic characteristics of the wines.

 In the samples analyzed, the average concentration of fatty acid was often close and sometimes higher than their ODT. Nevertheless, their odor activity value (OAV), which is the ratio between the concentration and ODT, was in the range of 1-4 for many of the compounds analyzed. The exception was octanoic acid, which is the most representative fatty acid in the

257 volatile composition of Ribolla wines (OAV  $\approx$  30). These results suggest that fatty acids might have minor relevance in the aromatic characteristics of Ribolla Gialla. Acetic acid (which is responsible for the typical vinegar off-flavor) was found to be lower than ODT in all samples analyzed.

 Interestingly, the variability among the samples (i.e., the CVs) for fatty acids and volatile compounds in general (Table 2) was higher than that detected for the basic parameters in Table 1. This highlights a certain level of differentiation among Ribolla Gialla DOC wines, which is probably related to a number of variables, such as those typically connected with the development of wine aromas. Concerning fatty acids, for instance, the differences among the samples might have resulted from the different origin of the grapes used, the amount of lipid substances in the musts, the diverse winemaking conditions, and the yeast strains used for fermentations [27].

aromas. Concerning fatty acids, for instant<br>sulted from the different origin of the gra<br>sts, the diverse winemaking conditions, a<br>are also important volatiles among the com<br>s during alcoholic fermentation as produce<br>ons to Higher alcohols (HA) are also important volatiles among the compounds listed in Table 2. They are produced by yeasts during alcoholic fermentation as products of amino acid metabolism [11]. Their contributions to wine aroma vary from honey, rose, and floral characters (2- phenylethyl alcohol and benzyl alcohol) to pungent and solvent-like smells (1-propanol, 1- butanol, 2- and 3-methyl-1-butanol) [28], and the effects depend on their concentration [9]. The average amounts of HA observed in the wines analyzed were generally found to be lower than 275 their ODT except for 2- and 3-methyl-1-butanol and 2-phenylethanol (ODT: 40 and 10 mg  $L^{-1}$ , respectively). This composition may be linked with the yeast strains used, as well as the winemaking conditions adopted and the amino acidic composition of the grapes. The presence of significant amounts of 2-phenylethanol is very interesting for its fresh rose-like odor [29], even if its OAV range was about 1-6.

 Esters are the most representative class of volatile compounds found in Ribolla Gialla sparkling wines. Isoamyl acetate (3-methyl-1-butanol acetate), ethyl hexanoate, ethyl lactate, ethyl octanoate, diethyl succinate, and 2-phenylethyl acetate showed the highest concentrations. The

 compounds eluted at higher retention times (e.g., diethyl malate) presumably had a minor impact on the aroma profile because of their low volatility. Some esters, such as ethyl butanoate and ethyl hexanoate, were close to or higher than their ODT (20 µg and 65 µg) in all the samples analyzed. Others such as isoamyl acetate, 2-phenylethyl acetate, and hexyl acetate were detected in only certain wines with concentrations higher than their ODT. Compounds such as ethyl hexanoate and isoamyl acetate were found to have significant OAVs (e.g., up to 18 for ethyl hexanoate, 85 for hexyl acetate, and approximately 200 for isoamyl acetate). This observation is very interesting from a sensory point of view because esters generally confer floral, fresh, and fruity notes (rose, banana, pear, green apple) to the wines [28, 30, 31], which are generally recognized as typical in Ribolla wines. For this reason, esters appeared to be an important component of the volatile profile of Ribolla Gialla.

r notes (rose, banana, pear, green apple) to<br>ed as typical in Ribolla wines. For this rea<br>of the volatile profile of Ribolla Gialla.<br>ctate and diethyl succinate are considered "<br>rally increases during ageing and after mal<br> Among esters, ethyl lactate and diethyl succinate are considered "ageing esters" (AE), and their content in wines generally increases during ageing and after malolactic fermentation [29, 32]. The same consideration is true for diethyl malate, but this compound is normally detected less frequently in wines. The concentration of AE was generally found to be lower than the ODT in most of the wines analyzed. However, in some samples, ethyl lactate had higher concentrations than its odor threshold (60 mg/L) in accordance with the amount of malic and lactic acid. In fact, the highest concentration of these esters was observed in the same products in which malolactic fermentation was hypothesized to have taken place (i.e., the wines where malic acid was not detected), as shown in Table 1. This may confirm that certain Ribolla Gialla sparkling wines are produced with a more or less prolonged period of *sur-lies* ageing or by refermenting them in bottle by the traditional method.

 Carbonyls, diols, and C6 compounds appeared not to have an important contribution to the aromatic characterization of the samples, and most of the cases, they were detected at levels below their ODT. Hexanal was found to overcome its ODT in some samples, but at the low concentrations detected, it may or may not contribute to the fresh vegetal notes of Ribolla wines.

Also  $\beta$ -citronellol, was found at lower level<br>owed similar average values to those of so<br>sylated forms, only geraniol was significa<br>prevalence of the latter (combined) form a<br>aniol (with minor traces of nerol) was the c 309 Table 3 reports the detected varietal aromas (terpenes and  $C_{13}$ -norisoprenoids) in free and bound form in the samples. Terpenes are well known to confer floral odors to wines [29]. In Ribolla, they were especially found in the free form, but in most cases, their average concentrations were below their ODT with a few exceptions for linalool and geraniol. Although it was present below its ODT, α-terpineol was the most abundant free terpenic alcohol in the wines. Interestingly, even if Ribolla Gialla cannot be considered as an aromatic variety, the concentrations detected for α-terpineol and linalool were higher than those normally observed in certain international white varieties, such as Pinot blanc, Pinot Gris, Chardonnay, and Sauvignon blanc [19]. Also  $\beta$ -citronellol, was found at lower levels with respect to its ODT (100 µg L-1) [33], it showed similar average values to those of some Riesling wines [19]. Considering the glycosylated forms, only geraniol was significantly detected as both free and bound terpenol with a prevalence of the latter (combined) form of the molecule. According to other experiments, geraniol (with minor traces of nerol) was the only terpenic alcohol normally found in Ribolla Gialla grapes [34]. For this reason, in Ribolla Gialla, linalool, α-terpineol, and 323 B-citronellol may be formed from such terpenic alcohols during grape processing. Citronellol, for instance, is reported to be produced by *S. cerevisiae* from geraniol and nerol during alcoholic fermentation, and other formation pathways have also been suggested for linalool (from geraniol) and α-terpineol (by cyclization of nerol) [35]. For both free and bound geraniol, the average values were generally lower than the ODT reported for the alcohol, but some of the samples showed significant levels of total geraniol (Table 3). Considering that the bound form generally prevails for this terpenol, one interesting way to improve the volatile profile of these wines might be the application of techniques to increase the release of free geraniol from its glycosides in the production process of Ribolla Gialla sparkling wines. In contrast to terpenes, which did not appear particularly characteristic for the aroma profile of

333 the wines, some  $C_{13}$ -norisoprenoids were detected in appreciable concentrations (Table 3).

334 TDN and B-damascenone, for instance, overcame their sensory thresholds in different samples.

Page 15 of 35

 $\mathbf{1}$ 

 Norisoprenoids are generally not connected to specific aromatic grapevine varieties and 336 generally have a low odor threshold. The ODT of  $\beta$ -damascenone is 0.04-0.06  $\mu$ g L<sup>-1</sup> in a model dilute alcohol solution [29], while a relatively wide range of ODTs was found in wine: Sefton 338 et al. [36] reported 0.14  $\mu$ g L<sup>-1</sup> in deodorized white wine,  $0.85 - 2.10 \mu$ g L<sup>-1</sup> in deodorized red 339 wine, and 7.00  $\mu$ g L<sup>-1</sup> in red wine. The presence of  $\beta$ -damascenone may be particularly interesting for Ribolla wines for not only the relatively high concentrations found but also because it is reported to modify the sensory perception of some esters. Escudero et al.[37] found 342 that the addition of low levels of  $\beta$ -damascenone (0.85 µg L<sup>-1</sup>) to a solution of esters increased 343 the fruity notes of the mixture, while higher levels  $(3.5 \text{ µg } L^{-1})$  accounted for the development of strong raisin/dry-plum odors.

mixture, while higher levels  $(3.5 \mu g L^{-1})$  as<br>almodors.<br>prenoids is also dependent on the winemak<br>ved at higher concentrations in young w<br>naphthalene (TDN), a positive correlation b<br>ed.[38] Their odor descriptors range fr 345 The amount of norisoprenoids is also dependent on the winemaking conditions. In general, β- damascenone is observed at higher concentrations in young wines [36], whereas for 1,1,6- trimethyl-1,2-dihydronaphthalene (TDN), a positive correlation between its occurrence in wine and ageing was reported.[38] Their odor descriptors range from floral (e.g., vitispiranes) and fruity-honey scents ( -damascenone) to notes of kerosene and petrol (for TDN) [39]. The presence of TDN among the compounds listed in Table 3 may represent further evidence about the ageing of certain Ribolla Gialla sparkling wines after refermentation.

 Another interesting aspect related to norisoprenoids is the high concentration observed for some 353 of them in bound form, particularly -oxo- $\alpha$ -ionol and  $3$ -hydroxy- $\beta$ -damascone, which are the most representative compounds. As for terpenes, Ribolla Gialla appeared to have an unexpressed aromatic potential for norisoprenoids. Thus, from a technical point of view, the optimization of winemaking process to maximize the release of this bound aromatic potential might be an interesting way to improve the quality of Ribolla Gialla sparkling wines.

#### *Metabolomic fingerprint of tryptophan metabolites and indoles*

 The second group of metabolites analyzed was the amino acid tryptophan and its catabolites, as shown in Table 4. There were twenty-eight compounds detected, among which the sulfonated derivates of indole 3-lactic acid, tryptophol, indole 3-lactic acid glucoside, and indole 3-acetic acid are presented. Sulfonation is the addition of a sulfonic acid group ( $-SO<sub>3</sub>H$ ) to an organic compound and is a widespread industrial process used in a diverse range of products. It also has a major function in modulating biological activities that are known to occur in wine, which involve several metabolites such as polyphenols and indoles [23].

blites such as polyphenols and indoles [23]<br>ell known to have much lower amounts of<br>to the second fermentation [23]. Tryptoph<br>.01 to 0.39 mg L<sup>-1</sup> with a mean value of 0.1<br>ester (0.44 mg L<sup>-1</sup> mean value). TOL-SO<sub>3</sub>l<br>n me Sparkling wines are well known to have much lower amounts of tryptophan than other wines, which is probably due to the second fermentation [23]. Tryptophan values for the commercial samples ranged from 0.01 to 0.39 mg L-1 with a mean value of 0.13 mg L-1, and which similarly 369 occurred for its ethyl ester  $(0.44 \text{ mg } L^{-1}$  mean value). TOL-SO<sub>3</sub>H had higher amounts among all the other tryptophan metabolites with a mean value of 3.73 mg L<sup>-1</sup>. The unsulfonated form of tryptophol varied from 0.01 to 1.08 mg L-1 in all samples. This could mean that the 372 sulfonated/unsulfonated TOL ratio that favored TOL-SO<sub>3</sub>H may also give rise to further products similar to 2-AAP.

374 The ILA  $(0.03 \text{ mg } L^{-1})$  and ILA-SO<sub>3</sub>H  $(8.53 \text{ mg } L^{-1})$  pairing showed similar behavior to the 375 previously described TOL and TOL-SO<sub>3</sub>H pairing. Arapitsas et al. [23], found that young red wines and rosé wines had a much higher concentration of ILA and especially ILA-GLU, which is synthesized by plants [40]. This was probably due to maceration with the skins during winemaking. In contrast, white and especially sparkling wines had the lowest concentrations due to soft pressing and secondary fermentation, which necessitate further nitrogen consumption.

381 Sulfonated indole 3-acetic acid was detected at concentrations between 0.13 and 0.39 mg  $L^{-1}$ , while its parent compounds were detectable at very low concentrations. Hoenicke et al. [41] showed that the sulfonation of indoles in a model wine solution could be responsible for their  $\mathbf{1}$  $\overline{2}$ 

 degradation and the formation of aromatic aminobenzenes such as 2-aminoacetophenone, which are responsible for some of the heavy aromatic characteristics of white wines. On other hand, the structural similarity of TOL to IAA could also cause the detection of 2-AAP or other similar aromatic compounds in wine. Because of the lack of anthocyanins and the low flavanol content in white wines, the sulfonation of indoles can increase the risk of developing an atypical off-flavor. Thus, the sparkling wines produced from the Ribolla Gialla variety might have a higher tendency to develop atypical ageing defects, which could also be promoted by inappropriate storage temperature.

*Metabolomic fingerprint of lipids*

int of lipids<br>nty-nine lipid compounds found in the same<br>HPLC-MS-MS. Most of the compounds for<br>ore than 12 carbon atoms), although the m<br>ir esters have a major influence on the org<br>that lipids are an integral part of solid Table 5 shows the twenty-nine lipid compounds found in the samples of commercial sparkling wines according to UHPLC-MS-MS. Most of the compounds found are saturated long-chain fatty acids (LCFAs, more than 12 carbon atoms), although the mid-chain fatty acids (MCFAs, 4–12 carbons) and their esters have a major influence on the organoleptic properties of wine. In addition to the fact that lipids are an integral part of solid grape tissues, they are also an important building block in wine yeasts, where the majority of LCFAs are esterified with glycerol or glycerophosphate to form mono-, di-, and tri-acylglycerides or glycerophospholipids, respectively [42].

 The two most abundant LCFAs detected in the samples were palmitic acid (C16:0), which 402 ranged from 5.21 to 11.49 mg  $L^{-1}$  with a mean value of 9.24 mg  $L^{-1}$ , and stearic acid (C18:0), 403 which ranged from 4.43 to 8.40 mg  $L^{-1}$  (mean value: 6.49 mg  $L^{-1}$ ). Previous studies showed that after the beginning of second fermentation, which is after the inoculation of the medium with yeast, the concentration of saturated fatty acids C16:0 slightly increased, while the proportion of C18:0 remained constant. However, after the end of the growth phase of the yeasts, the distributions of both saturated fatty acids decreased exponentially [43].

 Unlike saturated fatty acids (SFA), the values of unsaturated fatty acids (UFA) were lower in the wine samples (*cis*-11-eicosanoic acid 0.11 mg L-1, myristoleic acid 0.05 mg L-1, linoleic 410 acid 0.04 mg L<sup>-1</sup>, palmitoleic acid 0.02 mg L<sup>-1</sup>, and linolenic acid 0.01 mg L<sup>-1</sup>). UFAs also had a much smaller proportion than SFAs. Because of their membrane fluidity, wine yeasts are able to modify the UFA/SFA ratio in their membrane composition in response to stress conditions, such as the presence of ethanol or cooler fermentation conditions under which fermentation takes place, especially for white wines. Normally, the UFA/SFA ratio is close to 1:1, but a deficiency of UFA leads to impaired biosynthesis of phospholipids and consequently stuck

[44] studied the effect of the total contents<br>at best define foam stability in wines and<br>a positive correlation between these c<br>nat the wines with a greater amount of th<br>ore foam with high stability than those wit<br>in the p Similarly, Pueyo et al. [44] studied the effect of the total contents of linolenic acid and palmitic acid as compounds that best define foam stability in wines and foam height in the cavas, respectively. They found a positive correlation between these compounds and the measured properties, meaning that the wines with a greater amount of these fatty acids in the lipidic fraction could form more foam with high stability than those with lower concentrations. Since the analyzed samples in the present study contained a greater concentration of palmitic acid compared to the linolenic acid, it can be expected that foam formation of the Ribolla Gialla sparkling wines will be increased, which is a key parameter for determining the quality of

 In conclusion, Ribolla Gialla sparkling wines showed homogeneous characteristics concerning basic parameters (e.g., sugar content, alcoholic strength, pH, and titratable acidity) as well as their aroma composition. Sparkling Ribolla wines are generally characterized by low levels of free terpenols, and their aroma seems to be mostly characterized by the development of volatile esters and β-damascenone during fermentations, storage, and processing. The presence of these compounds is probably connected with the fresh and fruity notes that normally characterize the sparkling wines produced from Ribolla grapes.

Page 19 of 35

 $\mathbf{1}$ 

may lead to higher foam height in the state way quality features of sparkling wines in g<br>n tryptophan metabolites, Ribolla Gialla co<br>mas. Considering that this specific Italian p<br>rkets, further investigations should caref<br> A minor amount of malolactic fermentation was found, which reflects that two distinct wine styles coexist for Ribolla sparkling wines. The first characterizes young wines normally produced by refermentation in stainless-steel autoclaves (Martinotti/Charmat short refermentation method), and the second is based on a more or less prolonged ageing period on refermentation lees in an autoclave or bottles (according to the traditional refermentation method). The analysis of lipid molecules, tryptophan, and its metabolites highlighted other interesting features of Ribolla sparkling wines from a practical point of view. Ribolla Gialla shows a high ratio between saturated (e.g., palmitic acid) and unsaturated fatty acids (e.g., linolenic acid), which may lead to higher foam height in the sparkling wines produced and represents one of the key quality features of sparkling wines in general. In contrast, due to the high amounts of certain tryptophan metabolites, Ribolla Gialla could be prone to the formation of atypical ageing aromas. Considering that this specific Italian product has only recently been appearing in wine markets, further investigations should carefully consider these results to properly address the production practices and techniques towards producing high quality products, thus increasing the local and international competitiveness of Ribolla Gialla sparkling wines.

#### **Acknowledgments**

 This research was financially supported by Friuli Venezia Giulia Region (LR 29 Dec 2016, n. 25, art. 3, paragraphs 6-10. Project Title: Maturation parameters and optimization of the agronomic and enological techniques for the production of quality sparkling wines, Jul 2017 – Dec 2019) and by the ADP 2017 project and PhD scholarship (D.S.) funded by the Autonomous Province of Trento. The authors are grateful to Dr. Matteo Corradin for his cooperation in aroma analyses.

# **References**

- 1. Rusjan D, Jug T, Štajner N (2010) Evaluation of genetic diversity: Which of the varieties can be named "Rebula" (*Vitis vinifera* L.)? Vitis 49:189–192
- 2. De Lorenzis G, Imazio S, Rusjan D, Vouillamoz J F, Nikolaou N, Failla O, Scienza A
- (2013) Genetic investigation of grapevine varieties 'Ribolla Gialla' (Italy), 'Rebula' (Slovenia) and 'Robola' (Ionian Islands). Sci Hortic 150:425–431.
- 3. Imazio S, Scienza A, Brancadoro L, de Lorenzis G, Failla O (2016) Evidence for a Sympatric Origin of Ribolla gialla, Gouais Blanc and Schiava cultivars (*V. vinifera* L.). South African J Enol Vitic 35:149–156.
- m of Kibolia gialia, Gouais Blanc and Sch<br>Enol Vitic 35:149–156.<br>Prenzis G, Scienza A, Failla O, Vouillamoz<br>2014) 'Ribolla Gialla' from North Eastern<br>obola' from Ionian Islands; Do They Be<br>They Genetically Different? Acta 4. Imazio S, De Lorenzis G, Scienza A, Failla O, Vouillamoz J, Korosec-Koruza Z, Rusjan D, Nikolao N (2014) 'Ribolla Gialla' from North Eastern Italy, 'Rebula' from Northern Balkans and 'Robola' from Ionian Islands; Do They Belong to the Same Population Variety or Are They Genetically Different? Acta Hortic 1046:645–652.
- 5. Bavčar D, Baša Česnik H, Čuš F, Vanzo A, Gašperlin L, Košmerl T (2011) Impact of alternative skin contact procedures on the aroma composition of white wine. South African J Enol Vitic 32:190–203.
- 6. Bavčar D, Baša Česnik H, Čuš F, Košmerl T (2011) The influence of skin contact during alcoholic fermentation on the aroma composition of Ribolla Gialla and Malvasia Istriana Vitis vinifera (L.) grape wines. Int J Food Sci Technol 46:1801–1808.
- 7. Rapp A, Versini G (1995) Influence of nitrogen compounds in grapes on aroma compounds of wines. Dev Food Sci 37:1659–1694
- 8. Pueyo E, Martínez-Rodríguez A, Polo MC, Santa-María G, Bartolomé B (2000) Release of Lipids during Yeast Autolysis in a Model Wine System. J Agric Food Chem 46:116– 122.
	- 9. Swiegers JH, Bartowsky EJ, Henschke PA, Pretorius IS (2005) Yeast and bacterial modulation of wine aroma and flavour. Aust J Grape Wine Res 11:139-173.



Wine Foaming. J Agric Food Chem 50:704<br>
mat TJ, Steinhart H, Köhler HJ, Schwab A<br>
1 Indole-3-Acetic Acid, Tryptophan, and<br>
1 Wine. J Agric Food Chem 49:5494–5501<br>
114) Atypical Aging Defect: Sensory Discri<br>
Consequences. A 10. Callejón RM, Margulies B, Hirson GD, Ebeler SE (2012) Dynamic changes in volatile compounds during fermentation of Cabernet Sauvignon grapes with and without skins. Am J Enol Vitic 63:301–312. 11. Hirst MB, Richter CL (2016) Review of Aroma Formation through Metabolic Pathways of *Saccharomyces cerevisiae* in Beverage Fermentations. Am J Enol Vitic 67:361–370. 12. Waterhouse AL, Sacks GL, Jeffery DW (2016) Understanding Wine Chemistry. John Wiley & Sons, Chichester (UK). 13. Gallart M, Lopez-Tamames E, Suberbiola G, Buxaderas Susana (2002) Influence of Fatty Acids on Wine Foaming. J Agric Food Chem 50:7042–7045. 14. Hoenicke K, Simat TJ, Steinhart H, Köhler HJ, Schwab A (2001) Determination of Free and Conjugated Indole-3-Acetic Acid, Tryptophan, and Tryptophan Metabolites in Grape Must and Wine. J Agric Food Chem 49:5494–5501. 15. Schneider V (2014) Atypical Aging Defect: Sensory Discrimination, Viticultural Causes, and Enological Consequences. A Review. Am J Enol Vitic 65:277–284. 16. Dziadas M, Jeleń HH (2010) Analysis of terpenes in white wines using SPE-SPME- GC/MS approach. Anal Chim Acta 677:43–49. 17. Louw L, Tredoux AGJ, Rensburg PV, Kidd M, Naes T, Nieuwoudt HH (2010) Fermentation-derived Aroma Compounds in Varietal Young Wines from South Africa. South African J Enol Vitic 31:213–225. 18. D'Onofrio C, Matarese F, Cuzzola A (2017) Study of the terpene profile at harvest and during berry development of *Vitis vinifera* L. aromatic varieties Aleatico, Brachetto, Malvasia di Candia aromatica and Moscato bianco. J Sci Food Agric 97:2898–2907. 19. Vilanova M, Genisheva Z, Graña M, Oliveira JM (2013) Determination of Odorants in Varietal Wines from International Grape Cultivars (*Vitis vinifera*) Grown in NW Spain. South African J Enol Vitic 34:212–222.

20. Comuzzo P, Marconi M, Zanella G, Querzè M (2018) Pulsed electric field processing of

white grapes (cv. Garganega): Effects on wine composition and volatile compounds.

- Food Chem 264:16–23.
- 21. Loira I, Morata A, Comuzzo P, Callejo MJ, González C, Calderón F, Suárez-Lepe JA (2015) Use of *Schizosaccharomyces pombe* and *Torulaspora delbrueckii* strains in mixed and sequential fermentations to improve red wine sensory quality. Food Res Int 76:325–333.
- ed using solid-phase micro extraction<br>s chromatography coupled with time-of-fli<br>0.<br>ella G, Mattivi F (2018) The impact of SO<sub>2</sub><br>ne style and age. Sci Rep 8:858.<br>Chitarrini G, Di Gangi IM, Masuero D, Sc<br>LC–MS/MS method for 22. Carlin S, Vrhovsek U, Franceschi P, Lotti C, Bontempo L, Camin F, Toubiana D, Zottele F, Toller G, Fait A, Mattivi F (2016) Regional features of northern Italian sparkling wines, identified using solid-phase micro extraction and comprehensive two- dimensional gas chromatography coupled with time-of-flight mass spectrometry. Food Chem 208:68–80.
- 520 23. Arapitsas P, Guella G, Mattivi F (2018) The impact of  $SO_2$  on wine flavanols and indoles in relation to wine style and age. Sci Rep 8:858.
- 24. Della Corte A, Chitarrini G, Di Gangi IM, Masuero D, Soini E, Mattivi F, Vrhovsek U (2015) A rapid LC–MS/MS method for quantitative profiling of fatty acids, sterols, glycerolipids, glycerophospholipids and sphingolipids in grapes. Talanta 140:52–61.
- 25. Commission Regulation (EC) No 607/2009 (2009) Laying down certain detailed rules for the implementation of Council Regulation (EC) No 479/2008 as regards protected designations of origin and geographical indications, traditional terms, labeling and presentation of certain wine sector products. Off J Eur Communities 60–139.
- 26. Patynowski RJ, Jiranek V, Markides AJ (2002) Yeast viability during fermentation and sur lie ageing of a defined medium and subsequent growth of *Oenococcus oeni*. Aust J Grape Wine Res 8:62–69.
- 27. Lambrechts MG, Pretorius IS (2000) Yeast and its Importance to Wine Aroma A Review. South African J Enol Vitic 21:97-129.
	- 28. Fariña L, Villar V, Ares G, Carrau F, Dellacassa E, Boido E (2015) Volatile composition





 $\mathbf{1}$  $\overline{2}$ 



  For Perican

# **Figure Captions**

 **Figure 1.** Geographical position of Friuli Venezia Giulia region. Grey areas mark the municipalities in which the wineries that supplied wine samples are located. The number of samples/wineries for each municipality are also reported.

For Per Review 

 

 **Table 1. Chemical composition of Ribolla Gialla commercial wines. SD: standard deviation; MIN: minimum value; MAX: maximum value; CV: coefficient of variation.**



a n.d.: not detected

**Table 2. Non-varietal aroma compounds detected in Ribolla Gialla commercial sparkling wines by different analytical approaches. Concentrations** 

**are expressed in µg L-1 .** 



11

13

17

21

23 24 25

27

29 30 31

33 34 35

European Food Research and Technology





<sup>a</sup> AM: analytical method; <sup>b</sup> IM: identification method (S comparison of mass spectra and retention time with those of standard compounds; RI comparison of order of elution with those reported in literature; MS comparison of mass spectra with those reported in mass spectrum libraries); C SD: standard deviation; <sup>d</sup> MIN: minimum value; <sup>e</sup> MAX: maximum value; <sup>f</sup> CV: coefficient of variation; <sup>g</sup> ODT: odor detection threshold; <sup>h</sup> n.d.: not detected

**Table 3. Free and bound terpenes and norisoprenoids detected in Ribolla Gialla commercial sparkling wines by different analytical approaches.** 

**Concentrations are expressed in µg L-1 .** 





<sup>a</sup> AM: analytical method; <sup>b</sup> IM: identification method (S comparison of mass spectra and retention time with those of standard compounds; RI comparison of order of elution with those reported in literature; MS comparison of mass spectra with those reported in mass spectrum libraries); C SD: standard deviation; <sup>d</sup> MIN: minimum value; <sup>e</sup> MAX: maximum value; <sup>f</sup> CV: coefficient of variation; <sup>g</sup> ODT: odor detection threshold; <sup>h</sup> n.d.: not detected CV: coefficient of variation; <sup>g</sup>

SPE MS RI 16  $\pm$  2<br>
SPE MS RI 16  $\pm$  2<br>
method (S comparison of mass spectra and retention time<br>
rature; MS comparison of mass spectra with those repeatimum value; f CV: coefficient of variation; <sup>g</sup> ODT: odo:

606



607 **Table 4. Tryptophan metabolites detected in Ribolla Gialla commercial sparkling wines.** 

608 **Concentrations are expressed in mg L-1. SD: standard deviation; MIN: minimum value;** 

## 609 **MAX: maximum value; CV: coefficient of variation.**



**a** ILA-SO <sup>3</sup>H: indole-lactic acid-2-sulfonate; TOL-SO <sup>3</sup>H: tryptophol-2-sulfonate; TYL: tryptophol; TYR: tyrosine; ILA-GLU-SO <sup>3</sup>H: sulfonated indole-3-lactic acid; PHE: phenylalanine; TRP-EE: tryptophan-ethyl ester; TOL: tryptophol; ABA: abscisic acid; IAA-SO <sup>3</sup>H: sulfonated indole 3-acetic acid; ABA-GLU: glucoside of abscisic acid; TRP: tryptophan; ILA-GLU: indole 3-lactic acid glucoside; TYR-EE: tyrosine-ethyl ester; ILA: indole 3-lactic acid; KYNA: kynurenic acid; AA: anthranilic acid; IAA-ASP: indole 3 acetic acid conjugate with aspartic acid; N-TYR-EE: N-acetyl-tyrosine-ethyl ester; KYN: kynurenine; IAA: indole 3-acetic acid; N-TRP-EE: N-acetyl-tryptophan-ethyl ester.

b n.d.: not detected

610

611

**Table 5. Lipid compounds detected in Ribolla Gialla commercial sparkling wines.** 

**Concentrations are expressed in mg L-1. SD: standard deviation; MIN: minimum value;** 

**MAX: maximum value; CV: coefficient of variation.**



