



UNIVERSITÀ
DEGLI STUDI
DI UDINE

Università degli studi di Udine

Polish metric spaces with fixed distance set

Original

Availability:

This version is available <http://hdl.handle.net/11390/1191393> since 2021-03-22T18:25:00Z

Publisher:

Published

DOI:10.1016/j.apal.2020.102832

Terms of use:

The institutional repository of the University of Udine (<http://air.uniud.it>) is provided by ARIC services. The aim is to enable open access to all the world.

Publisher copyright

(Article begins on next page)

POLISH METRIC SPACES WITH FIXED DISTANCE SET

RICCARDO CAMERLO, ALBERTO MARCONE, AND LUCA MOTTO ROS

ABSTRACT. We study Polish spaces for which a set of possible distances $A \subseteq \mathbb{R}^+$ is fixed in advance. We determine, depending on the properties of A , the complexity of the collection of all Polish metric spaces with distances in A , obtaining also example of sets in some Wadge classes where not many natural examples are known. Moreover we describe the properties that A must have in order that all Polish spaces with distances in that set belong to a given class, such as zero-dimensional, locally compact, etc. These results lead us to give a fairly complete description of the complexity, with respect to Borel reducibility and again depending on the properties of A , of the relations of isometry and isometric embeddability between these Polish spaces.

CONTENTS

1. Introduction	1
2. Preliminaries	4
3. Distance sets of particular Polish metric spaces	8
4. The complexity of \mathcal{M}_A and \mathcal{M}_A^*	10
5. Isometry and isometric embeddability	15
5.1. Isometry	17
5.2. Isometric embeddability	21
6. Open problems	26
References	26

1. INTRODUCTION

A common problem in mathematics is to classify interesting objects up to some natural notion of equivalence. More precisely, one considers a class of objects X and an equivalence relation E on X , and tries to find a set of complete invariants I for (X, E) . To be of any use, such an assignment of invariants should be as simple as possible. In most cases, both X and I carry some intrinsic Borel structures, so that it is natural to ask the assignment to be a Borel measurable map.

Date: May 12, 2020.

2010 Mathematics Subject Classification. Primary: 03E15; Secondary: 54E50.

Key words and phrases. Borel reducibility; Polish spaces; isometry; isometric embeddability.

Camerlo's research was partially carried out while visiting the Équipe de logique at the Université de Lausanne. Marcone's research was supported by PRIN 2012 Grant "Logica, Modelli e Insiemi" and departmental PRID funding "HiWei — The higher levels of the Weihrauch hierarchy". Motto Ros was supported by the Young Researchers Program "Rita Levi Montalcini" 2012 through the project "New advances in Descriptive Set Theory".

A classical example is the problem of classifying separable complete metric spaces, called *Polish metric spaces*, up to isometry. In [Gro99] Gromov showed for instance that one can classify compact Polish metric spaces using (essentially) elements of \mathbb{R} as complete invariants; in this case, one says that the corresponding classification problem is smooth. However, as pointed out by Vershik in [Ver98] the problem of classifying arbitrary Polish metric spaces is «an enormous task», in particular it is far from being smooth. Thus it is natural to ask “how complicated” is such a classification problem.

A natural tool for studying the complexity of classification problems is the notion of Borel reducibility introduced in [FS89] and in [HKL90]: we say that a classification problem (X, E) is *Borel reducible* to another classification problem (Y, F) (in symbols, $E \leq_B F$) if there exists a Borel measurable function $f: X \rightarrow Y$ such that $x E x' \iff f(x) F f(x')$ for all $x, x' \in X$. Intuitively, this means that the classification problem (X, E) is not more complicated than (Y, F) : in fact, any assignment of complete invariants for (Y, F) may be turned into an assignment for (X, E) by composing with f . A comprehensive reference for the theory of Borel reducibility is [Gao09].

In the seminal [GK03] (see also [CGK01, Cle12]), Gao and Kechris were able to determine the exact complexity of the classification problem for isometry on arbitrary Polish metric spaces with respect to Borel reducibility: it is Borel bireducible with the most complex orbit equivalence relation (so every equivalence relation induced by a Borel action of a Polish group on a Polish space Borel reduces to it). However they left the open problems of establishing the complexity of isometry on locally compact ultrametric and zero-dimensional Polish spaces. We have been able to solve the first of these questions in [CMMR18] using an approach that goes back to Clemens [Cle07] and Gao and Shao [GS11]: Clemens studied the complexity of isometry on the collection of Polish metric spaces using only distances in a set $A \subseteq \mathbb{R}^+$ fixed in advance, while Gao and Shao considered the restriction of Clemens’ problem to ultrametric Polish spaces.

We answered the questions left open by Gao and Shao in [CMMR18], where we focused on the study of ultrametric Polish spaces with a fixed set of distances and, as a byproduct, we showed that isometry on locally compact (and even discrete) ultrametric Polish spaces is Borel bireducible with countable graph isomorphism. In this paper we instead settle various problems, or provide new proofs for known results, about *arbitrary* Polish metric spaces with a fixed set of distances.

Let $\mathbb{R}^+ = \{r \in \mathbb{R} \mid r \geq 0\}$. Let (X, d_X) be a Polish metric space, i.e. a separable space with a complete metric d_X (which often is left implicit). We denote by $D(X)$ the set of distances that are realized in X , i.e.

$$D(X) = \{r \in \mathbb{R}^+ \mid \exists x, y \in X (d_X(x, y) = r)\}.$$

All metric spaces X we consider are always assumed to be nonempty, so that $0 \in D(X)$.

Definition 1.1. We say that $A \subseteq \mathbb{R}^+$ is a *distance set* if $A = D(X)$ for some Polish metric space X . When $A = D(X)$ we say that A is *realized by X* . Let \mathcal{D} denote the set of all distance sets $A \subseteq \mathbb{R}^+$.

Clemens characterized the members of \mathcal{D} in his PhD thesis.

Theorem 1.2 ([Cle01, Theorem 4.3]). *Let $A \subseteq \mathbb{R}^+$. Then A is a distance set if and only if A is analytic, $0 \in A$, and either A is countable or 0 is a limit point of A .*

Clemens studied also for which $A \in \mathcal{D}$, given a property of Polish spaces (like being locally compact, or σ -compact, or discrete, or countable, and so on) some Polish metric space with this property has distance set A (see Theorem 3.1 below). Here we consider the following dual question:

Question 1. *For which $A \in \mathcal{D}$ every Polish metric space with distance set A has a given property?*

We answer this question in Section 3, and in particular in Theorem 3.6. Our results show in particular that lower bounds for the complexity of the restriction of isometry to zero-dimensional Polish metric spaces (one of the problems of Gao and Kechris) can be obtained by classifying the restriction of isometry to spaces with a fixed distance set which is dense in some right neighborhood of 0 but does not contain any such neighborhood.

Another natural question is the following:

Question 2. *Given $A \in \mathcal{D}$, what is the complexity of the collection \mathcal{M}_A^* of Polish metric spaces having distance set A ? What about the collection \mathcal{M}_A of Polish metric spaces having distance set included in A (in which case we can drop the requirement $A \in \mathcal{D}$)?*

This (and Question 3 below) requires to view Polish metric spaces as members of some hyperspace of all Polish metric spaces: we describe the set-up in Section 2 and answer quite satisfactorily Question 2 in Section 4. In particular Theorems 4.5(2) and 4.8(1) characterize when \mathcal{M}_A and \mathcal{M}_A^* are standard Borel. Tables 1 and 2 summarize our results for the complexity of \mathcal{M}_A and \mathcal{M}_A^* when $A \in \mathcal{D}$.

As a corollary, in Theorem 4.14 we also extend the characterization of the distance sets A which admit an A -Urysohn space obtained by Sauer [Sau13]¹.

The last main questions we deal with are the original motivation for this research:

Question 3. *Given $A \in \mathcal{D}$, what is the complexity with respect to Borel reducibility of isometry and isometric embeddability restricted to \mathcal{M}_A^* (denoted respectively \cong_A^* and \sqsubseteq_A^*)? What about the same problem for isometry and isometric embeddability restricted to \mathcal{M}_A (denoted respectively \cong_A and \sqsubseteq_A)?*

We study this question in Section 5, and our main results include the following:

- The fact that \sqsubseteq_A and \sqsubseteq_A^* have the same complexity for all $A \in \mathcal{D}$ (Corollary 5.13) and that \cong_A and \cong_A^* have the same complexity when A is countable (Theorem 5.10).
- The classification with respect to Borel reducibility of \cong_A^* , depending on the properties of A (Theorem 5.7). In particular, we characterize when countable graph isomorphism Borel reduces to \cong_A^* (Theorem 5.6).
- The exhaustive description of the complexity with respect to Borel reducibility of \sqsubseteq_A^* (Theorem 5.12).
- The fact that whenever \sqsubseteq_A^* is complete analytic, it has also the stronger property of being invariantly universal (Theorem 5.20).

¹We thank Joseph Zielinski for directing us to Sauer's paper.

The first item substantially enriches the picture obtained by Clemens in [Cle07], almost completely solving his original problem about isometry. We also answer some of the other questions contained in [Cle07] (Proposition 5.1, Theorem 5.8 and Theorem 5.10), and their analogues concerning isometric embeddability (Proposition 5.1 and Corollary 5.13).

2. PRELIMINARIES

If \mathcal{A} is a countably generated σ -algebra of subsets of X that separates points we refer to the members of \mathcal{A} as Borel sets (indeed, as shown e.g. in [Kec95, Proposition 12.1], in this case \mathcal{A} is the collection of Borel sets of some separable metrizable topology on X), and to (X, \mathcal{A}) as a Borel space. The Borel space (X, \mathcal{A}) is standard if \mathcal{A} is the collection of Borel sets of some Polish (i.e. separable and completely metrizable) topology on X . A map between two Borel spaces is Borel if the preimages of Borel sets of the target space are Borel sets of the domain.

We denote by $\Sigma_1^1(X)$ the family of subsets of the standard Borel spaces X which are Borel images of a standard Borel space. For $n > 0$, $\Pi_n^1(X)$ is the class of all complements of sets in $\Sigma_n^1(X)$, and $\Sigma_{n+1}^1(X)$ is the family of Borel images of a set in $\Pi_n^1(Y)$ for some standard Borel space Y . We have $\Sigma_n^1 \cup \Pi_n^1 \subseteq \Sigma_m^1 \cap \Pi_m^1$ whenever $n < m$, and for uncountable standard Borel spaces the inclusion is strict. This hierarchy is the *projective hierarchy*. Σ_1^1 and Π_1^1 sets are called resp. *analytic* and *coanalytic* sets. The class of differences of two analytic subsets (equivalently: of intersections of an analytic and a coanalytic subset) of X is denoted $D_2(\Sigma_1^1)(X)$.

We extend these notions also to Borel spaces X which are not necessarily standard. In particular we say that $A \subseteq X$ is analytic (or Σ_1^1) if there exists a standard Borel space $Y \supseteq X$ such that the Borel subsets of X are the intersections of the Borel subsets of Y with X and A is the intersection of some $B \in \Sigma_1^1(X)$ with X .

If Γ is a class of sets in Borel spaces closed under Borel preimages (like Σ_n^1 and Π_n^1), Y is a standard Borel space and $A \subseteq Y$, we say that A is *Borel Γ -hard* if for every $B \in \Gamma(X)$, where X is a standard Borel space, is *Borel Wadge reducible* to A , i.e. there exists a Borel function $f : X \rightarrow Y$ such that $f^{-1}(A) = B$. We say that A is *Borel Γ -complete* if, in addition, $A \in \Gamma(Y)$. If A is Borel Γ -hard and A is Borel Wadge reducible to B , then B is Borel Γ -hard as well: this is the typical way to prove hardness results.

The classes Γ we are interested in are closed under Borel preimages and such that either Γ or its dual $\check{\Gamma}$ (consisting of the complements of the elements of Γ) is closed under intersection with Π_1^1 sets. For these classes and any Polish topology compatible with the standard Borel spaces, Borel Γ -hardness can be witnessed by continuous functions: see [Kec97], where this fact is stated for the class Π_1^1 , but the argument actually works under our more general assumptions on Γ . Therefore Borel Γ -hardness and Borel Γ -completeness coincide with Γ -hardness and Γ -completeness, which are the notions used when dealing with Wadge reducibility. For this reason we drop Borel from this terminology.

Most results in Section 4 state that a collection of Polish metric spaces is Γ -complete for some Γ , and thus pinpoint the complexity of that particular collection by showing that it belongs to Γ and not to any simpler class. When $\Gamma \neq \check{\Gamma}$, this implies in particular that such a collection does not belong to $\check{\Gamma}$.

Borel Wadge reducibility can be generalized from sets to binary (and in fact, n -ary for any n) relations as follows. Let R and S be binary relations on Borel spaces X and Y , respectively. We say that R is *Borel reducible* to S , and we write $R \leq_B S$, if there is a Borel function $f : X \rightarrow Y$ such that $x R x'$ if and only if $f(x) S f(x')$ for all $x, x' \in X$. If $R \leq_B S$ and $S \leq_B R$ we say that R and S are *Borel bireducible* and we write $R \sim_B S$. If on the other hand we have $R \leq_B S$ and $S \not\leq_B R$ we write $R <_B S$.

If Γ is a class of binary relations on standard Borel spaces and $S \in \Gamma$, we say that S is *complete for Γ* if $R \leq_B S$ for all $R \in \Gamma$. Some relevant classes Γ one might consider are the collection of all analytic equivalence relations and the collection of all analytic quasi-orders. (Recall that a quasi-order is a reflexive and transitive binary relation.) An example of a complete quasi-order for the latter class is embeddability between countable graphs, see [LR05]. When Γ is the class of orbit equivalence relations (that is, those Borel reducible to a relation induced by a Borel action of a Polish group on a standard Borel space) a complete element for Γ is isometry on arbitrary Polish spaces, see [GK03]. Another important example is the class of equivalence relations classifiable by countable structures, that is those Borel reducible to isomorphism on countable structures. The canonical example of an equivalence relation complete for this class is countable graph isomorphism, see [FS89].

We reserve the term “complete for Γ ” for relations defined on some standard Borel space. In Section 5, we often consider analytic relations on Borel spaces which (by the results of Section 4) are not standard. In these cases we rather state that a relation is *Borel bireducible with a complete for Γ relation*.

We denote by \cong and \sqsubseteq the relations of isometry and isometric embeddability between metric spaces. Recall that a metric space is Polish if and only if it is isometric to an element of $F(\mathbb{U})$, the collection of all nonempty closed subsets of the Urysohn space \mathbb{U} (here we differ slightly from [Kec95], where $F(\mathbb{U})$ includes the empty set). The space $F(\mathbb{U})$ is endowed with the Effros Borel structure, which turns it into a standard Borel space: the hyperspace containing all Polish metric spaces up to isometry. Notice that \cong and \sqsubseteq are analytic relations on $F(\mathbb{U})$. We fix also a sequence of Borel functions $(\psi_n)_{n \in \omega}$ from $F(\mathbb{U})$ into \mathbb{U} such that $\{\psi_n(X) \mid n \in \omega\}$ is dense in X for every $X \in F(\mathbb{U})$, see [Kec95, Theorem 12.13].

Remark 2.1. Another possible coding for Polish metric spaces (used e.g. in [Cle07]) is sometimes convenient. In this approach a Polish metric space U is coded by an element M of a suitable \mathcal{M} , which is a closed subset of ${}^{\omega \times \omega} \mathbb{R}$: U is the completion of a set of points $\{x_i \mid i \in \omega\}$ such that the distance between x_i and x_j equals $M(i, j)$. As explained in [MR17, Section 2], this coding is equivalent to the one introduced above, in the sense that there are Borel functions $\Phi : F(\mathbb{U}) \rightarrow \mathcal{M}$ and $\Psi : \mathcal{M} \rightarrow F(\mathbb{U})$ such that $\Phi(X)$ codes a space isometric to X and $\Psi(M)$ is isometric to the space coded by M . Therefore the results can be transferred between the two settings. In particular, it is often easier to check that certain maps are Borel-measurable using \mathcal{M} rather than $F(\mathbb{U})$ (see e.g. the proof of Proposition 3.3).

Using the coding in $F(\mathbb{U})$ we have the following formalizations of the collections of Polish metric spaces using a prescribed set of distances.

Definition 2.2. Given $A \subseteq \mathbb{R}^+$, let

$$\mathcal{M}_A = \{X \in F(\mathbb{U}) \mid D(X) \subseteq A\} \text{ and } \mathcal{M}_A^* = \{X \in F(\mathbb{U}) \mid D(X) = A\}.$$

Let the equivalence relations \cong_A and \cong_A^* and the quasi-orders \sqsubseteq_A and \sqsubseteq_A^* be the restrictions of isometry and isometric embeddability to \mathcal{M}_A and \mathcal{M}_A^* .

The relations \cong_A , \cong_A^* , \sqsubseteq_A , and \sqsubseteq_A^* are defined on Borel spaces which are not necessarily standard (not even when A is countable, as we will show). We will discuss the complexity of \mathcal{M}_A and \mathcal{M}_A^* at length in Section 4.

In [CMMR18] we studied the restrictions of isometry and isometric embeddability to

$$\mathcal{U}_A = \{X \in \mathcal{M}_A \mid d_X \text{ is an ultrametric}\} \text{ and } \mathcal{U}_A^* = \{X \in \mathcal{M}_A^* \mid d_X \text{ is an ultrametric}\}.$$

By Theorem 3.1 below the latter is nonempty exactly when $A \in \mathcal{D}$ is countable. In contrast with the results of Section 4, \mathcal{U}_A and \mathcal{U}_A^* are both standard Borel spaces (see [CMMR18, Proposition 4.5]). We first considered the case of A ill-founded (with respect to the standard ordering of the reals) and, extending results of [GK03], proved the following.

Theorem 2.3 ([CMMR18, Corollary 5.7, and Theorems 6.3 and 6.4]). *Let $A \in \mathcal{D}$ be countable and ill-founded. Then:*

- (1) *Isometry on \mathcal{U}_A and on \mathcal{U}_A^* are both complete for equivalence relations classifiable by countable structures.*
- (2) *Isometric embeddability on \mathcal{U}_A and on \mathcal{U}_A^* are both complete for analytic quasi-orders.*

Then we dealt with well-founded A 's. Lemma 4.11 of [CMMR18] implies that the complexities of isometry and isometric embeddability on \mathcal{U}_A and \mathcal{U}_A^* for $A \in \mathcal{D}$ well-founded depend only on the order type of A . If α with $1 \leq \alpha < \omega_1$ is the order type of A we can then write, as in [CMMR18], \cong_α , \cong_α^* , \sqsubseteq_α , and \sqsubseteq_α^* in place of $\cong \upharpoonright \mathcal{U}_A$, $\cong \upharpoonright \mathcal{U}_A^*$, $\sqsubseteq \upharpoonright \mathcal{U}_A$ and $\sqsubseteq \upharpoonright \mathcal{U}_A^*$, respectively. Our results include the following.

Theorem 2.4 ([CMMR18, Lemma 5.8, and Theorems 5.15 and 6.11]).

- (1) *For every α such that $1 \leq \alpha < \omega_1$ we have $\cong_\alpha \sim_B \cong_\alpha^*$ and these equivalence relations are classifiable by countable structures.*
- (2) *The relations \cong_α , for $1 \leq \alpha < \omega_1$, form a strictly increasing chain under \leq_B of Borel equivalence relations which is cofinal below countable graph isomorphism (i.e. cofinal among Borel equivalence relations classifiable by countable structures).*
- (3) *For every α such that $1 \leq \alpha < \omega_1$ we have $\sqsubseteq_\alpha \sim_B \sqsubseteq_\alpha^*$.*
- (4) *Let $1 \leq \alpha < \omega_1$. Then*
 - (i) *if $\alpha \leq \omega$, then \sqsubseteq_α is Borel;*
 - (ii) *if $\alpha > \omega$, \sqsubseteq_α contains both upper and lower cones that are Σ_1^1 -complete, and hence \sqsubseteq_α is analytic non-Borel;*
 - (iii) *all classes of the equivalence relation induced by $\sqsubseteq_{\omega+1}$ are Borel, hence $\sqsubseteq_{\omega+1}$ is not complete for analytic quasi-orders;*
 - (iv) *for all $\alpha < \beta \leq \omega + 2$, $\sqsubseteq_\alpha <_B \sqsubseteq_\beta$.*

The problem of establishing the exact complexity of \sqsubseteq_α for $\alpha \geq \omega + 2$ is still open.

Some basic tools to change from a set of distances to another one are *metric preserving functions*, i.e. functions $f: A \rightarrow \mathbb{R}^+$ such that for every metric d on a space X with range contained in A we have that $f \circ d$ is still a metric on X . There is a vast literature about metric preserving functions defined on the whole

\mathbb{R}^+ (see [Dob95, Cor99] for surveys). Since we are dealing with Polish metric spaces we introduce the following definition, where we consider functions with a possibly proper subset of \mathbb{R}^+ as domain.

Definition 2.5. A function $f: A \rightarrow \mathbb{R}^+$ is *Polish metric preserving* if for every complete metric d with range contained in A on a Polish space X we have that $(X, f \circ d)$ is still a Polish metric space.

Proposition 2.6. A function $f: A \rightarrow \mathbb{R}^+$ is *Polish metric preserving if and only if it is metric preserving and for every sequence $(x_n)_{n \in \omega}$ in A ,*

$$(2.1) \quad \lim_{n \rightarrow \infty} x_n = 0 \quad \text{if and only if} \quad \lim_{n \rightarrow \infty} f(x_n) = 0.$$

Proof. Let f be Polish metric preserving. First we prove that f is metric preserving. Let (X, d) be a metric space with distances in A : if $(X, f \circ d)$ were not a metric space this would be witnessed on a subset $X' \subseteq X$ of size two or three; then (X', d) is Polish but $(X', f \circ d)$ is not even a metric space.

Fix now a sequence $(x_n)_{n \in \omega}$ in A . If $(x_n)_{n \in \omega}$ converges to 0 but $(f(x_n))_{n \in \omega}$ does not, then it can be assumed that $(x_n)_{n \in \omega}$ is strictly decreasing and $(f(x_n))_{n \in \omega}$ is bounded away from 0. Let d the metric on ${}^\omega\omega$ defined by letting, for distinct $\alpha, \beta \in {}^\omega\omega$, $d(\alpha, \beta) = x_n$ where n is least such that $\alpha(n) \neq \beta(n)$. Then $({}^\omega\omega, d)$ is a Polish metric space, while in $({}^\omega\omega, f \circ d)$ every point is isolated, so $({}^\omega\omega, f \circ d)$ is not separable and hence not Polish. Conversely, if $(x_n)_{n \in \omega}$ does not converge to 0 but $(f(x_n))_{n \in \omega}$ does, it can be assumed that $(x_n)_{n \in \omega}$ is bounded away from 0. Let $X = \{x_n \mid n \in \omega\}$, and define the distance d on X by letting $d(x_n, x_m) = \max\{x_n, x_m\}$ if $x_n \neq x_m$. Then (X, d) is a discrete Polish ultrametric space. Then the sequence $(x_n)_{n \in \omega}$ is a Cauchy sequence in $(X, f \circ d)$ but it does not converge, as $(X, f \circ d)$ is discrete.

Assume now that f is metric preserving and that condition (2.1) holds for every sequence $(x_n)_{n \in \omega}$ in A . This means that if (X, d) is a metric space with distances in A , then the identity is a homeomorphism between (X, d) and $(X, f \circ d)$. In particular, if (X, d) is Polish, then $(X, f \circ d)$ is separable; to conclude that $(X, f \circ d)$ is complete too, notice that a sequence in X is d -Cauchy if and only if it is $f \circ d$ -Cauchy. \square

A Polish metric preserving function $f: A \rightarrow A'$ transforms a space $(X, d) \in \mathcal{M}_A$ into the space $(X, f \circ d) \in \mathcal{M}_{A'}$, which is homeomorphic to (X, d) via the identity function by the previous characterization. The following proposition ensures that this transformation is always Borel.

Proposition 2.7. *Let $A \in \mathcal{D}$. Every Polish metric preserving $f: A \rightarrow \mathbb{R}^+$ is Borel-measurable.*

Proof. Fix $(X, d) \in \mathcal{M}_A^*$ and a countable dense $D = \{x_i \mid i \in \omega\} \subseteq X$. Let $r_{i,j} = d(x_i, x_j)$ and $r'_{i,j} = f(r_{i,j})$. Define $F \subseteq \mathbb{R}^+ \times \mathbb{R}^+$ by setting $(a, b) \in F$ if and only if

$$\begin{aligned} \exists (i_k)_{k \in \omega}, (j_k)_{k \in \omega} [(x_{i_k})_{k \in \omega} \text{ and } (x_{j_k})_{k \in \omega} \text{ are Cauchy sequences in } X \\ \wedge \lim r_{i_k, j_k} = a \wedge \lim r'_{i_k, j_k} = b]. \end{aligned}$$

The set F is clearly analytic. Since id_X is a homeomorphism between (X, d) and $(X, f \circ d)$, the set F is the graph of f . Using the fact that Souslin's Theorem holds for analytic spaces ([Kec95, Exercise 28.3]), we have that the proof of [Kec95,

Theorem 14.12] shows that functions on analytic spaces with analytic graphs are Borel. Therefore, since A is analytic by Theorem 1.2, we have that f is Borel. \square

3. DISTANCE SETS OF PARTICULAR POLISH METRIC SPACES

Beside proving Theorem 1.2 characterizing distance sets, Clemens also characterized the $A \in \mathcal{D}$ which can be realized by Polish metric spaces in a given class.

Theorem 3.1 ([Cle01]). *Let $A \in \mathcal{D}$. Then:*

- (1) \mathcal{M}_A^* always contains a zero-dimensional Polish metric space;
- (2) \mathcal{M}_A^* contains a Polish ultrametric space if and only if it contains a discrete Polish metric space, if and only if A is countable;
- (3) \mathcal{M}_A^* contains a connected Polish metric space if and only if it contains a path-connected Polish metric space if and only if A is an interval with left endpoint 0;
- (4) \mathcal{M}_A^* contains a compact Polish metric space if and only if A is compact and either it is finite or it has 0 as limit point;
- (5) \mathcal{M}_A^* contains a locally compact Polish metric space if and only if it contains a σ -compact Polish metric space, if and only if A is either countable or it is σ -compact and has 0 as limit point.

We now consider the dual problem of determining when $A \in \mathcal{D}$ is realized *only* by Polish metric spaces in a given class. We need the following construction, which will be used repeatedly throughout the paper.

Let (X, d_X) and (Y, d_Y) be metric spaces. Given two points $\bar{x} \in X$ and $\bar{y} \in Y$ and a real $r > 0$ we can extend the metrics d_X and d_Y to the disjoint union $Z = X \cup Y$ by setting $d_Z(x, y) = \max\{d_X(x, \bar{x}), d_Y(y, \bar{y}), r\}$ for $x \in X$ and $y \in Y$.

Lemma 3.2. *The function d_Z defined above is a metric. Moreover (Z, d_Z) is Polish whenever (X, d_X) and (Y, d_Y) are Polish.*

Proof. To prove that d_Z is a metric, we just need to check $d_Z(a, b) \leq d_Z(a, c) + d_Z(c, b)$ for distinct points $a, b, c \in X \cup Y$. If $a, b, c \in X$ or $a, b, c \in Y$ this is trivial.

Now assume $a, b \in X$ and $c \in Y$: then $d_Z(a, b) = d_X(a, b) \leq d_X(a, \bar{x}) + d_X(\bar{x}, b) \leq d_Z(a, c) + d_Z(c, b)$. The case $a, b \in Y$ and $c \in X$ is symmetric.

Assume now $a \in X$ and $b \in Y$ (the symmetric case is analogous). We distinguish three cases.

- Assume $d_Z(a, b) = r$. If $c \in X$ then $d_Z(c, b) \geq r$, while if $c \in Y$ then $d_Z(a, c) \geq r$. In both cases $d_Z(a, b) = r \leq d_Z(a, c) + d_Z(c, b)$.
- Assume $d_Z(a, b) = d_X(a, \bar{x})$. If $c \in X$ then $d_Z(a, b) = d_X(a, \bar{x}) \leq d_X(a, c) + d_X(c, \bar{x}) \leq d_Z(a, c) + d_Z(c, b)$. If instead $c \in Y$ then $d_X(a, \bar{x}) \leq d_Z(a, c)$, whence $d_Z(a, b) = d_X(a, \bar{x}) \leq d_Z(a, c) + d_Z(c, b)$.
- The case $d_Z(a, b) = d_Y(b, \bar{y})$ is similar to the previous one.

Polishness is preserved because every Cauchy sequence in Z is eventually contained either in X or in Y , so the construction does not add new limits of Cauchy sequences. \square

We denote the metric space (Z, d_Z) by $X \oplus_r Y$, omitting reference to \bar{x} and \bar{y} because the choice of these two points will be irrelevant in most of our applications. The following proposition shows that given $r > 0$, the map $F(\mathbb{U}) \times F(\mathbb{U}) \rightarrow F(\mathbb{U})$ sending $X, Y \in F(\mathbb{U})$ to $X \oplus_r Y$ may be construed as a Borel function.

Proposition 3.3. *There is a Borel-measurable function $f : F(\mathbb{U}) \times F(\mathbb{U}) \rightarrow F(\mathbb{U})$ such that $f(X, Y)$ is isometric to $X \oplus_r Y$ for some choice of the gluing points $\bar{x} \in X$ and $\bar{y} \in Y$.*

Proof. This is immediate using Remark 2.1, as it is easy to define a Borel counterpart of this function from $\mathcal{M} \times \mathcal{M}$ to \mathcal{M} . \square

The most important property of this construction is the following:

Fact 3.4. *For every r we have $D(X \oplus_r Y) = D(X) \cup D(Y) \cup \{r\}$.*

Definition 3.5. $A \in \mathcal{D}$ is *well-spaced* if $r < r'$ implies $2r < r'$ for all $r, r' \in A$.

Notice that if A is well-spaced and infinite then $A \setminus \{0\}$ is either a decreasing sequence converging to 0, an unbounded increasing sequence, or the union of these two. This follows from the fact that if A is well-spaced then for $n \in \mathbb{Z}$ the set $A \cap [2^n, 2^{n+1})$ contains at most one point.

Theorem 3.6. *Let $A \in \mathcal{D}$.*

- (1) *All spaces in \mathcal{M}_A^* are zero-dimensional if and only if A does not contain a right neighborhood of 0.*
- (2) *All spaces in \mathcal{M}_A^* are ultrametric if and only if A is well-spaced.*
- (3) *All spaces in \mathcal{M}_A^* are discrete if and only if they are all locally compact, if and only if they are all σ -compact, if and only if 0 is isolated in A .*
- (4) *All spaces in \mathcal{M}_A^* are connected if and only if they are all compact, if and only if they are all singletons, if and only if $A = \{0\}$.*

All the above characterizations remain true if we replace \mathcal{M}_A^ with \mathcal{M}_A .*

Proof. To prove all forward directions we construct spaces of the form $X \oplus_r Y$ with $r \in A$, $X \in \mathcal{M}_A^*$, and $Y \in \mathcal{M}_A$ lacking the relevant topological properties. We always have $X \oplus_r Y \in \mathcal{M}_A^*$ by Fact 3.4.

(1) Suppose that A contains an interval $[0, r]$ for some $r > 0$. Fix $X \in \mathcal{M}_A^*$ and let $Y = [0, r] \subseteq \mathbb{R}$ with the usual metric. Then $X \oplus_r Y$ belongs to \mathcal{M}_A^* but is not zero-dimensional.

Conversely, if 0 is a limit point of $\mathbb{R}^+ \setminus A$, then for any $X \in \mathcal{M}_A^*$ the collection of balls with radius in $\mathbb{R}^+ \setminus A$ is a clopen basis for X .

(2) Recall that a space is ultrametric if and only if every triangle is isosceles with legs not shorter than the base. Suppose that A is not well-spaced and pick $r, r' \in A$ with $r < r' \leq 2r$. Fix $X \in \mathcal{M}_A^*$ and let Y be a triangle with two sides of length r and one of length r' . Then $X \oplus_r Y$ belongs to \mathcal{M}_A^* but is not ultrametric.

Conversely, if $X \in \mathcal{M}_A^*$ is not ultrametric, then it must contain a triangle with sides of length $r'' \leq r < r'$. Then by the triangle inequality $r' \leq r'' + r \leq 2r$, and hence A is not well-spaced.

(3) Suppose that A contains a decreasing sequence $(r_n)_{n \in \omega}$ converging to 0. Fix again $X \in \mathcal{M}_A^*$ and let Y be the Baire space ${}^\omega\omega$ equipped with the metric defined by $d_Y(y, y') = r_n$ where n is least such that $y(n) \neq y'(n)$. Then the space $X \oplus_{r_0} Y$ belongs to \mathcal{M}_A^* but is not σ -compact (here we are also using the fact that Y is closed in $X \oplus_{r_0} Y$), and hence, by separability, neither locally compact nor discrete.

Conversely, if 0 is isolated in A , then any $X \in \mathcal{M}_A^*$ is discrete, and thus also locally compact and σ -compact.

(4) Suppose that $r > 0$ belongs to A and $X \in \mathcal{M}_A^*$. Let Y be the countable space with all distinct points at distance r : then $X \oplus_r Y$ belongs to \mathcal{M}_A^* but is neither connected nor compact. The other implications are obvious.

Finally, consider the statements for \mathcal{M}_A . The forward directions follow from $\mathcal{M}_A^* \subseteq \mathcal{M}_A$. For the backward directions, let $X \in \mathcal{M}_A$ and set $A' = D(X)$ so that $X \in \mathcal{M}_{A'}^*$: since $A' \subseteq A$ and all the stated conditions on A are inherited by subsets, the results follow from what we proved above. \square

Theorem 3.6(1) shows that restricting the attention to Polish metric spaces with a fixed set of distance $A \in \mathcal{D}$ may provide useful information on the complexity of the isometry relation on zero-dimensional Polish metric spaces: indeed, if A does not contain a right neighborhood of 0, then \cong_A^* is a lower bound for such a relation. In contrast, (3) and (4) of Theorem 3.6 imply that the approach of restricting isometry to Polish metric spaces using a specific distance set cannot provide interesting lower bounds for the complexity of locally compact or connected Polish metric spaces.

4. THE COMPLEXITY OF \mathcal{M}_A AND \mathcal{M}_A^*

We consider the problem of determining the complexity of the subspaces \mathcal{M}_A and \mathcal{M}_A^* of $F(\mathbb{U})$, in particular characterizing when they are standard Borel spaces. While it is worth studying \mathcal{M}_A for any $A \subseteq \mathbb{R}^+$ with $0 \in A$, we have $\mathcal{M}_A^* \neq \emptyset$ only when $A \in \mathcal{D}$.

Notice the following fact, immediate from the definitions:

Fact 4.1. *If A is analytic (in particular if $A \in \mathcal{D}$) then both \mathcal{M}_A and \mathcal{M}_A^* are $\mathbf{\Pi}_2^1$, while when A is Borel then \mathcal{M}_A is $\mathbf{\Pi}_1^1$. If A is countable then \mathcal{M}_A is $\mathbf{\Pi}_1^1$ and \mathcal{M}_A^* belongs to $D_2(\mathbf{\Sigma}_1^1)$.*

The following reductions are easy to prove and very general.

Proposition 4.2. *Let $0 \in A \subseteq \mathbb{R}^+$.*

- (1) *A Borel reduces to \mathcal{M}_A ;*
- (2) *if $A \in \mathcal{D}$ then \mathcal{M}_A Borel reduces to \mathcal{M}_A^* .*

Proof. (1) One can define a Borel function $f : \mathbb{R}^+ \rightarrow F(\mathbb{U})$ such that $f(0)$ is a singleton and otherwise $f(r)$ is a space consisting of two points at distance r : then $A = f^{-1}(\mathcal{M}_A)$.

(2) If $A = \{0\}$, then $\mathcal{M}_A = \mathcal{M}_A^*$. Otherwise, fix $Y \in \mathcal{M}_A^*$ and $r \in A \setminus \{0\}$. The Borel map $X \mapsto X \oplus_r Y$ reduces \mathcal{M}_A to \mathcal{M}_A^* . \square

To obtain sharper results we make extensive use of the following Borel construction of Polish metric spaces.

Definition 4.3. A triple $((r_n)_{n \in \omega}, (r'_n)_{n \in \omega}, x)$ is *tree-suitable* if

- $x > 0$;
- $(r_n)_{n \in \omega}$ is strictly decreasing and converges to 0;
- $(r'_n)_{n \in \omega}$ is strictly monotone and converges to x ;
- $r_0 < \min(x, r'_0)$;
- $\forall n \in \omega \ |r'_n - x| < r_n$, so that $\forall n, m \ |r'_n - r'_m| \leq \max\{r_n, r_m\}$.

In this case one can define an assignment $\Phi_{r_n r'_n}$ that sends a tree $T \subseteq \omega^{<\omega}$ to some $\Phi_{r_n r'_n}(T) \in F(\mathbb{U})$ which is isometric to the completion of $T \cup \{*\}$ under the metric d defined by setting $d(s, t) = r_n$ if $s, t \in T$ are distinct and n is largest such that $s \upharpoonright n = t \upharpoonright n$, and $d(s, *) = r'_{\text{lh}(s)}$ for $s \in T$. The last of the conditions in the definition of tree-suitability ensures that d satisfies the triangle inequality. Using Remark 2.1 and going through \mathcal{M} , we can assume that $\Phi_{r_n r'_n}$ is Borel.

The main property of $\Phi_{r_n r'_n}(T)$ is the following:

Fact 4.4. *If $((r_n)_{n \in \omega}, (r'_n)_{n \in \omega}, x)$ is tree-suitable then for any tree $T \subseteq \omega^{<\omega}$:*

- $D(\Phi_{r_n r'_n}(T)) \subseteq \{0\} \cup \{r_n, r'_n \mid n \in \omega\} \cup \{x\}$;
- $x \in D(\Phi_{r_n r'_n}(T))$ if and only if T is ill-founded.

We first study the complexity of \mathcal{M}_A .

Theorem 4.5. *Let $0 \in A \subseteq \mathbb{R}^+$.*

- (1) *If A is not closed and 0 is a limit point of A , then \mathcal{M}_A is $\mathbf{\Pi}_1^1$ -hard;*
- (2) *\mathcal{M}_A is Borel if and only if either A is closed, or A is Borel and 0 is not a limit point of A ;*
- (3) *if A is Borel but not closed and 0 is a limit point of A then \mathcal{M}_A is $\mathbf{\Pi}_1^1$ -complete;*
- (4) *if A is $\mathbf{\Sigma}_1^1$ -complete and 0 is a limit point of A then \mathcal{M}_A is $\mathbf{\Pi}_2^1$ -complete.*

Proof. (1) Fix $x \in \bar{A} \setminus A$, and sequences $(r_n)_{n \in \omega}$ and $(r'_n)_{n \in \omega}$ in A such that $((r_n)_{n \in \omega}, (r'_n)_{n \in \omega}, x)$ is tree-suitable. Then, by Fact 4.4, T is well-founded if and only if $\Phi_{r_n r'_n}(T) \in \mathcal{M}_A$.

(2) Assume first that A is a closed subset of \mathbb{R}^+ , or that A is Borel and 0 is not a limit point of A . Then, in either case, for any $X \in F(\mathbb{U})$ we have $X \in \mathcal{M}_A \Leftrightarrow \forall n, m \in \omega d(\psi_n(X), \psi_m(X)) \in A$: for the backwards implication when 0 is not a limit point of A , use the fact that the condition $\forall n, m \in \omega d(\psi_n(X), \psi_m(X)) \in A$ implies that X is discrete, so $X = \{\psi_n(X) \mid n \in \omega\}$.

Conversely, assume that \mathcal{M}_A is Borel. Then A is Borel by Proposition 4.2(1) and if A is not closed and 0 is a limit point of A we derive a contradiction from (1).

(3) is immediate from (1) and Fact 4.1.

To prove (4) first recall that \mathcal{M}_A is $\mathbf{\Pi}_2^1$ by Fact 4.1.

To show that \mathcal{M}_A is $\mathbf{\Pi}_2^1$ -hard we fix a strictly decreasing sequence $(\varepsilon_i)_{i \in \omega}$ in A with $\varepsilon_i < 2^{-i}$.

Let $P \subseteq 2^\omega$ be an arbitrary $\mathbf{\Pi}_2^1$ set. We assume first that $A \subseteq [0, 1]$. Since A is $\mathbf{\Sigma}_1^1$ -complete, there exists a continuous function $f : 2^\omega \times 2^\omega \rightarrow [0, 1]$ such that $P = \{\alpha \in 2^\omega \mid \forall \beta \in 2^\omega f(\alpha, \beta) \in A\}$. Let us define a Borel function g from 2^ω to the space of pruned trees on $\{0, 1\}$ as follows. Given $\alpha \in 2^\omega$ consider the compact set $C_\alpha = \{f(\alpha, \beta) \mid \beta \in 2^\omega\} \cup \{\varepsilon_i \mid i \in \omega\} \cup \{0\}$. Define $g(\alpha)$ to be the pruned tree such that $[g(\alpha)] = \{\gamma \in 2^\omega \mid \sum \gamma(i)2^{-(i+1)} \in C_\alpha\}$. The function g is Borel, using the fact that, given $s \in 2^{<\omega}$,

$$s \in g(\alpha) \iff \left[\sum_{i=0}^{\text{lh}(s)-1} \frac{s(i)}{2^{i+1}}, \sum_{i=0}^{\text{lh}(s)-1} \frac{s(i)}{2^{i+1}} + \frac{1}{2^{\text{lh}(s)}} \right] \cap C_\alpha \neq \emptyset$$

together with the continuity of the function $2^\omega \rightarrow K([0, 1])$, $\alpha \mapsto C_\alpha$ (using [Kec95, Exercise 4.29, iv) and vi]) and the fact that the relation of non-disjointness is closed in $(K([0, 1]))^2$ ([Kec95, Exercise 4.29, iii]).

We now apply to $g(\alpha)$ the construction used by Clemens in his proof of [Cle01, Theorem 4.7] using the sequence $(\varepsilon_i)_{i \in \omega}$ we fixed in advance. We thus obtain a function $h : 2^\omega \rightarrow \mathcal{M}_{[0,1]}$ such that $D(h(\alpha)) = C_\alpha$ for every $\alpha \in 2^\omega$. To see that h is Borel one needs to inspect Clemens' construction, keeping in mind Remark 2.1 since $h(\alpha)$ is introduced by defining the restriction of the distance function to a countable dense subset. It is immediate that $\alpha \in P$ if and only if $h(\alpha) \in \mathcal{M}_A$. Thus \mathcal{M}_A is $\mathbf{\Pi}_2^1$ -complete.

Properties of A	complexity of \mathcal{M}_A	Reference
A closed or 0 isolated in A	Borel	4.6
A Borel not closed and 0 not isolated in A	$\mathbf{\Pi}_1^1$ -complete	4.5(3)
A true analytic	neither $\mathbf{\Sigma}_1^1$ nor $\mathbf{\Pi}_1^1$	4.7(1)
A $\mathbf{\Sigma}_1^1$ -complete	$\mathbf{\Pi}_2^1$ -complete	4.7(2)

TABLE 1. Summary of the complexity of \mathcal{M}_A for $A \in \mathcal{D}$

When $A \subseteq [0, n]$, by rescaling we obtain a function $h_n : 2^\omega \rightarrow \mathcal{M}_{[0, n]}$ reducing P to \mathcal{M}_A .

If A is unbounded, let $\varphi : 2^\omega \rightarrow \mathbb{R}^+$ be a continuous function reducing a $\mathbf{\Sigma}_1^1$ -complete subset of 2^ω to A . Since the range of φ is bounded, it follows that $A_n = A \cap [0, n]$ is $\mathbf{\Sigma}_1^1$ -complete for some n and h_n still does the job. \square

If we further assume that $A \in \mathcal{D}$ we can draw the following corollaries. These, combined with Theorem 4.5(3), provide a complete picture of the complexity of \mathcal{M}_A for $A \in \mathcal{D}$ under analytic determinacy (which ensures that every analytic set which is not Borel is $\mathbf{\Sigma}_1^1$ -complete).

Corollary 4.6. *Let $A \in \mathcal{D}$. The following are equivalent:*

- (i) \mathcal{M}_A is Borel;
- (ii) \mathcal{M}_A is $\mathbf{\Sigma}_1^1$;
- (iii) A is closed or 0 is not a limit point of A .

Proof. (i) implies (ii) is obvious. (ii) implies (iii) follows from Theorem 4.5(1) because if \mathcal{M}_A is $\mathbf{\Sigma}_1^1$ then it is not $\mathbf{\Pi}_1^1$ -hard. To check that (iii) implies (i) notice that if A is closed then Theorem 4.5(2) applies, yielding immediately (i). If instead 0 is not a limit point of A then, since $A \in \mathcal{D}$, A is countable by Theorem 1.2. Thus A is Borel and Theorem 4.5(2) applies again. \square

Corollary 4.7. *Let $A \in \mathcal{D}$.*

- (1) *If A is not Borel then \mathcal{M}_A is neither analytic nor coanalytic;*
- (2) *if A is $\mathbf{\Sigma}_1^1$ -complete then \mathcal{M}_A is $\mathbf{\Pi}_2^1$ -complete.*

Proof. (1) If \mathcal{M}_A were analytic then it would be Borel by Corollary 4.6, and then A would be Borel by Proposition 4.2(1). Since A is analytic by Theorem 1.2, Proposition 4.2(1) implies also that \mathcal{M}_A is not coanalytic.

(2) Since any $\mathbf{\Sigma}_1^1$ -complete set is uncountable, the result follows immediately from Theorems 1.2 and 4.5(4). \square

For the reader's convenience we summarize in Table 1 our results for the complexity of \mathcal{M}_A when $A \in \mathcal{D}$.

We now show that the complexity of \mathcal{M}_A^* often depends on the limit points of A .

Theorem 4.8. *Let $A \in \mathcal{D}$.*

- (1) \mathcal{M}_A^* is Borel if and only if either 0 is not a limit point of A or 0 is the unique limit point of A .
- (2) *Suppose 0 is a limit point of A and A has other limit points (which may belong to A or not).*

- (a) If A is closed then \mathcal{M}_A^* is Σ_1^1 -hard;
 - (b) if A is not closed then \mathcal{M}_A^* is Π_1^1 -hard;
 - (c) if A is not closed and at least one of its limit points different from 0 belongs to A , then \mathcal{M}_A^* is $D_2(\Sigma_1^1)$ -hard.
- (3) if A is Σ_1^1 -complete then \mathcal{M}_A^* is Π_2^1 -complete.

Proof. We start from (2). For (a) pick $y \neq 0$ which is a limit point of A : obviously $y \in A$. Fix now sequences $(s_n)_{n \in \omega}, (s'_n)_{n \in \omega}$ in A such that the triple $((s_n)_{n \in \omega}, (s'_n)_{n \in \omega}, y)$ is tree-suitable. As $A \setminus \{y\} \in \mathcal{D}$ by Theorem 1.2, fix also a space $Y \in \mathcal{M}_{A \setminus \{y\}}^*$. Then, using Facts 4.4 and 3.4, the function $T \mapsto \Phi_{s_n s'_n}(T) \oplus_{s_0} Y$ is a Borel reduction from ill-founded trees to \mathcal{M}_A^* .

(b) follows from Proposition 4.2(2) and Theorem 4.5(1).

To prove (c) let $x \in \bar{A} \setminus A$, and let $y \in A \setminus \{0\}$ which is a limit point of A . Fix sequences $(r_n)_{n \in \omega}, (s_n)_{n \in \omega}, (t_n)_{n \in \omega}$ in $A \setminus \{y\}$ such that both $((r_n)_{n \in \omega}, (s_n)_{n \in \omega}, x)$ and $((r_n)_{n \in \omega}, (t_n)_{n \in \omega}, y)$ are tree-suitable. By Theorem 1.2 again, $A \setminus \{y\} \in \mathcal{D}$, so fix $X \in \mathcal{M}_{A \setminus \{y\}}^*$. Then, by Facts 4.4 and 3.4, the Borel function

$$(U, T) \mapsto \Theta(U, T) = (\Phi_{r_n s_n}(U) \oplus_{r_0} \Phi_{r_n t_n}(T)) \oplus_{r_0} X$$

is such that $\Theta(U, T) \in \mathcal{M}_A^*$ if and only if U is well-founded and T is ill-founded.

We now deal with (1). The forward direction follows from (2), since if the thesis were false one of (a) and (b) would apply.

Conversely, assume first that 0 is not a limit point of A . Then, by Theorem 1.2, A is countable and all members of \mathcal{M}_A^* are discrete. Then for any $X \in F(\mathbb{U})$ we have that $X \in \mathcal{M}_A^*$ if and only if $X \in \mathcal{M}_A \wedge \forall a \in A \exists m_1, m_2 \in \omega d(\psi_{m_1}(X), \psi_{m_2}(X)) = a$. Theorem 4.5(2) allows to conclude in this case.

Finally, suppose 0 is the unique limit point of A . Then A is closed and countable and all elements of A different from 0 are isolated in A . Thus for any $X \in F(\mathbb{U})$ we have again that $X \in \mathcal{M}_A^*$ if and only if $X \in \mathcal{M}_A \wedge \forall a \in A \exists m_1, m_2 \in \omega d(\psi_{m_1}(X), \psi_{m_2}(X)) = a$, which allows to conclude by applying Theorem 4.5(2).

(3) follows immediately from Proposition 4.2(2) and Corollary 4.7(2). \square

We now consider the case when A is countable. Then \mathcal{M}_A is either Borel or Π_1^1 -complete according to Theorem 4.5(2 and 3). We can obtain a complete classification of the complexity of \mathcal{M}_A^* as well.

Theorem 4.9. *Let A be a countable subset of \mathbb{R}^+ , with $0 \in A$.*

- (1) If 0 is not a limit point of A or 0 is the unique limit point of A , then \mathcal{M}_A^* is Borel;
- (2) if 0 is a limit point of A and A is closed having other limit points besides 0, then \mathcal{M}_A^* is Σ_1^1 -complete;
- (3) if 0 is a limit point of A , A is not closed, and all limit points of A different from 0 do not belong to A , then \mathcal{M}_A^* is Π_1^1 -complete;
- (4) if 0 is a limit point of A , A is not closed and contains a limit point different from 0, then \mathcal{M}_A^* is $D_2(\Sigma_1^1)$ -complete.

Proof. (1) follows from Theorem 4.8(1).

(2) In this case, by Theorem 4.5(2), \mathcal{M}_A is Borel. Since $X \in \mathcal{M}_A^*$ if and only if $X \in \mathcal{M}_A \wedge \forall a \in A \exists x, y \in X d(x, y) = a$, we have that \mathcal{M}_A^* is Σ_1^1 . Completeness follows from Theorem 4.8(2a).

Properties of A	complexity of \mathcal{M}_A^*	Reference
0 isolated in A or 0 unique limit point of A	Borel	4.8(1)
0 not isolated in A , A closed with other limit points	Σ_1^1 -hard	4.8(2)(a)
0 not isolated in A , A countable closed with other limit points	Σ_1^1 -complete	4.9(2)
0 not isolated in A , A not closed with other limit points	Π_1^1 -hard	4.8(2)(b)
0 not isolated in A , A countable not closed with all other limit points not in A	Π_1^1 -complete	4.9(3)
0 not isolated in A , A not closed with other limit points in A	$D_2(\Sigma_1^1)$ -hard	4.8(2)(c)
0 not isolated in A , A countable not closed with other limit points in A	$D_2(\Sigma_1^1)$ -complete	4.9(4)
A Σ_1^1 -complete	Π_2^1 -complete	4.8(3)

TABLE 2. Summary of the complexity of \mathcal{M}_A^* for $A \in \mathcal{D}$

(3) In this case, all points of A different from 0 are isolated in A so that, as at the end of the proof of Theorem 4.8(1), we have $X \in \mathcal{M}_A^*$ if and only if $X \in \mathcal{M}_A \wedge \forall a \in A \exists m_1, m_2 d(\psi_{m_1}(X), \psi_{m_2}(X)) = a$. Hence \mathcal{M}_A^* is Π_1^1 because \mathcal{M}_A is Π_1^1 by Fact 4.1. Completeness follows from Theorem 4.8(2b).

(4) By Fact 4.1 \mathcal{M}_A^* is $D_2(\Sigma_1^1)$ and completeness follows from Theorem 4.8(2c). \square

Remark 4.10. In the literature, there are very few “natural” examples of sets belonging to the class $D_2(\Sigma_1^1)$ but not to simpler ones. The set \mathcal{M}_A^* with A as in Theorem 4.9(4) is one of these. Other notable examples are: the collection of countable graphs whose automorphism group is isomorphic to ${}^\omega\mathbb{Z}_p$ for p a prime number [CK00]; the collection of countable linear orders which are not strongly surjective [CCM19]; and some collection of measurable sets generated using the density function on the Cantor space [AC19].

Table 2 summarizes our results for the complexity of \mathcal{M}_A^* when $A \in \mathcal{D}$.

Several sections of Gao and Shao’s paper [GS11] are devoted to different ways of constructing, for every countable $A \in \mathcal{D}$ (which, by Theorem 3.1(2), means for every A such that $\mathcal{U}_A^* \neq \emptyset$), a Polish A -ultrametric spaces X which is A -universal (i.e. such that $X \in \mathcal{U}_A^*$, and $Y \sqsubseteq X$ for every $Y \in \mathcal{U}_A$) and ultrahomogeneous. They call such a space A -ultrametric Urysohn. The analogous question for Polish spaces was considered by Sauer in [Sau13] (beware that Sauer calls homogeneous the spaces we call ultrahomogeneous, and that his definition of universality is equivalent to ours only for ultrahomogeneous and complete spaces).

Definition 4.11. We say that a metric space X is *ultrahomogeneous* if every isometry between finite subsets of X can be extended to an isometry of the whole X .

Definition 4.12. Let $A \in \mathcal{D}$. We say that $X \in \mathcal{M}_A$ is *Polish A -universal* if $Y \sqsubseteq X$ for every $Y \in \mathcal{M}_A$ (clearly $X \in \mathcal{M}_A^*$ must then hold). If additionally X is ultrahomogeneous then we say it is *A -Urysohn*.

Here we use Corollary 4.6 to extend Sauer's characterization ([Sau13, Theorem 4.13], which is the equivalence between (i) and (iv) in Theorem 4.14 below) and give a different proof of the necessity of the condition for the existence of A -Urysohn spaces. The following property was isolated in [DLPS07].

Definition 4.13. A triple (a, b, c) of elements of \mathbb{R}^+ is *metric* if $a \leq b+c$, $b \leq a+c$, and $c \leq a+b$. A set $A \subseteq \mathbb{R}^+$ satisfies the *4-values condition* if for all pairs of metric triples of numbers in A of the form (a, b, x) and (c, d, x) there exists $y \in A$ such that both (b, c, y) and (a, d, y) are metric triples.

Theorem 4.14. *Let $A \in \mathcal{D}$. The following are equivalent:*

- (i) *A satisfies the 4-values condition and either is closed or 0 is not a limit point of A ;*
- (ii) *A satisfies the 4-values condition and \mathcal{M}_A is Borel;*
- (iii) *A satisfies the 4-values condition and \mathcal{M}_A is Σ_1^1 ;*
- (iv) *there exists $X \in \mathcal{M}_A$ which is A -Urysohn;*
- (v) *A satisfies the 4-values condition and there exists $X \in \mathcal{M}_A$ which is Polish A -universal.*

Proof. The equivalence between (i), (ii), and (iii) follows immediately from Corollary 4.6.

(i) implies (iv) is obtained by Sauer repeating the classical construction of the Urysohn space (which in our terminology would be \mathbb{R}^+ -Urysohn) using only spaces in \mathcal{M}_A : we amalgamate (using the 4-values condition) the finite members of \mathcal{M}_A obtaining an ultrahomogeneous Z with $D(Z) = A$ which isometrically embeds all countable metric spaces using distances in A . We then let X to be the completion of Z , and we need to check that (i) guarantees that X does not use distances outside A . If 0 is not a limit point of A then Z is discrete so that $X = Z \in \mathcal{M}_A$. Otherwise $D(X) \subseteq \overline{D(Z)} = \overline{A}$, thus if A is closed we have $X \in \mathcal{M}_A$. Moreover X is still ultrahomogeneous by [Sau13, Theorem 4.4].

To prove (iv) implies (v) we need Theorem 3.9 of [Sau13], stating that the set of distances of an ultrahomogeneous universal metric space satisfies the 4-values condition.

We complete the proof by showing that (v) implies (iii). This is immediate because if a Polish A -universal X exists, then $Y \in \mathcal{M}_A$ if and only if $Y \sqsubseteq X$, and \sqsubseteq is analytic. \square

5. ISOMETRY AND ISOMETRIC EMBEDDABILITY

As a first step in our analysis of isometry and isometric embeddability restricted to \mathcal{M}_A and \mathcal{M}_A^* we prove the following proposition, which also answers the first part of [Cle07, Question 3] by showing that if $A, A' \in \mathcal{D}$ and $A \subseteq A'$, then $\cong_A^* \leq_B \cong_{A'}^*$.

Proposition 5.1. *Let $A, A' \in \mathcal{D}$ and assume $A \subsetneq A'$. Then $\cong_A^* \leq_B \cong_A \leq_B \cong_{A'}^* \leq_B \cong_{A'}$ and $\sqsubseteq_A^* \leq_B \sqsubseteq_A \leq_B \sqsubseteq_{A'}^* \leq_B \sqsubseteq_{A'}$.*

Proof. For any A we have $\mathcal{M}_A^* \subseteq \mathcal{M}_A$ and hence $\cong_A^* \leq_B \cong_A$ and $\sqsubseteq_A^* \leq_B \sqsubseteq_A$. Thus we need only to prove $\cong_A \leq_B \cong_{A'}^*$ and $\sqsubseteq_A \leq_B \sqsubseteq_{A'}^*$.

Suppose first that $A' = A \cup \{r_0\}$ and $a < r_0$ for all $a \in A$. Fix a space $Z \in \mathcal{M}_A^*$. Given $X \in \mathcal{M}_A$, define $X' = X \oplus_{r_0} Z \in \mathcal{M}_{A'}^*$ by Fact 3.4: this map is Borel by Proposition 3.3. We show that it is the required reduction. Notice that by case assumption $d_{X'}(x, z) = r_0$ whenever $x \in X$ and $z \in Z$. If $\psi: X \rightarrow Y$ is an isometric embedding (respectively, an isometry), then $\psi \cup \text{id}_Z: X' \rightarrow Y'$ is an isometric embedding (respectively, an isometry) as well. Conversely, suppose $\psi: X' \rightarrow Y'$ is an isometric embedding. Since the distance r_0 is never realized inside any of X , Y , and Z , either $\psi(X) \subseteq Y$ and $\psi(Z) \subseteq Z$, in which case $\psi \upharpoonright X$ witnesses $X \sqsubseteq Y$, or else $\psi(X) \subseteq Z$ and $\psi(Z) \subseteq Y$. In the latter case, $X \sqsubseteq Z$ and $Z \sqsubseteq Y$, so again $X \sqsubseteq Y$. If moreover ψ is onto, then we have either $\psi(X) = Y$, or else $\psi(X) = Z$ and $\psi(Z) = Y$, which allows us to conclude that $X \cong Y$.

Otherwise there exist $r_0 \in A' \setminus A$ and $r_1 \in A'$ with $r_0 < r_1$. Notice that there is $Z \in \mathcal{M}_{A'}^*$ with the property that $d_Z(z_0, z_1) = r_0$ for exactly one pair of points $\{z_0, z_1\} \subseteq Z$. Indeed $A' \setminus \{r_0\} \in \mathcal{D}$ by Theorem 1.2, so let $Z = W \oplus_{r_1} V$ where $W \in \mathcal{M}_{A' \setminus \{r_0\}}^*$ and $V = \{z_0, z_1\}$ consists of two points at distance r_0 , so that $d_Z(w, z_i) \geq r_1$ for every $w \in W$ and $i \in \{0, 1\}$.

We show that the mapping that associates to every $X \in \mathcal{M}_A$ the space $X \times Z$ with the product metric $d_{X \times Z}((x, z), (x', z')) = \max\{d_X(x, x'), d_Z(z, z')\}$ is the required Borel reduction. Borelness is proved using Remark 2.1.

If $\psi: X \rightarrow Y$ is an isometric embedding (respectively, an isometry), then $\psi \times \text{id}_Z: X \times Z \rightarrow Y \times Z$ is an isometric embedding (respectively, an isometry). Conversely, suppose $\psi: X \times Z \rightarrow Y \times Z$ is an isometric embedding. Then for every $x \in X$ there exists $y \in Y$ such that $\psi(x, z_0) \in \{(y, z_0), (y, z_1)\}$, since in both $X \times Z$ and $Y \times Z$ the points having second coordinate equal to z_0 or z_1 are the only points that realize the distance r_0 by our choice of Z . This defines a function $\varphi: X \rightarrow Y$ with the property that for every $x \in X$ we have $\psi(x, z_0) = (\varphi(x), z_i)$ for some $i \in \{0, 1\}$. In order to prove that φ is an isometric embedding, let $x, x' \in X$ and let $r = d_X(x, x') = d_{X \times Z}((x, z_0), (x', z_0)) = d_{Y \times Z}(\psi(x, z_0), \psi(x', z_0))$. Since we have $\psi(x, z_0) = (\varphi(x), z_i)$ and $\psi(x', z_0) = (\varphi(x'), z_j)$ for some $i, j \in \{0, 1\}$, we obtain $r = \max\{d_Y(\varphi(x), \varphi(x')), d_Z(z_i, z_j)\}$; recalling that $r \neq r_0 = d_Z(z_0, z_1)$, the equality $d_Y(\varphi(x), \varphi(x')) = r$ follows. So φ is an isometric embedding. Assume now ψ is surjective and let $y \in Y$. So, suppose $\psi(x, z) = (y, z_0)$ and $\psi(x', z') = (y, z_1)$. Recall that $r_0 \notin A$ is not realized in $X \in \mathcal{M}_A$ and is realized only by the pair $\{z_0, z_1\}$ in Z . Since $d_{Y \times Z}((y, z_0), (y, z_1)) = r_0$ we have that $\{z, z'\} = \{z_0, z_1\}$. If $z = z_0$ then $\varphi(x) = y$, while if $z' = z_0$ then $\varphi(x') = y$. So φ too is surjective. \square

Remark 5.2. One may be interested in analogues of Proposition 5.1 obtained by restricting the relations of isometry and isometric embeddability to a given class of Polish metric spaces.

The same proof shows that the conclusion of the proposition holds for ultrametric, zero-dimensional, countable, locally compact, σ -compact, and discrete spaces. This is the case because such classes are closed under finite products and the operations \oplus_r , and whenever they have an element in $\mathcal{M}_{A'}^*$, for every $r_0 \in A'$ they also have an element in $\mathcal{M}_{A' \setminus \{r_0\}}^*$ and contain a space consisting of two points at distance r_0 . For the classes mentioned above the latter property follows from Theorem 3.1: this is clear for ultrametric, zero-dimensional, countable, and discrete spaces; for the classes of locally compact and σ -compact spaces notice that removing a point from a σ -compact subset of \mathbb{R} yields a σ -compact set. For contrast, this

argument does not work for compact metric spaces² and we do not know whether Proposition 5.1 holds restricted to this class.

One can also restrict attention to spaces of a fixed dimension different from 0 and obtain the same results even if these classes are not closed under finite products: this is because in the last paragraph of the proof of Proposition 5.1 we can require (by Theorem 3.1(1)) Z to be zero-dimensional, so that $X \times Z$ has the same dimension of X .

We will use the following folklore construction to turn a countable graph into a discrete metric space. Fix $r, r' \in \mathbb{R}$ with $r < r' \leq 2r$. To each graph G on ω associate the metric space $X_G = (G, d_G)$ by letting $d_G(a, b) = r$ if (a, b) is an edge in G , and $d(a, b) = r'$ if $a \neq b$ and (a, b) is not an edge in G . The following Lemma is straightforward.

Lemma 5.3. *The map $G \mapsto X_G$ Borel reduces countable graph isomorphism and countable graph embeddability to $\cong_{\{0, r, r'\}}$ and $\sqsubseteq_{\{0, r, r'\}}$, respectively. Moreover, if we restrict the map to nontrivial graphs (i.e. different from the empty graph and from the countable clique), we get a reduction to $\cong_{\{0, r, r'\}}^*$ and $\sqsubseteq_{\{0, r, r'\}}^*$.*

5.1. Isometry. The study of the complexity of \cong_A^* was started by Clemens in [Cle07], where such relation is called E_A . Clemens' main results about \cong_A^* are summarized in the following theorem.

Theorem 5.4 ([Cle07, Theorem 23]). *Let $A \in \mathcal{D}$.*

- (1) *If A contains a right neighborhood of 0, then \cong_A^* is Borel bireducible with any complete orbit equivalence relation.*
- (2) *If A is dense in some right neighborhood of 0 but does not contain any of them, then both the action of the density ideal on ω^2 and countable graph isomorphism are Borel reducible to \cong_A^* . In particular, \cong_A^* is strictly above countable graph isomorphism with respect to \leq_B .*
- (3) *If A is not dense in any right neighborhood of 0 and either 0 is a limit point of A or A is not well-spaced, then \cong_A^* is Borel bireducible with countable graph isomorphism.*
- (4) *If $A = \{0\} \cup \{r_i \mid i \in \omega\}$ with $0 < r_i < r_{i+1}$ is well-spaced, then \cong_A^* is Borel bireducible with isomorphism between reverse trees (as defined in [Cle07]).*
- (5) *If $A = \{0, r_0, \dots, r_{n-1}\}$ with $0 < r_i < r_{i+1}$ is well-spaced, then \cong_A^* is Borel bireducible with isomorphism between trees of height n . Thus these relations form a \leq_B -strictly increasing chain of equivalence relations classifiable by countable structures and they are strictly below countable graph isomorphism.*

Notice that the conditions considered in Theorem 5.4 are exhaustive. In fact, if 0 is not isolated in A or A is not well-spaced then we are in either case (1), (2), or (3). If 0 is isolated in A and A is well-spaced then A is well-founded, since strictly decreasing sequences in a well-spaced set converge to 0. Since a well-founded and well-spaced set has order type $\leq \omega$, we are either in case (4) or in case (5).

Proposition 5.1 may be used to give a simpler proof of part (1) of Theorem 5.4. Let r be such that $[0, r] \subseteq A$. To any Polish metric space (X, d) , associate the

²In a previous version of the paper we claimed that this was the case: we thank the referee for pointing out our mistake.

space (X, d') , where $d'(x, y) = r \cdot \frac{d(x, y)}{1+d(x, y)}$. This reduces isometry on all Polish metric spaces to $\cong_{[0, r)}$, which in turn reduces to \cong_A^* by Proposition 5.1. Since isometry on arbitrary Polish metric spaces is Borel bireducible with the complete orbit equivalence relation by [GK03, Theorem 1], we are done.

Theorem 5.4 yields the following sufficient condition for countable graph isomorphism being Borel reducible to \cong_A^* .

Corollary 5.5. *Let $A \in \mathcal{D}$. If A is ill-founded or not well-spaced, then countable graph isomorphism Borel reduces to \cong_A^* .*

We will now show how the results from [CMMR18] cited in Section 2 can be used to complete the description of the behaviour of \cong_A^* . We begin by proving the converse of Corollary 5.5, i.e. we characterize when countable graph isomorphism is Borel reducible to \cong_A^* .

Theorem 5.6. *Let $A \in \mathcal{D}$. Countable graph isomorphism Borel reduces to \cong_A^* if and only if A is either ill-founded or not well-spaced.*

Proof. One direction is Corollary 5.5, but for the reader's convenience we give here an alternative and simpler proof. If A is ill-founded, let $(r_n)_{n \in \omega}$ be a decreasing sequence in A . Since countable graph isomorphism Borel reduces to $\cong_{(r_n)_{n \in \omega}}^*$ by Theorem 2.3(1), it suffices to use Proposition 5.1 when $(r_n)_{n \in \omega} \subsetneq A$. If instead A is not well-spaced fix $r, r' \in A$ with $r < r' \leq 2r$. Lemma 5.3 gives a Borel reduction of isomorphism between nontrivial countable graphs to $\cong_{\{0, r, r'\}}^*$. Then apply Proposition 5.1 if $\{0, r, r'\} \subsetneq A$.

Finally, assume that A is well-founded and well-spaced. Then $\mathcal{M}_A^* = \mathcal{U}_A^*$ by Theorem 3.6(2), and countable graph isomorphism does not Borel reduce to \cong_A^* because the latter is Borel by Theorem 2.4(2). \square

We now have the following fairly complete picture of the structure of the relations \cong_A^* . Notice that conditions (1)–(4) exhaust all possible cases for A .

Theorem 5.7. *Let $A \in \mathcal{D}$.*

- (1) *The relations \cong_A^* with A well-founded and well-spaced form a strictly increasing chain of order type $\omega + 1$ under \leq_B , consisting of Borel equivalence relations, and they are Borel reducible to all the other $\cong_{A'}^*$, with $A' \in \mathcal{D}$.*
- (2) *If A is either ill-founded or not well-spaced, and moreover A is not dense in any right neighborhood of 0, then \cong_A^* is Borel bireducible with countable graph isomorphism.*
- (3) *If A is dense in some right neighborhood of 0 but does not contain any such neighborhood, then \cong_A^* is strictly above countable graph isomorphism and Borel reducible to any complete orbit equivalence relation.*
- (4) *If A contains a right neighborhood of 0, then \cong_A^* is Borel bireducible with any complete orbit equivalence relation.*

Proof. Parts (2)–(4) follow from Theorem 5.4 and the observations following it, so let us prove (1).

Let α be the order type of A . The fact that A is well-spaced implies $1 \leq \alpha \leq \omega$ and $\mathcal{M}_A^* = \mathcal{U}_A^*$ (by Theorem 3.6(2)). Thus \cong_A^* and \cong_α^* are the same relation. By Theorem 2.4(2), when α varies between 1 and ω these equivalence relations form a strictly increasing chain of length $\omega + 1$ under \leq_B and they are Borel equivalence relations. Finally, since \cong_α^* Borel reduces to countable graph

isomorphism by Theorem 2.4(1), it follows from Theorem 5.6 that \cong_A^* reduces to any other $\cong_{A'}^*$, for A' not satisfying the conditions of (1). \square

We will now partially answer also the second part of [Cle07, Question 3], which asked whether $\cong_A \sim_B \cong_A^*$ for every $A \in \mathcal{D}$.

Theorem 5.8. *Let $A \in \mathcal{D}$ satisfy at least one of the following conditions:*

- (i) *A is not dense in any right neighborhood of 0;*
- (ii) *A has maximum;*
- (iii) *there exists $f: A \rightarrow A$ which is Polish metric preserving, injective, and non-surjective.*

Then $\cong_A \sim_B \cong_A^$.*

Proof. For any $A \in \mathcal{D}$ we have $\cong_A^* \leq_B \cong_A$ because $\mathcal{M}_A^* \subseteq \mathcal{M}_A$.

Assume first that A is not dense in any right neighborhood of 0. In this case \cong_A is classifiable by countable structures. In fact the argument of [Cle07, Proposition 18] applies not only to \cong_A^* but also to \cong_A . We distinguish two cases. If A is either ill-founded or not well-spaced, then \cong_A^* is Borel bireducible with countable graph isomorphism by Theorem 5.7(2), and hence $\cong_A \leq_B \cong_A^*$. If instead A is well-founded and well-spaced then $\mathcal{M}_A = \mathcal{U}_A$ and $\mathcal{M}_A^* = \mathcal{U}_A^*$, so that we can use Theorem 2.4(1).

Now assume that $r = \max A$. Since $\mathcal{M}_{\{0\}} = \mathcal{M}_{\{0\}}^*$, we may assume that A has more than one element. Fix $Z \in \mathcal{M}_{A \setminus \{r\}}^*$ and consider the map sending $X \in \mathcal{M}_A$ to $X' = X \oplus_r Z \in \mathcal{M}_A^*$, so that by case assumption $d_{X'}(x, z) = r$ whenever $x \in X$ and $z \in Z$. The map is Borel and we claim that it witnesses $\cong_A \leq_B \cong_A^*$. To show this we use an argument similar to the one employed in the first part of the proof of Theorem 2.4(1) in [CMMR18]. Fix $X, Y \in \mathcal{M}_A$. First assume that $\varphi: X \rightarrow Y$ witnesses $X \cong_A Y$: then $\varphi \cup \text{id}_Z$ is a witness of $X' \cong_A^* Y'$.

Conversely, let ψ be an isometry between X' and Y' , and let $X_0 = \psi^{-1}(Z)$, so that $\psi(X' \setminus X_0) = Y$. Notice that $d_{X'}(x_0, x_1) < r$ for every $x_0, x_1 \in X_0$ since $\psi(x_0), \psi(x_1) \in Z \in \mathcal{M}_{A \setminus \{r\}}^*$. Hence, by construction of X' , either $X_0 \subseteq Z$ or $X_0 \subseteq X$. In the former case $X_0 = Z$ because $\psi(Z)$ cannot intersect both Y and Z , since any two points of Z are less than r apart. Thus $\psi(X) = \psi(X' \setminus X_0) = Y$ and $\psi \upharpoonright X$ is an isometry between X and Y . If instead $X_0 \subseteq X$, we claim that $\varphi = \psi \upharpoonright (X \setminus X_0) \cup (\psi \circ \psi) \upharpoonright X_0$ witnesses $X \cong_A Y$. Notice that φ is well-defined by the fact that by definition $\psi(X_0) = Z \subseteq X'$. For the same reason, the range of φ equals the range of $\psi \upharpoonright (X \setminus X_0) \cup \psi \upharpoonright Z = \psi \upharpoonright (X' \setminus X_0)$, thus φ is a surjection from X onto Y . Finally, we check that φ preserves distances. It is clearly enough to show that for $x \in X \setminus X_0$ and $x' \in X_0$ we have $d_X(x, x') = d_Y(\varphi(x), \varphi(x'))$. Since $\psi(x) \in Y$ and $\psi(x') \in Z$, we have $d_{Y'}(\psi(x), \psi(x')) = r$, whence $d_X(x, x') = d_{X'}(x, x') = d_{Y'}(\psi(x), \psi(x')) = r$. Since $\psi(x') \in Z \subseteq X'$ and $x \in X$, $d_{X'}(x, \psi(x')) = r$, therefore $d_{Y'}(\psi(x), \psi(\psi(x'))) = r$. Since $\psi(x) = \varphi(x)$ and $\psi(\psi(x')) = \varphi(x')$ by definition of φ , we have $d_Y(\varphi(x), \varphi(x')) = d_{Y'}(\varphi(x), \varphi(x')) = r = d_X(x, x')$, as required.

Finally, assume that f is as in (iii) and let $A' \subsetneq A$ be the range of f . We map any $X \in \mathcal{M}_A$ to $X' \in \mathcal{M}_{A'}$ by composing the metric d_X with f : the fact that f is Polish metric preserving makes sure that $d_{X'} = f \circ d_X$ is still a Polish metric space. Consider the map $X \mapsto X'$. It is Borel because f is such by Proposition

2.7, and, since f is injective, witnesses $\cong_A \leq_B \cong_{A'}$. By Proposition 5.1 we have $\cong_{A'} \leq_B \cong_{A'}^*$, and thus $\cong_A \leq_B \cong_A^*$. \square

It is not obvious when Condition (iii) of Theorem 5.8 holds. Notice that a sufficient condition for a nondecreasing $f: A \rightarrow \mathbb{R}$ to be metric preserving is that for all $r, s, t \in A$, if $s \leq t < r \leq s+t$ then $f(r) \leq f(s) + f(t)$. Using this we see that Condition (iii) holds for instance when $A = \mathbb{Q}^+$ or $A = (\mathbb{R}^+ \setminus \mathbb{Q}) \cup \{0\}$, as witnessed by the map $f(r) = r/(1+r)$. The same is true when A contains an interval $[a, b]$, as witnessed by

$$f(r) = \begin{cases} r & \text{if } r < a; \\ a + (b-a) \frac{r-a}{1+(r-a)} & \text{if } r \geq a. \end{cases}$$

(Notice that b does not belong to the range of f .) On the other hand, Condition (iii) can fail even for countable sets: if $A = \{0\} \cup \{2^k \mid k \in \mathbb{Z}\}$ then every injective Polish metric preserving function $f: A \rightarrow A$ satisfies $f(2^k) = 2^{k+z}$ for some $z \in \mathbb{Z}$, and hence is surjective. Notice however that this particular A satisfies Condition (i) of Theorem 5.8.

In fact, we do not know if there exists $A \in \mathcal{D}$ which does not satisfy any of the conditions of Theorem 5.8. However if such an A exists, it must be uncountable. To show this, we need the following fact, which might be of independent interest.

Proposition 5.9. *Let $A \in \mathcal{D}$ be such that A is dense in (a, b) and $A \cap (a, +\infty)$ is countable for some a, b with $0 < a < b$. Then Condition (iii) of Theorem 5.8 holds, i.e. there exists $f: A \rightarrow A$ which is Polish metric preserving, injective, and non-surjective.*

Proof. We may assume that $a, b \in A$. We will define f strictly increasing which is the identity up to a and maps $A \cap (a, +\infty)$ into (a, b) , so that $b \in A$ is not in the range of f .

Let $(a_n)_{n \in \omega}$ be an enumeration without repetitions of $A \cap (a, +\infty)$. Since A is dense in (a, b) we can recursively define $f(a_n)$ so that:

- $f(a_n) < a_n$;
- if $c_0 = a < c_1 < \dots < c_{n+1}$ enumerate in increasing order $\{a\} \cup \{a_m \mid m \leq n\}$ then the slope of the segment with endpoints $(c_i, f(c_i))$ and $(c_{i+1}, f(c_{i+1}))$ is larger than the slope of the segment with endpoints $(c_{i+1}, f(c_{i+1}))$ and $(c_{i+2}, f(c_{i+2}))$ for each $i < n$.

It remains to prove that f is Polish metric preserving. Since f is nondecreasing, by Proposition 2.6 and the observation after Theorem 5.8, it suffices to show that for all $r, s, t \in A$, if $s \leq t < r \leq s+t$ then $f(r) \leq f(s) + f(t)$. By construction we have that if $x, y, z \in A$ with $x < y < z$ then $S(y, z) \leq S(x, z) \leq S(x, y)$ where $S(v, w)$ is the slope of the segment with endpoints $(v, f(v))$ and $(w, f(w))$. From this it follows that for r, s, t as above $S(t, r) \leq S(0, s)$, whence

$$f(r) = f(t) + (r-t) \cdot S(t, r) \leq f(t) + s \cdot S(t, r) \leq f(t) + s \cdot S(0, s) = f(t) + f(s).$$

\square

Theorem 5.10. *If $A \in \mathcal{D}$ is countable then $\cong_A \sim_B \cong_A^*$.*

Proof. If A is not dense in any right neighborhood of 0 we are in case (i) of Theorem 5.8. Otherwise by Proposition 5.9 we are in case (iii) of Theorem 5.8. \square

5.2. Isometric embeddability. Recall again that if A is well-founded and well-spaced, then $\mathcal{M}_A^* = \mathcal{U}_A^*$ and the order type of A is $\leq \omega$. Hence in this case the structure of the relations \sqsubseteq_A^* is described by (i) and (iv) of Theorem 2.4(4). The remaining case is settled by the following proposition.

Proposition 5.11. *Let $A \in \mathcal{D}$. If A is either ill-founded or not well-spaced, then \sqsubseteq_A^* is Borel bireducible with a complete analytic quasi-order.*

Proof. First notice that, even if \mathcal{M}_A^* may not be a standard Borel space, the relation \sqsubseteq_A^* still Borel reduces to isometric embeddability on all Polish metric spaces, which is a complete analytic quasi-order on a standard Borel space (in fact, Louveau and Rosendal [LR05] showed that isometric embeddability restricted to ultrametric Polish spaces is a complete analytic quasi-order, and we strengthened this in [CMMR13] by showing that it has the stronger property of being invariantly universal).

If A is ill-founded, let $(r_n)_{n \in \omega}$ be a decreasing sequence in A , and let $A' = \{0\} \cup \{r_n \mid n \in \omega\}$. Then isometric embeddability on $\mathcal{U}_{A'}^*$ is a complete analytic quasi-order by Theorem 2.3(2). Hence so is $\sqsubseteq_{A'}^*$ because $\mathcal{U}_{A'}^* \subseteq \mathcal{M}_{A'}^*$. By applying Proposition 5.1 in case $A' \subsetneq A$, we get the desired result.

Suppose now that there are $r, r' \in A$ with $r < r' \leq 2r$. Then Lemma 5.3 yields a Borel reduction of embeddability between nontrivial graphs on ω to $\sqsubseteq_{\{0, r, r'\}}^*$. Now apply Proposition 5.1 again in case $\{0, r, r'\} \subsetneq A$. \square

Summing up, we have the following full description of the relations \sqsubseteq_A^* for $A \in \mathcal{D}$.

Theorem 5.12. *Let $A \in \mathcal{D}$.*

- (1) *The relations \sqsubseteq_A^* with A well-founded and well-spaced form a strictly increasing chain of order type $\omega + 1$ consisting of Borel quasi-orders, i.e. the quasi-orders \sqsubseteq_α^* for $\alpha \leq \omega$ from (i) and (iv) of Theorem 2.4(4). These relations are Borel reducible to all remaining $\sqsubseteq_{A'}^*$ for $A' \in \mathcal{D}$.*
- (2) *The relations \sqsubseteq_A^* when A is either ill-founded or not well-spaced are Borel bireducible with a complete analytic quasi-order.*

The following is the analogue of Theorem 5.8, but in this case the result is unconditional.

Corollary 5.13. *If $A \in \mathcal{D}$ then $\sqsubseteq_A \sim_B \sqsubseteq_A^*$.*

Proof. For any A we have $\sqsubseteq_A^* \leq_B \sqsubseteq_A$ because $\mathcal{M}_A^* \subseteq \mathcal{M}_A$.

If we are in case (1) of the previous Theorem, $\sqsubseteq_A \leq_B \sqsubseteq_A^*$ follows from Theorem 2.4(3) because in this case $\mathcal{M}_A = \mathcal{U}_A$ and $\mathcal{M}_A^* = \mathcal{U}_A^*$ by Theorem 3.6(2). If we are in case (2), notice that \sqsubseteq_A is Borel reducible to isometric embeddability on arbitrary Polish spaces. The latter, being an analytic quasi-order on a standard Borel space, is Borel reducible to any complete analytic quasi-order and hence to \sqsubseteq_A^* . \square

We observe that Theorem 5.12 holds even if we further restrict the relation \sqsubseteq_A^* to zero-dimensional spaces. In fact in case (1) all elements of \mathcal{M}_A^* are discrete and hence zero-dimensional, while in case (2), which is based on Proposition 5.11, we use Remark 5.2. For any other topological dimension we have that \mathcal{M}_A^* contains such spaces if and only if A includes a right neighborhood of 0 (by minor modifications of the proof of Theorem 3.6(1)). Theorem 5.12 holds also in this case by Remark 5.2, but only case (2) can occur. Similar observations apply to Corollary 5.13.

The analogous problems about isometry between spaces of fixed dimension (different from ∞) appear to be more delicate and are discussed more in detail in [CMMR18, Question 7.1 and the ensuing discussion]. The only result about isometry that we know still holds after fixing dimension is Theorem 5.8, because we can still use Remark 5.2.

Proposition 5.11 can be strengthened by replacing completeness with invariant universality (the notion originates in [FMR11], and was formally introduced in [CMMR13]).

Definition 5.14. Let the pair (S, E) consist of an analytic quasi-order S and an analytic equivalence relation $E \subseteq S$, with both relations defined on the same standard Borel space X . Then (S, E) is *invariantly universal* (for analytic quasi-orders) if for any analytic quasi-order R there is a Borel $B \subseteq X$ invariant under E such that $R \sim_B S \upharpoonright B$.

When E is isometry and S is isometric embeddability on some class of metric spaces, we just say that S is invariantly universal.

Notice that if (S, E) is invariantly universal, then S is complete for analytic quasi-orders.

A notion strictly connected to invariant universality is the following (see [CMMR18]). Given a pair (S, E) as above, we denote by S/E the E -quotient of S , i.e. the quasi-order on X/E induced by S . If F and E are equivalence relations on sets X and Y and $f: X/F \rightarrow Y/E$, then a *lifting* of f is a function $\hat{f}: X \rightarrow Y$ such that $[\hat{f}(x)]_E = f([x]_F)$ for every $x \in X$.

Definition 5.15. Let (R, F) and (S, E) be pairs consisting of a quasi-order and an equivalence relation on some standard Borel spaces, with $F \subseteq R$ and $E \subseteq S$.

We say that (R, F) is *classwise Borel isomorphic* to (S, E) , in symbols $(R, F) \simeq_{cB} (S, E)$, if there is an isomorphism of quasi-orders f between R/F and S/E such that both f and f^{-1} admit Borel liftings.

When the equivalence relations F and E are clear from the context we just say that R is classwise Borel isomorphic to S , and write $R \simeq_{cB} S$.

It is easy to see that if (R, F) is invariantly universal and for some Borel E -invariant B we have $(R, F) \simeq_{cB} (S \upharpoonright B, E \upharpoonright B)$ then (S, E) is invariantly universal as well.

Lemma 5.16. *Let $A = \{0, r, r'\}$ with $r < r' \leq 2r$. Then \mathcal{M}_A^* is Borel in $F(\mathbb{U})$ and \sqsubseteq_A^* is invariantly universal.*

Proof. The set \mathcal{M}_A^* is Borel by Theorem 4.8. We now show that embeddability between nontrivial countably infinite graphs is classwise Borel isomorphic to the restriction of \sqsubseteq_A^* to the class of infinite spaces. This suffices because embeddability between these graphs is invariantly universal by [FMR11]. First notice that the class of infinite spaces in \mathcal{M}_A^* is Borel because is the collection of $X \in \mathcal{M}_A^*$ satisfying

$$\forall n \in \omega \exists m \in \omega \forall i \leq n (\psi_m(X) \neq \psi_i(X)).$$

The classwise Borel isomorphism between the quotient quasi-orders is the quotient of the map introduced before Lemma 5.3. The inverse of this map is induced by the Borel function $X \mapsto G_X$ where (n, m) is an edge in G_X if and only if $d_X(\psi_n(X), \psi_m(X)) = r$. \square

Remark 5.17. The proof of [FMR11, Theorem 3.9] actually shows that the embeddability relation is already invariantly universal when restricted to the (Borel) class of connected graphs on ω : this ensures that the restriction of \sqsubseteq_A^* (for A as in the hypothesis of Lemma 5.16) to spaces in which every point realizes the distance r is still invariantly universal.

Lemma 5.18. *Let $A = \{0\} \cup \{r_n \mid n \in \omega\}$ be well-spaced with $(r_n)_{n \in \omega}$ a strictly decreasing sequence converging to 0. Then \mathcal{M}_A^* is Borel in $F(\mathbb{U})$ and \sqsubseteq_A^* is invariantly universal.*

Proof. The fact that under the hypotheses of the lemma the set \mathcal{M}_A^* is Borel follows from Theorem 3.6(2): since A is well-spaced, then $\mathcal{M}_A^* = \mathcal{U}_A^*$, and the latter is a Borel subset of $F(\mathbb{U})$. The fact that \sqsubseteq_A^* is invariantly universal is proved in [CMMR13, Theorem 5.19] for the special case $r_n = 2^{-n}$, but notice that the same proof works with any other choice for the r_n 's as well. \square

Remark 5.19. The proof of [CMMR13, Theorem 5.19] actually shows that \sqsubseteq_A^* (for A as in the hypothesis of Lemma 5.18) is already invariantly universal when restricted to spaces $X \in \mathcal{M}_A^*$ such that for all $n \in \omega$ there is $m \in \omega$ with $d_X(\psi_n(X), \psi_m(X)) = r_0$.

Theorem 5.20. *Let $A \in \mathcal{D}$, and assume that A is either ill-founded or not well-spaced. Then \sqsubseteq_A^* is invariantly universal, meaning that:³ For every analytic quasi-order R on a standard Borel space there is a Borel subset C' of $F(\mathbb{U})$ invariant under isometry such that $C' \subseteq \mathcal{M}_A^*$ and $R \sim_B \sqsubseteq \upharpoonright C'$.*

Proof. By Lemma 5.16 and Lemma 5.18, we can assume that A is neither of the form $\{0, r, r'\}$ with $r < r' \leq 2r$, nor of the form $\{0\} \cup \{r_n \mid n \in \omega\}$ with $(r_n)_{n \in \omega}$ a strictly decreasing sequence converging to 0 such that $2r_{n+1} < r_n$.

Claim 5.20.1. *There exist $A' \subseteq A$, $B \subseteq \mathcal{M}_{A'}^*$, $r_0 \in A'$, and $r_1, \bar{r} \in A$ such that:*

- (1) B is a Borel subset of $F(\mathbb{U})$ which is invariant under isometry;
- (2) $\sqsubseteq \upharpoonright B$ is invariantly universal;
- (3) $r_0, r_1 < \bar{r}$, $r_0 \neq r_1$, and $r \leq \bar{r}$ for all $r \in A'$;
- (4) for every $X \in B$ and for every countable dense $D \subseteq X$ it holds

$$\forall x \in X \exists y \in D (d_X(x, y) = r_0).$$

Proof. Suppose first that A is not well-spaced, and let $r, r' \in A$ be such that $r < r' \leq 2r$. Set $A' = \{0, r, r'\}$. Set $r_0 = r$, and pick $\bar{r} \in A \cap (r', +\infty)$ if such set is nonempty, and $\bar{r} = r'$ otherwise. Finally, let r_1 be any element of A distinct from r_0 and smaller than \bar{r} ; indeed, if $\bar{r} \neq r'$ we can just take $r_1 = r'$, otherwise the existence of such an r_1 is guaranteed by the fact that $\bar{r} = r' = \max A$ and $A \neq A'$. These choices ensure that (3) is satisfied.

Since $\mathcal{M}_{A'}^*$ is Borel by Lemma 5.16 the set $B \subseteq F(\mathbb{U})$ consisting of the $X \in \mathcal{M}_{A'}^*$ such that

$$\forall n \exists m \forall i \leq n (\psi_m(X) \neq \psi_i(X)) \wedge \forall n \exists m (d_X(\psi_n(X), \psi_m(X)) = r)$$

is Borel. Notice that every $X \in B$ is discrete and infinite, so that the ψ_n 's enumerate the entire X and no distances outside A' are possible. It easily follows that B is

³Formally, the definition of invariant universality is given only for quasi-orders with a standard Borel domain (Definition 5.15). Here we are considering the natural generalization of this concept to quasi-orders whose domain is an arbitrary Borel space.

invariant under isometry so that (1) is satisfied. Condition (4) is satisfied as well by the second part of the definition of B and the fact that we set $r_0 = r$. Finally, condition (2) follows from Lemma 5.16 and Remark 5.17.

Assume now that A is well-spaced and ill-founded. Notice that any strictly decreasing sequence $(r_n)_{n \in \omega}$ in A must converge to 0. Also we may assume without loss of generality that there is $\bar{r} \in A$ with $r_0 < \bar{r}$ (otherwise we shift the decreasing sequence by one). Then setting $A' = \{0\} \cup \{r_n \mid n \in \omega\}$ we get that (3) is satisfied. Moreover, A' is well-spaced by case assumption, hence $\mathcal{M}_{A'}^*$ is Borel by Lemma 5.18.

Let B be the collection of those $X \in \mathcal{M}_{A'}^*$ such that

$$\forall n \exists m (d_X(\psi_n(X), \psi_m(X)) = r_0).$$

The set B is clearly Borel in $F(\mathbb{U})$. We will now show that (4) is satisfied for such B : since condition (4) is preserved by isometry, it will also follow that B is invariant under isometry, i.e. that (1) is satisfied. So let $X \in B$, let D be dense in X , and let $x \in X$ be arbitrary. Let $n \in \omega$ be such that $d_X(\psi_n(X), x) < r_0$, and let $m \in \omega$ be such that $d_X(\psi_n(X), \psi_m(X)) = r_0$, which exists because $X \in B$. Then since X is ultrametric by Theorem 3.6(2) and the fact that A' is well-spaced, we also get $d_X(x, \psi_m(X)) = r_0$. Using the density of D , pick $y \in D$ such that $d_X(\psi_m(X), y) < r_0$: using again the fact that X is ultrametric, we get $d_X(x, y) = r_0$, as required. Finally, part (2) follows from Lemma 5.18 and Remark 5.19. \square

Claim 5.20.2. *Let $A', B \subseteq \mathcal{M}_{A'}^*$, r_0, r_1 , and \bar{r} be as in Claim 5.20.1. Then there is a Borel map $f: B \rightarrow \mathcal{M}_A^*$ such that*

- (1) $C = [f(B)]_{\cong} = \{Y \in F(\mathbb{U}) \mid \exists X \in B (f(X) \cong Y)\}$ is Borel (and obviously invariant under isometry);
- (2) f reduces $\sqsubseteq \upharpoonright B$ to \sqsubseteq_A^* and $\cong \upharpoonright B$ to \cong_A^* ;
- (3) there is a Borel $g: C \rightarrow B$ such that $(g \circ f)(X) \cong X$ for all $X \in B$ and $(f \circ g)(Y) \cong Y$ for all $Y \in C$.

Proof. Fix $W \in \mathcal{M}_{A \setminus \{r_0, r_1, \bar{r}\}}^*$, and let $Z = W \oplus_{\bar{r}} V$, where V consists of two points z_0 and z_1 at distance r_1 . Notice that by Fact 3.4 $Z \in \mathcal{M}_{A \setminus \{r_0\}}^*$ and that z_0, z_1 are the unique points of Z which realize the distance r_1 . Moreover, they are isolated in Z , so they belong to any dense subset of Z . Finally, notice that $d_Z(z_0, z) = d_Z(z_1, z)$ for any $z \in W$: it follows that if $X \in B \subseteq \mathcal{M}_{A'}^*$, the choice of the gluing point in X (because of Claim 5.20.1(3)) and of z_0 or z_1 as gluing point in Z does not change the space $X \oplus_{\bar{r}} Z$.

Let now f be the Borel map

$$f: B \rightarrow \mathcal{M}_A^*, \quad X \mapsto \tilde{X} = X \oplus_{\bar{r}} Z,$$

where X and Z are glued using one of z_0, z_1 . Notice that $\tilde{X} \in \mathcal{M}_A^*$ because $r_0 \in A'$, $X \in \mathcal{M}_{A'}^*$, and $Z \in \mathcal{M}_{A \setminus \{r_0\}}^*$, so that f is well-defined.

It is not hard to see that f satisfies (2). Indeed, if φ is an isometry (respectively, an isometric embedding) between $X, X' \in B$, then $\varphi \cup \text{id}_Z$ is an isometry (respectively, an isometric embedding) between \tilde{X} and \tilde{X}' . Conversely, if ψ is an isometry (respectively, an isometric embedding) between \tilde{X} and \tilde{X}' , then $\psi \upharpoonright X$ is an isometry (respectively, an isometric embedding) between X and X' because of Claim 5.20.1(4) and the fact that $r_0 \notin D(Z)$.

We now check that $C = [f(B)]_{\cong}$ is Borel in $F(\mathbb{U})$. First observe that since topologically \tilde{X} is a direct sum of X and Z , then for any dense subset D of \tilde{X} one

has $\forall x \in X \exists y \in D d_X(x, y) = r_0$. Together with the facts that $r_0 \notin D(Z)$ and $r_0 \neq \bar{r}$, this allows us to recover X from the space \tilde{X} as the completion (i.e. closure in \mathbb{U}) of

$$\{\psi_n(\tilde{X}) \mid \exists m(d(\psi_n(\tilde{X}), \psi_m(\tilde{X}))) = r_0\},$$

and Z as the completion of

$$\{\psi_n(\tilde{X}) \mid \neg \exists m(d(\psi_n(\tilde{X}), \psi_m(\tilde{X}))) = r_0\}.$$

We now generalize this process to an arbitrary $Y \in F(\mathbb{U})$. Let $\text{Rlz}(n, Y, r)$ be an abbreviation for the Borel condition

$$\exists m(d(\psi_n(Y), \psi_m(Y)) = r),$$

and set

$$X(Y) = \text{cl}(\{\psi_n(Y) \mid \text{Rlz}(n, Y, r_0)\})$$

and

$$Z(Y) = \text{cl}(\{\psi_n(Y) \mid \neg \text{Rlz}(n, Y, r_0)\}).$$

Notice that the maps $Y \mapsto X(Y)$ and $Y \mapsto Z(Y)$ are Borel functions from $F(\mathbb{U})$ into itself, and for $X \in B$ we have $X(\tilde{X}) \cong X$ and $Z(\tilde{X}) \cong Z$. Then for an arbitrary $Y \in F(\mathbb{U})$ we have that $Y \in C$ if and only if $\exists X \in B (Y \cong \tilde{X})$, if and only if $X(Y) \in B$, $Z(Y) \cong Z$, and $Y \cong X(Y) \oplus_{\bar{r}} Z(Y)$ (where the two spaces are glued via one of the only two points in $Z(Y)$ realizing r_1 in $Z(Y)$), if and only if

$$\begin{aligned} & \tilde{X}(Y) \in B \wedge Z(Y) \cong Z \wedge \exists n[\psi_n(Y) \in Z(Y) \wedge \text{Rlz}(n, Y, r_1) \wedge \\ & \forall m, k(\psi_m(Y) \in X(Y) \wedge \psi_k(Y) \in Z(Y) \Rightarrow d(\psi_m(Y), \psi_k(Y)) = \max\{\bar{r}, d(\psi_k(Y), \psi_n(Y))\})]. \end{aligned}$$

This is a Borel condition (here we are also using the fact that since isometry on $F(\mathbb{U})$ is Borel bi-reducible with an orbit equivalence relation, then the isometry class of a fixed element Z is Borel), whence C is a Borel subset of $F(\mathbb{U})$.

Finally, let

$$g: C \rightarrow B, \quad Y \mapsto X(Y),$$

where $X(Y)$ is as above. As already noticed, such map is Borel and $(g \circ f)(X) = g(\tilde{X}) = X(\tilde{X}) \cong X$ for every $X \in B$. Conversely, for every $Y \in C$ we have $(f \circ g)(Y) = \widetilde{X(Y)} \cong X(Y) \oplus_{\bar{r}} Z$: since $Y \cong X(Y) \oplus_{\bar{r}} Z(Y)$ where $Z(Y) \cong Z$ and $X(Y)$ and $Z(Y)$ are glued using one of the only two points of $Z(Y)$ which realize the distance r_1 in $Z(Y)$, it follows that $(f \circ g)(Y) \cong X(Y) \oplus_{\bar{r}} Z \cong X(Y) \oplus_{\bar{r}} Z(Y) \cong Y$. \square

Notice that conditions (2) and (3) of Claim 5.20.2 imply that g reduces $\sqsubseteq \upharpoonright C$ and $\cong \upharpoonright C$ to $\sqsubseteq \upharpoonright B$ and $\cong \upharpoonright B$, respectively.

Let now R be an arbitrary analytic quasi-order on a standard Borel space. By Claim 5.20.1 there is $B' \subseteq B$ Borel (in both B and $F(\mathbb{U})$) and invariant under isometry such that $R \sim_B \sqsubseteq \upharpoonright B'$; let $h_1: \text{dom}(R) \rightarrow B'$ and $h_2: B' \rightarrow \text{dom}(R)$ be witnesses of this last fact. Let $C' = [f(B')]_{\cong} = \{Y \in F(\mathbb{U}) \mid \exists X \in B'(f(X) \cong Y)\}$. Then C' is invariant under isometry by definition, and it is a Borel subset of $F(\mathbb{U})$: indeed, this follows from the fact that $C' = g^{-1}(B')$ by Claim 5.20.2(3), and that C is Borel in $F(\mathbb{U})$ by Claim 5.20.2(1). Moreover, $C' \subseteq C \subseteq \mathcal{M}_A^*$. Finally, $f \circ h_1$ witnesses $R \leq_B \sqsubseteq \upharpoonright C'$, while $h_2 \circ (g \upharpoonright C')$ witnesses $\sqsubseteq \upharpoonright C' \leq_B R$ by parts (2) and (3) of Claim 5.20.2. \square

6. OPEN PROBLEMS

Theorem 5.7 gives a fairly neat picture of the behaviour of the relations \cong_A^* , except for the following case.

Question 6.1. *If $A \in \mathcal{D}$ is dense in some right neighborhood of 0 but does not contain entirely any of them, can something more precise be said about the complexity of \cong_A^* ?*

Since for such A we have that \mathcal{M}_A^* consists only of zero-dimensional spaces by Theorem 3.6(1), this question is linked to one of the main questions of [GK03] which is still open (this question is discussed more at depth in [CMMR18], where it is labeled Question 7.1).

Question 6.2. *What is the complexity of isometry between zero-dimensional Polish metric spaces?*

A step to get some insight into Question 6.1 would be to answer the following:

Question 6.3. *Which A and A' as in Question 6.1 are such that $\cong_A^* \leq_B \cong_{A'}^*$?*

This question has a positive answer for a given pair A, A' whenever there exists $f: A \rightarrow A'$ which is injective and Polish metric preserving. In fact, arguing as in the proof of case (iii) of Theorem 5.8 one can build a witness for $\cong_A^* \leq_B \cong_{f(A)}^*$ by changing the distances of the spaces using f , and, if $f(A) \subsetneq A'$, successively applying Proposition 5.1 to obtain $\cong_{f(A)}^* \leq_B \cong_{A'}^*$. For example, in this way one sees that $\cong_{\mathbb{Q}^+}^* \leq_B \cong_{(\mathbb{R}^+ \setminus \mathbb{Q}) \cup \{0\}}^*$ via the map $f(q) = \sqrt{2}q$.

Another problem related to Question 6.1 is the following:

Question 6.4. *Is it the case that $\cong_A \sim_B \cong_A^*$ for every $A \in \mathcal{D}$?*

Theorem 5.8 and the results following it give a positive answer to this question under a wide range of hypotheses on A , but we do not know if there are distance sets which do not satisfy any of those hypotheses.

REFERENCES

- [AC19] Alessandro Andretta and Riccardo Camerlo. Analytic sets of reals and the density function in the Cantor space. *Eur. J. Math.*, 5(1):49–80, 2019.
- [CCM19] Riccardo Camerlo, Raphaël Carroy, and Alberto Marcone. Linear orders: when embeddability and epimorphism agree. *J. Math. Log.*, 19(1):1950003, 32, 2019.
- [CGK01] John D. Clemens, Su Gao, and Alexander S. Kechris. Polish metric spaces: their classification and isometry groups. *Bull. Symbolic Logic*, 7(3):361–375, 2001.
- [CK00] Riccardo Camerlo and Alexander S. Kechris. Countable structures with a fixed group of homeomorphisms. *Isr. J. Math.*, 117:105–124, 2000.
- [Cle01] John D. Clemens. *Descriptive Set Theory, Equivalence Relations, and Classification Problems in Analysis*. PhD thesis, University of California, Berkeley, 2001.
- [Cle07] John D. Clemens. Isometry of Polish metric spaces with a fixed set of distances. Preprint downloaded from <http://citeseerx.ist.psu.edu/viewdoc/summary?doi=10.1.1.434.8685>, 2007.
- [Cle12] John D. Clemens. Isometry of Polish metric spaces. *Ann. Pure Appl. Logic*, 163(9):1196–1209, 2012.
- [CMMR13] Riccardo Camerlo, Alberto Marcone, and Luca Motto Ros. Invariantly universal analytic quasi-orders. *Trans. Amer. Math. Soc.*, 365(4):1901–1931, 2013.
- [CMMR18] Riccardo Camerlo, Alberto Marcone, and Luca Motto Ros. On isometry and isometric embeddability between ultrametric Polish spaces. *Adv. Math.*, 329:1231–1284, 2018.

- [Cor99] Paul Corazza. Introduction to metric-preserving functions. *Amer. Math. Monthly*, 106(4):309–323, 1999.
- [DLPS07] Christian Delhommé, Claude Laflamme, Maurice Pouzet, and Norbert Sauer. Divisibility of countable metric spaces. *European J. Combin.*, 28(6):1746–1769, 2007.
- [Dob95] Jozef Doboš. A survey of metric preserving functions. *Questions Answers Gen. Topology*, 13(2):129–134, 1995.
- [FMR11] Sy-David Friedman and Luca Motto Ros. Analytic equivalence relations and bi-embeddability. *J. Symbolic Logic*, 76(1):243–266, 2011.
- [FS89] Harvey Friedman and Lee Stanley. A Borel reducibility theory for classes of countable structures. *J. Symbolic Logic*, 54(3):894–914, 1989.
- [Gao09] Su Gao. *Invariant descriptive set theory*, volume 293 of *Pure and Applied Mathematics*. CRC Press, Boca Raton, 2009.
- [GK03] Su Gao and Alexander S. Kechris. On the classification of Polish metric spaces up to isometry. *Mem. Amer. Math. Soc.*, 161(766):viii+78, 2003.
- [Gro99] Misha Gromov. *Metric structures for Riemannian and non-Riemannian spaces*, volume 152 of *Progress in Mathematics*. Birkhäuser Boston, Inc., Boston, MA, 1999.
- [GS11] Su Gao and Chuang Shao. Polish ultrametric Urysohn spaces and their isometry groups. *Topology Appl.*, 158(3):492–508, 2011.
- [HKL90] Leo A. Harrington, Alexander S. Kechris, and Alain Louveau. A Glimm-Effros dichotomy for Borel equivalence relations. *J. Amer. Math. Soc.*, 3(4):903–928, 1990.
- [Kec95] Alexander S. Kechris. *Classical descriptive set theory*, volume 156 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1995.
- [Kec97] Alexander S. Kechris. On the concept of Π_1^1 -completeness. *Proc. Amer. Math. Soc.*, 125(6):1811–1814, 1997.
- [LR05] Alain Louveau and Christian Rosendal. Complete analytic equivalence relations. *Trans. Amer. Math. Soc.*, 357(12):4839–4866, 2005.
- [MR17] Luca Motto Ros. Can we classify complete metric spaces up to isometry? *Boll. Unione Mat. Ital.*, 10(3):369–410, 2017.
- [Sau13] Norbert W. Sauer. Distance sets of Urysohn metric spaces. *Canad. J. Math.*, 65(1):222–240, 2013.
- [Ver98] Anatoly M. Vershik. The universal Uryson space, Gromov’s metric triples, and random metrics on the series of natural numbers. *Uspekhi Mat. Nauk*, 53(5(323)):57–64, 1998.

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI GENOVA, VIA DODECANESO 35, 16146 GENOVA — ITALY

Email address: `camerlo@dima.unige.it`

DIPARTIMENTO DI SCIENZE MATEMATICHE, INFORMATICHE E FISICHE, UNIVERSITÀ DI UDINE, VIA DELLE SCIENZE 208, 33100 UDINE — ITALY

Email address: `alberto.marcone@uniud.it`

DIPARTIMENTO DI MATEMATICA «GIUSEPPE PEANO», UNIVERSITÀ DI TORINO, VIA CARLO ALBERTO 10, 10123 TORINO — ITALY

Email address: `luca.mottoros@unito.it`