

Università degli studi di Udine

GLOBAL EXISTENCE OF SOLUTIONS FOR A MULTI-PHASE FLOW: A BUBBLE IN A LIQUID TUBE AND RELATED CASES

Original

Availability: This version is available http://hdl.handle.net/11390/1072077 since 2021-03-25T11:16:31Z

Publisher:

Published DOI:10.1016/S0252-9602(15)30024-2

Terms of use:

The institutional repository of the University of Udine (http://air.uniud.it) is provided by ARIC services. The aim is to enable open access to all the world.

Publisher copyright

(Article begins on next page)



Available online at www.sciencedirect.com ScienceDirect Acta Mathematica Scientia 2015,35B(4):1–23



http://actams.wipm.ac.cn

GLOBAL EXISTENCE OF SOLUTIONS FOR A MULTI-PHASE FLOW: A BUBBLE IN A LIQUID TUBE AND RELATED CASES*

Dedicated to Professor Tai-Ping Liu on the occasion of his 70th birthday

Debora AMADORI¹ Paolo BAITI² Andrea CORLI³ Edda DAL SANTO¹

1. Department of Engineering and Computer Science and Mathematics, University of L'Aquila, Italy

2. Department of Mathematics and Computer Science, University of Udine, Italy

3. Department of Mathematics and Computer Science, University of Ferrara, Italy E-mail: amadori@univaq.it; paolo.bciti@uniud.it; crl@unife.it

Abstract In this paper we study the problem of the global existence (in time) of weak, entropic solutions to a system of three hyperbolic conservation laws, in one space dimension, for large initial data. The system models the dynamics of phase transitions in an isothermal fluid; in Lagrangian coordinates, the phase interfaces are represented as stationary contact discontinuities. We focus on the persistence of solutions consisting in three bulk phases separated by two interfaces. Under some stability conditions on the phase configuration and by a suitable front tracking algorithm we show that, if the **BV**-norm of the initial data is less than an explicit (large) threshold, then the Cauchy problem has global solutions.

Key words hyperbolic systems of conservation laws; phase transitions; wave-front tracking algorithm

2010 MR Subject Classification 35L65; 35L60; 35L67; 76T99

1 Introduction

This paper concludes a long analysis, begun in [1, 2], concerning the global existence in time of weak entropy solutions for a system of conservation laws modeling phase transitions in a fluid. More precisely, the focus of the analysis is on the persistence of solutions with stationary interfaces.

The system under consideration consists of three equations, namely,

$$\begin{cases} v_t - u_x = 0, \\ u_t + p(v, \lambda)_x = 0, \\ \lambda_t = 0, \end{cases}$$
(1.1)

where the state variables $(v, u, \lambda) \in \Omega :=]0, +\infty[\times \mathbb{R} \times [0, 1]]$ denote the specific volume, the velocity and the mass-density fraction of the vapor in the fluid, respectively. System (1.1) is

^{*}Received February 21, 2015.

the conservative part of a more complex model first introduced in [11], one of whose novelties lies in the pressure p, which depends not only on v but also on λ . More precisely, the pressure is prescribed by the law

$$p(v,\lambda) = \frac{a^2(\lambda)}{v},\tag{1.2}$$

where the **C**¹ function *a* is assumed to be strictly positive in [0, 1]. For example, we can take $p = (1 + \lambda)k/v$ for some positive constant *k*.

A related example of pressure law occurs in the theory of (non-isothermal) ionized gases. In that case, we have $p = (1 + \alpha)RT/mv$, where R, m and T are the universal gas constant, the molecular mass and the temperature, respectively, while α is the ionization degree; if $\alpha = 0$ the gas is not ionized. In that model, however, α is usually assigned as a given function of both p and T by Saha's law [12] while, in the case under consideration, the mass-density fraction λ is understood as an independent variable.

System (1.1) is strictly hyperbolic in Ω ; the eigenvalues $\pm a/v$ are genuinely nonlinear while 0 is linearly degenerate. As a consequence, the field associated to this latter eigenvalue supports contact discontinuities, which are understood in the model as phase interfaces.

The Cauchy problem for (1.1) includes the initial data

$$(v_o(x), u_o(x), \lambda_o(x)), \qquad v_o(x) \ge \underline{v} > 0, \ x \in \mathbb{R}.$$
(1.3)

The global existence (in time) of solutions to the initial-value problem for any strictly hyperbolic system of conservation laws whose eigenvalues do not change type is well-known and can be proved either with the Glimm scheme or with a front tracking algorithm, see [7, 8]. The case of large initial data is a challenging problem and can be tackled only for special systems; then, the issue is to find classes of initial data with (large) total variation for which global solutions to the Cauchy problem exist. This topic was studied in [18] and [19] in the case of the isothermal, respectively isentropic, *p*-system, see also [9, 10]; these results were extended in the seminal papers [16, 17] to the case of nonisentropic gas dynamics. We also refer to [14] for the extension of Nishida's result to the initial-value problem in Special Relativity.

As far as (1.1)-(1.3) is concerned, a positive answer was first given in [3] and then in [5]. In particular, in the former paper an explicit threshold of the **BV**-norm of the initial data was provided in order to have the global existence of solutions. Moreover, both papers require a sort of balance of the **BV**-norm of the initial data: the larger the variation of (v_o, u_o) , the smaller the variation of λ_o and vice-versa. It would be interesting to prove whether or not, for general **BV**-data, the solution exists globally in time. Motivated by the techniques introduced in [3], we considered in [1] the special case of initial data (1.3) where

$$\lambda_o(x) = \begin{cases} \lambda_\ell & \text{if } x < 0, \\ \lambda_r & \text{if } x > 0, \end{cases}$$

which models the dynamics of a two-phases fluid. In order to deal with this particular framework, an original Riemann solver was proposed by which we proved the global existence of solutions for a wide class of large initial data. Such results improved by far those of [3] when adapted to that setting. We briefly mention that the problem of smooth perturbations of Riemann-type solutions has been studied by many authors, see for instance [15].

3

The next step regarded the case of initial data with two phase interfaces, namely,

$$\lambda_o(x) = \begin{cases} \lambda_\ell & \text{if } x < a, \\ \lambda_m & \text{if } a < x < b, \\ \lambda_r & \text{if } x > b, \end{cases}$$
(1.4)

where $\lambda_{\ell}, \lambda_m, \lambda_r$ are constant in [0, 1] and $\mathbf{a} < \mathbf{b}$ are the location of the waves; denote $a_{\ell} = a(\lambda_{\ell})$, $a_m = a(\lambda_m)$, $a_r = a(\lambda_r)$. Clearly, this case is much more complicated than the previous one, because of the possible bouncing back and forward of the waves in the middle region $[\mathbf{a}, \mathbf{b}]$. In [2] we answered in the positive to the above issue by assuming the condition $a_m < \min\{a_{\ell}, a_r\}$; if v is fixed, then the pressure in the middle region is lower than the pressure in the outside regions. We also recovered the results of [1] by passing to the limit when either $a_m \to a_{\ell}$ or $a_m \to a_r$. To give a physical flavor to the problem, assume for a moment that the function $a(\lambda)$ is increasing with λ , as is the interesting case in modeling. Then the condition above also means that the mass-density fraction of vapor inside $[\mathbf{a}, \mathbf{b}]$ is less than outside: for brevity, we address to this case as the drop case. We refer to [1–3] for more details on the model and references.

In this paper, we deal with the two remaining cases, namely,

either
$$a_m > \min\{a_\ell, a_r\}$$
 or $a_\ell < a_m < a_r$,

since the case $a_{\ell} > a_m > a_r$ can be deduced by the latter. Reminding of the previous physical interpretation, we shall loosely address to these cases as the bubble case and the increasingpressure case, respectively. As in [2], we introduce two special Riemann solvers. Both of them replace the standard non-physical waves [7] by waves defined through integral curves; this allows us to attach these waves to the phase waves. Then, following again the lines of [2], we introduce an "asymmetrical" Glimm functional F, in the sense that its interaction potential Q takes into account only certain shock waves approaching the phase waves while, on the contrary, all rarefaction waves approaching the phase waves are included. We point out that the definition of F differs in each of the three cases mentioned above.

A key feature of both [2] and the current paper is the possible occurrence of a stability condition, depending on the case under consideration, to control interactions in the middle zone. More precisely, such a condition is needed both in the drop and in the increasing-pressure cases and imposes bounds to the strengths of the phase waves. However, in the bubble case we require no stability condition and, as a consequence, any phase wave is admitted.

The plan of the paper is the following. The main result is stated in Section 2. In Section 3 we introduce the Riemann solvers, the composite waves and, at last, we define the functional F; we also recall some background facts from [1–3]. Sections 4 and 5 focus on the bubble and on the monotone-pressure cases, respectively; in particular, we prove there that the functional F decreases at every interaction. The last Section 6 deals with the two cases at the same time; by showing the convergence of the front tracking algorithm we conclude the proof of the main result.

2 Main Results

In this section we state the main results of this paper, which concerns the existence of solutions to the Cauchy problem (1.1)-(1.3), (1.4). Under the notation defined in the Introduction,

1

we set

$$\gamma = 2 \frac{a_m - a_\ell}{a_m + a_\ell}, \qquad \zeta = 2 \frac{a_r - a_m}{a_r + a_m}.$$
 (2.1)

The quantities η and ζ range over]-2, 2[; they are the strengths of the two contact discontinuities (see (3.4) below) that support the phase interfaces of the model, located at $x = \mathbf{a}$ and $x = \mathbf{b}$, respectively. We denote by $\mathcal{L}, \mathcal{M}, \mathcal{R}$ the three regions separated by η and ζ in the (x, t)-plane, see Figure 1.

As already mentioned in the Introduction, we focus on the bubble case, which corresponds to consider $\eta > 0$ and $\zeta < 0$, and on the increasing-pressure case, which corresponds to both $\eta > 0$ and $\zeta > 0$.



Fig.1 The two phase waves with strengths η and ζ in the (x, t)-plane and the regions $\mathcal{L}, \mathcal{M}, \mathcal{R}$.

In order to state the existence theorem, we introduce some threshold functions. First, as in [1, 2], we define the strictly decreasing function

$$\mathcal{K}(r) := \frac{2}{1+r} \log \left(1 + \frac{2}{r} \left(1 + \sqrt{1+r} \right) \right), \qquad r \in \mathbb{R}^+,$$
(2.2)

which plays a key role in the main results; we notice that $\lim_{r \to 0^+} \mathcal{K}(r) = +\infty$ and $\lim_{r \to +\infty} \mathcal{K}(r) = 0$. Moreover, as in [2], we need another function related to the stability of the two phase waves configuration, which differs from that in [2]. In the bubble case it is

$$\mathcal{H}_{b}(|\eta|,|\zeta|) := \frac{4}{4 - |\eta\zeta|} \max\left\{ |\eta| \frac{2 + |\zeta|}{2 - |\zeta|}, |\zeta| \frac{2 + |\eta|}{2 - |\eta|} \right\},$$
(2.3)



Fig.2 The domain D_c in the $(|\eta|, |\zeta|)$ -plane

for $(|\eta|, |\zeta|) \in D_b := [0, 2[\times[0, 2[$. As for the increasing-pressure case, the definition of the function \mathcal{H}_c as well as that of its domain D_c is more complicated though explicit; we refer to (5.23) and (5.22) below, respectively, and to Figure 2 for a picture of D_c . We can immediately observe that in the bubble case the pair $(|\eta|, |\zeta|)$ can vary inside the whole square $[0, 2[\times[0, 2[$, while in the increasing-pressure case they can cover only a portion of it.

We denote $p_o(x) = p(v_o(x), \lambda_o(x))$. The following theorem states the global in time existence of solutions in the bubble and in the increasing-pressure case. Notice that the statement is the same in both cases.

Theorem 2.1 Assume (1.2) and consider initial data (1.3), (1.4). Let $D = D_b$ and $\mathcal{H} = \mathcal{H}_b$ in the bubble case, $D = D_c$ and $\mathcal{H} = \mathcal{H}_c$ in the increasing-pressure case. Moreover, assume that for η, ζ as in (2.1), the pair $(|\eta|, |\zeta|)$ belongs to D. If

$$\operatorname{TV}\left(\log(p_o)\right) + \frac{1}{\min\{a_\ell, a_m, a_r\}} \operatorname{TV}\left(u_o\right) < \mathcal{K}\left(\mathcal{H}(|\eta|, |\zeta|)\right)$$
(2.4)

holds, then the Cauchy problem (1.1)–(1.3), (1.4) has a weak entropic solution (v, u, λ) defined for $t \in [0, +\infty)$. If $\eta = \zeta = 0$ the same conclusion holds with $\mathcal{K}(\mathcal{H}(|\eta|, |\zeta|))$ replaced by $+\infty$ in (2.4).

Moreover, the solution is valued in a compact set and $(v(t, \cdot), u(t, \cdot)) \in L^{\infty}([0, \infty[; \mathbf{BV}(\mathbb{R})))$.

The sub-level sets $S_h = \{(|\eta|, |\zeta|) \in \mathcal{D} : \mathcal{H}(|\eta|, |\zeta|) < h\}, h > 0$, of the function \mathcal{H} play an important role in condition (2.4), see for instance Figure 3 in the bubble case. Indeed, condition (2.4) holds for every $(|\eta|, |\zeta|) \in S_h$ if

$$\operatorname{TV}\left(\log(p_o)\right) + \frac{1}{\min\{a_\ell, a_m, a_r\}} \operatorname{TV}\left(u_o\right) < \mathcal{K}(h),$$

since \mathcal{K} is decreasing. When h = 2, we have $\mathcal{K}(2) = 2\log(2+\sqrt{3})/3$ and the case is particularly significative; let us consider, for example, the bubble framework. As in the drop case [2], the domain \mathcal{S}_2 includes the segments [0, 2[on each axis, but the 2-level set of \mathcal{H}_b is no more the graph of the function $\zeta(|\eta|) = 2(2-|\eta|)/(2+|\eta|)$, see Figure 3.



Fig.3 Sets of level c of the function \mathcal{H}_b : cases c = 1, 2, 3; the thin line is the curve $\zeta = \zeta(|\eta|)$

3 Preliminaries and Functionals

In this section we collect some preliminary results from [1-3], focusing on the front tracking algorithm used to construct the approximate solutions. Moreover, we introduce some functionals needed to estimate the total variation of such solutions.

First, we recall some basic notions. System (1.1) is strictly hyperbolic with two genuinely nonlinear characteristic fields (of family 1 and 3) and a linearly degenerate one of family 2. For i = 1, 3, the *i*-th right Lax curves through the point $\hat{U} = (\hat{v}, \hat{u}, \hat{\lambda}) \in \Omega$ are

$$v \mapsto \left(v, \hat{u} + 2a(\hat{\lambda})h(\varepsilon_i), \hat{\lambda}\right), \qquad v > 0,$$

$$(3.1)$$

where ε_i denotes the strength of an *i*-wave,

$$\varepsilon_1 = \frac{1}{2} \log\left(\frac{v}{\hat{v}}\right), \qquad \varepsilon_3 = \frac{1}{2} \log\left(\frac{\hat{v}}{v}\right),$$
(3.2)

and h is the function defined by

$$h(\varepsilon) = \begin{cases} \varepsilon & \text{if } \varepsilon \ge 0 \,,\\ \sinh \varepsilon & \text{if } \varepsilon < 0 \,. \end{cases}$$
(3.3)

Rarefaction waves have positive strength, while shock waves have negative strength. The wave curve for the second characteristic field through $\hat{U} \in \Omega$ is given by

$$\lambda \mapsto \left(\hat{v} \frac{a^2(\lambda)}{a^2(\hat{\lambda})}, \hat{u}, \lambda \right), \qquad \lambda \in [0, 1],$$

and the strength of a 2-wave is

$$\delta = 2 \frac{a(\lambda) - a(\hat{\lambda})}{a(\lambda) + a(\hat{\lambda})}.$$
(3.4)

We solve the Riemann problems by means of some Pre-Riemann solvers, which are introduced in the following proposition. We use the symbols 'L' and 'I' to denote Lax and Integral curves, respectively. For i = 1, 3 and $\theta_i \in \{L, I\}$, we define the functions

$$\Theta_i = \begin{cases} h & \text{if } \theta_i = L, \\ Id & \text{if } \theta_i = I. \end{cases}$$
(3.5)

Proposition 3.1 (Pre-Riemann solvers) Fix $\theta_i \in \{L, I\}$, for i = 1, 3. There exists a map $R_{\theta_1\theta_3} : \Omega \times \Omega \to \mathbb{R} \times] - 2, 2[\times \mathbb{R}$ such that for any two states $U_- = (v_-, u_-, \lambda_-), U_+ = (v_+, u_+, \lambda_+) \in \Omega$ we have

$$R_{\theta_1\theta_3}(U_-, U_+) = (\varepsilon_1, \delta, \varepsilon_3), \qquad (3.6)$$

where $\varepsilon_1, \delta, \varepsilon_3$ represent waves of family 1, 2, 3, respectively, satisfying the following relations:

$$\varepsilon_3 - \varepsilon_1 = \frac{1}{2} \log\left(\frac{p_+}{p_-}\right), \quad a_-\Theta_1(\varepsilon_1) + a_+\Theta_3(\varepsilon_3) = \frac{u_+ - u_-}{2}, \quad \delta = 2\frac{a_+ - a_-}{a_+ + a_-}.$$
 (3.7)

We denoted $a_{\pm} = a(\lambda_{\pm}), p_{\pm} = p(v_{\pm}, \lambda_{\pm}).$

We refer to [2, Proposition 3.1] for a proof of the previous result. In a few words, Proposition 3.1 states the existence of four Pre-Riemann solvers R_{LL} , R_{II} , R_{LI} and R_{IL} , that prescribe how to solve a Riemann problem with *i*-waves taken along (Lax or integral) *i*-curves, for i = 1, 3. In particular, R_{LL} is the solver of [1, 3] that employs Lax curves.

No.4

We use a front tracking algorithm [7] to build up the approximate solutions to (1.1)-(1.3), (1.4). The first step in the construction consists in taking a sequence $(v_{\sigma}^{\nu}, u_{\sigma}^{\nu})_{\nu \in \mathbb{N}}$ of piecewiseconstant functions with a finite number of jumps, that approximate the initial data (1.4) in the sense of [2, Section 4]. We choose two parameters $\sigma = \sigma_{\nu} > 0$, $\rho = \rho_{\nu} > 0$ and proceed as follows. At time t = 0 we apply the solver R_{LL} at each jump of the approximated initial data; we split rarefactions into a finite number of rarefaction shocks, each of size $\leq \sigma$, whose speed equals the characteristic speed at the right state (see [2, Section 4] for more details). Then, an approximate solution $(v^{\nu}, u^{\nu}, \lambda)(\cdot, t)$ is defined until the first time two wave fronts interact and a new Riemann problem arises. In the case of interaction between two fronts of families 1 or 3, we again use R_{LL} and adopt the following strategy to approximate outgoing rarefaction, otherwise they are split into a fan of waves as before. On the other hand, when solving an interaction of a wave of family 1 or 3 with a 2-wave δ , we possibly make use of a method that attaches certain reflected waves to δ ; the outcome is a wave of the same family of the incoming one and a stationary composite wave, which is defined below.

Definition 3.2 (Composite wave [2]) Consider two states $U_{-} = (v_{-}, u_{-}, \lambda_{-})$ and $U_{+} = (v_{+}, u_{+}, \lambda_{+})$ of Ω , with $\lambda_{-} \neq \lambda_{+}$. The composite wave $\delta_{0} = (\delta_{0}^{1}, \delta, \delta_{0}^{3})$ connecting U_{-} to U_{+} is the stationary wave defined by $\delta_{0} = R_{II}(U_{-}, U_{+})$. We write $|\delta_{0}| = |\delta_{0}^{1}| + |\delta_{0}^{3}|$.



Fig.4 A composite wave δ_0 in the (x, t)-plane: in (a) it is drawn as three parallel close lines, while the auxiliary picture (b) is used to determine the states in the interactions

Notice that in Definition 3.2 the waves δ_0^1, δ_0^3 are given zero speed and δ_0 reduces to a 2-wave as long as $\delta_0^1 = \delta_0^3 = 0$. Hence, instead of dealing with 2-waves, we are left with composite waves belonging to a fictitious 0-family.

When a wave front of family 1 or 3 with strength δ_i interacts with a composite wave $\delta_0 = (\delta_0^1, \delta, \delta_0^3)$ at t > 0, we exploit two different procedures to solve the emerging Riemann problem of states U_-, U_+ . We indicate by $I_i(\varepsilon)(U)$ the integral curves of family i = 1, 3, parametrized by ε and of origin $U \in \Omega$; then, we define $\widetilde{U}_- = I_1(\delta_0^1)(U_-)$ and $\widetilde{U}_+ = I_3(-\delta_0^3)(U_+)$. Here follows a description of the Riemann solvers; for a proof of the following facts see [2, Proposition 3.7].

(1) Accurate Riemann solver. If $|\delta_i| \ge \rho$, then the solution is formed by waves $\varepsilon_1, \varepsilon_0, \varepsilon_3$, where $(\varepsilon_1, \delta, \varepsilon_3) = R_{LL}(\widetilde{U}_-, \widetilde{U}_+)$ and $\varepsilon_0 = \delta_0$.

(2) Simplified Riemann solver. If $|\delta_i| < \rho$, then we distinguish case i = 1 and i = 3:

i) for i = 1, the solution is formed by waves $\varepsilon_1, \varepsilon_0$ such that $(\varepsilon_1, \delta, \varepsilon_3) = R_{LI}(\widetilde{U}_-, \widetilde{U}_+)$ and $\varepsilon_0 = (\delta_0^1, \delta, \delta_0^3 + \varepsilon_3);$

ii) for i = 3, the solution is formed by waves $\varepsilon_0, \varepsilon_3$ such that $(\varepsilon_1, \delta, \varepsilon_3) = R_{IL}(\widetilde{U}_-, \widetilde{U}_+)$ and $\varepsilon_0 = (\delta_0^1 + \varepsilon_1, \delta, \delta_0^3)$.

We emphasize that $\Theta_i = h$ in all cases and the following relations are verified:

$$\varepsilon_3 - \varepsilon_1 = \begin{cases} -\delta_1 & \text{if } i = 1, \\ \delta_3 & \text{if } i = 3, \end{cases} \quad a_-\Theta_1(\varepsilon_1) + a_+\Theta_3(\varepsilon_3) = \begin{cases} a_+\Theta_1(\delta_1) & \text{if } i = 1, \\ a_-\Theta_3(\delta_3) & \text{if } i = 3, \end{cases}$$
(3.8)

$$\operatorname{sgn} \varepsilon_{i} = \operatorname{sgn} \delta_{i}, \qquad \operatorname{sgn} \varepsilon_{j} = \begin{cases} \operatorname{sgn} \delta \cdot \operatorname{sgn} \delta_{i} & \text{if } i = 1, \\ -\operatorname{sgn} \delta \cdot \operatorname{sgn} \delta_{i} & \text{if } i = 3. \end{cases}$$
(3.9)

We recall the interaction estimates [2, Lemma 5.2 and 5.4]. For i = 1, 3, we denote by ε_i the strength of the transmitted wave and by ε_j , j = 1, 3, $j \neq i$, the strength of the reflected one (even in case of interaction with a composite wave treated by the Simplified solver).

Lemma 3.3 (Interaction estimates) For the interaction between two waves at time t > 0 we have the following; let $i, j = 1, 3, j \neq i$.

(1) Assume that an *i*-wave δ_i interacts with a composite wave $\delta_0 = (\delta_0^1, \delta, \delta_0^3)$. If $|\delta_i| \ge \rho$, we have

$$|\varepsilon_i - \delta_i| = |\varepsilon_j| \le \frac{1}{2} |\delta_i \delta|$$
 and $|\varepsilon_0 - \delta_0| = 0,$ (3.10)

while, if $|\delta_i| < \rho$, it holds

$$|\varepsilon_i - \delta_i| = |\varepsilon_0 - \delta_0| = |\varepsilon_j| \le \begin{cases} \frac{C_o}{2} |\delta_i \delta| & \text{if } \delta_i < 0 \text{ and either } (i = 1, \, \delta > 0) \text{ or } (i = 3, \, \delta < 0), \\ \frac{1}{2} |\delta_i \delta| & \text{otherwise,} \end{cases}$$

$$(3.11)$$

where

$$C_o = C_o(\rho) = \frac{\sinh \rho}{\rho} \,. \tag{3.12}$$

(2) If two waves α_i and β_j of different families interact with each other, then,

$$|\varepsilon_i| = |\alpha_i|, \qquad |\varepsilon_j| = |\beta_j|.$$
(3.13)

(3) Assume that two waves α_i and β_i of the same family interact. If both α_i and β_i are shocks, then the reflected wave ε_j is a rarefaction, while the transmitted wave ε_i is a shock and satisfies

$$|\varepsilon_i| > \max\{|\alpha_i|, |\beta_i|\}.$$
(3.14)

If α_i and β_i have different signs, e.g. $\alpha_i < 0 < \beta_i$, then the reflected wave is a shock, both the amounts of shocks and rarefactions of the *i*-th family decrease across the interaction and one has

$$|\varepsilon_j| \le c(\alpha_i) \cdot \min\{|\alpha_i|, |\beta_i|\}, \qquad c(z) = \frac{\cosh z - 1}{\cosh z + 1}.$$
(3.15)

About the term in C_o in (3.12), we notice that $C_o(\rho) > 1$ for any $\rho > 0$ and $C_o(\rho) \to 1^+$ for $\rho \to 0^+$.

Now, we introduce some functionals needed to prove the boundedness of the total variation of the approximate solutions. Using indices ℓ, m, r to refer to waves in $\mathcal{L}, \mathcal{M}, \mathcal{R}$, respectively, we define

$$L^{\ell,m,r} = \sum_{\substack{i=1,3,\,\delta_i > 0\\\delta_i \in \mathcal{L},\mathcal{M},\mathcal{R}}} |\delta_i| + \xi \sum_{\substack{i=1,3,\,\delta_i < 0\\\delta_i \in \mathcal{L},\mathcal{M},\mathcal{R}}} |\delta_i| \,,$$

where $\xi > 1$ is a parameter to be determined. We denote

$$L = L^{\ell} + L^m + L^r$$
, $L^0 = |\eta_0| + |\zeta_0|$

and

No.4

$$\bar{L} = \bar{L}^{\ell} + \bar{L}^{m} + \bar{L}^{r} = \sum_{\substack{i=1,3\\\delta_{i} \in \mathcal{L}}} |\delta_{i}| + \sum_{\substack{i=1,3\\\delta_{i} \in \mathcal{M}}} |\delta_{i}| + \sum_{\substack{i=1,3\\\delta_{i} \in \mathcal{R}}} |\delta_{i}| = \frac{1}{2} \operatorname{TV} \left(\log p(t, \cdot)\right) - |\eta_{0}| - |\zeta_{0}|.$$

In the following sections we also introduce the interaction potential

$$Q = Q^{\ell} + Q^m + Q^r \tag{3.16}$$

and specify $Q^{\ell,m,r}$, that differ in the bubble case and in the increasing-pressure case. The resulting functional

$$F = L + L^0 + Q \tag{3.17}$$

is equivalent to the total variation of the approximate solutions and has an asymmetrical character in a double sense: firstly, it depends on the phases and, secondly, shocks and rarefactions play a different role. As for the latter, not only shocks are weighted by ξ (a procedure of [4] also exploited in [1, 2]), but the contributions of certain shock waves from Q are dropped. This is due to the fact that in the interaction of some shocks with a 0-wave we have that ΔL is already nonpositive; more precisely, from (3.8) it follows that such unnecessary waves are either 3-shocks interacting with a 2-wave $\delta > 0$ or 1-shocks interacting with a 2-wave $\delta < 0$. On the contrary, rarefaction waves are always counted in Q if they approach a phase wave. As shown in Figure 5, the total variation of the solutions increases going towards the more liquid regions.



Fig.5 How the total variation varies for interactions with the phase waves in the bubble case (a) and in the increasing-pressure case (b)

In the potentials $Q^{\ell,m,r}$ we insert some positive weights $K_{\eta,\zeta}^{\ell,m,r}$ that keep track of the regions of provenience $(\mathcal{L}, \mathcal{M}, \mathcal{R})$ of the approaching waves and of the 0-wave approached $(\eta_0 \text{ or } \zeta_0)$, see Figure 6.

In the next sections we show that the functional F decreases across any interaction, under suitable conditions on the parameters ξ , $K_{\eta,\zeta}^{\ell,m,r}$ and ρ , which differ in the bubble case and in the increasing-pressure case. In the proof we fix $m_o > 0$ and assume that, for i = 1, 3, the size δ_i of any *i*-shock satisfies

$$|\delta_i| \le m_o \,. \tag{3.18}$$

Such parameter is necessary in our proof to relate the total variation of the initial data u_o, p_o to the sizes of the phase waves.



Fig.6 The parameters $K_{\eta,\zeta}^{\ell,m,r}$

4 The Bubble Case

In this section, we prove the decreasing of the functional F in (3.17) in the bubble case, where the phase-dependent interaction potentials are defined by

$$\begin{split} Q^{\ell} &= \left(K_{\eta}^{\ell} |\eta| + K_{\zeta}^{\ell} |\zeta|\right) \sum_{\substack{\delta_{3} > 0 \\ \delta_{3} \in \mathcal{L}}} |\delta_{3}| + \xi K_{\zeta}^{\ell} \sum_{\substack{\delta_{3} < 0 \\ \delta_{3} \in \mathcal{L}}} |\delta_{3}\zeta| \,, \\ Q^{m} &= K_{\eta}^{m} \left(\sum_{\substack{\delta_{1} > 0 \\ \delta_{1} \in \mathcal{M}}} |\delta_{1}\eta| + \xi \sum_{\substack{\delta_{1} < 0 \\ \delta_{1} \in \mathcal{M}}} |\delta_{1}\eta|\right) + K_{\zeta}^{m} \left(\sum_{\substack{\delta_{3} > 0 \\ \delta_{3} \in \mathcal{M}}} |\delta_{3}\zeta| + \xi \sum_{\substack{\delta_{3} < 0 \\ \delta_{3} \in \mathcal{M}}} |\delta_{3}\zeta|\right), \\ Q^{r} &= \left(K_{\eta}^{r} |\eta| + K_{\zeta}^{r} |\zeta|\right) \sum_{\substack{\delta_{1} > 0 \\ \delta_{1} \in \mathcal{R}}} |\delta_{1}| + \xi K_{\eta}^{r} \sum_{\substack{\delta_{1} < 0 \\ \delta_{1} \in \mathcal{R}}} |\delta_{1}\eta| \,. \end{split}$$

As previously mentioned, the interaction potential Q in (3.16) lacks certain shock waves: precisely, 3-shocks interacting with η_0 and 1-shocks interacting with ζ_0 , see Figure 7. In the next two propositions we list the conditions on the parameters ξ , $K_{\eta,\zeta}^{\ell,m,r}$ and ρ needed for the decrease of F.



Fig.7 The waves considered in Q for the bubble case

Proposition 4.1 Assume that at time t > 0 a wave δ_i , i = 1, 3, interacts with one of the composite waves η_0 or ζ_0 . Then, $\Delta F(t) \leq 0$ provided that

$$\xi \ge 1, \qquad K^m_{\zeta}, K^m_{\eta} \ge 1, \qquad \frac{\xi - 1}{2} \le K^r_{\zeta}, K^\ell_{\eta}, \qquad K^m_{\eta} \le K^r_{\eta}, \qquad K^m_{\zeta} \le K^\ell_{\zeta}, \tag{4.1}$$

$$1 + K_{\eta}^{m} \frac{|\eta|}{2} - K_{\zeta}^{m} \le 0, \qquad 1 + K_{\zeta}^{m} \frac{|\zeta|}{2} - K_{\eta}^{m} \le 0, \tag{4.2}$$

$$C_o(\rho) \le \frac{2\xi}{\xi+1} \min\{K_{\zeta}^m, K_{\eta}^m\}.$$
 (4.3)



Fig.8 Interactions of 1- and 3-waves with ζ_0 solved by means of the Accurate solver. The fronts carrying the composite waves are represented by a single line

Proof Since the two cases give symmetric conditions, we only analyze interactions involving ζ_0 ; see Figure 8. By (3.8) and (3.9) we have

$$\begin{cases} \varepsilon_3 - \varepsilon_1 = -\delta_1, & |\varepsilon_1| - |\delta_1| = -|\varepsilon_3|, & \text{if } i = 1, \\ \varepsilon_3 - \varepsilon_1 = \delta_3, & |\varepsilon_3| - |\delta_3| = |\varepsilon_1|, & \text{if } i = 3. \end{cases}$$

i = 1. If the interacting wave is a rarefaction, then by (3.10) we have

$$\Delta L + \Delta L^{0} = \begin{cases} \xi |\varepsilon_{3}| + |\varepsilon_{1}| - |\delta_{1}| = (\xi - 1)|\varepsilon_{3}| \leq \frac{\xi - 1}{2}|\delta_{1}\zeta| & \text{if } |\delta_{1}| \geq \rho, \\ |\varepsilon_{3}| + |\varepsilon_{1}| - |\delta_{1}| = 0 & \text{if } |\delta_{1}| < \rho, \end{cases}$$

and $\Delta Q = K_{\eta}^{m} |\varepsilon_{1}\eta| - K_{\eta}^{r} |\delta_{1}\eta| - K_{\zeta}^{r} |\delta_{1}\zeta|$. Therefore, since $|\varepsilon_{1}| \leq |\delta_{1}|$ by the interaction estimates,

$$\Delta F \leq \begin{cases} \left(K_{\eta}^{m} - K_{\eta}^{r}\right)|\delta_{1}\eta| + \left[\frac{\xi - 1}{2} - K_{\zeta}^{r}\right]|\delta_{1}\zeta| & \text{if } |\delta_{1}| \ge \rho, \\ \left(K_{\eta}^{m} - K_{\eta}^{r}\right)|\delta_{1}\eta| - K_{\zeta}^{r}|\delta_{1}\zeta| & \text{if } |\delta_{1}| < \rho, \end{cases}$$

which is nonpositive by $(4.1)_{3,4}$. Instead, if the interacting wave is a shock, then in both the accurate and simplified cases we have $\Delta L + \Delta L^0 = |\varepsilon_3| + \xi |\varepsilon_1| - \xi |\delta_1| = -(\xi - 1)|\varepsilon_3|$ and $\Delta Q = \xi (K_\eta^m |\varepsilon_1| - K_\eta^r |\delta_1|) |\eta|$. As a consequence, $\Delta F \leq -(\xi - 1)|\varepsilon_3| + \xi (K_\eta^m - K_\eta^r) |\delta_1\eta|$, which is ≤ 0 by $(4.1)_{1,4}$.

i = 3. If the interacting wave is a rarefaction, then by the interaction estimates (3.10) $\Delta L + \Delta L^0 = |\varepsilon_3| + |\varepsilon_1| - |\delta_3| = 2|\varepsilon_1| \le |\delta_3\zeta|$ and

$$\Delta Q = \begin{cases} K_{\eta}^{m} |\varepsilon_{1}\eta| - K_{\zeta}^{m} |\delta_{3}\zeta| & \text{if } |\delta_{3}| \ge \rho, \\ -K_{\zeta}^{m} |\delta_{3}\zeta| & \text{if } |\delta_{3}| < \rho. \end{cases}$$

Then,

$$\Delta F \leq \begin{cases} \left[1 + K_{\eta}^{m} \frac{|\eta|}{2} - K_{\zeta}^{m}\right] |\delta_{3}\zeta| & \text{if } |\delta_{3}| \geq \rho, \\ \left[1 - K_{\zeta}^{m}\right] |\delta_{3}\zeta| & \text{if } |\delta_{3}| < \rho, \end{cases}$$

which is nonpositive by $(4.1)_2$, $(4.2)_1$. On the other hand, if the interacting wave is a shock, then

$$\Delta L + \Delta L^{0} = \begin{cases} \xi |\varepsilon_{1}| + \xi |\varepsilon_{3}| - \xi |\delta_{3}| = 2\xi |\varepsilon_{1}| \le \xi |\delta_{3}\zeta| & \text{if } |\delta_{3}| \ge \rho, \\ |\varepsilon_{1}| + \xi |\varepsilon_{3}| - \xi |\delta_{3}| = (\xi + 1)|\varepsilon_{1}| \le (\xi + 1)\frac{C_{o}}{2}|\delta_{3}\zeta| & \text{if } |\delta_{3}| < \rho, \end{cases}$$

and

$$\Delta Q = \begin{cases} \xi \left(K_{\eta}^{m} |\varepsilon_{1}\eta| - K_{\zeta}^{m} |\delta_{3}\zeta| \right) & \text{if } |\delta_{3}| \ge \rho, \\ -\xi K_{\zeta}^{m} |\delta_{3}\zeta| & \text{if } |\delta_{3}| < \rho. \end{cases}$$

Therefore,

$$\Delta F \leq \begin{cases} \xi \left[1 + K_{\eta}^{m} \frac{|\eta|}{2} - K_{\zeta}^{m} \right] |\delta_{3}\zeta| & \text{if } |\delta_{3}| \geq \rho, \\ \left[(\xi + 1) \frac{C_{o}}{2} - \xi K_{\zeta}^{m} \right] |\delta_{3}\zeta| & \text{if } |\delta_{3}| < \rho, \end{cases}$$

which is nonpositive by $(4.2)_1$, $(4.3)_1$.

Proposition 4.2 Consider the interaction at time t > 0 of two waves of the same family 1 or 3 and assume (3.18). Then, $\Delta F(t) \leq 0$ provided that

$$1 \le \xi \le \frac{1}{c(m_o)}, \qquad K_{\zeta}^m \le \frac{\xi - 1}{|\zeta|}, \qquad K_{\eta}^m \le \frac{\xi - 1}{|\eta|},$$
(4.4)

$$K_{\eta}^{r}|\eta| + K_{\zeta}^{r}|\zeta| \le \xi - 1, \qquad K_{\eta}^{\ell}|\eta| + K_{\zeta}^{\ell}|\zeta| \le \xi - 1.$$
 (4.5)

Proof First, we consider the interactions taking place in \mathcal{M} , see Figure 9. For brevity, we only deal with the case of interactions between two 3-waves α_3 and β_3 giving rise to ε_1 and ε_3 (the 1-waves case is analogous).



Fig.9 Interactions of 3-waves in \mathcal{M}

If both α_3 and β_3 are shocks, then ε_1 is a rarefaction by Lemma 3.3 and, as in [1, Proposition 5.8], we have

$$\Delta L + |\varepsilon_1|(\xi - 1) = 0, \qquad (4.6)$$

for any $\xi \geq 1$. Moreover, we have

$$\Delta Q = K_{\eta}^{m} |\varepsilon_{1}\eta| - \xi K_{\zeta}^{m} |\varepsilon_{1}\zeta| \le K_{\eta}^{m} |\varepsilon_{1}\eta|, \qquad \Delta F \le \left[-(\xi - 1) + K_{\eta}^{m} |\eta|\right] |\varepsilon_{1}|$$

and F is non-increasing by $(4.4)_{1,3}$. On the other hand, when the two interacting waves are of different type, for example $\alpha_3 < 0 < \beta_3$, as in [1, Proposition 5.8], by $(4.4)_1$ one can deduce that

$$\Delta L + \xi(\xi - 1)|\varepsilon_1| \le 0.$$
(4.7)

If ε_3 is a rarefaction, then $\Delta Q = \xi K_{\eta}^m |\varepsilon_1 \eta| + K_{\zeta}^m (|\varepsilon_3| - \xi |\alpha_3| - |\beta_3|) |\zeta|$; on the other hand, if ε_3 is a shock, $\Delta Q = \xi K_{\eta}^m |\varepsilon_1 \eta| + K_{\zeta}^m (\xi |\varepsilon_3| - \xi |\alpha_3| - |\beta_3|) |\zeta|$. Therefore, by Lemma 3.3 in both cases it holds

$$\Delta Q \le \xi K_{\eta}^{m} |\varepsilon_{1}\eta|, \qquad \Delta F \le \xi \left[-(\xi - 1) + K_{\eta}^{m} |\eta|\right] |\varepsilon_{1}|$$

and F decreases by $(4.4)_{1,3}$. The analysis of the interactions between 1-waves requires symmetrically the condition $K_{\zeta}^m \leq (\xi - 1)/|\zeta|$.

Next, we analyze the case of interactions taking place in \mathcal{R} , the case of interactions in \mathcal{L} being analogous. By $(4.4)_1$ it is easy to verify that F decreases when two 1-waves interact.



Fig.10 Interactions of 3-waves in \mathcal{R}

Now, we consider the interactions between 3-waves (see Figure 10). When the interacting waves α_3, β_3 are both shocks we have (4.6), while in the other two cases estimate (4.7) still holds under condition (4.4)₁. If $\alpha_3, \beta_3 < 0$ we have

$$\Delta Q = K_{\eta}^{r} |\varepsilon_{1}\eta| + K_{\zeta}^{r} |\varepsilon_{1}\zeta|, \qquad \Delta F = \left[-(\xi - 1) + K_{\eta}^{r} |\eta| + K_{\zeta}^{r} |\zeta|\right] |\varepsilon_{1}|,$$

while if, for example, $\alpha_3 < 0 < \beta_3$ we have

$$\Delta Q = K_{\eta}^{r} \xi |\varepsilon_{1}\eta|, \qquad \Delta F \leq \xi \left[-(\xi - 1) + K_{\eta}^{r}|\eta|\right] |\varepsilon_{1}|.$$

Consequently, F is non-increasing by $(4.5)_1$. The condition $K_{\eta}^{\ell}|\eta| + K_{\zeta}^{\ell}|\zeta| \leq \xi - 1$ is required for the interactions occurring in the region \mathcal{L} .

Now, we can determine the order of choice of the parameters. To simplify the analysis, we can let $K_{\eta}^{m} = K_{\eta}^{r}$ and $K_{\zeta}^{m} = K_{\zeta}^{\ell}$, since the final result does not change otherwise. Once η, ζ have been fixed, we choose in turn: m_{o}, ξ, K_{η}^{m} and $K_{\zeta}^{m}, K_{\zeta}^{r}$ and K_{η}^{ℓ} ; finally, we choose ρ (i.e. C_{o}). First, notice that the conditions in (4.2) identify the set in the $(K_{\eta}^{m}, K_{\zeta}^{m})$ -plane represented in Figure 11. Hence, by (4.2) we deduce

$$K_{\eta}^{m} \ge 1 + K_{\zeta}^{m} \frac{|\zeta|}{2} \ge 1 + \frac{|\zeta|}{2} \left(1 + K_{\eta}^{m} \frac{|\eta|}{2}\right) \quad \text{and} \quad K_{\zeta}^{m} \ge 1 + K_{\eta}^{m} \frac{|\eta|}{2} \ge 1 + \frac{|\eta|}{2} \left(1 + K_{\zeta}^{m} \frac{|\zeta|}{2}\right),$$

that imply

$$K_{\eta}^{m} \ge \frac{1+|\zeta|/2}{1-|\eta\zeta|/4} \quad \text{and} \quad K_{\zeta}^{m} \ge \frac{1+|\eta|/2}{1-|\eta\zeta|/4}.$$
 (4.8)

In particular, by replacing the inequality sign by equality in (4.8) we get the coordinates of the intersection point V between the two lines of Figure 11. Notice also that (4.8) implies $(4.1)_2$.

Since we have chosen $K_{\eta}^{m} = K_{\eta}^{r}$ and $K_{\zeta}^{m} = K_{\zeta}^{\ell}$, conditions (4.5) imply (4.4)_{2,3}. By (4.1)₃ and (4.5)₁, we get $K_{\eta}^{m} |\eta| + (\xi - 1)|\zeta|/2 \leq \xi - 1$, which is equivalent to

$$K_{\eta}^{m} \frac{|\eta|}{1 - |\zeta|/2} \le \xi - 1;$$
(4.9)



Fig.11 Graphical representation of conditions (4.2) for $|\eta| = 1/2$ and $|\zeta| = 3/2$

similarly, by $(4.1)_3$ and $(4.5)_2$ we get

$$K_{\zeta}^{m} \frac{|\zeta|}{1 - |\eta|/2} \le \xi - 1.$$
(4.10)

By (4.8), (4.9) and (4.10) it follows that ξ must satisfy the inequality

$$\xi \ge 1 + \max\left\{\frac{1+|\zeta|/2}{1-|\zeta|/2} \, \frac{|\eta|}{1-|\eta\zeta|/4}, \, \frac{1+|\eta|/2}{1-|\eta|/2} \, \frac{|\zeta|}{1-|\eta\zeta|/4}\right\}$$

This condition must match with $(4.4)_1$; then, recalling (2.3) we must require

$$1 + \mathcal{H}_b(|\eta|, |\zeta|) \le \xi \le \frac{1}{c(m_o)}, \qquad (4.11)$$

which is a condition that relates m_o to $|\eta|, |\zeta|$. When one of the phase waves tends to zero, $\mathcal{H}_b(|\eta|, |\zeta|)$ tends to the other one and we completely recover the results of [1].

Summarizing, we choose the parameters as follows and keep strict inequalities for later need; let $(|\eta|, |\zeta|) \in D_b$ be given.

• First, we fix m_o such that

$$1 + \mathcal{H}_b(|\eta|, |\zeta|) < \frac{1}{c(m_o)} \tag{4.12}$$

and take ξ in the interior of the interval given by (4.11).

• In the $(K_{\eta}^m, K_{\zeta}^m)$ -plane we choose a point in the affine cone defined by (4.2) and sufficiently close to V; moreover, we require

$$K_{\eta}^{m} \frac{|\eta|}{1 - |\zeta|/2} < \xi - 1 \quad \text{and} \quad K_{\zeta}^{m} \frac{|\zeta|}{1 - |\eta|/2} < \xi - 1.$$
 (4.13)

• We choose $K^r_{\eta} = K^m_{\eta}, K^{\ell}_{\zeta} = K^m_{\zeta}$ and, then, by (4.13) we choose K^r_{ζ} and K^{ℓ}_{η} such that

$$\frac{\xi - 1}{2} < K_{\zeta}^{r} < \frac{\xi - 1}{|\zeta|} - K_{\eta}^{m} \frac{|\eta|}{|\zeta|} \quad \text{and} \quad \frac{\xi - 1}{2} < K_{\eta}^{\ell} < \frac{\xi - 1}{|\eta|} - K_{\zeta}^{m} \frac{|\zeta|}{|\eta|}.$$
(4.14)

• Finally, we choose ρ such that (4.3) holds.

Now, we can prove the global in time decreasing of the functional F.

Proposition 4.3 Let $m_o > 0$ satisfy (4.12). Moreover, assume that ξ , $K_{\eta,\zeta}^{\ell,m,r}$ and ρ satisfy (4.11)–(4.14) and (4.3). Then, the following two statements hold.

i) Local Decreasing. For any interaction at time t > 0 between two waves satisfying (3.18), it holds

$$\Delta F(t) \le 0 \,.$$

ii) Global Decreasing. If

$$L(0) \le m_o c(m_o) \tag{4.15}$$

and the approximate solution is defined in [0, T], then $F(0) \le m_o$, $\Delta F(t) \le 0$ for every $t \in (0, T]$ and (3.18) is satisfied.

Proof The first statement has been proved above. As for the second assertion, let us denote by $L_{iR}^{\ell,m,r}$ and $L_{iS}^{\ell,m,r}$ the partial sums in $L^{\ell,m,r}$ due to *i*-rarefaction waves (iR) and *i*-shock waves (iS), respectively. By $(4.4)_{2,3}$ we have

$$F^{m}(0) = L^{m}(0) + Q^{m}(0) \le L^{m}(0) \left(1 + \max\{K_{\eta}^{m}|\eta|, K_{\zeta}^{m}|\zeta|\}\right) \le \xi^{2} \bar{L}^{m}(0).$$

Moreover, from (4.5) it follows

$$\begin{split} F^{\ell}(0) &\leq L_{1R}^{\ell}(0) + L_{1S}^{\ell}(0) + L_{3R}^{\ell}(0) \left(1 + K_{\eta}^{\ell} |\eta| + K_{\zeta}^{\ell} |\zeta|\right) + L_{3S}^{\ell}(0) \left(1 + K_{\zeta}^{\ell} |\zeta|\right) \leq \xi^{2} \bar{L}^{\ell}(0) ,\\ F^{r}(0) &\leq L_{3R}^{r}(0) + L_{3S}^{r}(0) + L_{1R}^{r}(0) \left(1 + K_{\eta}^{r} |\eta| + K_{\zeta}^{r} |\zeta|\right) + L_{1S}^{r}(0) \left(1 + K_{\eta}^{r} |\eta|\right) \leq \xi^{2} \bar{L}^{r}(0) . \end{split}$$

Then,

No.4

$$F(0) = F^{\ell}(0) + F^{m}(0) + F^{r}(0) \le \xi^{2} \bar{L}(0).$$

For a fixed $t \leq T$, suppose by induction that $F(\tau) \leq m_o$ and $\Delta F(\tau) \leq 0$ for every $0 < \tau < t$, interaction time. Then, the inequality $\Delta F(t) \leq 0$ implies that

$$F(t) \le F(0) \le \xi^2 \bar{L}(0) \,.$$

Hence, by (4.15) the size of a shock δ_i (i = 1, 3) at time t satisfies

$$|\delta_i| \le \frac{1}{\xi} F(t) \le \xi \bar{L}(0) \le \frac{1}{c(m_o)} \bar{L}(0) \le m_o$$

and (3.18) is verified.

5 The Increasing-Pressure Case

In this section, we prove the decreasing of the functional F in (3.17) in the increasingpressure case. As before, we first introduce the following interaction potentials

$$\begin{split} Q^{\ell} &= \left(K_{\eta}^{\ell} |\eta| + K_{\zeta}^{\ell} |\zeta| \right) \sum_{\substack{\delta_{3} > 0 \\ \delta_{3} \in \mathcal{L}}} |\delta_{3}| ,\\ Q^{m} &= K_{\eta}^{m} |\eta| \bigg(\sum_{\substack{\delta_{1} > 0 \\ \delta_{1} \in \mathcal{M}}} |\delta_{1}| + \xi \sum_{\substack{\delta_{1} < 0 \\ \delta_{1} \in \mathcal{M}}} |\delta_{1}| \bigg) + K_{\zeta}^{m} |\zeta| \sum_{\substack{\delta_{3} > 0 \\ \delta_{3} \in \mathcal{M}}} |\delta_{3}| ,\\ Q^{r} &= \left(K_{\eta}^{r} |\eta| + K_{\zeta}^{r} |\zeta| \right) \bigg(\sum_{\substack{\delta_{1} > 0 \\ \delta_{1} \in \mathcal{R}}} |\delta_{1}| + \xi \sum_{\substack{\delta_{1} < 0 \\ \delta_{1} \in \mathcal{R}}} |\delta_{1}| \bigg) . \end{split}$$

The interaction potential Q in (3.16) lacks the 3-shocks interacting with η_0 and with ζ_0 , see Figure 12. The next proposition, which is analogous to Proposition 4.1, gives a first list of conditions that guarantee the decrease of F.



Fig.12 The waves considered in Q for the increasing-pressure case

Proposition 5.1 Assume that at time t > 0 a wave δ_i , i = 1, 3, interacts with one of the composite waves η_0 or ζ_0 . Then, $\Delta F(t) \leq 0$ provided that

$$\xi \ge 1, \quad K_{\eta}^{m} \ge 1, \quad K_{\eta}^{m} \le \frac{\xi - 1}{|\eta|},$$
(5.1)

$$\left(\frac{\xi - 1}{2} - K_{\eta}^{\ell}\right)|\eta| + \left(K_{\zeta}^{m} - K_{\zeta}^{\ell}\right)|\zeta| \le 0, \quad (1 - K_{\zeta}^{r})|\zeta| + \left(K_{\eta}^{m}\left(1 + \frac{|\zeta|}{2}\right) - K_{\eta}^{r}\right)|\eta| \le 0, \quad (5.2)$$

$$\frac{\xi - 1}{2} + K_{\eta}^{m} \xi \frac{|\eta|}{2} - K_{\zeta}^{m} \le 0, \quad 1 + K_{\zeta}^{m} \frac{|\zeta|}{2} - K_{\eta}^{m} \le 0, \quad (5.3)$$

$$\left((\xi+1)\frac{C_o}{2} - \xi K_{\zeta}^r \right) |\zeta| + \xi \left(K_{\eta}^m \left(1 + \frac{C_o}{2} |\zeta| \right) - K_{\eta}^r \right) |\eta| \le 0, \quad (\xi+1)\frac{C_o}{2} - \xi K_{\eta}^m \le 0.$$
(5.4)

Proof Both in the case of interaction with η_0 and ζ_0 , by (3.8), (3.9) we have

$$\begin{cases} \varepsilon_3 - \varepsilon_1 = -\delta_1, & |\varepsilon_1| - |\delta_1| = |\varepsilon_3|, & \text{if } i = 1, \\ \varepsilon_3 - \varepsilon_1 = \delta_3, & |\varepsilon_3| - |\delta_3| = -|\varepsilon_1|, & \text{if } i = 3. \end{cases}$$

However, in this case we have to treat separately the interactions with the phase waves, since the interaction potential Q is not symmetric with respect to η_0 and ζ_0 .

Interactions with η_0 . Assume i = 1. If δ_1 is a rarefaction, then $\Delta L + \Delta L^0 = 2|\varepsilon_3| \le |\delta_1 \eta|$ and

$$\Delta Q = \begin{cases} K_{\zeta}^{m} |\varepsilon_{3}\zeta| - K_{\eta}^{m} |\delta_{1}\eta| & \text{if } |\delta_{1}| \ge \rho, \\ -K_{\eta}^{m} |\delta_{1}\eta| & \text{if } |\delta_{1}| < \rho. \end{cases}$$

Hence,

$$\Delta F \leq \begin{cases} \left[1 + K_{\zeta}^{m} \frac{|\zeta|}{2} - K_{\eta}^{m} \right] |\delta_{1}\eta| & \text{if } |\delta_{1}| \geq \rho, \\ \left[1 - K_{\eta}^{m} \right] |\delta_{1}\eta| & \text{if } |\delta_{1}| < \rho. \end{cases}$$

Then, $\Delta F \leq 0$ by $(5.3)_2$ and $(5.1)_2$. On the other hand, if δ_1 is a shock then

$$\Delta L + \Delta L^{0} = \begin{cases} 2\xi |\varepsilon_{3}| \leq \xi |\delta_{1}\eta| & \text{if } |\delta_{1}| \geq \rho, \\ (\xi + 1)|\varepsilon_{3}| \leq \frac{C_{o}}{2}(\xi + 1)|\delta_{1}\eta| & \text{if } |\delta_{1}| < \rho \end{cases}$$

and $\Delta Q = -\xi K_{\eta}^{m} |\delta_{1}\eta|$. Hence,

$$\Delta F \leq \begin{cases} \xi \left[1 - K_{\eta}^{m} \right] |\delta_{1}\eta| & \text{if } |\delta_{1}| \geq \rho, \\ \left[(\xi + 1) \frac{C_{o}}{2} - \xi K_{\eta}^{m} \right] |\delta_{1}\eta|, & \text{if } |\delta_{1}| < \rho, \end{cases}$$

which is nonpositive by $(5.1)_2$ and $(5.4)_2$.

Now, let i = 3. If δ_3 is a rarefaction, then

$$\Delta L + \Delta L^{0} = \begin{cases} (\xi - 1)|\varepsilon_{1}| \leq \frac{\xi - 1}{2}|\delta_{3}\eta| & \text{if } |\delta_{3}| \geq \rho, \\ 0 & \text{if } |\delta_{3}| < \rho, \end{cases}$$

and $\Delta Q = K_{\zeta}^{m} |\varepsilon_{3}\zeta| - K_{\eta}^{\ell} |\delta_{3}\eta| - K_{\zeta}^{\ell} |\delta_{3}\zeta|$. Then, since $|\varepsilon_{3}| \leq |\delta_{3}|$ we have

$$\Delta F \leq \begin{cases} \left[\left(\frac{\xi - 1}{2} - K_{\eta}^{\ell}\right) |\eta| + (K_{\zeta}^{m} - K_{\zeta}^{\ell}) |\zeta| \right] |\delta_{3}| & \text{if } |\delta_{3}| \geq \rho, \\ (K_{\zeta}^{m} - K_{\zeta}^{\ell}) |\delta_{3}\zeta| - K_{\eta}^{\ell} |\delta_{3}\eta| & \text{if } |\delta_{3}| < \rho, \end{cases}$$

and F decreases by (5.2)₁. If δ_3 is a shock, then in any case we get $\Delta F = -(\xi - 1)|\varepsilon_1| \leq 0$ by (5.1)₁.

Interactions with ζ_0 . Assume i = 1. If δ_1 is a rarefaction, then $\Delta L + \Delta L^0 = 2|\varepsilon_3| \leq |\delta_1\zeta|$ and $\Delta Q = K_{\eta}^m |\varepsilon_1\eta| - K_{\eta}^r |\delta_1\eta| - K_{\zeta}^r |\delta_1\zeta|$ in both the Accurate and the Simplified case. Hence, by $(5.2)_2$ we have

$$\Delta F \leq \left[(1 - K_{\zeta}^r) |\zeta| + \left(K_{\eta}^m \left(1 + \frac{|\zeta|}{2} \right) - K_{\eta}^r \right) |\eta| \right] |\delta_1| \leq 0.$$

On the other hand, if δ_1 is a shock, then

$$\Delta L + \Delta L^{0} = \begin{cases} 2\xi |\varepsilon_{3}| \leq \xi |\delta_{1}\zeta| & \text{if } |\delta_{1}| \geq \rho, \\ \\ (\xi + 1)|\varepsilon_{3}| \leq \frac{C_{o}}{2}(\xi + 1)|\delta_{1}\zeta| & \text{if } |\delta_{1}| < \rho, \end{cases}$$

and $\Delta Q = \xi K_{\eta}^{m} |\varepsilon_{1}\eta| - \xi K_{\eta}^{r} |\delta_{1}\eta| - \xi K_{\zeta}^{r} |\delta_{1}\zeta|$. Thus,

$$\Delta F \leq \begin{cases} \xi \left[(1 - K_{\zeta}^{r})|\zeta| + \left(K_{\eta}^{m} \left(1 + \frac{|\zeta|}{2} \right) - K_{\eta}^{r} \right) |\eta| \right] |\delta_{1}| & \text{if } |\delta_{1}| \geq \rho, \\ \left[\left((\xi + 1) \frac{C_{o}}{2} - \xi K_{\zeta}^{r} \right) |\zeta| + \xi \left(K_{\eta}^{m} \left(1 + \frac{C_{o}}{2} |\zeta| \right) - K_{\eta}^{r} \right) |\eta| \right] |\delta_{1}| & \text{if } |\delta_{1}| < \rho, \end{cases}$$

which is nonpositive by $(5.2)_2$ and $(5.4)_1$.

Assume i = 3. If δ_3 is a rarefaction, then

$$\Delta L + \Delta L^{0} = \begin{cases} (\xi - 1)|\varepsilon_{1}| \leq \frac{\xi - 1}{2}|\delta_{3}\zeta| & \text{if } |\delta_{3}| \geq \rho_{4}\\ 0 & \text{if } |\delta_{3}| < \rho_{4} \end{cases}$$

and

$$\Delta Q = \begin{cases} \xi K_{\eta}^{m} |\varepsilon_{1}\eta| - K_{\zeta}^{m} |\delta_{3}\zeta| & \text{if } |\delta_{3}| \ge \rho, \\ \\ -K_{\zeta}^{m} |\delta_{3}\zeta| & \text{if } |\delta_{3}| < \rho. \end{cases}$$

Hence,

18

$$\Delta F \leq \begin{cases} \left[\frac{\xi - 1}{2} + K_{\eta}^{m} \xi \frac{|\eta|}{2} - K_{\zeta}^{m}\right] |\delta_{3}\zeta| & \text{if } |\delta_{3}| \geq \rho, \\ -K_{\zeta}^{m} |\delta_{3}\zeta| & \text{if } |\delta_{3}| < \rho, \end{cases}$$

which is nonpositive by $(5.3)_1$. If, instead, δ_3 is a shock, then $\Delta L + \Delta L^0 = -(\xi - 1)|\varepsilon_1|$ and

$$\Delta Q = \begin{cases} K_{\eta}^{m} |\varepsilon_{1}\eta| & \text{if } |\delta_{3}| \ge \rho, \\ 0 & \text{if } |\delta_{3}| < \rho. \end{cases}$$

Then,

$$\Delta F \leq \begin{cases} -(\xi - 1)|\varepsilon_1| + K_{\eta}^m|\varepsilon_1\eta| & \text{if } |\delta_3| \ge \rho, \\ \\ -(\xi - 1)|\varepsilon_1| & \text{if } |\delta_3| < \rho, \end{cases}$$

which is nonpositive by $(5.1)_{1,3}$.

As for an interaction between two waves of the same family, Proposition 4.2 still holds with the current functional F. Therefore, the conditions required on the various parameters are (4.4), (4.5). We omit the proof, since it can be carried out as above.

Here, we make some comments on the conditions (5.1)–(5.4) and (4.4), (4.5); finally, we establish the order in which we can choose the parameters.

First, notice that $(5.1)_2$ is implied by $(5.3)_2$. Secondly, we can rewrite (5.2) as

$$\frac{\xi - 1}{2} |\eta| + K_{\zeta}^{m} |\zeta| \le K_{\eta}^{\ell} |\eta| + K_{\zeta}^{\ell} |\zeta|, \quad K_{\eta}^{m} \left(1 + \frac{|\zeta|}{2} \right) |\eta| + |\zeta| \le K_{\eta}^{r} |\eta| + K_{\zeta}^{r} |\zeta|.$$
(5.5)

Putting together $(5.5)_2$ with $(4.5)_2$ and $(5.5)_1$ with $(4.5)_1$, we have

$$K_{\eta}^{m} \left(1 + \frac{|\zeta|}{2}\right) |\eta| + |\zeta| \le K_{\eta}^{r} |\eta| + K_{\zeta}^{r} |\zeta| \le \xi - 1, \qquad (5.6)$$

$$\frac{\xi - 1}{2} |\eta| + K_{\zeta}^{m} |\zeta| \le K_{\eta}^{\ell} |\eta| + K_{\zeta}^{\ell} |\zeta| \le \xi - 1, \qquad (5.7)$$

then $(4.4)_{2,3}$ are implied by (5.2) and (4.5). Moreover, by (5.3) we have

$$K_{\eta}^{m} \ge 1 + K_{\zeta}^{m} \frac{|\zeta|}{2}, \qquad K_{\zeta}^{m} \ge \frac{\xi - 1}{2} + \xi K_{\eta}^{m} \frac{|\eta|}{2},$$
(5.8)

that give the following lower bounds on K_{η}^{m} and K_{ζ}^{m} :

$$K_{\eta}^{m} \ge \frac{1 + (\xi - 1)|\zeta|/4}{1 - \xi|\eta\zeta|/4}, \qquad K_{\zeta}^{m} \ge \frac{(\xi - 1) + \xi|\eta|}{2(1 - \xi|\eta\zeta|/4)}.$$
(5.9)

Remark that (5.8) represents an affine cone in the $(K_{\eta}^{m}, K_{\zeta}^{m})$ -plane under the condition

$$\xi \le \frac{4}{|\eta\zeta|} \,. \tag{5.10}$$

The vertex is the point whose coordinates are given by (5.9). Hence, K_{η}^{m} and K_{ζ}^{m} must be chosen in the non-empty intervals identified by (5.6), (5.9)₁ and (5.7), (5.9)₂, respectively. This means that (5.7), (5.9)₂ give the condition

$$\frac{\xi - 1}{2} |\eta| + \frac{(\xi - 1) + \xi |\eta|}{2(1 - \xi |\eta\zeta|/4)} |\zeta| \le \xi - 1,$$
(5.11)

while (5.6), $(5.9)_1$ give the condition

$$\frac{1 + (\xi - 1)|\zeta|/4}{1 - \xi|\eta\zeta|/4} \left(1 + \frac{|\zeta|}{2}\right)|\eta| + |\zeta| \le \xi - 1.$$
(5.12)

We introduce the notation $|\eta| = x, |\zeta| = y$ and $\xi - 1 = z$. Then, by (5.10) we rewrite (5.11) and (5.12) as, respectively,

$$\frac{xy}{4}(2-x)z^2 + \left[y(x+1) - (2-x)\left(1 - \frac{xy}{4}\right)\right]z + xy \le 0,$$
(5.13)

$$\frac{xy}{4}z^2 + \left[\frac{xy}{8}(4-y) - 1\right]z + \left(1 + \frac{y}{2}\right)x + y\left(1 - \frac{xy}{4}\right) \le 0.$$
(5.14)

We also denote a(x, y) = xy(2 - x)/4, b(x, y) = y(x + 1) - (2 - x)(1 - xy/4), c(x, y) = xy, d(x, y) = xy/4, e(x, y) = xy(4 - y)/8 - 1 and f(x, y) = (1 + y/2)x + y(1 - xy/4) so that (5.13) and (5.14) become, respectively,

$$a(x,y)z^{2} + b(x,y)z + c(x,y) \le 0, \qquad (5.15)$$

$$d(x,y)z^{2} + e(x,y)z + f(x,y) \le 0.$$
(5.16)

Notice that the coefficients a, c, d, f are positive, e is negative and b may change sign. In order that each equations associated to (5.15) and (5.16) have distinct solutions, the discriminants $b^2 - 4ac$ and $e^2 - 4df$ must be strictly positive. If b < 0, such solutions are positive. Thus, about (5.13) we require

$$y(x+1) - (2-x)\left(1 - \frac{xy}{4}\right) + xy\sqrt{2-x} < 0, \qquad (5.17)$$

while about (5.14) we impose

$$\left[\frac{xy}{8}(4-y)-1\right]^2 - xy\left[\left(1+\frac{y}{2}\right)x+y\left(1-\frac{xy}{4}\right)\right] > 0.$$
(5.18)

By a numerical comparison, we see that the set defined by (5.17) is included in that defined by (5.18). Under (5.17) and (5.18), we denote by

$$z_{1,2}(x,y) = \frac{(2-x)(1-xy/4) - y(x+1) \pm \sqrt{[y(x+1) - (2-x)(1-xy/4)]^2 - x^2y^2(2-x)}}{xy(2-x)/2}$$
(5.19)

the two positive solutions of the equation associated to (5.15) and by

$$z_{3,4}(x,y) = \frac{1 - xy(4-y)/8 \pm \sqrt{[xy(4-y)/8 - 1]^2 - xy[(1+y/2)x + y(1-xy/4)]}}{xy/2}$$
(5.20)

the solutions of the equation associated to (5.16). Hence, by (5.10), (5.13) and (5.14) we get

$$1 + \max\left\{z_1(x,y), z_3(x,y)\right\} < \xi < 1 + \min\left\{z_2(x,y), z_4(x,y), \frac{4}{xy} - 1\right\}.$$
 (5.21)

Therefore, we can define the domain D_c represented in Figure 2 as

 $D_c = \{ (|\eta|, |\zeta|) = (x, y) : (5.21) \text{ holds} \}$ (5.22)

and the function

$$\mathcal{H}_{c}(|\eta|, |\zeta|) = \max\left\{z_{1}(|\eta|, |\zeta|), z_{3}(|\eta|, |\zeta|)\right\}.$$
(5.23)

By $(4.4)_1$ we find the condition that relates m_o to $|\eta|, |\zeta|$, i.e.,

$$1 + \mathcal{H}_c(|\eta|, |\zeta|) < \frac{1}{c(m_o)}$$
 (5.24)

As a final remark, we notice that $(5.4)_1$ is equivalent to

$$\left(\frac{\xi+1}{2\xi}C_o - K_{\zeta}^r\right)|\zeta| + \left(K_{\eta}^m\left(1 + \frac{C_o}{2}|\zeta|\right) - K_{\eta}^r\right)|\eta| \le 0.$$
(5.25)

Then, by taking ρ sufficiently small (since $C_o(\rho) \to 1$ if $\rho \to 0^+$) and $\xi > 1$, (5.25) is implied by $(5.2)_2$.

For the choice of the parameters we proceed as follows.

• We fix $|\eta|, |\zeta|$ such that

$$1 + \max\left\{z_1(|\eta|, |\zeta|), z_3(|\eta|, |\zeta|)\right\} < 1 + \min\left\{z_2(|\eta|, |\zeta|), z_4(|\eta|, |\zeta|), \frac{4}{|\eta\zeta|} - 1\right\}.$$
 (5.26)

Then, we fix m_o such that (5.24) hold and, in turn, we choose ξ satisfying both (5.21) and

$$1 + \mathcal{H}_c(|\eta|, |\zeta|) < \xi < \frac{1}{c(m_o)},$$
(5.27)

so that $(4.4)_1$ holds.

• We choose $K_{\eta}^{m}, K_{\zeta}^{m}$ such that (5.3) holds; in particular, we take $(K_{\eta}^{m}, K_{\zeta}^{m})$ sufficiently close to the vertex of the cone and satisfying (5.6)–(5.9), i.e., such that

$$\frac{1+(\xi-1)|\zeta|/4}{1-\xi|\eta\zeta|/4} \le K_{\eta}^{m} < \frac{\xi-1-|\zeta|}{(1+|\zeta|/2)|\eta|},$$

$$\frac{(\xi-1)+\xi|\eta|}{2(1-\xi|\eta\zeta|/4)} \le K_{\zeta}^{m} < \frac{\xi-1-(\xi-1)|\eta|/2}{|\zeta|}.$$
(5.28)

Then, we choose $K^\ell_\eta = K^\ell_\zeta,\, K^r_\eta = K^r_\zeta$ such that

$$\frac{(\xi-1)|\eta|/2 + K_{\zeta}^{m}|\zeta|}{|\eta| + |\zeta|} \le K_{\eta}^{\ell} = K_{\zeta}^{\ell} < \frac{\xi - 1}{|\eta| + |\zeta|},$$

$$\frac{K_{m}^{\eta}(1 + |\zeta|/2)|\eta| + |\zeta|}{|\eta| + |\zeta|} \le K_{\eta}^{r} = K_{\zeta}^{r} < \frac{\xi - 1}{|\eta| + |\zeta|}.$$
(5.29)

Thus, (4.5) and (5.2) hold; hence, also $(4.4)_{2,3}$ are verified.

• Finally, we choose ρ such that $C_o(\rho)$ verifies (5.4).

In the next proposition, we claim the global in time decreasing of the functional F. The proof is omitted since it is analogous to that of Proposition 4.3.

Proposition 5.2 Let $m_o > 0$ satisfy (5.24). Moreover, assume that ξ , $K_{\eta,\zeta}^{\ell,m,r}$ and ρ satisfy (5.21)–(5.29) and (5.4). Then, the following two statements hold.

• Local Decreasing. For any interaction at time t > 0 between two waves satisfying (3.18), it holds

$$\Delta F(t) \le 0$$
.

• Global Decreasing. If

$$\bar{L}(0) \le m_o c(m_o) \tag{5.30}$$

and the approximate solution is defined in [0, T], then $F(0) \le m_o$, $\Delta F(t) \le 0$ for every $t \in (0, T]$ and (3.18) is satisfied.

6 End of the Proof of Theorem 2.1

In this last section, we conclude the proof of Theorem 2.1 and add some final comments. The proof fits in the general framework detailed in [2]; hence, we only outline the most important changes.

First, as in [2, Section 6] the front tracking algorithm used to construct the approximate solutions is well-defined and converges. Moreover, it is consistent in the sense that the total size of the non-physical waves carried by the composite waves vanishes with ν . In brief, to estimate this quantity we use the notion of generation order, i.e., we attach an index $k \geq 1$ to each wave generated in the construction. Then, according to each k, we introduce suitable functionals L_k, Q_k, F_k simply by referring the functionals L, Q, F to waves with order k. In particular, following the same steps as in [1, 2], in both cases we can prove that

$$\tilde{F}_k(t) = \sum_{j \ge k} F_j(t) \le \mu^{k-1} F_1(0), \tag{6.1}$$

where $\mu \in [0, 1]$ is either μ_b or μ_c . More precisely, in the bubble case we find

$$\mu_{b} = \max\left\{\frac{1+K_{\zeta}^{m}|\zeta|}{2K_{\eta}^{m}-1}, \frac{1+K_{\eta}^{m}|\eta|}{2K_{\zeta}^{m}-1}, \frac{\xi}{1+2K_{\eta}^{\ell}}, \frac{\xi}{1+2K_{\zeta}^{r}}, \frac{1+K_{\eta}^{m}|\eta|}{\xi}, \frac{1+K_{\zeta}^{m}|\zeta|}{\xi}, \frac{1+K_{\zeta}^{m}|\zeta|}{\xi}, \frac{1+K_{\eta}^{r}|\eta|+K_{\zeta}^{r}|\zeta|}{\xi}, \frac{C_{o}}{\xi(2K_{\eta}^{m}-C_{o})}, \frac{C_{o}}{\xi(2K_{\zeta}^{m}-C_{o})}\right\};$$

while in the increasing-pressure case we have

$$\mu_{c} = \max\left\{\frac{1+K_{\zeta}^{m}|\zeta|}{2K_{\eta}^{m}-1}, \frac{\xi(1+K_{\eta}^{m}|\eta|)}{1+2K_{\zeta}^{m}}, \frac{1+K_{\eta}^{m}|\eta|}{\xi}, \frac{1+K_{\eta}^{\ell}(|\eta|+|\zeta|)}{\xi}, \frac{1+K_{\eta}^{r}(|\eta|+|\zeta|)}{\xi} \right\}$$

$$\frac{\xi|\eta|/2}{(K_{\eta}^{\ell}-1/2)|\eta|+(K_{\zeta}^{\ell}-K_{\zeta}^{m})|\zeta|}, \frac{|\zeta|/2}{(K_{\zeta}^{r}-1/2)|\zeta|+[K_{\eta}^{r}-K_{\eta}^{m}(1+|\zeta|/2)]|\eta|}, \frac{C_{o}|\zeta|/2}{\xi(K_{\zeta}^{r}-C_{o}/2)|\zeta|+\xi[K_{\eta}^{r}-K_{\eta}^{m}(1+C_{o}|\zeta|/2)]|\eta|}, \frac{C_{o}}{\xi(2K_{\eta}^{m}-C_{o})}\right\}.$$

In both cases, simple calculations show that $\mu < 1$ because of our choice of keeping strict inequalities in the final parts of Sections 4 and 5. We exploit formula (6.1) to show that the total size of the composite waves tends to zero, as follows. We have

[total size of composite waves]

 $\leq [\text{size of composite waves of order} \geq k] + [\text{size of composite waves of order} < k] \\ \leq \mu^{k-1} \cdot F_1(0) + \frac{\rho}{2} C_o(\rho)(|\eta| + |\zeta|) \text{ [number of fronts of order} < k] \\ \leq \mu^{k-1} \cdot m_o + \frac{\rho}{2} C_o(\rho)(|\eta| + |\zeta|) \text{ [number of fronts of order} < k] .$ (6.2)

Then, (6.2) is less than $1/\nu$ if we choose k sufficiently large to have the first term less than $1/(2\nu)$ and $\rho = \rho_{\nu}(m_o)$ small enough to have the second term less than $1/(2\nu)$.

End of the proof of Theorem 2.1 As in the proof of [2, Theorem 2.1], in the bubble case (increasing-pressure case) by (2.4) and (4.15) ((5.30), respectively) we prove that

$$\bar{L}(0) \le \frac{1}{2} \operatorname{TV}(\log(p_o)) + \frac{1}{2\min\{a_\ell, a_m, a_r\}} \operatorname{TV}(u_o) .$$
(6.3)

Now, by (4.12) ((5.24), respectively) and (6.3) we look for an m_o satisfying both inequalities below:

$$\mathcal{H}(|\eta|, |\zeta|) < \frac{1}{c(m_o)} - 1 = \frac{2}{\cosh m_o - 1} =: w(m_o), \qquad (6.4)$$

$$\Gamma V\left(\log(p_o)\right) + \frac{1}{\min\{a_\ell, a_m, a_r\}} \operatorname{TV}(u_o) < 2m_o c(m_o) =: z(m_o).$$
(6.5)

Recall from [2] that w(r) is strictly decreasing and z(r) is strictly increasing, for $r \in \mathbb{R}^+$. Moreover, we have $\mathcal{K}(r) = z(w^{-1}(r))$, see (2.2). Hence, by (2.4) one can choose m_o such that (6.4), (6.5) hold in both the cases. Therefore, we can conclude as in [7] and Theorem 2.1 is completely proved.

As in [2], we want to compare the results obtained here in the bubble casewith that of [3]. More precisely, we set $x = |\eta|, y = |\zeta|$ and we claim that

$$\mathcal{H}_b(x,y) \le x+y \qquad \text{for } 0 \le x+y < 1/2.$$
(6.6)

Since \mathcal{H}_b is a symmetric function of x and y, it suffices to verify that in the common domain it holds

$$\frac{(2+x)4y}{(2-x)(4-xy)} < x+y.$$
(6.7)

By simplifying expression (6.7), we find that it is equivalent to

$$x^2y + xy^2 - (2xy + 2y^2 + 4x + 8y) + 8 > 0 \,,$$

which will be satisfied if $xy + y^2 + 2x + 4y < 4$. Since x < 1/2 - y, this last inequality is verified if

$$\left(\frac{1}{2} - y\right)y + y^2 + 2\left(\frac{1}{2} - y\right) + 4y < 4,$$

that is when y < 6/5. Therefore, (6.6) holds and, since \mathcal{K} is decreasing, we have

$$\mathcal{K}\left(\mathcal{H}_b(|\eta|, |\zeta|)\right) > \mathcal{K}\left(|\eta| + |\zeta|\right)$$

in the common domain $|\eta| + |\zeta| < 1/2$. Hence, Theorem 2.1 improves [3, Theorem 2.2] in the bubble case.

References

- Amadori D, Baiti P, Corli A, Dal Santo E. Global weak solutions for a model of two-phase flow with a single interface. submitted, 2014
- [2] Amadori D, Baiti P, Corli A, Dal Santo E. Global existence of solutions for a multi-phase flow: a drop in a gas-tube. submitted, 2015
- [3] Amadori D, Corli A. On a model of multiphase flow. SIAM J Math Anal, 2008, 40(1): 134–166
- [4] Amadori D, Guerra G. Global BV solutions and relaxation limit for a system of conservation laws. Proc Roy Soc Edinburgh Sect A, 2001, 131(1): 1–26
- [5] Asakura F, Corli A. Global existence of solutions by path decomposition for a model of multiphase flow. Quart Appl Math, 2013, 71(1): 135–182
- [6] Baiti P, Dal Santo E. Front tracking for a 2 × 2 system of conservation laws. Electron J Differ Equ, 2012, (220): 14
- [7] Bressan A. Hyperbolic Systems of Conservation Laws. The One-dimensional Cauchy Problem. Oxford University Press, 2000
- [8] Dafermos C M. Hyperbolic Conservation Laws in Continuum Physics. 3rd ed. Berlin: Springer-Verlag, 2010
- [9] DiPerna R. Existence in the large for quasilinear hyperbolic conservation laws. Arch Rational Mech Anal, 1973, 52: 244–257

- [10] DiPerna R. Global solutions to a class of nonlinear hyperbolic systems of equations. Comm Pure Appl Math, 1973, 26: 1–28
- [11] Fan H. On a model of the dynamics of liquid/vapor phase transitions. SIAM J Appl Math, 2000, 60(4): 1270–1301
- [12] Fermi E. Thermodynamics. Dover Publications, 1956
- [13] Godlewski E. Coupling fluid models. Exploring some features of interfacial coupling//Finite Volumes for Complex Applications V, London: ISTE, 2008: 87–102
- [14] Groah J, Smoller J, Temple B. Shock Wave Interactions in General Relativity. Springer Monographs in Mathematics. New York: Springer, 2007
- [15] Li T -T, Yu W -C. Boundary Value Problems for Quasilinear Hyperbolic Systems. Duke University, 1985
- [16] Liu T -P. Initial-boundary value problems for gas dynamics. Arch Rational Mech Anal, 1977, 64: 137–168
- [17] Liu T -P. Solutions in the large for the equations of nonisentropic gas dynamics. Indiana Univ Math J, 1977, 26: 147–177
- [18] Nishida T. Global solution for an initial boundary value problem of a quasilinear hyperbolic system. Proc Japan Acad, 1968, 44: 642–646
- [19] Nishida T, Smoller J A. Solutions in the large for some nonlinear hyperbolic conservation laws. Comm Pure Appl Math, 1973, 26: 183–200