



UNIVERSITÀ  
DEGLI STUDI  
DI UDINE

Università degli studi di Udine

## A Corrected Sadowsky Functional for Inextensible Elastic Ribbons

*Original*

*Availability:*

This version is available <http://hdl.handle.net/11390/1081370> since 2021-03-15T13:28:53Z

*Publisher:*

*Published*

DOI:10.1007/s10659-015-9551-4

*Terms of use:*

The institutional repository of the University of Udine (<http://air.uniud.it>) is provided by ARIC services. The aim is to enable open access to all the world.

*Publisher copyright*

(Article begins on next page)

# A CORRECTED SADOWSKY FUNCTIONAL FOR INEXTENSIBLE ELASTIC RIBBONS

LORENZO FREDDI, PETER HORNUNG, MARIA GIOVANNA MORA, AND ROBERTO PARONI

ABSTRACT. The classical theory of ribbons, developed by Sadowsky and Wunderlich, has recently received renewed attention. Here, by means of  $\Gamma$ -convergence, we re-examine the derivation of the limit energy of an inextensible, isotropic, elastic strip as the width goes to zero. We find that this rigorously derived functional agrees with the classical Sadowsky functional only for “large” curvature of the centerline of the strip.

## 1. INTRODUCTION

In 1930 Sadowsky [12] established the existence of a developable Möbius strip and stated that the configuration assumed by the strip can be computed by minimizing the bending energy. He further argued that the bending energy density is proportional to the square of the mean curvature of the surface.

The (bending) energy of an inextensible elastic strip  $S_\varepsilon = (-\ell/2, \ell/2) \times (-\varepsilon/2, \varepsilon/2)$ , where  $\ell > 0$  and  $\varepsilon > 0$  are the length and the thickness of the strip, is

$$u \mapsto \frac{1}{\varepsilon} \int_{S_\varepsilon} Q(A_u(x)) dx, \quad (1.1)$$

where  $Q$  is the bending energy density and  $A_u$  denotes the second fundamental form of  $u : S_\varepsilon \rightarrow \mathbb{R}^3$ . For isotropic strips the energy density  $Q$  depends only on the determinant and the trace of  $A_u$ : the Gaussian curvature  $K_u := \det A_u$  and the mean curvature  $H_u := 1/2 \operatorname{tr} A_u$ . Since developable surfaces have null Gaussian curvature we have that  $Q$  only depends on  $H_u$ . By assuming, as tacitly done by Sadowsky [12], the material to be isotropic and the energy density to be quadratic, the bending energy is proportional to

$$u \mapsto \frac{1}{\varepsilon} \int_{S_\varepsilon} |H_u(x)|^2 dx. \quad (1.2)$$

Sadowsky, still in [12], also argued that as  $\varepsilon \rightarrow 0$  the energy of the ribbon reduces to

$$y \mapsto \int_{-\ell/2}^{\ell/2} \frac{(\kappa^2 + \tau^2)^2}{\kappa^2} ds \quad (1.3)$$

where  $\kappa$  and  $\tau$  are the curvature and the torsion, respectively, of the centerline  $y$  of the strip, and  $s$  is its arc length. This functional is now known as Sadowsky’s functional. The seminal paper [12] and also the subsequent paper by Sadowsky [13] have recently been translated into English [8, 9].

Wunderlich in [17] gave a formal justification of the energy (1.3), see [16] for a translation. His analysis is based on the fact that every smooth developable surface is a ruled surface, in particular, as pointed out in [14], it follows that if the centerline  $y$  is smooth and with *non vanishing curvature*  $\kappa$ , then the surface  $u(S_\varepsilon)$  has a parametrization of the form

$$(s, z) \mapsto y(s) + z[b(s) + \eta(s)t(s)],$$

where  $t$  and  $b$  are the tangent and the binormal of the centerline  $y$  and  $\eta := \tau/\kappa$ . With this parametrization one may compute the mean curvature  $H_u$  and then rewrite the energy (1.2) as

$$y \mapsto \int_{-\ell/2}^{\ell/2} \frac{(\kappa^2 + \tau^2)^2}{\kappa^2} \frac{1}{\varepsilon \eta'} \ln \frac{1 + \varepsilon \eta'/2}{1 - \varepsilon \eta'/2} ds. \quad (1.4)$$

The energy (1.3) is then obtained as the pointwise limit, as  $\varepsilon \rightarrow 0$ , of this sequence of energies. Wunderlich's analysis makes it clear that the Sadowsky functional is derived under the assumption of non vanishing curvature of the centerline of the strip. Kirby and Fried [10] recently investigated if also the  $\Gamma$ -limit, under an appropriate topology, of the sequence of functionals (1.4) is the Sadowsky functional. They gave a positive answer after restricting the domain of the functional (1.4) to a space of curves with non vanishing curvature and with certain regularity.

Other interesting papers addressing the Sadowsky functional are [2, 3, 4, 11, 15].

In this paper we study the  $\Gamma$ -limit, with respect to a topology that ensures the convergence of the minimizers, of the sequence of energies (1.2): we therefore re-examine the same problem studied by Sadowsky in [12]. In our analysis, however, we make no a priori assumptions on the curvature of the centerline. The obtained limit functional depends on three orthonormal vectors  $d_1$ ,  $d_2$ , and  $d_3$ . The first director  $d_1$  is the tangent to the centerline  $y$ , whereas  $d_2$  represents the "transversal" orientation of the strip, and  $d_3 = d_1 \wedge d_2$ . The limit problem describes an inextensible beam (since  $y' = d_1$ ) which cannot bend within the plane of the strip (since the directors have to satisfy the constraint  $d_1' \cdot d_2 = 0$ ). Its energy is given by

$$J(d_1, d_2, d_3) = \int_{-\ell/2}^{\ell/2} \bar{Q}(d_1' \cdot d_3, d_2' \cdot d_3) ds,$$

where the energy density  $\bar{Q}$  is

$$\bar{Q}(\alpha, \beta) = \begin{cases} \frac{(\alpha^2 + \beta^2)^2}{\alpha^2} & \text{if } |\alpha| > |\beta|, \\ 4\beta^2 & \text{if } |\alpha| \leq |\beta|. \end{cases} \quad (1.5)$$

We also show that if the curvature  $\kappa$  of the centerline  $y$  is everywhere different from zero, then  $\kappa = |d_1' \cdot d_3|$  and the torsion  $\tau = d_2' \cdot d_3$ , so that the limit functional can be rewritten, using the notation adopted earlier, as

$$y \mapsto \int_{-\ell/2}^{\ell/2} \bar{Q}(\kappa, \tau) ds.$$

In view of this expression, we see that Sadowsky's functional (1.3) only agrees with the rigorously derived asymptotic functional when the curvature exceeds the absolute value of the torsion.

The techniques employed in our analysis make a strong use of the isotropy assumption. A Sadowsky type of functional for anisotropic elastic ribbons will be derived, with a more involved argument, in a forthcoming paper [7].

## 2. MAIN RESULTS

Let  $\ell > 0$  and  $I := (-\ell/2, \ell/2)$ . For  $0 < \varepsilon \ll 1$ , we take  $S_\varepsilon = I \times (-\varepsilon/2, \varepsilon/2)$  as the reference configuration of an inextensible elastic narrow strip. For a smooth deformation  $u : S_\varepsilon \rightarrow \mathbb{R}^3$  we have, due to the inextensibility constraint, that  $\partial_i u \cdot \partial_j u = \delta_{ij}$ , where  $\delta_{ij}$  denotes the Kronecker delta. The second fundamental form of  $u$ , denoted by  $A_u : S_\varepsilon \rightarrow \mathbb{R}_{\text{sym}}^{2 \times 2}$ , is defined by

$$(A_u)_{ij} = n_u \cdot \partial_i \partial_j u,$$

where

$$n_u = \partial_1 u \wedge \partial_2 u$$

is the unit normal to  $u$ . We recall that by Gauss's Theorema Egregium we have  $\det A_u = 0$ , because the Gaussian curvature is everywhere equal to zero. We assume the energy density of the strip,  $Q : \mathbb{R}_{\text{sym}}^{2 \times 2} \rightarrow [0, +\infty)$ , to be an isotropic and quadratic function of the second fundamental form. Under these assumptions we may take

$$Q(A) = |A|^2 \text{ for every } A \in \mathbb{R}_{\text{sym}}^{2 \times 2}.$$

We note that  $Q(A_u) = |A_u|^2 = (\text{tr } A_u)^2$  since for every  $A \in \mathbb{R}_{\text{sym}}^{2 \times 2}$  we have that  $|A|^2 + 2 \det A = (\text{tr } A)^2$ ; hence, the energy density considered here is equivalent with that discussed in the introduction.

Denoting the space of  $W^{2,2}$  isometries of  $S_\varepsilon$  by  $W_{iso}^{2,2}(S_\varepsilon; \mathbb{R}^3)$ , that is,

$$W_{iso}^{2,2}(S_\varepsilon; \mathbb{R}^3) := \{u \in W^{2,2}(S_\varepsilon; \mathbb{R}^3) : \partial_i u \cdot \partial_j u = \delta_{ij} \text{ a.e. in } S_\varepsilon\},$$

we have that the bending energy of the strip is

$$E_\varepsilon(u) = \frac{1}{\varepsilon} \int_{S_\varepsilon} |A_u(x)|^2 dx,$$

for any  $u \in W_{iso}^{2,2}(S_\varepsilon; \mathbb{R}^3)$ .

We now change variables in order to rewrite the energy over the fixed domain

$$S = I \times \left(-\frac{1}{2}, \frac{1}{2}\right).$$

We introduce the rescaled version  $y : S \rightarrow \mathbb{R}^3$  of  $u$  by setting

$$y(x_1, x_2) = u(x_1, \varepsilon x_2).$$

With the scaled gradient defined by

$$\nabla_\varepsilon \cdot = (\partial_1 \cdot \mid \varepsilon^{-1} \partial_2 \cdot),$$

we have that

$$\nabla_\varepsilon y(x) = \nabla u(x_1, \varepsilon x_2).$$

In particular, if  $u \in W_{iso}^{2,2}(S_\varepsilon; \mathbb{R}^3)$ , the map  $y$  belongs to the space of *scaled isometries*

$$W_{iso,\varepsilon}^{2,2}(S; \mathbb{R}^3) := \{y \in W^{2,2}(S; \mathbb{R}^3) : |\partial_1 y| = \varepsilon^{-1} |\partial_2 y| = 1, \partial_1 y \cdot \varepsilon^{-1} \partial_2 y = 0 \text{ a.e. in } S\}.$$

Defining the scaled unit normal to  $y$  by

$$n_{y,\varepsilon} = \partial_1 y \wedge \varepsilon^{-1} \partial_2 y$$

and the scaled second fundamental form of  $y$  by

$$A_{y,\varepsilon} = \begin{pmatrix} n_{y,\varepsilon} \cdot \partial_1 \partial_1 y & \varepsilon^{-1} n_{y,\varepsilon} \cdot \partial_1 \partial_2 y \\ \varepsilon^{-1} n_{y,\varepsilon} \cdot \partial_1 \partial_2 y & \varepsilon^{-2} n_{y,\varepsilon} \cdot \partial_2 \partial_2 y \end{pmatrix},$$

we have  $A_u(x_1, \varepsilon x_2) = A_{y,\varepsilon}(x_1, x_2)$ . Introducing

$$J_\varepsilon(y) = \int_S |A_{y,\varepsilon}(x)|^2 dx,$$

we have  $J_\varepsilon(y) = E_\varepsilon(u)$ .

Our first result, whose proof will be postponed to the next section, is about compactness.

**Lemma 2.1.** *Let  $(y_\varepsilon) \subset W_{iso,\varepsilon}^{2,2}(S; \mathbb{R}^3)$  be a sequence of scaled isometries such that*

$$\sup_\varepsilon J_\varepsilon(y_\varepsilon) < \infty. \quad (2.1)$$

*Then, up to a subsequence and additive constants, there exist a deformation  $y \in W^{2,2}(I; \mathbb{R}^3)$  and an orthonormal frame field  $(d_1 \mid d_2 \mid d_3) \in W^{1,2}(I; SO(3))$  satisfying*

$$d_1 = y' \quad \text{and} \quad d'_1 \cdot d_2 = 0 \quad \text{a.e. in } I, \quad (2.2)$$

*such that*

$$y_\varepsilon \rightharpoonup y \text{ in } W^{2,2}(S; \mathbb{R}^3), \quad \nabla_\varepsilon y_\varepsilon \rightharpoonup (d_1 \mid d_2) \text{ in } W^{1,2}(S; \mathbb{R}^{3 \times 2}), \quad (2.3)$$

*and*

$$A_{y_\varepsilon, \varepsilon} \rightharpoonup \begin{pmatrix} d'_1 \cdot d_3 & d'_2 \cdot d_3 \\ d'_2 \cdot d_3 & \gamma \end{pmatrix} \text{ in } L^2(S; \mathbb{R}_{\text{sym}}^{2 \times 2}) \quad (2.4)$$

*with  $\gamma \in L^2(S)$ .*

Hence a sequence  $(y_\varepsilon)$  of scaled isometries with equi-bounded energy weakly converges in  $W^{2,2}$  to a deformation  $y$  that depends only on  $x_1$ . The orthonormal vectors  $d_1$ ,  $d_2$ , and  $d_3$  are the directors of the “limit beam”, with  $d_1$  tangent to the deformation  $y$ ,  $d_2$  representing the “transversal” orientation of the strip, and  $d_3 = d_1 \wedge d_2$ . The limiting values of the 11 and 12 components of the second fundamental form are measures of flexure and twist, respectively, cf. [1]. The 22 component instead cannot be expressed in terms of the directors. We also note that  $A_{y_\varepsilon, \varepsilon}$  has null determinant while its limit, i.e., the limit matrix appearing in (2.4), does not need to have null determinant. The first constraint in (2.2) states that the “limit beam” is inextensible, while the second states that the strip does not bend within its plane. The same constraints were also obtained in [5, 6].

The previous lemma motivates the next definition. We set

$$\mathcal{A} = \{(d_1, d_2, d_3) : (d_1 | d_2 | d_3) \in W^{1,2}(I; SO(3)) \text{ and } d'_1 \cdot d_2 = 0 \text{ a.e. in } I\}.$$

In order to state our next result we need to first introduce some definitions. Let  $\overline{Q} : \mathbb{R} \times \mathbb{R} \rightarrow [0, +\infty)$  and  $J : \mathcal{A} \rightarrow \mathbb{R}$  be defined by

$$\overline{Q}(\kappa, \tau) := \min_{\gamma \in \mathbb{R}} \left\{ |M|^2 + 2|\det M| : M = \begin{pmatrix} \kappa & \tau \\ \tau & \gamma \end{pmatrix} \right\}$$

and

$$J(d_1, d_2, d_3) := \int_I \overline{Q}(d'_1 \cdot d_3, d'_2 \cdot d_3) dx_1.$$

A simple computation shows that  $\overline{Q}$  can be written as in (1.5).

**Theorem 2.2.** *As  $\varepsilon \rightarrow 0$ , the functionals  $J_\varepsilon$  are  $\Gamma$ -converging to the functional  $J$  in the following sense:*

- (i) (liminf inequality) *for every sequence  $(y_\varepsilon) \subset W^{2,2}_{iso,\varepsilon}(S; \mathbb{R}^3)$ ,  $y \in W^{2,2}(I; \mathbb{R}^3)$ , and  $(d_1, d_2, d_3) \in \mathcal{A}$  such that  $y' = d_1$  a.e. in  $I$ ,  $y_\varepsilon \rightharpoonup y$  in  $W^{2,2}(S; \mathbb{R}^3)$ , and  $\nabla_\varepsilon y_\varepsilon \rightharpoonup (d_1 | d_2)$  in  $W^{1,2}(S; \mathbb{R}^{3 \times 2})$ , we have that*

$$\liminf_{\varepsilon \rightarrow 0} J_\varepsilon(y_\varepsilon) \geq J(d_1, d_2, d_3);$$

- (ii) (recovery sequence) *for every  $(d_1, d_2, d_3) \in \mathcal{A}$  there exists a sequence  $(y_\varepsilon) \subset W^{2,2}_{iso,\varepsilon}(S; \mathbb{R}^3)$  such that  $y_\varepsilon \rightharpoonup y$  in  $W^{2,2}(S; \mathbb{R}^3)$ ,  $\nabla_\varepsilon y_\varepsilon \rightharpoonup (d_1 | d_2)$  weakly in  $W^{1,2}(S; \mathbb{R}^{3 \times 2})$ , and*

$$\limsup_{\varepsilon \rightarrow 0} J_\varepsilon(y_\varepsilon) \leq J(d_1, d_2, d_3),$$

where  $y$  is defined up to a constant by  $y' = d_1$  a.e. in  $I$ .

We conclude the section by comparing the obtained  $\Gamma$ -limit with Sadowsky’s functional. For  $(d_1, d_2, d_3) \in \mathcal{A}$ , let  $y$  be the function defined, up to a constant, by  $y' = d_1$ . Until the end of the section we suppose that  $y$  is smooth enough and the curvature

$$\kappa = |d'_1| = |y''|$$

is a strictly positive function. Under this assumption, the normal  $n = d'_1/\kappa$ , the binormal  $b = t \wedge n$ , and the torsion

$$\tau = -n \cdot b'$$

are well defined at every point of  $I$ . Set  $\tilde{\kappa} = d'_1 \cdot d_3$  and  $\tilde{\tau} = d'_2 \cdot d_3$ . We now study the relation between  $\kappa, \tau, \tilde{\kappa}$ , and  $\tilde{\tau}$ . Since  $d'_1$  is orthogonal to  $d_1$  and  $d_2$  it follows that  $d'_1 = \tilde{\kappa}d_3$ . Thus

$$\kappa = |d'_1| = |\tilde{\kappa}|, \quad n = \frac{\tilde{\kappa}}{\kappa}d_3 = \text{sgn}(\tilde{\kappa})d_3, \quad b = d_1 \wedge n = \text{sgn}(\tilde{\kappa})d_1 \wedge d_3 = -\text{sgn}(\tilde{\kappa})d_2,$$

where  $\text{sgn}$  is the sign function. Since  $n$  and  $d_3$  are continuous functions, the second equality implies that  $\text{sgn}(\tilde{\kappa})$  is continuous, hence constant. Thus, differentiating the third equality above we have that  $b' = -\text{sgn}(\tilde{\kappa})d'_2$ , hence

$$\tau = -n \cdot b' = -\text{sgn}(\tilde{\kappa})d_3 \cdot (-\text{sgn}(\tilde{\kappa})d'_2) = d_3 \cdot d'_2 = \tilde{\tau}.$$

Thus, if the curvature  $\kappa$  of  $y$  is strictly positive everywhere, then the  $\Gamma$ -limit can be rewritten as

$$J(d_1, d_2, d_3) = \int_I \bar{Q}(\kappa, \tau) dx_1,$$

since  $\kappa = |\tilde{\kappa}|$  and  $\tau = \tilde{\tau}$ . Therefore, the  $\Gamma$ -limit coincides with Sadowsky's functional for  $\kappa > |\tau|$  and  $\kappa > 0$ .

### 3. PROOFS

Here we prove the results stated in the previous section.

*Proof of Lemma 2.1.* Let  $(y_\varepsilon) \subset W_{iso,\varepsilon}^{2,2}(S; \mathbb{R}^3)$  be a sequence satisfying (2.1), that is

$$\sup_\varepsilon \|A_{y_\varepsilon, \varepsilon}\|_{L^2(S)} < +\infty.$$

Since  $y_\varepsilon$  is a scaled isometry, we have that

$$\partial_1 \partial_1 y_\varepsilon = (A_{y_\varepsilon, \varepsilon})_{11} n_{y_\varepsilon, \varepsilon}, \quad \varepsilon^{-1} \partial_1 \partial_2 y_\varepsilon = (A_{y_\varepsilon, \varepsilon})_{12} n_{y_\varepsilon, \varepsilon}, \quad \varepsilon^{-2} \partial_2 \partial_2 y_\varepsilon = (A_{y_\varepsilon, \varepsilon})_{22} n_{y_\varepsilon, \varepsilon},$$

where  $n_{y_\varepsilon, \varepsilon}$  is the scaled unit normal to  $y_\varepsilon$ . For instance, the last equation can be checked by differentiating with respect to  $x_1$  and  $x_2$  the identities  $\varepsilon^{-1} |\partial_2 y_\varepsilon| = 1$  and  $\partial_1 y_\varepsilon \cdot \varepsilon^{-1} \partial_2 y_\varepsilon = 0$ . Thus

$$\sup_\varepsilon (\|\partial_1 \partial_1 y_\varepsilon\|_{L^2(S)} + \|\varepsilon^{-1} \partial_1 \partial_2 y_\varepsilon\|_{L^2(S)} + \|\varepsilon^{-2} \partial_2 \partial_2 y_\varepsilon\|_{L^2(S)}) < +\infty. \quad (3.1)$$

Moreover,  $(y_\varepsilon) \subset W_{iso,\varepsilon}^{2,2}(S; \mathbb{R}^3)$  also implies that

$$\|\partial_1 y_\varepsilon\|_{L^\infty(S)} = 1, \quad \|\partial_2 y_\varepsilon\|_{L^\infty(S)} = \varepsilon, \quad (3.2)$$

and hence, up to additive constants, the sequence  $(y_\varepsilon)$  is uniformly bounded in  $W^{2,2}(S; \mathbb{R}^3)$ . Therefore, up to subsequences, we have that  $y_\varepsilon \rightharpoonup y$  in  $W^{2,2}(S; \mathbb{R}^3)$  and strongly in  $W^{1,p}(S; \mathbb{R}^3)$  for every  $p < \infty$ . Identities (3.2) imply that  $y$  is independent of  $x_2$  and  $|y'| = 1$  a.e. in  $I$ .

By the previous bounds the sequence  $(\varepsilon^{-1} \partial_2 y_\varepsilon)$  is bounded in  $L^\infty(S; \mathbb{R}^3)$  and  $(\nabla_\varepsilon(\varepsilon^{-1} \partial_2 y_\varepsilon))$  is bounded in  $L^2(S; \mathbb{R}^{3 \times 2})$ . Hence, up to subsequences,  $\varepsilon^{-1} \partial_2 y_\varepsilon \rightharpoonup d_2$  weakly in  $W^{1,2}(S; \mathbb{R}^3)$  and strongly in  $L^p(S; \mathbb{R}^3)$  for every  $p < \infty$ , with  $d_2$  independent of  $x_2$  and  $|d_2| = 1$  a.e. in  $S$ . Moreover, by passing to the limit in the relation  $\partial_1 y_\varepsilon \cdot (\varepsilon^{-1} \partial_2 y_\varepsilon) = 0$ , we deduce that  $y' \cdot d_2 = 0$  a.e. in  $I$ . Since  $n_{y_\varepsilon, \varepsilon} = \partial_1 y_\varepsilon \wedge \varepsilon^{-1} \partial_2 y_\varepsilon$  we have that  $n_{y_\varepsilon, \varepsilon} \rightarrow d_3$  in  $L^p(S; \mathbb{R}^3)$  for every  $p < \infty$ , where  $d_3 = y' \wedge d_2$ .

By differentiating the equality  $\partial_1 y_\varepsilon \cdot \partial_1 y_\varepsilon = 1$  with respect to  $x_2$  and scaling by  $\varepsilon$ , we obtain

$$\partial_1(\varepsilon^{-1} \partial_2 y_\varepsilon) \cdot \partial_1 y_\varepsilon = 0.$$

By letting  $\varepsilon$  go to zero we find  $d'_2 \cdot y' = 0$ , from which, setting  $d_1 := y'$  and using that  $d_1 \cdot d_2 = 0$ , we deduce that  $d'_1 \cdot d_2 = 0$ .

Finally, up to subsequences, we have that  $A_{y_\varepsilon, \varepsilon}$  weakly converges to a matrix field  $A$  in  $L^2(S; \mathbb{R}_{\text{sym}}^{2 \times 2})$ . By using the convergences established above, it follows that  $A_{11} = y'' \cdot d_3$  and  $A_{12} = d'_2 \cdot d_3$ . The entry  $A_{22}$ , that cannot be identified in terms of  $y$ ,  $d_2$ , and  $d_3$ , is denoted by  $\gamma$  in the statement.  $\square$

We now prove the liminf inequality in Theorem 2.2.

*Proof of Theorem 2.2-(i), liminf inequality.* We may suppose that  $\liminf_{\varepsilon \rightarrow 0} J_\varepsilon(y_\varepsilon) < \infty$ , since otherwise there is nothing to prove. Then, by passing to a subsequence, we may suppose that  $\sup_\varepsilon J_\varepsilon(y_\varepsilon) < \infty$ . By Lemma 2.1 we have that  $A_{y_\varepsilon, \varepsilon} \rightharpoonup A$  in  $L^2(S; \mathbb{R}_{\text{sym}}^{2 \times 2})$ , where

$$A = \begin{pmatrix} d'_1 \cdot d_3 & d'_2 \cdot d_3 \\ d'_2 \cdot d_3 & \gamma \end{pmatrix}$$

with  $\gamma \in L^2(S)$ . In the rest of the proof, to simplify the notation, we set  $A^\varepsilon := A_{y_\varepsilon, \varepsilon}$ . We note that  $|A^\varepsilon|^2 = (\text{tr } A^\varepsilon)^2$ , since  $|B|^2 + 2 \det B = (\text{tr } B)^2$  for every  $B \in \mathbb{R}_{\text{sym}}^{2 \times 2}$  and  $\det A^\varepsilon = 0$ , and also that

$$(\text{tr } A^\varepsilon)^2 = (A_{11}^\varepsilon - A_{22}^\varepsilon)^2 + 4A_{11}^\varepsilon A_{22}^\varepsilon = (A_{11}^\varepsilon - A_{22}^\varepsilon)^2 + 4(A_{12}^\varepsilon)^2.$$

Let  $S^+ = S \cap \{\det A \geq 0\}$  and  $S^- = S \cap \{\det A < 0\}$ , then

$$\begin{aligned} \liminf_{\varepsilon \rightarrow 0} J_\varepsilon(y_\varepsilon) &= \liminf_{\varepsilon \rightarrow 0} \int_S (\operatorname{tr} A^\varepsilon)^2 dx = \liminf_{\varepsilon \rightarrow 0} \left\{ \int_{S^+} (\operatorname{tr} A^\varepsilon)^2 dx + \int_{S^-} (A_{11}^\varepsilon - A_{22}^\varepsilon)^2 + 4(A_{12}^\varepsilon)^2 dx \right\} \\ &\geq \int_{S^+} (\operatorname{tr} A)^2 dx + \int_{S^-} (A_{11} - A_{22})^2 + 4(A_{12})^2 dx, \end{aligned}$$

where in the last inequality we used the lower semicontinuity of convex functionals with respect to  $L^2$ -weak convergence. Noticing again that  $(\operatorname{tr} A)^2 = |A|^2 + 2\det A$  and that

$$(A_{11} - A_{22})^2 + 4(A_{12})^2 = (A_{11})^2 + (A_{22})^2 + 2(A_{12})^2 - 2(A_{11}A_{22} - (A_{12})^2) = |A|^2 - 2\det A,$$

we deduce that

$$\begin{aligned} \liminf_{\varepsilon \rightarrow 0} J_\varepsilon(y_\varepsilon) &\geq \int_{S^+} |A|^2 + 2\det A dx + \int_{S^-} |A|^2 - 2\det A dx = \int_S |A|^2 + 2|\det A| dx \\ &\geq \int_S \overline{Q}(d'_1 \cdot d_3, d'_2 \cdot d_3) dx = J(d_1, d_2, d_3), \end{aligned}$$

where the last inequality follows from the definitions of  $A$  and of  $\overline{Q}$ .  $\square$

The following lemma plays a crucial role in the construction of the recovery sequence of Theorem 2.2.

**Lemma 3.1.** *Let  $M \in L^2(I; \mathbb{R}_{\text{sym}}^{2 \times 2})$ . There exists a sequence  $(M_n) \subset L^2(I; \mathbb{R}_{\text{sym}}^{2 \times 2})$  such that  $\det M_n = 0$  for every  $n$ ,  $M_n \rightharpoonup M$  in  $L^2(I; \mathbb{R}_{\text{sym}}^{2 \times 2})$ , and*

$$\int_I |M_n|^2 dx \rightarrow \int_I |M|^2 + 2|\det M| dx.$$

*Proof.* If  $\det M = 0$ , then we can choose all  $M_n$  to be identically equal to  $M$ . Thus, hereafter we assume  $\det M \neq 0$ . We subdivide the proof into three steps.

*Step 1: assume  $M$  is constant.* Further assume, for the moment, that there exist  $\lambda_1, \lambda_2 \in \mathbb{R} \setminus \{0\}$  such that  $M = \operatorname{diag}(\lambda_1, \lambda_2) = \lambda_1 e_1 \otimes e_1 + \lambda_2 e_2 \otimes e_2$ . Let

$$\theta = \frac{|\lambda_1|}{|\lambda_1| + |\lambda_2|} \in (0, 1),$$

so that

$$\frac{\lambda_1^2}{\theta} + \frac{\lambda_2^2}{1-\theta} = \lambda_1^2 + \lambda_2^2 + 2|\lambda_1 \lambda_2| = |M|^2 + 2|\det M|. \quad (3.3)$$

Let  $\chi : \mathbb{R} \rightarrow \{0, 1\}$  denote the 1-periodic extension of the characteristic function of the interval  $(0, \theta)$ . Define  $M_n : I \rightarrow \mathbb{R}_{\text{sym}}^{2 \times 2}$  by setting

$$M_n(x) = \chi(nx_1) \frac{\lambda_1}{\theta} e_1 \otimes e_1 + (1 - \chi(nx_1)) \frac{\lambda_2}{1-\theta} e_2 \otimes e_2.$$

Clearly  $\det M_n = 0$  for every  $n$  and  $M_n$  converges weakly\* in  $L^\infty(I; \mathbb{R}_{\text{sym}}^{2 \times 2})$  to the constant matrix  $M$ , since  $\chi(n \cdot) \rightharpoonup \theta$  weakly\* in  $L^\infty(I)$ .

On the other hand, using (3.3), we compute

$$\int_I |M_n|^2 dx = \int_I \theta \cdot \frac{\lambda_1^2}{\theta^2} + (1 - \theta) \cdot \frac{\lambda_2^2}{(1-\theta)^2} dx = \int_I |M|^2 + 2|\det M| dx. \quad (3.4)$$

This concludes the proof of the step in the case that  $M$  is diagonal.

For an arbitrary constant matrix  $M \in \mathbb{R}_{\text{sym}}^{2 \times 2}$  with  $\det M \neq 0$  there exists  $Q \in O(2)$  such that  $Q^T M Q$  is diagonal. Applying the construction above to  $Q^T M Q$  we obtain a sequence  $\widetilde{M}_n$  with the properties stated in the lemma for  $Q^T M Q$ , and setting  $M_n = Q \widetilde{M}_n Q^T$  we find the desired sequence.

*Step 2: assume  $M$  is piecewise constant.* It suffices to apply Step 1 on each interval on which  $M$  is constant.

*Step 3: assume  $M \in L^2(I; \mathbb{R}_{\text{sym}}^{2 \times 2})$ .* It suffices to approximate  $M$  by a sequence  $(M_k)$  of piecewise constant matrices in the strong topology of  $L^2(I; \mathbb{R}_{\text{sym}}^{2 \times 2})$ . For each  $M_k$  we apply Step 2 and obtain

a sequence  $(M_{k,n})$  with the required properties and such that  $\|M_{k,n}\|_{L^2}^2 \leq 2\|M_k\|_{L^2}^2$  for every  $k$  and  $n$ , in view of (3.4). This allows one to apply a diagonal argument and conclude the proof.  $\square$

*Proof of Theorem 2.2-(ii), recovery sequence.* Let  $(d_1, d_2, d_3) \in \mathcal{A}$  and let  $y \in W^{2,2}(I; \mathbb{R}^3)$  be such that  $y' = d_1$  a.e. in  $I$ . We set  $R := (y' \mid d_2 \mid d_3) \in SO(3)$  a.e. in  $I$  and

$$M := \begin{pmatrix} y'' \cdot d_3 & d_2' \cdot d_3 \\ d_2' \cdot d_3 & \gamma \end{pmatrix},$$

where  $\gamma \in L^2(I)$  is such that

$$\overline{Q}(y'' \cdot d_3, d_2' \cdot d_3) = |M|^2 + 2|\det M| \quad \text{a.e. in } I.$$

Such a  $\gamma$  can indeed be chosen measurable. Moreover,  $\gamma \in L^2(I)$  because by minimality, comparing  $M$  to the same matrix with 0 instead of  $\gamma$ , we have

$$\gamma^2 \leq |M|^2 + 2|\det M| \leq M_{11}^2 + 4M_{12}^2 \quad \text{a.e. in } I,$$

and the right-hand side is in  $L^1(I)$ .

By Lemma 3.1 there exist  $M_n \in L^2(I; \mathbb{R}_{\text{sym}}^{2 \times 2})$  with  $\det M_n = 0$  for every  $n$  and such that  $M_n \rightharpoonup M$  weakly in  $L^2(I; \mathbb{R}_{\text{sym}}^{2 \times 2})$  and

$$\mathcal{F}(M_n) := \int_I |M_n|^2 dx_1 \rightarrow \overline{\mathcal{F}}(M) := \int_I |M|^2 + 2|\det M| dx_1,$$

as  $n \rightarrow \infty$ . Denote by  $\lambda_n \in L^2(I)$  the trace of  $M_n$ . Since  $M_n$  is symmetric with  $\det M_n = 0$ , there exists  $\beta_n(x_1) \in (-\pi/2, \pi/2]$  such that

$$M_n = \begin{pmatrix} \cos \beta_n & -\sin \beta_n \\ \sin \beta_n & \cos \beta_n \end{pmatrix} \begin{pmatrix} \lambda_n & 0 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} \cos \beta_n & \sin \beta_n \\ -\sin \beta_n & \cos \beta_n \end{pmatrix},$$

and  $\beta_n$  is uniquely determined if  $\lambda_n \neq 0$ . When  $\lambda_n(x_1) = 0$ , we set  $\beta_n(x_1) = 0$ . After truncating  $\lambda_n$  in modulus by  $n$ , we may assume without loss of generality that  $\lambda_n \in L^\infty(I)$ , while  $M_n$  still enjoys the same properties as before. By mollification, we can find  $\lambda_{n,k} \in C^\infty(\bar{I})$  and  $\beta_{n,k} \in C^\infty(\bar{I})$  such that

- $\beta_{n,k}(x_1) \in (-\pi/2, \pi/2)$  for every  $x_1 \in \bar{I}$  and every  $n, k$  (this condition is achieved after possibly multiplying each mollification by a constant smaller than 1);
- $\lambda_{n,k} \rightarrow \lambda_n$  in  $L^p(I)$  for every  $p < +\infty$ , as  $k \rightarrow \infty$ ;
- $\beta_{n,k} \rightarrow \beta_n$  in  $L^p(I)$  for every  $p < +\infty$ , as  $k \rightarrow \infty$ .

Set

$$M_{n,k} := \begin{pmatrix} \cos \beta_{n,k} & -\sin \beta_{n,k} \\ \sin \beta_{n,k} & \cos \beta_{n,k} \end{pmatrix} \begin{pmatrix} \lambda_{n,k} & 0 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} \cos \beta_{n,k} & \sin \beta_{n,k} \\ -\sin \beta_{n,k} & \cos \beta_{n,k} \end{pmatrix}.$$

Then,  $\det M_{n,k} = 0$  for every  $n, k$  and  $M_{n,k} \rightarrow M_n$  in  $L^2(I; \mathbb{R}_{\text{sym}}^{2 \times 2})$ , as  $k \rightarrow \infty$ .

Thus, by a diagonal argument, we may assume that there exist  $\lambda^j \in C^\infty(\bar{I})$  and  $\beta^j \in C^\infty(\bar{I})$  such that  $|\beta^j| < \pi/2$  on  $\bar{I}$ , and with

$$\begin{aligned} M^j &:= \begin{pmatrix} \cos \beta^j & -\sin \beta^j \\ \sin \beta^j & \cos \beta^j \end{pmatrix} \begin{pmatrix} \lambda^j & 0 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} \cos \beta^j & \sin \beta^j \\ -\sin \beta^j & \cos \beta^j \end{pmatrix} \\ &= \lambda^j \begin{pmatrix} \cos^2 \beta^j & \sin \beta^j \cos \beta^j \\ \sin \beta^j \cos \beta^j & \sin^2 \beta^j \end{pmatrix} \end{aligned}$$

we have that  $\det M^j = 0$  for every  $j$ ,  $M^j \rightharpoonup M$  in  $L^2(I; \mathbb{R}_{\text{sym}}^{2 \times 2})$ , and  $\mathcal{F}(M_j) \rightarrow \overline{\mathcal{F}}(M)$ , as  $j \rightarrow \infty$ .

Set  $\vartheta^j = \frac{\pi}{2} + \beta^j$  and define

$$\tilde{b}^j(\xi_1) := \cos \vartheta^j(\xi_1) e_1 + \sin \vartheta^j(\xi_1) e_2$$

and  $\Phi^j(\xi_1, \xi_2) := \xi_1 e_1 + \xi_2 \tilde{b}^j(\xi_1)$ . Since  $\beta^j \in (-\pi/2, \pi/2)$ , we can argue as in [7] to see that, for  $\varepsilon$  small enough,  $(\Phi^j)^{-1} : S_\varepsilon \rightarrow \mathbb{R}^2$  is well defined.



Let  $R^j : I \rightarrow SO(3)$  be the solution of the ODE

$$(R^j)' = R^j \begin{pmatrix} 0 & 0 & -M_{11}^j \\ 0 & 0 & -M_{12}^j \\ M_{11}^j & M_{12}^j & 0 \end{pmatrix} \quad (3.5)$$

with  $R^j(0) = R(0) = (y'(0) \mid d_2(0) \mid d_3(0))$ . Since  $M^j$  is smooth, so is  $R^j$  and, since  $R(0) \in SO(3)$ ,  $R^j$  attains values in  $SO(3)$ . We set

$$d_k^j(t) = R^j(t)e_k \text{ for } k = 1, 2, 3, \quad y^j(t) = y(0) + \int_0^t d_1^j(s) ds.$$

Then  $y^j \rightharpoonup y$  weakly in  $W^{2,2}(I; \mathbb{R}^3)$ , see, for instance, the proof of Lemma 4.2 of [5]. It follows from (3.5) that

$$\begin{aligned} (d_1^j)' \cdot d_2^j &= 0, \\ (d_2^j)' \cdot d_3^j &= M_{12}^j = \lambda^j \sin \beta^j \cos \beta^j = -\lambda^j \sin \vartheta^j \cos \vartheta^j, \\ (d_1^j)' \cdot d_3^j &= M_{11}^j = \lambda^j \cos^2 \beta^j = \lambda^j \sin^2 \vartheta^j. \end{aligned} \quad (3.6)$$

Define

$$\begin{aligned} b^j(\xi_1) &= \cos \vartheta^j(\xi_1) d_1^j(\xi_1) + \sin \vartheta^j(\xi_1) d_2^j(\xi_1), \\ v^j(\xi_1, \xi_2) &= y^j(\xi_1) + \xi_2 b^j(\xi_1), \\ u^j(x_1, x_2) &= v^j((\Phi^j)^{-1}(x_1, x_2)). \end{aligned}$$

Then

$$\nabla v^j = ((y^j)' + \xi_2(b^j)' \mid b^j), \quad \nabla \Phi^j = \begin{pmatrix} 1 - \xi_2(\vartheta^j)' \sin \vartheta^j & \cos \vartheta^j \\ \xi_2(\vartheta^j)' \cos \vartheta^j & \sin \vartheta^j \end{pmatrix}, \quad (\nabla u^j)(\Phi^j) \nabla \Phi^j = \nabla v^j.$$

By means of (3.6) we can check that

$$(b^j)' \cdot d_3^j = 0, \quad |(b^j)'| = |(\vartheta^j)'|.$$

With these identities we can show that  $(\nabla v^j)^T \nabla v^j = (\nabla \Phi^j)^T \nabla \Phi^j$ , that is,  $(\nabla u^j)^T \nabla u^j = I$ . Clearly  $u^j(x_1, 0) = y^j(x_1)$  and  $\partial_1 u^j(x_1, 0) = (y^j)'(x_1) = d_1^j(x_1)$ . Moreover, since  $\partial_2 \Phi^j = \tilde{b}^j$ , we have

$$\nabla u^j(\Phi^j) \tilde{b}^j = \partial_2 v^j = b^j. \quad (3.7)$$

From this one readily deduces that

$$\nabla u^j(\cdot, 0) = (d_1^j \mid d_2^j). \quad (3.8)$$

Taking  $\xi_2$ -derivatives on both sides of (3.7), we see that  $\nabla^2 u(\Phi^j)(\tilde{b}^j, \tilde{b}^j) = 0$ , and therefore

$$A_{u^j}(\Phi^j)(\tilde{b}^j, \tilde{b}^j) = 0. \quad (3.9)$$

Taking derivatives in (3.8), we see that

$$(A_{u^j}(\cdot, 0))_{11} = d_3^j \cdot \partial_1 \partial_1 u^j(\cdot, 0) = d_3^j \cdot (y^j)'' = M_{11}^j$$

and similarly that  $(A_{u^j}(\cdot, 0))_{12} = M_{12}^j$ . Combining these with (3.9) and with the fact that  $M^j \tilde{b}^j = 0$ , and recalling that  $\tilde{b}^j \cdot e_2 \neq 0$ , we conclude that  $A_{u^j}(x_1, 0) = M^j(x_1)$ , because both  $A_{u^j}$  and  $M^j$  are symmetric and have zero determinant.

For  $\varepsilon$  small enough, the maps  $y_\varepsilon^j : S \rightarrow \mathbb{R}^3$  given by  $y_\varepsilon^j(x_1, x_2) = u^j(x_1, \varepsilon x_2)$  are well-defined scaled  $C^2$  isometries of  $S$ , such that

$$\nabla_\varepsilon y_\varepsilon^j = (\nabla u^j)(T_\varepsilon) \rightarrow \nabla u^j(\cdot, 0) = (d_1^j \mid d_2^j) \quad \text{strongly in } W^{1,2}(S; \mathbb{R}^{3 \times 2}),$$

as  $\varepsilon \rightarrow 0$ ; here  $T_\varepsilon x = (x_1, \varepsilon x_2)$ . Set  $A_\varepsilon^j := A_{y_\varepsilon^j}$ . Then since  $A_{u^j}(x_1, 0) = M^j(x_1)$ , we see that  $A_\varepsilon^j \rightarrow M^j$  strongly in  $L^2(S; \mathbb{R}_{\text{sym}}^{2 \times 2})$ , as  $\varepsilon \rightarrow 0$ . Hence,

$$\lim_{\varepsilon \rightarrow 0} J_\varepsilon(y_\varepsilon^j) = \lim_{\varepsilon \rightarrow 0} \int_S |A_\varepsilon^j|^2 dx = \int_S |M^j|^2 dx = \mathcal{F}(M^j).$$

Therefore, taking diagonal sequences we obtain the desired maps.  $\square$

**Acknowledgements.** The work of LF has been supported by the GNAMPA-INdAM project: “Metodi asintotici per problemi di fratture, delaminazioni e giunzioni”. PH was supported by the Deutsche Forschungsgemeinschaft. MGM acknowledges support by GNAMPA-INdAM and by the European Research Council under Grant No. 290888 “Quasistatic and Dynamic Evolution Problems in Plasticity and Fracture”. RP is grateful to Regione Autonoma della Sardegna for the support given under the project “Modellazione Multiscala della Meccanica dei Materiali Compositi (M4C)”.

## REFERENCES

- [1] S.S. Antman: *Nonlinear Problems of Elasticity*. Springer, New York, 2005.
- [2] S. Bartels, P. Hornung: Bending Paper and the Möbius Strip, *J. Elasticity* **119** (2015), 113–136.
- [3] M.A. Dias, B. Audoly: “Wunderlich, Meet Kirchhoff”: A General and Unified Description of Elastic Ribbons and Thin Rods, *J. Elasticity* **119** (2015), 49–66.
- [4] E. Efrati: Non-Euclidean Ribbons, *J. Elasticity* **119** (2015), 251–261.
- [5] L. Freddi, M.G. Mora, R. Paroni: Nonlinear thin-walled beams with a rectangular cross-section – Part I. *Math. Models Methods Appl. Sci.* **22** (2012), 1150016 (34 pp).
- [6] L. Freddi, M.G. Mora, R. Paroni: Nonlinear thin-walled beams with a rectangular cross-section – Part II. *Math. Models Methods Appl. Sci.* **23** (2013), 743–775.
- [7] L. Freddi, P. Hornung, M.G. Mora, R. Paroni, *in preparation*.
- [8] D.F. Hinz, E. Fried: Translation of Michael Sadowsky’s Paper “An Elementary Proof for the Existence of a Developable Möbius Band and the Attribution of the Geometric Problem to a Variational Problem”, *J. Elasticity* **119** (2015), 3–6.
- [9] D.F. Hinz, E. Fried: Translation and Interpretation of Michael Sadowsky’s Paper “Theory of Elastically Bendable Inextensible Bands with Applications to the Möbius Band”, *J. Elasticity* **119** (2015), 7–17.
- [10] N.O. Kirby, E. Fried: Gamma-Limit of a Model for the Elastic Energy of an Inextensible Ribbon, *J. Elasticity* **119** (2015), 35–47.
- [11] T. Randrup, P. Røgen: Sides of the Möbius strip. *Arch. Math.* **66** (1996), 511–521.
- [12] M. Sadowsky: Ein elementarer Beweis für die Existenz eines abwickelbaren Möbiuschen Bandes und die Zurückführung des geometrischen Problems auf ein Variationsproblem. *Sitzungsber. Preuss. Akad. Wiss.* 1930 – Mitteilung vom 26. Juni, 412–415.
- [13] M. Sadowsky: Theorie der elastisch biegsamen undehnbaren Bänder mit Anwendungen auf das Möbiussche Band. *Verhandl. des 3. Intern. Kongr. f. Techn. Mechanik* **2** (1930), 444–451.
- [14] E.L. Starostin, G.H.M. van der Heijden: Equilibrium Shapes with Stress Localisation for Inextensible Elastic Möbius and Other Strips, *J. Elasticity* **119** (2015), 67–112.
- [15] E.L. Starostin, G.H.M. van der Heijden: The equilibrium shape of an elastic developable Möbius strip, *Proc. Appl. Math. Mech.* **7** (2007), 2020115–2020116
- [16] R.E. Todres: Translation of W. Wunderlich’s “On a Developable Möbius Band”, *J. Elasticity*, (2015) **119** (2015), 23–34.
- [17] W. Wunderlich: Über ein abwickelbares Möbiusband, *Monatshefte für Mathematik* **66** (1962), 276–289.

(L. Freddi) DIPARTIMENTO DI MATEMATICA E INFORMATICA, VIA DELLE SCIENZE 206, 33100 UDINE, ITALY  
*E-mail address:* `lorenzo.freddi@uniud.it`

(P. Hornung) FACHRICHTUNG MATHEMATIK, TU DRESDEN, 01062 DRESDEN, GERMANY  
*E-mail address:* `peter.hornung@tu-dresden.de`

(M.G. Mora) DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI PAVIA, VIA FERRATA 1, 27100 PAVIA, ITALY  
*E-mail address:* `mariagiovanna.mora@unipv.it`

(R. Paroni) DADU, UNIVERSITÀ DEGLI STUDI DI SASSARI, PALAZZO DEL POU SALIT, 07041 ALGHERO (SS), ITALY  
*E-mail address:* `paroni@uniss.it`