



UNIVERSITÀ
DEGLI STUDI
DI UDINE

Università degli studi di Udine

The iterability hierarchy above I3

Original

Availability:

This version is available <http://hdl.handle.net/11390/1142080> since 2021-03-16T17:21:22Z

Publisher:

Published

DOI:10.1007/s00153-018-0624-5

Terms of use:

The institutional repository of the University of Udine (<http://air.uniud.it>) is provided by ARIC services. The aim is to enable open access to all the world.

Publisher copyright

(Article begins on next page)

THE ITERABILITY HIERARCHY ABOVE I3

ALESSANDRO ANDRETTA AND VINCENZO DIMONTE

ABSTRACT. In this paper we introduce a new hierarchy of large cardinals between I3 and I2, the iterability hierarchy, and we prove that every step of it strongly implies the ones below.

1. INTRODUCTION

In the late 70's and early 80's there was a flurry of activity around rank-into-rank axioms, a new kind of large cardinal hypotheses at the top of the hierarchy. They were I3 (the existence of an elementary embedding $j: V_\lambda \prec V_\lambda$), I2 (the existence of an elementary embedding $j: V \prec M$ with $V_\lambda \subseteq M$) and I1 (the existence of an elementary embedding $j: V_{\lambda+1} \prec V_{\lambda+1}$). The initial belief of the majority of set theorists was that their existence was eventually going to be disproved in ZFC, and hence the I suggesting inconsistency, but a result by Martin started to change the mood: he proved in [?] that a hypothesis strictly below I2 and above I3 implies the determinacy of $\mathbf{\Pi}_2^1$ sets. The hypothesis used by Martin was the existence of an *iterable* (see Section ??) $j: V_\lambda \prec V_\lambda$ —we call this large cardinal axiom $I3_\infty$. The excitement grew when Woodin, a few years later, proved the consistency of the Axiom of Determinacy using an axiom stronger than I1, called I0. But in the following years Woodin, building on work of Martin and Steel, showed that determinacy had much lower consistency strength, so $I3_\infty$ slowly fell into oblivion. Interest in rank-into-rank axioms re-emerged twenty years ago, thanks to Laver's results on the algebra of the I3-embeddings, and more recently after Woodin's extensive work on I0. Moving to the present age, the researcher that wants to know more about $I3_\infty$, however, is going to be disappointed: the only reference is Martin's original paper, that is very terse and lacking in details. For example, it says that I2 strictly implies $I3_\infty$, but it provides no proof for that. Even the very definition of iterable embedding is not fully satisfactory, as it is founded on operations whose validity has not been fully provided in print (as in Lemma ??). Also, is $I3_\infty$ strictly stronger than I3?


The aim of this paper is to approach iterability of I3-embeddings in all the details and in a modern way, thanks to the better understanding of rank-into-rank axioms that decades of work has given us. In Section ?? the current knowledge of

Date: December 14, 2017.

2010 Mathematics Subject Classification. 03E55.

Key words and phrases. Rank-into-rank embeddings, linear iterations.

rank-into-rank axioms is described, introducing the concept of “strong implication”, more suitable for such axioms than the usual notion of strict implication. In Section ?? we define exactly what it is an iterable $\mathfrak{I}3$ -embedding, and we introduce a new hierarchy of axioms, depending on how long the $\mathfrak{I}3$ -embedding can be iterable. It turns out that, like for iterations of a single measure or an extender, if an $\mathfrak{I}3$ -embedding is ω_1 -iterable, then it is iterable for any length, and this is strictly (in fact: strongly) weaker than $\mathfrak{I}2$. Finally, in Section ?? we prove that every step in the iterability hierarchy strongly implies the ones below, with even one more step at limit points. The final picture is therefore the following:



Acknowledgments. The second author would like to thank the program “Rita Levi Montalcini 2013” for the support, and the Department of Mathematics of the University of Turin for its hospitality.

2. PRELIMINARIES

To avoid confusion or misunderstandings, all notations and standard basic results are collected here.

If M and N are sets or classes, $j: M \prec N$ denotes that j is an elementary embedding from M to N , that is a function such that for any formula φ and any $x \in M$, $M \models \varphi(x)$ iff $N \models \varphi(j(x))$; when this holds only for Σ_n formulæ, we write

$j: M \prec_n N$. The case in which j is the identity, i.e., if M is an elementary (or Σ_n -elementary) submodel of N , is simply written as $M \prec N$ (or $M \prec_n N$).

If $M \models \text{AC}$ or $N \subseteq M$ and $j: M \prec N$ is not the identity, then it moves at least one ordinal; the least such ordinal is the *critical point* of j and it is denoted by $\text{crt}(j)$. Let j be an elementary embedding and $\kappa = \text{crt}(j)$. Define $\kappa_0 = \kappa$ and $\kappa_{n+1} = j(\kappa_n)$. Then $\langle \kappa_n : n \in \omega \rangle$ is the *critical sequence* of j .

Kunen [?] proved that if $M = N = V_\eta$ for some ordinal η , and λ is the supremum of the critical sequence, then η cannot be bigger than $\lambda + 1$ (and of course cannot be smaller than λ). Kunen's result actually does not say anything about the cases $\eta = \lambda$ or $\eta = \lambda + 1$. Therefore we can introduce the following hypotheses without fearing an immediate inconsistency:

- I3: There exists $j: V_\lambda \prec V_\lambda$, where λ is the supremum of the critical sequence of j .
- I1: There exists $j: V_{\lambda+1} \prec V_{\lambda+1}$, where λ is the supremum of the critical sequence of j .

We will be flexible in handling this and other rank-into-rank notations, but the meaning will be always clear. For example, I3(λ) means that there is a $j: V_\lambda \prec V_\lambda$ (so it is a property of λ), while I3(j) indicates that $j: V_\lambda \prec V_\lambda$. Sometimes we write I3(j, λ) to underline the role of λ , etc.

Another way to reach the apogee of the large cardinal hierarchy is via the usual template asserting the existence of an elementary embedding $j: V \prec M \subseteq V$ with M resembling V .

Definition 2.1. A cardinal κ is:

- *superstrong* iff there exists $j: V \prec M$ such that $\text{crt}(j) = \kappa$ and $V_{\kappa_1} \subseteq M$, where κ_1 is the second element of the critical sequence of j ;
- *n -superstrong* iff there exists $j: V \prec M$ such that $\text{crt}(j) = \kappa$ and $V_{\kappa_n} \subseteq M$, where κ_n is the $n + 1$ -th element of the critical sequence of j ;
- *ω -superstrong* iff there exists $j: V \prec M$ such that $\text{crt}(j) = \kappa$ and $V_\lambda \subseteq M$, where λ is the supremum of the critical sequence of j .

If κ is ω -superstrong as witnessed by j, λ , then $j(\lambda) = \lambda$, and therefore I3 holds. Moreover, like other large cardinals, it can be formulated as the existence of an extender (see [?]), in this case a (κ, λ) -extender E such that $V_\lambda \subseteq \text{Ult}(V, E)$, where λ is the supremum of the κ_n 's.

It is possible to pinpoint exactly how much ω -superstrongness is stronger than I3, but for this we need to clarify what we mean by "being stronger":

Definition 2.2. Let $\Phi(j, \lambda)$ and $\Psi(j, \lambda)$ be two large cardinal properties as above. Then

- Φ implies Ψ iff $\text{ZFC} \vdash \Phi(j, \lambda) \rightarrow \Psi(j, \lambda)$;
- Φ strictly implies Ψ iff Φ implies Ψ and $\text{ZFC} \vdash \Phi(j, \lambda) \rightarrow \exists \lambda' \exists j' (\Psi(j', \lambda') \wedge \neg(\Phi(j', \lambda')))$;

- Φ strongly implies Ψ iff Φ implies Ψ and $\text{ZFC} \vdash \Phi(j, \lambda) \rightarrow \exists \lambda' < \lambda \exists j' \Psi(j', \lambda')$.

Note that strong implication yields strict implication: If Φ strongly implies Ψ , let λ be the smallest such that there is a j such that $\Phi(j, \lambda)$ holds. Then there are j' and $\lambda' < \lambda$ such that $\Psi(j', \lambda')$ holds, and since λ was the smallest for Φ , then $\Phi(j', \lambda')$ does not hold. The difference between “strict” versus “strong” is a consequence of the peculiar nature of rank-into-rank axioms. For weaker axioms, usually the focal cardinal is the critical point of an elementary embedding, and such cardinal is measurable. If, assuming some property Φ of κ , we can find a $\kappa' < \kappa$ that satisfies a property Ψ , the reasoning is as follows: let κ be the smallest cardinal that satisfies Φ ; find $\kappa' < \kappa$ that satisfies Ψ ; then κ' must not satisfy Φ ; then V_{κ} is a model of ZFC where no cardinal satisfy Φ , but some cardinal satisfy Ψ , so the consistency strength of Φ is stronger than that of Ψ . So actually strict implication and strong implication are the same.

In the rank-into-rank case, the focal cardinal is λ , as for the same λ there can be many different embeddings j and critical points of them (see discussion after Lemma ??). Therefore, given an embedding $j: V_{\lambda} \prec V_{\lambda}$ satisfying a property Φ , if we find another embedding $j': V_{\lambda} \prec V_{\lambda}$ with $\text{crt}(j') < \text{crt}(j)$ that satisfies a property Ψ , this would prove that the two properties are actually different, so it would prove that Φ strictly implies Ψ : Let j, λ satisfy Φ with $\text{crt}(j)$ the smallest possible. Then if we can find $j' : V_{\lambda} \prec V_{\lambda}$ with $\text{crt}(j') < \text{crt}(j)$, and so $\Phi(j', \lambda)$ does not hold. But this falls short of actually proving that the consistency strength of one is stronger than the other. As $V_{\lambda} \models \text{ZFC}$, if λ is least for Φ and Φ strongly implies Ψ , then V_{λ} is a model for Ψ and for $\neg\Phi$.

There is an alternative way to look at **I1**, as a higher-order **I3**-embedding. A second-order language is formally a two-sorted language, where one sort is interpreted by elements of the model, and the other sort is interpreted by subsets of the model (usually the first sort is lowercase and the second is uppercase). Both variables and parameters have two sorts. So, for example, if φ has no quantifiers, $V_{\lambda} \models \forall X \forall x \varphi(x, X, a, A)$ means “ $\forall X \subseteq V_{\lambda} \forall x \in V_{\lambda} \varphi(x, X, a, A)$ ”, where $a \in V_{\lambda}$ and $A \subseteq V_{\lambda}$. If a second-order formula does not have quantifiers with uppercase variables, then it is Δ_0^1 (or Σ_0^1 or Π_0^1). In a similar way to first-order formulæ, a formula is Σ_n^1 if there are n alternations of \exists and \forall quantifiers with uppercase variable, the first one being a \exists .

Now, as $V_{\lambda+1}$ is just $\mathcal{P}(V_{\lambda})$, if there is a $j: V_{\lambda} \prec V_{\lambda}$, for any $X \in V_{\lambda+1}$ we can define $j^+(X) = \bigcup_{\alpha < \lambda} j(X \cap V_{\alpha})$, so that j extends to $j^+: V_{\lambda+1} \rightarrow V_{\lambda+1}$. With this in mind, it makes sense now to ask whether j preserves second-order formulæ, i.e., whether $V_{\lambda} \models \varphi(a, A)$ iff $V_{\lambda} \models \varphi(j(a), j^+(A))$. If j preserves all second-order formulæ then, clearly, $j^+: V_{\lambda+1} \prec V_{\lambda+1}$. We say that j is a Σ_n^1 -elementary embedding if it preserves Σ_n^1 formulæ. If j is a Σ_n^1 -elementary embedding for any $n \in \omega$, then j^+ witnesses **I1**. The other direction also holds: if $j: V_{\lambda+1} \prec V_{\lambda+1}$, then the extension of $j \upharpoonright V_{\lambda}$ is j itself (see Lemma 3.4 in [?]), so every **I3**-embedding, if it can be extended to a **I1**-embedding, it can be extended in a unique way. It is a standard fact that

if j witnesses I3, then it is a Σ_0^1 -elementary embedding (this is a consequence of the more general Lemma ??). The general concept of Σ_n^1 -elementary embeddings appears first in the paper by Laver [?], building on Martin's seminal work.

In order to state and prove the results that follow, we introduce the following

Definition 2.3. $\mathcal{E}(\lambda)$ is the set of all $j: V_\lambda \prec V_\lambda$, and $\mathcal{E}_n(\lambda)$ is the set of all $j \in \mathcal{E}(\lambda)$ that are Σ_n^1 -elementary.

Therefore $\mathcal{E}(\lambda) = \mathcal{E}_0(\lambda) \supseteq \mathcal{E}_1(\lambda) \supseteq \mathcal{E}_2(\lambda) \supseteq \dots$, I3(λ) means $\mathcal{E}(\lambda) \neq \emptyset$, and I1(λ) is $\bigcap_n \mathcal{E}_n(\lambda) \neq \emptyset$.

Now we can characterize ω -superstrongness within this template:

Theorem 2.4 (Martin, [?]). *Any $j \in \mathcal{E}_1(\lambda)$ can be extended to an $i: V \prec M$ such that $V_\lambda \subseteq M$. Conversely, if $i: V \prec M$ is such that $V_\lambda \subseteq M$, with λ supremum of the critical sequence, then $i \upharpoonright V_\lambda \in \mathcal{E}_1(\lambda)$.*

In view of Theorem ??, we write I2(λ) for one of the two following equivalent statements:

- there is a $j: V \prec M$ such that $V_\lambda \subseteq M$, where λ is the supremum of the critical sequence of j ;
- $\mathcal{E}_1(\lambda) \neq \emptyset$.

Going back to Σ_n^1 -embeddings, there is a hierarchy of hypotheses of length $\omega + 1$, that starts with I3, I2, and then Σ_2^1 -, Σ_3^1 -, ... elementarity up to I1. Do they really form a proper hierarchy?

Theorem 2.5 (Laver, Martin, [?], [?]). *For every $n \in \omega$*

- (1) $\mathcal{E}_{2n+1}(\lambda) = \mathcal{E}_{2n+2}(\lambda)$;
- (2) if $j \in \mathcal{E}_{2n+1}(\lambda)$ and $\kappa = \text{crt}(j)$, then $C_n = \{\lambda' < \kappa : \mathcal{E}_{2n}(\lambda') \neq \emptyset\}$ is ω -club in κ .

Thus the existence of a Σ_{2n+1}^1 -embedding from V_λ to itself strongly implies the existence of a Σ_{2n}^1 -embedding from V_λ to itself.

The following result is Corollary 5.24 in [?]. Since the proof in that paper follows from a long-winded argument, for the reader's convenience we present a self-contained one.

Corollary 2.6. *I1(λ) strongly implies $\forall n \mathcal{E}_n(\lambda) \neq \emptyset$.*

Proof. If $j: V_{\lambda+1} \prec V_{\lambda+1}$ witnesses I1, then $j \upharpoonright V_\lambda$ is in $\mathcal{E}_{2n+1}(\lambda)$ for any $n \in \omega$. Let $C_n \subseteq \text{crt}(j)$ be as in Theorem ??; as $\text{crt}(j)$ is regular, $\bigcap_{n \in \omega} C_n \neq \emptyset$, therefore there is a $\lambda' < \text{crt}(j) < \lambda$ such that $\mathcal{E}_{2n}(\lambda') \neq \emptyset$ for any $n \in \omega$. \square

Figure ?? summarizes the situation until now, where all vertical arrows are strong implications.

$$\begin{array}{c}
\mathbf{I1}(\lambda) \leftrightarrow \bigcap_n \mathcal{E}_n(\lambda) \neq \emptyset \\
\downarrow \\
\forall n \mathcal{E}_n(\lambda) \neq \emptyset \\
\downarrow \\
\vdots \\
\downarrow \\
\mathcal{E}_4(\lambda) = \mathcal{E}_3(\lambda) \neq \emptyset \\
\downarrow \\
\mathbf{I2}(\lambda) \leftrightarrow \mathcal{E}_2(\lambda) = \mathcal{E}_1(\lambda) \neq \emptyset \\
\downarrow \\
\mathbf{I3}(\lambda) \leftrightarrow \mathcal{E}(\lambda) = \mathcal{E}_0(\lambda) \neq \emptyset
\end{array}$$

FIGURE 1.

3. ITERATIONS OF I3

If E is an extender in a transitive model M of ZFC, the iteration of length ν is a commutative system of transitive models, extenders, and elementary embeddings $\langle (M_\alpha, E_\alpha, j_{\beta,\alpha}) : \beta \leq \alpha \in \nu \rangle$ defined as follows:

- $M_0 = M$, $E_0 = E$, and $j_{0,0}$ is the identity on M ,
- $j_{\alpha,\alpha+1}: M_\alpha \prec \text{Ult}(M_\alpha, E_\alpha)^{M_\alpha} = M_{\alpha+1}$ is the ultrapower embedding, and $\text{Ult}(M_\alpha, E_\alpha)^{M_\alpha}$ is the ultrapower of M_α via E_α computed in M_α ,
- $E_{\alpha+1} = j_{\alpha,\alpha+1}(E_\alpha)$, and for $\beta \leq \alpha$ we set $j_{\beta,\alpha+1} = j_{\alpha,\alpha+1} \circ j_{\beta,\alpha}$,
- for γ limit M_γ is the direct limit of the M_α s for $\alpha < \gamma$.

If $\nu \subseteq M$ one verifies by induction on $\alpha \in \nu$ that M_α is well-founded—see the proof of Lemma 19.5 in [?]. In particular, if M is a proper class then ν can be replaced by Ord.

In 1978 Martin showed that the determinacy of $\mathbf{\Pi}_2^1$ sets followed from the existence of an elementary embedding $j: V_\lambda \prec V_\lambda$ that is iterable [?]. Before considering iterability, we need to define this concept rigorously. The crux of the matter is that j is a proper class of V_λ , and it cannot be a definable class by a generalization of Kunen's Theorem by Suzuki [?]. So the hypothetical $j_{1,2}$ cannot be calculated directly as a $j(j)$, since j is not an element of V_λ , but it cannot even be calculated indirectly, as j is not an ultrapower embedding via some extender. The idea is then to exploit the fact that λ has cofinality ω , defining the first iterate as $j^+(j)$, and at limit stages finding a way to define $j_{0,\omega}^+(j)$. In this process, however, we do not have the assurance that we can always prolong the iterate, and in fact we will see in Theorem ?? that there are cases where it cannot even reach the ω -limit.

It is worthwhile noticing that climbing the hierarchy of rank-into-rank axioms up to I0 and beyond, this problem disappears again. In fact: I2(λ) is witnessed by ω -superstrong embeddings defined by an extender, which is iterable by the argument at the beginning of this section; I1(λ) is taken care by Proposition ??

below; $I0(\lambda)$ can be witnessed by *proper embeddings* (see [?, ?, ?]) which are in fact iterable, as they are defined from a normal ultrafilter. It seems therefore that iterability is a problem peculiar to I3.

Proposition 3.1. *Every $j: V_{\lambda+1} \prec V_{\lambda+1}$ is iterable.*

Proof. Let $j: V_{\lambda+1} \prec V_{\lambda+1}$. In particular $j \upharpoonright V_\lambda = k$ is a Σ_1^1 -embedding, therefore it can be extended to an embedding $i: V \prec M$ with $V_\lambda \subseteq M$. Now, i is iterable, so $i_\alpha = i_{\alpha, \alpha+1}$, the α -th iterate, and $i_{0, \alpha}$, the limit embedding, are defined for any α ordinal, and M_α , the α -th model of the iteration, is well-founded. We want to define an iterate for j . Since $j(\lambda) = \lambda$, we have that $j(k): V_\lambda \prec V_\lambda$, so we can define $j_2 = j(k)^+$, i.e., the extension of $j(k)$ to $V_{\lambda+1}$, $j_3 = j_2(j_2 \upharpoonright V_\lambda)^+$ and so on. Now, as the extension of an I3-embedding is uniquely defined, $i_n \upharpoonright V_{\lambda+1} = j_n$, and the ω -th model of the iteration of j is $(V_{i_{0, \omega}(\lambda)+1})^{M_\omega}$ (see proof of Proposition ??). We can define therefore $j_\omega = j_{0, \omega}(k)^+$, with $j_{0, \omega}(k): (V_{i_{0, \omega}(\lambda)})^{M_\omega} \prec (V_{i_{0, \omega}(\lambda)})^{M_\omega}$ and $i_\omega \upharpoonright (V_{i_{0, \omega}(\lambda)+1})^{M_\omega} = j_\omega$. Finally, “ k is Σ_n^1 ” is a Σ_{n+2}^1 -formula, (see Lemma 2.1 in [?]) and therefore by elementarity of the $j_{0, \omega}$, $j_{0, \omega}(k)$ is a Σ_n^1 -embedding for any $n \in \omega$, and therefore an I1-embedding. So we can continue indefinitely the construction, and j is iterable. \square

The following lemmas provide the key construction for iterates of rank-into-rank embeddings:

Lemma 3.2. *Let $M, N \models \text{ZFC}$ be transitive sets, $\pi: M \prec N$, and $X \subseteq M$. If*

- π is cofinal, i.e., $\forall \beta \in N \cap \text{Ord} \exists \alpha \in M \cap \text{Ord} \pi(\alpha) > \beta$;
- X is amenable for M , i.e., $\forall \alpha \in M \cap \text{Ord} X \cap (V_\alpha)^M \in M$,

define $\pi^+(X) = \bigcup_{\alpha \in M \cap \text{Ord}} \pi(X \cap (V_\alpha)^M)$. Then $\pi^+(X)$ is amenable for N and $\pi: (M, X) \prec_1 (N, \pi^+(X))$.

Proof. Well known. A simple induction proves that $\pi: (M, X) \prec_0 (N, \pi^+(X))$, and every cofinal Σ_0 -embedding is a Σ_1 -embedding. \square

The next result shows that with an assumption on the cofinality of $M \cap \text{Ord}$ we can have full elementarity:

Lemma 3.3. *Let $M, N \models \text{ZFC}$ be transitive sets. Suppose $\pi: M \prec N$ is cofinal, that $\gamma = \text{cof}(M \cap \text{Ord}) \in M$, and that $M^{<\gamma} \subseteq M$. Then $\pi: (M, X) \prec (N, \pi^+(X))$ for any $X \subseteq M$ amenable in M .*

Proof. We prove by induction on n that $\pi: (M, X) \prec_n (N, \pi^+(X))$ for any $X \subseteq M$ amenable in M . The case $n = 1$ holds by Lemma ??, so we may assume that the result holds for some $n \geq 1$ towards proving the result for $n + 1$.

Fix $F = \langle \kappa_\alpha : \alpha < \gamma \rangle$ cofinal in $M \cap \text{Ord}$. As $M^{<\gamma} \subseteq M$ then F is amenable for M . Let Fml_n be the set of (codes for) Σ_n -formulas in the language of set theory augmented with a 1-ary predicate \dot{X} . Let

$$B = \{(\exists y \psi(x, y), \alpha) \in \text{Fml}_n \times \gamma : \psi \text{ is } \Pi_{n-1} \wedge (M, X) \models \exists y \in V_{\kappa_\alpha} \psi(x, y)\}.$$

Note that B is amenable for M . If $\varphi(x) \in \text{Fml}_n$ is $\exists y\psi(x, y)$, then

$$\begin{aligned} \forall x(\varphi(x) \rightarrow \exists \alpha < \gamma(\varphi(x), \alpha) \in \mathring{B}) \\ \wedge \forall x \forall \kappa \forall \alpha < \gamma((\varphi(x), \alpha) \in \mathring{B} \wedge (\alpha, \kappa) \in \mathring{F} \rightarrow \exists y \in V_\kappa \psi(x, y)) \end{aligned}$$

is a Π_n -formula $\Psi_{\varphi(x)}$ in the language of set theory augmented with predicates $\mathring{X}, \mathring{F}, \mathring{B}$ that holds true in (M, X, F, B) . (The assumption that the cofinality of $M \cap \text{Ord}$ is singular is used to bound the quantifier $\exists \alpha < \gamma$ so that $\Psi_{\varphi(x)}$ is indeed a Π_1 formula when $n = 1$.) The formulas $\Psi_{\varphi(x)}$ with $\varphi(x) \in \text{Fml}_n$ describe that B is exactly as defined, therefore we can say that the Π_n -theory of (M, X, F, B) “knows” the definition of B . Since

$$\pi: (M, X, F, B) \prec_n (N, \pi^+(X), \pi^+(F), \pi^+(B)),$$

then $\pi^+(B)$ is as expected, i.e.,

$$\begin{aligned} \pi^+(B) = \{(\exists y\psi(x, y), \alpha) \in \text{Fml}_n \times \pi(\gamma) : \psi \text{ is } \Pi_{n-1} \\ \wedge (N, \pi^+(X)) \models \exists y \in V_{\pi(F(\alpha))} \psi(x, y)\}, \end{aligned}$$

so $(N, \pi^+(X), \pi^+(B)) \models \exists \alpha < \pi(\gamma) (\varphi(x), \alpha) \in \mathring{B}$ iff $(N, \pi^+(X)) \models \varphi(x)$.

We are now ready to show that π preserves all Π_{n+1} formulas and hence it is Σ_{n+1} -elementary. If φ is a Σ_n formula then

$$\begin{aligned} (M, X) \models \forall x \varphi(x) &\leftrightarrow (M, X, B) \models \forall x \exists \alpha < \gamma (\varphi(x), \alpha) \in \mathring{B} \\ &\leftrightarrow (N, \pi^+(X), \pi^+(B)) \models \forall x \exists \alpha < \pi(\gamma) (\varphi(x), \alpha) \in \mathring{B} \\ &\leftrightarrow (N, \pi^+(X)) \models \forall x \varphi(x) \end{aligned}$$

where the second equivalence follows from $\pi: (M, X, B) \prec_1 (N, \pi^+(X), \pi^+(B))$ by Lemma ?? and therefore it preserves Π_1 formulas. \square

When $M = N = V_\lambda$ then $\text{cof}(M \cap \text{Ord}) = \omega$ so that the hypothesis $M^{<\gamma} \subseteq M$ in the statement of Lemma ?? holds automatically. Lemma ?? for $M = N = V_\lambda$ appears in several places without proof (e.g., [?, ?]), but only in [?] there is a proof of that. Unfortunately, as it is written in [?] there is a small gap: the proof is based on defining j^+ first on Skolem functions, but it is not clear why $j^+(f)$ should be total for any f Skolem function. This problem is solved as Claim 3.7 in [?]. The proof above is instead an argument by Woodin found on MathOverflow [?].

Lemma ?? shows how to extend an elementary embedding to amenable subsets. A simple calculation proves that such extensions behave as expected between each other:

Lemma 3.4. *Let M_1, M_2, N_1, N_2 be transitive sets and models of ZFC.*

- (1) *If $j: M_1 \prec N_1$ and $\pi: N_1 \prec N_2$ are cofinal, and $X \subseteq M_1$ is amenable for M_1 , then $\pi^+(j^+(X)) = (\pi^+(j))^+(\pi^+(X))$.*

- (2) If $\pi_1: M_1 \prec N_1$, $\pi_2: M_2 \prec N_2$, $j_1: M_1 \prec M_2$, and $j_2: N_1 \prec N_2$ are cofinal, then $\pi_2^+ \circ j_1^+ = j_2^+ \circ \pi_1^+$ on the sets amenable for M_1 .

$$\begin{array}{ccc} N_1 & \xrightarrow{j_2} & N_2 \\ \pi_1 \uparrow & & \uparrow \pi_2 \\ M_1 & \xrightarrow{j_1} & M_2 \end{array}$$

The next result is folklore.

Lemma 3.5. *If $j, k: V_\lambda \prec V_\lambda$, then $j^+(k): V_\lambda \prec V_\lambda$.*

Proof. Let Fml be the set of all (codes of) first order formulas in the language of set theory. Note that “ k is an elementary embedding from V_λ to itself” amounts to say that $V_\lambda \models \Upsilon_{\varphi(x)}$ for every $\varphi(x) \in \text{Fml}$, where $\Upsilon_{\varphi(x)}$ is

$$\forall x(\varphi(x) \rightarrow \exists y((x, y) \in \mathring{k} \wedge \varphi(y)))$$

with \mathring{k} a binary predicate for k . Since $j: (V_\lambda, k) \rightarrow (V_\lambda, j^+(k))$ by Lemma ??, it follows that $j^+(k): V_\lambda \prec V_\lambda$. \square

In particular $j^+(j) = j_1$ will be an embedding with critical point $\text{crt}(j^+(j)) = j(\text{crt}(j)) = \kappa_1$. Letting $j_0 = j$ and $j_{n+1} = j^+(j_n)$ one proves by induction on n that $j_{n+1} = j_n^+(j_n)$. Note that $\text{crt}(j_n) = \kappa_n$. By induction on n it follows that $j_n(\kappa_m) = \kappa_{m+1}$ when $m \geq n$:

$$j_n(\kappa_m) = j_{n-1}(j_{n-1})(j_{n-1}(\kappa_{m-1})) = j_{n-1}(j_{n-1}(\kappa_{m-1})) = j_{n-1}(\kappa_m) = \kappa_{m+1}.$$

Let M_ω be the direct limit of the system $\langle (V_\lambda, j_{n,m}) : n, m \in \omega, n < m \rangle$, where $j_{n,n+1} = j_n$ and $j_{n,m} = j_m \circ j_{m-1} \circ \dots \circ j_n$. If M_ω is well-founded, then it is defined $j_{n,\omega}: V_\lambda \prec M_\omega$ for any $n \in \omega$. Now, $j_{0,\omega}$ is cofinal: Let $\alpha \in M_\omega$. Then there exist $n \in \omega$ and $\beta \in \lambda$ such that $\alpha = j_{n,\omega}(\beta)$. Let m be such that $\beta < \kappa_m$ and $m > n$. Then $j_{0,n}(\kappa_{m-n}) = \kappa_m > \beta$, and

$$j_{0,\omega}(\kappa_{m-n}) = j_{n,\omega}(j_{0,n}(\kappa_{m-n})) > j_{n,\omega}(\beta) = \alpha.$$

Therefore we can define $j_\omega = j_{0,\omega}^+(j): M_\omega \prec M_\omega$, and then $j_{\omega+1} = (j_\omega)^+(j_\omega): M_\omega \prec M_\omega$, and so on. At each limit point we ask whether the direct limit is well-founded, and if so we continue, otherwise we stop.

Note that, differently than in the case $j: V \prec M$, the model M_α is the same as $M_{\alpha+1}$, so they are either both well-founded or not. In other words, the construction can stop only at limit ordinals. We say that j is α -iterable, then, if the construction does not stop at the $\omega \cdot \alpha$ -th step, i.e, if $M_{\omega \cdot \alpha}$ is well-founded, and $<\alpha$ -iterable if $M_{\omega \cdot \beta}$ is well-founded for any $\beta < \alpha$. As usual, we identify M_β with its transitive collapse, when well-founded. We write $\mathbf{I3}_\alpha$ to indicate the existence of an α -iterable embedding from V_λ to itself, and $\mathbf{I3}_{<\alpha}$ for the existence of a $<\alpha$ -iterable embedding. We say that j is iterable, and we indicate the relative hypothesis with $\mathbf{I3}_\infty$, if it is α -iterable for any α ordinal.

If j is 1-iterable, as $j_m(\kappa_n) = \kappa_n$ for any $m > n$, we have that $\text{crt}(j_{n,\omega}) = \kappa_n$, so if $x \in V_{\kappa_n}$, $j_{n+1,\omega}(x) = x \in M_\omega$. This means that $j_{0,\omega}(\kappa_0) = \lambda$ and $V_\lambda \subseteq M_\omega$, so M_ω is actually “taller” than V_λ . As $(V_\lambda)^{M_\omega} = V_\lambda$, and this implies that $V_\lambda \in M_\omega$. In the same way, if j is 2-iterable then $M_\omega \in M_{\omega \cdot 2}$, and, more generally, if j is β -iterable then $M_{\omega \cdot \alpha} \in M_{\omega \cdot \beta}$ for any $\alpha < \beta$.

Suppose E is an extender in a transitive model M and let M_α denote the α th model of the iteration. It is a standard result in inner model theory that if M_α is well-founded for every $\alpha < \omega_1$, then every M_α is well-founded. This holds also in our situation.

Proposition 3.6. *For $j: V_\lambda \rightarrow V_\lambda$, the following are equivalent:*

- $\mathbf{I3}_\infty(j)$, that is M_α is well-founded for any α ordinal, i.e., j is iterable,
- $\mathbf{I3}_{\omega_1}(j)$, that is M_{ω_1} is well-founded, i.e., j is ω_1 -iterable,
- $\forall \beta < \omega_1 \mathbf{I3}_\beta(j)$, that is M_β is well-founded for any $\beta < \omega_1$, i.e., j is $< \omega_1$ -iterable.

Proof. Only one direction is not obvious. So suppose that $\forall \beta < \omega_1 \mathbf{I3}_\beta(j)$ and that there exists θ such that M_θ is ill-founded. Without loss of generality, we can assume that θ is least, limit and $\geq \omega_1$. We will prove that this leads to a contradiction.

Pick α large enough so that $\langle M_\nu : \nu < \theta \rangle \in V_\alpha$ together with witnesses of the ill-foundedness of M_θ . Let $\pi: \mathcal{P} \rightarrow V_\alpha$ the inverse of the collapse such that j , $\langle M_\nu : \nu < \theta \rangle$, V_λ , θ , κ_n for all $n \in \omega$ and the witnesses of the ill-foundedness of M_θ are all in the range of π , with \mathcal{P} countable. Let $\bar{M}_0 = \pi^{-1}(V_\lambda)$, $\bar{j}_0 = \pi^{-1}(j)$ and $\bar{\theta} = \pi^{-1}(\theta)$, and let $\langle \bar{M}_\nu : \nu \leq \bar{\theta} \rangle$ be the iteration of (\bar{M}_0, \bar{j}_0) . Then $\bar{M}_{\bar{\theta}}$ is ill-founded in \mathcal{P} .

We want all the models of the iterates of j and \bar{j} to satisfy the hypothesis of Lemma ??, so to have singular height. But note that, as $\langle \kappa_n : n \in \omega \rangle$ is cofinal in V_λ , for any $\nu < \theta$ we have that $\langle j_{0,\nu}(\kappa_n) : n \in \omega \rangle$ is cofinal in M_ν , as $j_{0,\nu}$ is a Σ_0^1 -elementary embedding by Lemma ?? and $\langle j_{0,\nu}(\kappa_n) : n \in \omega \rangle = j_{0,\nu}^+(\langle \kappa_n : n \in \omega \rangle)$. Let $\mu_n = \pi^{-1}(\kappa_n)$. Then also $\langle \mu_n : n \in \omega \rangle$ is cofinal in $\bar{M}_0 \cap \text{Ord}$ and for every $\nu < \bar{\theta}$ $\langle \bar{j}_{0,\nu}(\mu_n) : n \in \omega \rangle$ is cofinal in $\bar{M}_\nu \cap \text{Ord}$. So $\bar{M}_\nu \cap \text{Ord}$ has cofinality ω , for every $\nu \leq \bar{\theta}$. As $\bar{\theta}$ is countable, all the M_α and \bar{M}_α are well-founded for $\beta < \bar{\theta}$ for case assumption.

We build by induction π_ν , for every $\nu \leq \bar{\theta}$, such that:

- (1) $\pi_\nu: \bar{M}_\nu \prec M_\nu$ and it is cofinal;
- (2) $\pi_\nu \circ \bar{j}_{\delta,\nu} = j_{\delta,\nu} \circ \pi_\delta$ for every $\delta < \nu$;
- (3) $\pi_\nu^+(\bar{j}_\nu) = j_\nu$.

$$\begin{array}{ccccccc}
& & & \overset{j_{\delta,\nu}}{\curvearrowright} & & & \\
V_\lambda & \xrightarrow{j} & V_\lambda & \cdots & M_\delta & \cdots & M_\nu \xrightarrow{j_\nu} M_\nu & \cdots & M_{\bar{\theta}} & \cdots & M_\theta \\
\pi_0 \uparrow & \pi_1 \uparrow & & & \pi_\delta \uparrow & & \pi_\nu \uparrow & \pi_{\nu+1} \uparrow & \pi_{\bar{\theta}} \uparrow & & \nearrow \pi \\
\bar{M}_0 & \xrightarrow{j_0} & \bar{M}_0 & \cdots & \bar{M}_\delta & \cdots & \bar{M}_\nu \xrightarrow{j_\nu} \bar{M}_\nu & \cdots & \bar{M}_{\bar{\theta}} & & \\
& & & \underset{\bar{j}_{\delta,\nu}}{\curvearrowright} & & & & & & &
\end{array}$$

If this can be achieved, we easily reach a contradiction: In M_θ there is a sequence that witnesses that M_θ is ill-founded, and by construction of π such witnesses are in the range of π , therefore also $\bar{M}_{\bar{\theta}}$ is ill-founded. But then, by elementarity via $\pi_{\bar{\theta}}$, also $M_{\bar{\theta}}$ is ill-founded, a contradiction since $\bar{\theta}$ is countable and we assumed that all the M_α with α countable are well-founded.

For $\nu = 0$, let $\pi_0 = \pi \upharpoonright \bar{M}_0$. Then, of course, $\pi_0: \bar{M}_0 \prec V_\lambda$. It is cofinal because for all n , $\kappa_n \in \text{ran}(\pi_0)$. Note that by elementarity $\bar{M}_0 = (V_\eta)^{\mathcal{P}}$ for some η ; so if $X \subseteq M_0$ and $X \in \mathcal{P}$, then X is amenable in \bar{M}_0 , and $\pi_0^+(X) = \pi(X)$, therefore $\pi_0^+(\bar{j}_0) = \pi(\bar{j}_0) = j$.

Let ν be a limit ordinal. Let $x \in \bar{M}_\nu$. Then there exist $\delta < \nu$ and $y \in \bar{M}_\delta$ such that $x = \bar{j}_{\delta,\nu}(y)$. We define then $\pi_\nu(x) = j_{\delta,\nu}(\pi_\delta(y))$. It is easy to see that it is elementary and well defined. For any $n \in \omega$, then $\pi_\nu(\bar{j}_{0,\nu}(\mu_n)) = j_{0,\nu}(\pi_0(\mu_n)) = j_{0,\nu}(\kappa_n)$, therefore π_ν is cofinal. Let $\delta < \nu$ and let $x \in \bar{M}_\delta$. Then $\pi_\nu(\bar{j}_{\delta,\nu}(x)) = j_{\delta,\nu}(\pi_\delta(x))$ by definition of π_ν , therefore (2) holds. Also, $\pi_\nu^+(\bar{j}_\nu) = \pi_\nu^+(\bar{j}_{0,\nu}(\bar{j}_0)) = j_{0,\nu}^+(\pi_0^+(\bar{j}_0)) = j_{0,\nu}^+(\bar{j}_0) = j_\nu$, so π_ν is as desired.

Finally, let $\nu = \mu + 1$. Then define $\pi_\nu = \pi_\mu$. (1) is immediate. For (2), we prove it for $\delta = \mu$, and the rest is by easy induction. Note that $\bar{j}_{\mu,\nu} = \bar{j}_\mu$ and $j_{\mu,\nu} = j_\mu$. Let $x \in \bar{M}_\nu$. Then

$$\pi_\nu(\bar{j}_{\mu,\nu}(x)) = \pi_\nu(\bar{j}_\mu(x)) = \pi_\mu(\bar{j}_\mu(x)) = \pi_\mu^+(\bar{j}_\mu)(\pi_\mu(x)) = j_\mu(\pi_\mu(x)) = j_{\mu,\nu}(\pi_\mu(x)).$$

For (3), $\pi_\nu^+(\bar{j}_\nu) = \pi_\nu^+(\bar{j}_\mu(\bar{j}_\mu)) = \pi_\mu^+(\bar{j}_\mu)(\pi_\mu^+(\bar{j}_\mu)) = j_\mu(j_\mu) = j_\nu$.

But now there is $\pi_{\bar{\theta}}: \bar{M}_{\bar{\theta}} \prec M_{\bar{\theta}}$, with $M_{\bar{\theta}}$ well-founded because $\bar{\theta}$ is countable and $\bar{M}_{\bar{\theta}}$ ill-founded in \mathcal{P} , and therefore in V , contradiction. \square

We say that $j: V_\lambda \prec V_\lambda$ is iterable if j satisfies one of the three equivalent conditions of Proposition ??, and we denote with $\langle (M_\alpha, j_\alpha) : \alpha \in \text{Ord} \rangle$ the iteration of (V_λ, j) .

Lemma 3.7. *Suppose N is a transitive model of ZFC and that $N \models E$ is a (κ, λ) -extender witnessing κ is ω -superstrong. Let $i_E: N \rightarrow \text{Ult}(N, E)^N = N'$ be the ultrapower embedding, let $\kappa' = i_E(\kappa)$, and let $E' = i_E(E)$ so that E' is a (κ', λ) -extender witnessing in N' that κ' is ω -superstrong. Then $j^+(j) = i_{E'} \upharpoonright V_\lambda$, where $j = i_E \upharpoonright V_\lambda$.*

Proof. The result follows from the fact that a (κ, λ) -extender F is completely determined by $i_F \upharpoonright V_\lambda$. \square

The next result shows $\mathbf{I3}_\infty$ sits between $\mathbf{I3}$ and $\mathbf{I2}$.

Proposition 3.8. $\mathbf{I2}(j)$ implies $\mathbf{I3}_\infty(j)$, for all $j: V_\lambda \rightarrow V_\lambda$.

Proof. Suppose $\mathbf{I2}(j)$ and let $\kappa = \text{crt}(j)$. By Martin's Theorem ?? let E be a (κ, λ) -extender such that $i: V \prec \text{Ult}(V, E) \supseteq V_\lambda$ and $i \upharpoonright V_\lambda = j$. Let $\langle (N_\alpha, E_\alpha, i_{\beta, \alpha}) : \beta \leq \alpha \in \text{Ord} \rangle$ be the iteration of V via E . As argued at the beginning of this section, every N_α is well-founded. It is enough to show that $\langle (M_\alpha, j_\alpha) : \alpha \in \text{Ord} \rangle$ is the iteration of (V_λ, j) , where

$$\lambda_\alpha = i_{0, \alpha}(\lambda), \quad M_\alpha = N_\alpha \cap V_{\lambda_\alpha}, \quad j_\alpha = i_{\alpha, \alpha+1} \upharpoonright M_\alpha.$$

This boils-down to show that $j_\alpha^+(j_\alpha) = j_{\alpha+1}$, which follows from Lemma ?? and an easy induction on α . \square

Therefore all the iterable embeddings are in consistency strength between $\mathbf{I3}$ and $\mathbf{I2}$. Are they strictly or strongly between them? The tools developed in [?] by Laver will be essential to prove that $\mathbf{I3}_\infty$ is strongly between $\mathbf{I3}$ and $\mathbf{I2}$:

Proposition 3.9 (Laver, Square root of elementary embeddings, [?]). *Let $j: V_\lambda \prec V_\lambda$ and let $\kappa = \text{crt}(j)$.*

- (1) *If j is Σ_1^1 -elementary (so $\mathbf{I2}(j)$) and $\beta < \kappa$, then there exists $k: V_\lambda \prec V_\lambda$ such that $k^+(k) = j$ and $\beta < \text{crt}(k) < \kappa$.*
- (2) *If j is Σ_{n+2}^1 , then for any $B \subseteq V_\lambda$ there exists $\lambda' < \kappa$ and $J: V_{\lambda'} \prec V_\lambda$ that is Σ_n^1 such that $B \in \text{ran}(J^+)$.*

Proposition ??(1) is enough to prove that $\mathbf{I2}$ strictly implies $\mathbf{I3}_\infty$: Let $j: V_\lambda \prec V_\lambda$ be a Σ_1^1 -elementary embedding with least critical point, so that j witnesses $\mathbf{I2}$. Then by Proposition ??(1) there is a $k: V_\lambda \prec V_\lambda$ such that $\text{crt}(j) < \text{crt}(k)$. Since j was chosen with least critical point, k cannot be Σ_1^1 . But k is iterable, as $k_1 = j$ and therefore $k_{n+1} = j_n$ and their limit iterations are the same. But we can do better:

Proposition 3.10. *Let $j: V_\lambda \prec V_\lambda$ be a Σ_1^1 -elementary embedding. Then there is a $\lambda' < \lambda$ and a $k: V_{\lambda'} \prec V_\lambda$ that is iterable. In other words, $\mathbf{I2}$ strongly implies $\mathbf{I3}_\infty$.*

Proof. Let $j: V_\lambda \prec V_\lambda$ be Σ_1^1 . Use Proposition ??(2) above with $B = j$, and let k with $\lambda' < \lambda$ and $J^+(k) = j$ (remember that being Σ_1^1 is the same as being Σ_2^1). Note that “ $j: V_\lambda \prec V_\lambda$ ” is Δ_1^1 in V_λ , so $k: V_{\lambda'} \prec V_\lambda$. Let $\langle (\bar{M}_\alpha, k_\alpha) : \alpha < \gamma \rangle$ be an iteration of k of length $\gamma < \omega_1$, and let $\langle (M_\alpha, j_\alpha) : \alpha < \omega_1 \rangle$ be the iteration of length ω_1 of j . Now the proof is the same as in Proposition ??: define for any $\nu \leq \gamma$ $J_\nu: \bar{M}_\nu \prec M_\nu$, cofinal, such that $J_\nu \circ k_{\alpha, \nu} = j_{\alpha, \nu} \circ J_\alpha$ for any $\alpha < \nu$ and

$$J_\nu^+(k_\nu) = j_\nu.$$

$$\begin{array}{ccccccc}
& & & & \xrightarrow{j_{\alpha,\nu}} & & \\
V_\lambda & \xrightarrow{j} & V_\lambda & \cdots & M_\alpha & \cdots & M_\nu \xrightarrow{j_\nu} M_\nu & & M_\gamma \\
J_0 \uparrow & J_1 \uparrow & & & J_\alpha \uparrow & & J_\nu \uparrow & J_{\nu+1} \uparrow & J_\gamma \uparrow \\
V_{\lambda'} & \xrightarrow{k} & V_{\lambda'} & \cdots & \bar{M}_\alpha & \cdots & \bar{M}_\nu \xrightarrow{k_\nu} \bar{M}_\nu & \cdots & \bar{M}_\gamma \\
& & & & \xleftarrow{k_{\alpha,\nu}} & & & &
\end{array}$$

If \bar{M}_γ were ill-founded, then because of the elementarity of $J_\gamma: \bar{M}_\gamma \prec M_\gamma$ also M_γ would be ill-founded, but j is iterable, so \bar{M}_γ is well-founded. This holds for any $\gamma < \omega_1$, and therefore by Proposition ?? k is iterable. \square

Therefore the iterability hypotheses are not only between I3 and I2, but strongly below I2. But this is where Laver's tools stop, as they are too coarse to actually be useful in investigating gaps under I3 $_\infty$. For this, we need tools that are partially borrowed from the "classic" iterability.

4. THE ITERABILITY HIERARCHY

Recall from Definition ?? that $\mathcal{E}(\lambda)$ is the set of all $j: V_\lambda \prec V_\lambda$. Then

$$\mathcal{W}_\alpha(\lambda) = \{j \in \mathcal{E}(\lambda) : M_{w_\alpha} \text{ is well-founded}\}$$

is the set of all $j: V_\lambda \prec V_\lambda$ that are α -iterable. Therefore

$$\mathcal{E}(\lambda) = \mathcal{W}_0(\lambda) \supseteq \mathcal{W}_1(\lambda) \supseteq \cdots \supseteq \mathcal{W}_\alpha(\lambda) \supseteq \dots$$

With this notations Propositions ?? and ?? become

$$\mathcal{E}_1(\lambda) \subset \mathcal{W}_{\omega_1}(\lambda) = \bigcap_{\alpha < \omega_1} \mathcal{W}_\alpha(\lambda) = \mathcal{W}_\beta(\lambda)$$

for any $\beta \geq \omega_1$. We will prove that there is a strong hierarchy below I3 $_\infty$:

Theorem 4.1.

- If $\alpha < \omega_1$, then I3 $_{\alpha+1}(\lambda)$ strongly implies I3 $_\alpha(\lambda)$, i.e., if $j \in \mathcal{W}_{\alpha+1}(\lambda)$ then there is a $\lambda' < \lambda$ and an $e \in \mathcal{W}_\alpha(\lambda')$.
- If $\nu < \omega_1$ is limit, then I3 $_\nu(\lambda)$ strongly implies I3 $_{<\nu}(\lambda)$, i.e., if $j \in \mathcal{W}_\nu(\lambda)$, then there are a $\lambda' < \lambda$ and an $e \in \bigcap_{\alpha < \nu} \mathcal{W}_\alpha(\lambda')$.
- If $\nu \leq \omega_1$ is limit, then I3 $_{<\nu}(\lambda)$ strongly implies $\forall \alpha < \nu$ I3 $_\alpha(\lambda)$, i.e., if $j \in \bigcap_{\alpha < \nu} \mathcal{W}_\alpha(\lambda)$, then there is a $\lambda' < \lambda$ such that $\mathcal{W}_\alpha(\lambda') \neq \emptyset$ for all $\alpha < \nu$.

Moreover, for any instance of the above the λ' that witnesses the strong implication can be cofinally high under λ , so for any $\eta < \lambda$ there exists $\eta < \lambda' < \lambda$ that witnesses the strong implication. We will call this cofinal strong implication.

The hierarchy just above I3 will therefore look like this, where every vertical arrow is a cofinal strong implication:

$$\begin{array}{c}
\vdots \\
\downarrow \\
\mathcal{W}_{\omega+1}(\lambda) \neq \emptyset \\
\downarrow \\
\mathcal{W}_\omega(\lambda) \neq \emptyset \\
\downarrow \\
\bigcap_n \mathcal{W}_n(\lambda) \neq \emptyset \\
\downarrow \\
\forall n \in \omega \mathcal{W}_n(\lambda) \neq \emptyset \\
\downarrow \\
\vdots \\
\downarrow \\
\mathcal{W}_2(\lambda) \neq \emptyset \\
\downarrow \\
\mathcal{W}_1(\lambda) \neq \emptyset \\
\downarrow \\
\mathcal{W}_0(\lambda) = \mathcal{E}(\lambda) \neq \emptyset
\end{array}$$

FIGURE 2.

Proof. We prove it gradually, as going-up the iterability hierarchy will introduce more and more problems. As usual, if $j: V_\lambda \prec V_\lambda$, then $\langle \kappa_n : n \in \omega \rangle$ is the critical sequence. When it exists, we call λ_α the height of M_α , i.e., $M_\alpha \cap \text{Ord}$. Therefore $\lambda_0 = \lambda$, $\lambda_\omega = M_\omega \cap \text{Ord}$ and so on. At the same time, we define $\kappa_\alpha = j_{0,\alpha}(\kappa_0)$. These two sequences overlap often, as $\kappa_{\alpha+\omega} = \lambda_\alpha$ for any α limit, but there is a slight difference at limit of limit stages: the κ_α sequence is in fact continuous at limit points, so for example $\kappa_{\omega \cdot \omega} = \sup_{n \in \omega} \kappa_{\omega \cdot n}$, while the λ_α sequence is discontinuous, for example $\lambda_{\omega \cdot \omega} > \sup_{n \in \omega} \lambda_{\omega \cdot n} = \kappa_{\omega \cdot \omega}$, as $\kappa_{\omega \cdot \omega} \in M_{\omega \cdot \omega}$. In a certain sense, the κ_α sequence is finer than the λ_α sequence and continuously completes it.

Claim 4.2. \mathfrak{I}_1 cofinally strongly implies \mathfrak{I}_3 , i.e., for any 1-iterable $j: V_\lambda \prec V_\lambda$ there exists $\lambda' < \lambda$ and $k: V_{\lambda'} \prec V_{\lambda'}$, and for any $\eta < \lambda$ we can find such λ' to be larger than η .

Proof. Suppose there exists $j: V_\lambda \prec V_\lambda$ that is 1-iterable and let M_ω be the ω -th iterated model, which is well-founded by assumption. Note that $j_{0,\omega}(\kappa_0) = \lambda$, therefore λ is regular in M_ω .

As M_ω is well-founded we build in M_ω a descriptive set-theoretic tree of approximations of an \mathfrak{I}_3 -embedding. So let T_1 be defined as:

$$T_1 = \{ \langle \gamma_0, (e^0, \gamma_1), \dots, (e^n, \gamma_{n+1}) \rangle : \forall i < n \ e^i \subseteq e^{i+1}, \forall i \leq n \ e^i : V_{\gamma_i} \prec V_{\gamma_{i+1}} \},$$

as defined in M_ω . Note that T_1 is a tree on V_λ and $V_\lambda \in M_\omega$, so $T_1 \in M_\omega$. Now, for any $n \in \omega$, $j \upharpoonright V_{\kappa_n} \in V_{\kappa_{n+1}} \subseteq V_\lambda \subseteq M_\omega$, therefore $\langle \kappa_0, (j \upharpoonright V_{\kappa_0}, \kappa_1), \dots \rangle$ is a

branch of T_1 in V . By absoluteness of well-foundedness there is therefore a branch of T_1 in M_ω . Let $\langle (e^n, \gamma_{n+1}) : n \in \omega \rangle$ together with γ_0 be a branch of T_1 in M_ω . Let $\gamma_\omega = \sup_{n \in \omega} \gamma_n$ and $e = \bigcup_{n \in \omega} e^n$. Then $e: V_{\gamma_\omega} \prec V_{\gamma_\omega}$, and $\gamma_\omega < \lambda$, since λ is regular in M_ω . So $M_\omega \models \exists \lambda' < \lambda \exists e: V_{\lambda'} \prec V_{\lambda'}$, but then this is true also in V . This proved that 1-iterability strongly implies I3.

Let now $\eta < \lambda$, and let $n \in \omega$ such that $\kappa_n > \eta$. Define a revised version of T_1 , adding the condition that $\gamma_0 > \eta$. Then the sequence $\langle \kappa_n, (j_n \upharpoonright V_{\kappa_n}, \kappa_{n+1}), \dots \rangle$ is a branch of the revised T_1 , so there is a branch also in V , and the γ_ω defined by the branch will be such that $\lambda' = \gamma_\omega > \gamma_0 > \eta$. \square

Claim 4.3. *I3₂ cofinally strongly implies I3₁, i.e., for any 2-iterable $j: V_\lambda \prec V_\lambda$ there exists $\lambda' < \lambda$ and a 1-iterable $k: V_{\lambda'} \prec V_{\lambda'}$, and for any $\eta < \lambda$ we can find such λ' to be larger than η .*

Proof. Suppose now that $j: V_\lambda \prec V_\lambda$ is 2-iterable, so $M_{\omega+\omega}$ is well-founded. Again, we are going to define a tree T_2 whose branches are going to bring us I3-embeddings. But we want more, since we want such embeddings to be 1-iterable. The solution is to build in T_2 at the same time a family of embeddings $\langle k_m : m \in \omega \rangle$ that commutes with the iterates of e , so that the ω -limit of such family will witness that the ω -limit of e is going to be well-founded.

Let $\langle \gamma_0, (e^0, \gamma_1), \dots, (e^n, \gamma_{n+1}), \dots \rangle$ be a branch of T_1 , in other words a sequence of approximations of an I3-embedding e . We can define then approximations also for the iterates of e : for example

$$e_1 \upharpoonright V_{\gamma_1} = e_1^1 = e^1(e^0), \quad e_1 \upharpoonright V_{\gamma_2} = e_1^2 = e^2(e^1), \quad e_2 \upharpoonright V_{\gamma_2} = e_2^2 = e_1^2(e_1^1),$$

where the subscript indicates the iteration number, the superscript the level up to which the iterate is approximate. Of course, e_1^0 and e_2^1 are the identity. In general, we define $e_{m+1}^{n+m+1} = e_m^{n+m+1}(e_m^{n+m})$ for any $n, m \in \omega$, where $e_0^n = e^n$. So if we know e up to V_{γ_n} , then we know any finite iterate up to V_{γ_n} .

Working in $M_{\omega+\omega}$ we define a tree T_2 on the set $M_\omega \in M_{\omega+\omega}$ in the following way:

(1) the nodes of T_2 are of the form

$$\langle (\gamma_0, \eta_0), (e^0, k_0^0, \gamma_1, \eta_1), (e^1, k_0^1, k_1^1, \gamma_2, \eta_2), \dots, (e^n, k_0^n, \dots, k_n^n, \gamma_{n+1}, \eta_{n+1}) \rangle;$$

(2) $e^0 \subseteq e^1 \subseteq \dots \subseteq e^n$, i.e., $\forall i < n \ e^i \subseteq e^{i+1}$;

(3) $e^i: V_{\gamma_i} \prec V_{\gamma_{i+1}}$ for any $i \leq n$;

(4) note that k_m^l is defined only for $m \leq l \leq n$, and we want $k_m^m \subseteq k_m^{m+1} \subseteq \dots \subseteq k_m^n$ for any $m \leq n$, i.e. $\forall i < n, \forall l < n - i \ k_i^{i+l} \subseteq k_i^{i+l+1}$;

(5) for any $m \leq l \leq n$, $k_m^l: V_{\gamma_l} \prec (V_{\eta_{l-m}})^{M_\omega}$;

(6) for any $m \leq l < n$, $k_{m+1}^{l+1} \circ e_m^l = k_m^l$.

Consider now

$$\langle (\kappa_0, \kappa_\omega), (j \upharpoonright V_{\kappa_0}, j_{0,\omega} \upharpoonright V_{\kappa_0}, \kappa_1, \kappa_{\omega+1}), (j \upharpoonright V_{\kappa_1}, j_{0,\omega} \upharpoonright V_{\kappa_1}, j_{1,\omega} \upharpoonright V_{\kappa_1}, \kappa_2, \kappa_{\omega+2}), \dots \rangle.$$

We want to prove that this sequence is a branch of T_2 . We need to prove that every element of the sequence is in M_ω . Clearly $j \upharpoonright V_{\kappa_n} \in V_\lambda \in M_\omega$ for any $n \in \omega$. Moreover, also $j_{m,l} \upharpoonright V_{\kappa_n} \in V_\lambda$ for any $m \leq l \in \omega$ and $n \in \omega$. But then for any $m \leq n \in \omega$

$$j_{m,\omega} \upharpoonright V_{\kappa_n} = j_{n+1,\omega} \circ j_{m,n+1} \upharpoonright V_{\kappa_n} = j_{n+1,\omega}(j_{m,n+1} \upharpoonright V_{\kappa_n}) \in M_\omega$$

because $\text{crt}(j_{n+1,\omega}) = \kappa_{n+1} > \kappa_n$. Points (2), (3), (4) are immediate. For point (5),

$$j_{m,\omega}(\kappa_n) = j_{m,\omega}(j_{0,m}(\kappa_{n-m})) = j_{0,\omega}(\kappa_{n-m}) = \kappa_{\omega+n-m} \quad \text{for any } m \leq n \in \omega,$$

so by elementarity $j_{m,\omega} \upharpoonright V_{\kappa_n} : V_{\kappa_n} \prec (V_{\kappa_{\omega+n-m}})^{M_\omega}$ for any $m \leq n \in \omega$. Finally, for point (6), notice that the iterate j_m is $j_{m,m+1}$ for any $m \in \omega$, so if $x \in V_{\kappa_n}$ for some $n \leq m$, $n \in \omega$, $j_{m+1,\omega}(j_m(x)) = j_{m,\omega}(x)$, and since $j_m \upharpoonright V_{\kappa_n} : V_{\kappa_n} \prec V_{\kappa_{n+1}}$, we have that

$$j_{m+1,\omega} \upharpoonright V_{\kappa_{n+1}} \circ j_m \upharpoonright V_{\kappa_n} = j_{m+1,\omega} \circ j_m \upharpoonright V_{\kappa_n} = j_{m,\omega} \upharpoonright V_{\kappa_n}.$$

So T_2 has a branch in V , and therefore it has a branch in $M_{\omega+\omega}$. Consider such a branch, and let $\gamma_\omega = \sup_{n \in \omega} \gamma_n$, $\eta_\omega = \sup_{n \in \omega} \eta_n$, $e = \bigcup_{n \in \omega} e^n$ and $k_m = \bigcup_{n \in \omega} k_m^n$ for any $m \in \omega$. Then, as before, $e : V_{\gamma_\omega} \prec V_{\gamma_\omega}$ and $k_m : V_{\gamma_\omega} \prec (V_{\eta_\omega})^{M_\omega}$ for any $m \in \omega$. Note that $M_\omega = (V_{\lambda_\omega})^{M_{\omega+\omega}}$, therefore $\lambda = \kappa_\omega$ is regular also in $M_{\omega+\omega}$ and $\gamma_\omega < \lambda$. Also $\lambda_\omega = \kappa_{\omega+\omega}$ is regular in $M_{\omega+\omega}$, so $\eta_\omega < \lambda_\omega$. Consider now N_ω the ω -iterated model of e . The picture is the following:

$$\begin{array}{c} \begin{array}{ccccccc} V_{\gamma_\omega} & \xrightarrow{e} & V_{\gamma_\omega} & \rightarrow \cdots & \longrightarrow & V_{\gamma_\omega} & \xrightarrow{e_m} & V_{\gamma_\omega} & \rightarrow \cdots & N_\omega \\ & & & & & & & & & \uparrow k_\omega \\ & & & & & & & & & (V_{\eta_\omega})^{M_\omega} \end{array} \\ \begin{array}{c} \nearrow k_m \\ \nearrow k_{m+1} \end{array} \end{array}$$

By the properties of the direct limit, as the family of k_m commutes with the iterates of e , there exists a $k_\omega : N_\omega \prec (V_{\eta_\omega})^{M_\omega}$. Then, by elementarity and well-foundedness of M_ω , also N_ω is well-founded, and e is a 1-iterable embedding below λ . Note that it is not necessary that $\gamma_\omega = \eta_\omega$, like in the branch in V .

Again, for any $\eta < \kappa_n < \lambda$ we can add to the definition of T_2 the condition $\gamma_0 > \eta$, and the branch generated by j_n instead of j will make the proof work, so that we will find a 1-iterable $e : V_{\lambda'} \prec V_{\lambda'}$ with $\eta < \gamma_0 < \gamma_\omega = \lambda'$. \square

Claim 4.4. \mathfrak{I}_3 cofinally strongly implies \mathfrak{I}_2 .

Proof. This adds another layer of complexity. The tree T_2 as calculated in $M_{\omega,3}$ would give a 1-iterable embedding, but our aim is to build a 2-iterable embedding. The strategy of adding more witnesses to well-foundedness (so further $k_{\omega+n}$) cannot work in the same way, as k_ω is defined on N_ω , and the initial segments of N_ω are known only when the whole e is known, so it is not possible to build at the same time small approximations of e and k_ω . The solution is instead to build a 1-iterable

embedding h via $(T_2)^{M_\omega}$, so that $k_\omega^+(e_\omega) = h$, and an iteration argument will show that this is enough to prove that e is 2-iterable.

Define T_3 in $M_{\omega \cdot 3}$ on $M_{\omega \cdot 2} \in M_{\omega \cdot 3}$ in the following way:

(1) the nodes of T_3 are of the form

$$\langle (\gamma_0, \eta_0, \delta_0), (e^0, k_0^0, h^0, g_0^0, \gamma_1, \eta_1, \delta_1), (e^1, k_0^1, k_1^1, h^1, g_0^1, g_1^1, \gamma_2, \eta_2, \delta_2), \dots, \\ (e^n, k_0^n, \dots, k_n^n, h^n, g_0^n, \dots, g_n^n, \gamma_{n+1}, \eta_{n+1}, \delta_{n+1}) \rangle;$$

(2) the sequence

$$\langle (\gamma_0, \eta_0), (e^0, k_0^0, \gamma_1, \eta_1), (e^1, k_0^1, k_1^1, \gamma_2, \eta_2), \dots, (e^n, k_0^n, \dots, k_n^n, \gamma_{n+1}, \eta_{n+1}) \rangle$$

is a node of T_2 ;

(3) the sequence

$$\langle (\eta_0, \delta_0), (h^0, g_0^0, \eta_1, \delta_1), (h^1, g_0^1, g_1^1, \eta_2, \delta_2), \dots, (h^n, g_0^n, \dots, g_n^n, \eta_{n+1}, \delta_{n+1}) \rangle$$

is a node of $(T_2)^{M_\omega}$, that is defined as T_2 but in M_ω and with all instances of M_ω replaced by $M_{\omega \cdot 2}$, so for example $h^n: (V_{\eta_n})^{M_\omega} \prec (V_{\eta_n})^{M_\omega}$ and $g_m^l: (V_{\eta_l})^{M_\omega} \prec (V_{\delta_{l-m}})^{M_{\omega \cdot 2}}$;

(4) for any $m \leq l < n$, $k_m^{l+1}(e_m^l) = h^{l-m}$.


As before, we can find a branch of T_3 in V in the natural way, i.e., assign $\gamma_n = \kappa_n$, $\eta_n = \kappa_{\omega+n}$, $\delta_n = \kappa_{\omega+\omega+n}$, $e^n = j \upharpoonright V_{\kappa_n}$, $k_m^n = j_{m,\omega} \upharpoonright V_{\kappa_n}$, $h^n = j_\omega \upharpoonright (V_{\kappa_{\omega+n}})^{M_\omega}$ and $g_m^n = j_{\omega,\omega+m} \upharpoonright (V_{\kappa_{\omega+n}})^{M_\omega}$. As j is 3-iterable and

$$j_{m,\omega}(j_m) = j_{m,\omega}(j_{0,m}(j)) = j_{0,\omega}(j) = j_\omega,$$

everything works.

Consider a branch of T_3 in $M_{\omega \cdot 3}$. Then, calling $\gamma_\omega = \sup_{n \in \omega} \gamma_n$, $\eta_\omega = \sup_{n \in \omega} \eta_n$, $\delta_\omega = \sup_{n \in \omega} \delta_n$, $e = \bigcup_{n \in \omega} e^n$, $k_m = \bigcup_{n \in \omega} k_m^n$ for any $m \in \omega$, $h = \bigcup_{n \in \omega} h^n$, $g_m = \bigcup_{n \in \omega} g_m^n$ for any $m \in \omega$, by the previous results we have that $e: V_{\gamma_\omega} \prec V_{\gamma_\omega}$, $k_m: V_{\eta_\omega} \prec (V_{\eta_\omega})^{M_\omega}$ for any $m \in \omega$, $h: (V_{\eta_\omega})^{M_\omega} \prec (V_{\eta_\omega})^{M_\omega}$ and $g_m: (V_{\eta_\omega})^{M_{\omega \cdot 2}} \prec (V_{\delta_\omega})^{M_{\omega \cdot 2}}$. As before, by the regularity of λ , λ_ω and $\lambda_{\omega \cdot 2}$ in $M_{\omega \cdot 3}$, $\gamma_\omega < \lambda$, $\eta_\omega < \lambda_\omega$ and $\delta_\omega < \lambda_{\omega \cdot 2}$. Moreover, e and h are 1-iterable, and point (4) of the definition of T_3 guarantees that $k_m^+(e_m) = h$ for any $m \in \omega$. We want to prove that $k_\omega^+(e_\omega) = h$. But $e_\omega = e_{m,\omega}^+(e_m)$ for all $m \in \omega$, so $k_\omega^+(e_{m,\omega}^+(e_m))$ is, by definition, $k_m^+(e_m)$.


Now, if $e_\omega = h$, then we have that the ω -iterate of e is 1-iterable, and so e is 2-iterable. Otherwise this is the picture, where $(V_{\eta_\omega})^{M_\omega} = \bar{N}_0$ and $(V_{\delta_\omega})^{M_{\omega \cdot 2}} = \bar{N}_0$:



By the usual reasoning, there exists $k_{\omega \cdot 2}: N_{\omega \cdot 2} \prec \bar{N}_\omega$. We know that $(V_{\delta_\omega})^{M_{\omega \cdot 2}}$ is well-founded because j is 3-iterable, but then by elementarity of g_ω we have that \bar{N}_ω is well-founded, and so by elementarity of $k_{\omega \cdot 2}$ also $N_{\omega \cdot 2}$ is well-founded.

Note that the usual remark on the cofinality of the possible λ' under λ still holds, with the same proof. \square

The techniques used for T_3 can be used now to define T_4, T_5 , and so on, therefore proving the first part of the theorem for α finite. For example this is the diagram generated by a branch of T_4 .




It is immediate now to see that if for all $n \in \omega$ there is a $j: V_\lambda \prec V_\lambda$ that has $M_{\omega \cdot n}$ well-founded, in particular there is a j that is $(n+1)$ -iterable, and this cofinally reflects the existence of an e that is n -iterable.

We postpone the proof that $<\omega$ -iterability cofinally strongly implies the existence of n -iterable embeddings for any $n \in \omega$ to the end of the proof of the theorem because it uses different techniques.

Claim 4.5. *If there exists $j: V_\lambda \prec V_\lambda$ that is ω -iterable (therefore $M_{\omega \cdot \omega}$ is well-founded), then there exists $\lambda' < \lambda$ and $e: V_{\lambda'} \prec V_{\lambda'}$ that is n -iterable for any $n \in \omega$. Moreover, the set of such λ' is cofinal in λ .*

Proof. We want to build in $M_{\omega \cdot \omega}$ a tree T_ω that “glues” together all the T_n trees, so that its branches will generate for any $n \in \omega$ a 1-iterable embedding e^n in $M_{\omega \cdot n}$ and

families of k_m^n that witness that e^n is 1-iterable and such that $(k_\omega^n)^+(e^n) = e^{n+1}$.



It is important to define T_ω so that it is in $M_{\omega \cdot \omega}$, so that the argument of the absoluteness of well-foundedness gives a branch that is in $M_{\omega \cdot \omega}$, and therefore bounded below λ , so T_ω should be a subset of a set in $M_{\omega \cdot \omega}$. Note that $(V_{\kappa_{\omega \cdot \omega}})^{M_{\omega \cdot \omega}} = \bigcup_{n \in \omega} M_{\omega \cdot n}$, because of the properties of the direct limit and because $\text{crt}(j_{\omega \cdot n, \omega \cdot \omega}) = \text{crt}(j_{\omega \cdot n}) = \kappa_{\omega \cdot n} = \lambda_{\omega \cdot (n-1)}$ for any $n > 0$.

The most immediate approach would be to build all the embeddings e^n and k_m^n at the same time, step by step. At every passage, each approximation of the e^n and k_m^n will be actually in $M_{\omega \cdot n}$, as we have seen in the previous claims, therefore they would be all in $M_{\omega \cdot \omega}$. This approach, however, has a problem: in the finite cases, each node is a finite sequence of finite sequences, so it suffices to know that all its elements are in $M_{\omega \cdot n}$ to say that the whole tree is contained in it. In a T_ω defined in such a way we have instead infinite sequences, for example the first step would be to decide the critical points of all the e^n , and it is not clear why this sequence should be in $M_{\omega \cdot \omega}$. If we restrict ourselves only to the sequences that are in $M_{\omega \cdot \omega}$ then we are in trouble, as we possibly cannot then build a branch in V : for example the “natural” branch generated by j will have the following first element:

$$\langle \kappa_0, \kappa_\omega, \kappa_{\omega+\omega}, \dots \rangle.$$

and this cannot be in $M_{\omega \cdot \omega}$, as the supremum of it is exactly $\kappa_{\omega \cdot \omega}$, that is regular in $M_{\omega \cdot \omega}$. The solution is to rearrange the pace at which the approximations are introduced, so that every sequence that appears in the revised tree T_ω is finite. We leave the details to the reader. Now the tree T_ω is on $M_{\omega \cdot \omega}$, and the proof is as before, cofinally strong implication included. \square

As we have sufficiently analyzed the case of successor and limit of successors, the techniques just presented are enough to go up the hierarchy of the countable ordinals. The following claim completes the proof:

Claim 4.6. *Let $\nu \leq \omega_1$ be a limit ordinal. If there exists a $j: V_\lambda \prec V_\lambda$ that is α -iterable for any $\alpha < \nu$, then there exists a $\lambda' < \lambda$ such that for any $\alpha < \nu$ there is an $e: V_{\lambda'} \prec V_{\lambda'}$ that is α -iterable.*

Proof. Let $j: V_\lambda \prec V_\lambda$ that is α -iterable for any $\alpha < \nu$. Then, by the claims above, for any $\alpha < \nu$ there is a $\lambda' < \lambda$ and an $e: V_{\lambda'} \prec V_{\lambda'}$ that is α -iterable. We want to prove that there is a single $\lambda' < \lambda$ that works for all the $\alpha < \nu$.

For any $\omega \cdot \alpha < \nu$ let

$$E_\alpha = \{\lambda' < \lambda : \exists k: V_{\lambda'} \prec V_{\lambda'} \text{ that is } \alpha\text{-iterable}\}.$$

Then $E_\alpha \neq \emptyset$, and we need to prove that $\bigcap_{\omega \cdot \alpha < \nu} E_\alpha \neq \emptyset$. Since all the strong implications above are cofinal, E_α is cofinal in λ . We want to prove that E_α is definable in V_λ using only α as a parameter. If $\lambda' \in E_\alpha$, let e witness that, and let $n \in \omega$ be such that $e \in V_{\kappa_n}$. Then N_ω , its ω -iterate, is the set of the $e_{m,\omega}(y)$ with $m \in \omega$ and $y \in V_{\lambda'}$, therefore $|N_\omega| = |V_{\lambda'}|$. By induction, $|N_{\omega \cdot \beta}| = |V_{\lambda'}|$ for any $\beta \leq \alpha$, and as $N_{\omega \cdot \beta}$ is transitive this means that $N_{\omega \cdot \beta} \in V_{\kappa_n}$. Therefore e is α -iterable iff $V_\lambda \models e$ is α -iterable, i.e., V_λ computes correctly the iterability of embeddings inside it. Therefore E_α is definable in V_λ , so $j^+(E_\alpha) = E_\alpha$, and if $\eta < \text{crt}(j)$ and β is the η -th element of E_α , then $j(\beta)$ is the η -th element of E_α , i.e., β , so $\beta < \text{crt}(j)$. But then the ordertype of E_α must be bigger than $\text{crt}(j)$, otherwise E_α would be all inside $\text{crt}(j)$ and not cofinal, and the first $\text{crt}(j)$ elements of E_α are smaller than $\text{crt}(j)$. This means that $\bigcup_{\alpha < \beta} (E_\alpha \cap \text{crt}(j)) \neq \emptyset$, as $\text{crt}(j)$ is regular, and the proposition is proved. As E_α is cofinal under κ_0 , by elementarity it is cofinal also under λ , therefore also cofinal strong implication is proved. \square

\square

REFERENCES

- [1] V. Dimonte, *Totally non-proper ordinals beyond $L(V_{\lambda+1})$* , Archive for Mathematical Logic **50** (2011) 565–584
- [2] V. Dimonte, *A Partially Non-Proper Ordinal*, Annals of Pure and Applied Logic **163** (2012) 1309–1321.
- [3] V. Dimonte, *I0 and rank-into-rank axioms*, Bollettino dell’Unione Matematica Italiana, (2017) <https://doi.org/10.1007/s40574-017-0136-y>
- [4] T. Jech, *Set theory. The third millennium edition, revised and expanded*. Springer Monographs in Mathematics, Springer-Verlag, Berlin, 2003.
- [5] A. Kanamori, *The higher infinite. Large cardinals in set theory from their beginnings. Second edition*. Springer Monographs in Mathematics, Springer-Verlag, Berlin, 2003.
- [6] K. Kunen, *Elementary embeddings and infinite combinatorics*. Journal of Symbolic Logic **36** (1971), 209–231.
- [7] R. Laver, *On the algebra of elementary embeddings of a rank into itself*. Advances in Mathematics **110** (1995), 334–346.
- [8] R. Laver, *Implications between strong large cardinal axioms*. Annals of Pure and Applied Logic **90** (1997), 79–90.
- [9] D.A. Martin, *Infinite games*. Proceedings of the ICM, Helsinki, 1978, Acad. Sci. Fennica, Helsinki, 269–273.
- [10] <https://mathoverflow.net/questions/191742/a-question-on-rank-to-rank-embeddings>
- [11] A. Suzuki, *No elementary embedding from V into V is definable from parameters*. Journal of Symbolic Logic **64** (1999), 1591–1594.

- [12] W. H. Woodin, *Suitable extender models II: beyond ω -huge*. Journal of Mathematical Logic **11** (2011), 115–436.

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI TORINO, VIA CARLO ALBERTO 10, 10123 TORINO—ITALY

E-mail address: `alessandro.andretta@unito.it`

UNIVERSITÀ DEGLI STUDI DI UDINE, DIPARTIMENTO DI SCIENZE MATEMATICHE, INFORMATICHE E FISICHE, VIA DELLE SCIENZE, 206, 33100 UDINE—ITALY

E-mail address: `vincenzo.dimonte@gmail.com`

URL: `https://users.dimi.uniud.it/~vincenzo.dimonte/`