



UNIVERSITÀ  
DEGLI STUDI  
DI UDINE

Università degli studi di Udine

One-dimensional von Karman models for elastic ribbons

*Original*

*Availability:*

This version is available <http://hdl.handle.net/11390/1123988> since 2021-03-19T07:34:43Z

*Publisher:*

*Published*

DOI:10.1007/s11012-017-0666-5

*Terms of use:*

The institutional repository of the University of Udine (<http://air.uniud.it>) is provided by ARIC services. The aim is to enable open access to all the world.

*Publisher copyright*

(Article begins on next page)

## ONE-DIMENSIONAL VON KÁRMÁN MODELS FOR ELASTIC RIBBONS

LORENZO FREDDI, PETER HORNUNG, MARIA GIOVANNA MORA, AND ROBERTO PARONI

ABSTRACT. By means of a variational approach we rigorously deduce three one-dimensional models for elastic ribbons from the theory of von Kármán plates, passing to the limit as the width of the plate goes to zero. The one-dimensional model found starting from the “linearized” von Kármán energy corresponds to that of a linearly elastic beam that can twist but can deform in just one plane; while the model found from the von Kármán energy is a non-linear model that comprises stretching, bendings, and twisting. The “constrained” von Kármán energy, instead, leads to a new Sadowsky type of model.

KEYWORDS: Elastic ribbons, von Kármán plates, Sadowsky functional, Gamma-convergence

MATHEMATICS SUBJECT CLASSIFICATION : 49J45, 49S05, 74B20, 74K20, 74K99

## 1. INTRODUCTION

Geometrically a ribbon is a body with three length scales: it is a parallelepiped whose length  $\ell$  is much larger than the width  $\varepsilon$ , which, in turn, is much larger than the thickness  $h$ . That is,  $\ell \gg \varepsilon \gg h$ . Since two characteristic dimensions are much smaller than the length, ribbons can be efficiently modelled as a one-dimensional continuum, see [14]. In the literature, two types of one-dimensional models are found: rod models and “Sadowsky type” models. We shall mainly discuss the latter, since we work within that framework; for rod type models we refer to [7] and the references therein.

So far, “Sadowsky type” models have been deduced starting from a plate model, that is, from a two-dimensional model obtained from a three-dimensional problem by letting the thickness  $h$  go to zero. Starting from a Kirchhoff plate model, a one-dimensional model for an isotropic elastic ribbon was proposed by Sadowsky in 1930, [15, 20]. The model was formally justified in 1962 by Wunderlich, [23, 26], by considering the Kirchhoff model for a plate of length  $\ell$  and width  $\varepsilon$ , and by letting  $\varepsilon$  go to zero. The justification given was only formal, since it was based on an ansatz on the deformation. Wunderlich’s technique is quite ingenious, but it leads to a singular energy density; we refer to [17] for a rigorous analysis of the so-called Wunderlich energy. A corrected Sadowsky type of energy was derived in [9] and generalized in [1, 10].

A third approach, which partly justifies the two approaches mentioned above, is to let the width  $\varepsilon$  and the thickness  $h$  go to zero simultaneously. By appropriately tuning the rates at which  $\varepsilon$  and  $h$  converge to zero, one obtains a hierarchy of one-dimensional models: in [11, 12] several rod models have been deduced, and in a forthcoming paper we will show that also “Sadowsky type” models can be obtained.

Before describing the contents of the present paper, we point out that the literature on ribbons is really blooming in several interesting directions, see, for instance, [2, 3, 4, 6, 8, 18, 21, 22].

Our starting point are the von Kármán plate models, whereas the papers quoted above have the Kirchhoff plate model as a starting point. The von Kármán model for plates has been successfully used in [5] to describe the plethora of morphological instabilities observed in a stretched and twisted ribbon.

The von Kármán plate equations, formulated more than a hundred years ago [24], have been recently justified by Friesecke, James, and Müller [13]. These authors consider a three-dimensional non-linear hyper-elastic material in a reference configuration  $\Omega_h = S_\varepsilon \times (-\frac{h}{2}, \frac{h}{2})$  with a stored energy density  $W : \mathbb{R}^{3 \times 3} \rightarrow [0, +\infty)$  satisfying standard regularity and growth conditions. In [13] the set  $S_\varepsilon$  is quite general, but in this introduction, in order to be consistent with the previous discussion, we take  $S_\varepsilon = (-\ell/2, \ell/2) \times (-\varepsilon/2, \varepsilon/2)$ . Then, the energy associated with a deformation

$y : \Omega_h \rightarrow \mathbb{R}^3$  is given by

$$\mathcal{E}^h(y) = \int_{\Omega_h} W(\nabla y) dx.$$

By scaling the elastic energy per unit volume  $\mathcal{E}^h/h \sim h^\beta$ , with  $\beta$  a positive real parameter, in [13] a hierarchy of plate models has been derived (by letting  $h$  go to zero) by means of  $\Gamma$ -convergence theory. The larger  $\beta$  is, the smaller the energy becomes. Therefore, heuristically, for large  $\beta$  the limit of the rescaled energy should produce the linear plate equation. This is indeed corroborated in [13]. Still in the same paper it is shown that for  $\beta = 2\alpha - 2$  and for the regimes  $\alpha > 3$ ,  $\alpha = 3$ , and  $2 < \alpha < 3$  three different  $\Gamma$ -limits are obtained that correspond to von Kármán type of energies.

Precisely, denoting by  $u : S_\varepsilon \rightarrow \mathbb{R}^2$  and  $v : S_\varepsilon \rightarrow \mathbb{R}$  the in-plane and the out-of-plane displacement fields, respectively, the three asymptotic energies are as follows (see [13, Theorem 2]):

$(LvK)_\varepsilon$  for  $\alpha > 3$  we have the “linearized” von Kármán theory, where  $u = 0$  and  $v$  minimizes the functional

$$I_\varepsilon^{LvK}(v) := \frac{1}{24} \int_{S_\varepsilon} Q_2(\nabla^2 v) dx,$$

with  $Q_2 : \mathbb{R}_{\text{sym}}^{2 \times 2} \rightarrow [0, +\infty)$  the positive definite quadratic form of linearized elasticity, see Remark 2.5 for a precise definition;

$(vK)_\varepsilon$  for  $\alpha = 3$  we have the von Kármán theory, where the in-plane and the out-of-plane displacements  $u$  and  $v$  minimize the functional

$$I_\varepsilon^{vK}(u, v) := \frac{1}{2} \int_{S_\varepsilon} Q_2\left(\frac{1}{2}[\nabla u + (\nabla u)^T + \nabla v \otimes \nabla v]\right) dx + \frac{1}{24} \int_{S_\varepsilon} Q_2(\nabla^2 v) dx;$$

$(CvK)_\varepsilon$  for  $2 < \alpha < 3$  we have the “constrained” von Kármán theory, in which the functional

$$I_\varepsilon^{CvK}(v) := \frac{1}{24} \int_{S_\varepsilon} Q_2(\nabla^2 v) dx$$

has to be minimized under the non-linear constraint

$$\nabla u + (\nabla u)^T + \nabla v \otimes \nabla v = 0, \quad (1.1)$$

or, equivalently, the functional  $I_\varepsilon^{CvK}$  has to be minimized under the constraint

$$\det(\nabla^2 v) = 0$$

(which is, in turn, necessary and sufficient for the existence of a map  $u$  satisfying (1.1)).

The existence of minimizers and the characterization of the Euler equations for constrained von Kármán plates have been studied in [16].

By letting  $h$  go to zero, the three-dimensional domain  $\Omega_h = S_\varepsilon \times (-\frac{h}{2}, \frac{h}{2})$  is “squeezed” to become  $S_\varepsilon$ . In this paper, we consider the von Kármán energies and we let  $\varepsilon$  go to zero, still by means of  $\Gamma$ -convergence, to find one-dimensional models for elastic ribbons in the von Kármán regimes. In this way, the two-dimensional domain  $S_\varepsilon = (-\ell/2, \ell/2) \times (-\varepsilon/2, \varepsilon/2)$  is “squeezed” to the segment  $I = (-\ell/2, \ell/2)$ , which we parametrize with the coordinate  $x_1$ . In the limit, the in-plane displacement  $u : S_\varepsilon \rightarrow \mathbb{R}^2$  generates two displacements: an axial displacement  $\xi_1 : I \rightarrow \mathbb{R}$ , and an orthogonal “in-plane” displacement  $\xi_2 : I \rightarrow \mathbb{R}$ . The out-of-plane displacement  $v : S_\varepsilon \rightarrow \mathbb{R}$ , in turn, generates an “out-of-plane” displacement  $w : I \rightarrow \mathbb{R}$  and the derivative of  $v$  in the direction orthogonal to the axis leads to a rotation  $\vartheta : I \rightarrow \mathbb{R}$ . The limit energies that we find in the three regimes are the following:

$(LvK)$  the limit of the “linearized” von Kármán energy is

$$J^{LvK}(w, \vartheta) := \frac{1}{24} \int_I Q_1(w'', \vartheta') dx_1;$$

$(vK)$  the limit of the von Kármán energy is

$$J^{vK}(\xi, w, \vartheta) := \frac{1}{2} \int_I Q_0\left(\xi_1' + \frac{|w'|^2}{2}\right) dx_1 + \frac{1}{24} \int_I (Q_0(\xi_2'') + Q_1(w'', \vartheta')) dx_1;$$

$(CvK)$  the limit of the “constrained” von Kármán energy is

$$J^{CvK}(w, \vartheta) := \frac{1}{24} \int_I \overline{Q}(w'', \vartheta') dx_1.$$

Here  $Q_1, Q_0$ , and  $\overline{Q}$  are energy densities whose precise definition can be found in Section 2; see Remark 2.5 for the specialization of these energies in the isotropic case.

We note that, since  $Q_1$  is quadratic, the functional  $(LvK)$  corresponds to the energy of a linearly elastic “three-dimensional beam” in which the section  $S_\varepsilon$  is unstretchable: the energy is simply due to the “out-of-plane” bending of the axis and to the torsion of the cross-section orthogonal to the axis. The limit functional  $(vK)$  is non-linear and penalizes stretching and both bendings of the axis, as well as the torsion of the cross-section. The functional  $(CvK)$  is sometimes called the energy of a beam with large deflections, see [25]. Despite the appearance, the energy functional  $(CvK)$  is very different from that of  $(LvK)$ . Indeed, in contrast to  $Q_1$ , the energy density  $\overline{Q}$  is not quadratic. It incorporates into its definition the non-linear constraint (1.1) that appears into the two-dimensional model  $(CvK)_\varepsilon$ . The energy density  $\overline{Q}$  agrees with the corrected Sadowsky energy density found in [9] in the isotropic case, and with that found in [10] for the general anisotropic case. To the best of our knowledge, the model  $(CvK)$  is new.

We conclude this introduction by pointing out that the statements of the results and the precise definitions are given in Section 2, while Section 3 is exclusively devoted to the proofs of these results.

## 2. NARROW STRIPS

Let  $\ell > 0$ , let  $I$  denote the interval  $(-\ell/2, \ell/2)$ , and let  $S_\varepsilon = I \times (-\varepsilon/2, \varepsilon/2)$  with  $\varepsilon > 0$ . For  $u \in W^{1,2}(S_\varepsilon; \mathbb{R}^2)$  and  $v \in W^{2,2}(S_\varepsilon)$  we consider the scaled von Kármán extensional and bending energies

$$\mathcal{J}_\varepsilon^{ext}(u, v) = \frac{1}{\varepsilon} \frac{1}{2} \int_{S_\varepsilon} Q_2 \left( Eu + \frac{1}{2} \nabla v \otimes \nabla v \right) dx, \quad \mathcal{J}_\varepsilon^{ben}(v) = \frac{1}{\varepsilon} \frac{1}{24} \int_{S_\varepsilon} Q_2(\nabla^2 v) dx,$$

where  $Eu = \frac{1}{2}(\nabla u + \nabla u^T)$  is the symmetric part of the gradient of the in-plane displacement  $u$ , while  $\nabla^2 v$  denotes the Hessian matrix of the out-of-plane displacement  $v$ . The energy density  $Q_2 : \mathbb{R}_{\text{sym}}^{2 \times 2} \rightarrow [0, +\infty)$  is assumed to be a positive definite quadratic form.

To simplify our analysis we rewrite the energies over the domain  $S := S_1 = I \times (-1/2, 1/2)$ . More precisely, we introduce the scaled versions  $y : S \rightarrow \mathbb{R}^2$  and  $w : S \rightarrow \mathbb{R}$  of  $u$  and  $v$ , respectively, by setting

$$y_1(x_1, x_2) := u_1(x_1, \varepsilon x_2), \quad y_2(x_1, x_2) := \varepsilon u_2(x_1, \varepsilon x_2), \quad w(x_1, x_2) := v(x_1, \varepsilon x_2),$$

and define the scaled differential operators

$$E^\varepsilon y := \begin{pmatrix} \partial_1 y_1 & \frac{1}{2\varepsilon}(\partial_1 y_2 + \partial_2 y_1) \\ \frac{1}{2\varepsilon}(\partial_2 y_1 + \partial_1 y_2) & \frac{1}{\varepsilon^2} \partial_2 y_2 \end{pmatrix},$$

$$\nabla_\varepsilon w := \left( \partial_1 w, \frac{1}{\varepsilon} \partial_2 w \right), \quad \nabla_\varepsilon^2 w := \begin{pmatrix} \partial_{11}^2 w & \frac{1}{\varepsilon} \partial_{12}^2 w \\ \frac{1}{\varepsilon} \partial_{21}^2 w & \frac{1}{\varepsilon^2} \partial_{22}^2 w \end{pmatrix},$$

so that

$$E^\varepsilon y(x) = Eu(x_1, \varepsilon x_2), \quad \nabla_\varepsilon w(x) = \nabla v(x_1, \varepsilon x_2), \quad \nabla_\varepsilon^2 w(x) = \nabla^2 v(x_1, \varepsilon x_2).$$

By performing the change of variables in the energy integrals we have that  $\mathcal{J}_\varepsilon^{ext}(u, v) = \mathcal{J}_\varepsilon^{ext}(y, w)$  and  $\mathcal{J}_\varepsilon^{ben}(v) = \mathcal{J}_\varepsilon^{ben}(w)$ , where

$$\mathcal{J}_\varepsilon^{ext}(y, w) := \frac{1}{2} \int_S Q_2 \left( E^\varepsilon y + \frac{1}{2} \nabla_\varepsilon w \otimes \nabla_\varepsilon w \right) dx, \quad \mathcal{J}_\varepsilon^{ben}(w) := \frac{1}{24} \int_S Q_2(\nabla_\varepsilon^2 w) dx. \quad (2.1)$$

Since we do not impose boundary conditions, we require the displacements to have zero average and, for the out-of plane component, also zero average gradient. That is, we shall work in the following spaces: for every open set  $\Omega \subset \mathbb{R}^\alpha$  with  $\alpha = 1, 2$ , we consider

$$\begin{aligned} W_{(0)}^{1,2}(\Omega) &:= \left\{ g \in W^{1,2}(\Omega) : \int_{\Omega} g(x) dx = 0 \right\}, \\ W_{(0)}^{2,2}(\Omega) &:= \left\{ g \in W^{2,2}(\Omega) : \int_{\Omega} g(x) dx = 0 \text{ and } \int_{\Omega} \nabla g(x) dx = 0 \right\}, \end{aligned}$$

and similarly we define  $W_{(0)}^{1,2}(\Omega; \mathbb{R}^2)$ .

Our first result is about compactness of sequences with bounded energy; the limit of the in-plane displacements will belong to the space of two-dimensional Bernoulli-Navier functions defined by

$$\begin{aligned} BN_{(0)}(S; \mathbb{R}^2) &:= \{ g \in W_{(0)}^{1,2}(S; \mathbb{R}^2) : (Eg)_{12} = (Eg)_{22} = 0 \} \\ &= \{ g \in W_{(0)}^{1,2}(S; \mathbb{R}^2) : \exists \xi_1 \in W_{(0)}^{1,2}(I) \text{ and } \xi_2 \in W_{(0)}^{1,2}(I) \cap W^{2,2}(I) \text{ such that} \\ &\quad g_1(x) = \xi_1(x_1) - x_2 \xi_2'(x_1), g_2(x) = \xi_2(x_1) \}, \end{aligned}$$

where the second characterization can be obtained by arguing as in [19, Section 4.1].

**Lemma 2.1.** *Let  $(w_\varepsilon) \subset W_{(0)}^{2,2}(S)$  be a sequence such that*

$$\sup_{\varepsilon} J_{\varepsilon}^{ben}(w_{\varepsilon}) < \infty. \quad (2.2)$$

*Then, up to a subsequence, there exist a vertical displacement  $w \in W_{(0)}^{2,2}(I)$  and a twist function  $\vartheta \in W_{(0)}^{1,2}(I)$  such that*

$$w_{\varepsilon} \rightharpoonup w \text{ in } W^{2,2}(S), \quad \nabla_{\varepsilon} w_{\varepsilon} \rightharpoonup (w', \vartheta) \text{ in } W^{1,2}(S; \mathbb{R}^2), \quad (2.3)$$

*and*

$$\nabla_{\varepsilon}^2 w_{\varepsilon} \rightharpoonup \begin{pmatrix} w'' & \vartheta' \\ \vartheta' & \gamma \end{pmatrix} \text{ in } L^2(S; \mathbb{R}_{\text{sym}}^{2 \times 2}) \quad (2.4)$$

*for a suitable  $\gamma \in L^2(S)$ .*

*Moreover, if  $(y_{\varepsilon}) \subset W_{(0)}^{1,2}(S; \mathbb{R}^2)$  is a further sequence such that*

$$\sup_{\varepsilon} J_{\varepsilon}^{ext}(y_{\varepsilon}, w_{\varepsilon}) < \infty, \quad (2.5)$$

*then, up to a subsequence, there exists  $y \in BN_{(0)}(S; \mathbb{R}^2)$  such that*

$$y_{\varepsilon} \rightharpoonup y \text{ in } W^{1,2}(S; \mathbb{R}^2).$$

*Also,*

$$E^{\varepsilon} y_{\varepsilon} \rightharpoonup E \text{ in } L^2(S; \mathbb{R}_{\text{sym}}^{2 \times 2})$$

*for a suitable  $E \in L^2(S; \mathbb{R}_{\text{sym}}^{2 \times 2})$  such that  $E_{11} = \partial_1 y_1$ .*

The rest of this section is devoted to state the  $\Gamma$ -convergence results starting from the simpler case of the linearized theory  $(LvK)_{\varepsilon}$ , and proceeding in the order of increasing difficulty to consider the standard and the constrained models  $(vK)_{\varepsilon}$  and  $(CvK)_{\varepsilon}$ , respectively.

**2.1. The linearized von Kármán model.** In order to state our first convergence result we need to introduce some definitions. Let  $Q_1 : \mathbb{R} \times \mathbb{R} \rightarrow [0, +\infty)$  be defined by

$$Q_1(\kappa, \tau) := \min_{\gamma \in \mathbb{R}} \left\{ Q_2(M) : M = \begin{pmatrix} \kappa & \tau \\ \tau & \gamma \end{pmatrix} \right\}.$$

Let  $J^{LvK} : W_{(0)}^{2,2}(I) \times W_{(0)}^{1,2}(I) \rightarrow \mathbb{R}$  be defined by

$$J^{LvK}(w, \vartheta) := \frac{1}{24} \int_I Q_1(w'', \vartheta') dx_1.$$

**Theorem 2.2.** *As  $\varepsilon \rightarrow 0$ , the functionals  $J_{\varepsilon}^{ben}$   $\Gamma$ -converge to the functional  $J^{LvK}$  in the following sense:*

- (i) (liminf inequality) for every sequence  $(w_\varepsilon) \subset W_{(0)}^{2,2}(S)$ ,  $w \in W_{(0)}^{2,2}(I)$ , and  $\vartheta \in W_{(0)}^{1,2}(I)$  such that  $w_\varepsilon \rightharpoonup w$  in  $W^{2,2}(S)$ , and  $\nabla_\varepsilon w_\varepsilon \rightharpoonup (w', \vartheta)$  in  $W^{1,2}(S; \mathbb{R}^2)$ , we have that

$$\liminf_{\varepsilon \rightarrow 0} J_\varepsilon^{ben}(w_\varepsilon) \geq J^{LvK}(w, \vartheta);$$

- (ii) (recovery sequence) for every  $w \in W_{(0)}^{2,2}(I)$  and  $\vartheta \in W_{(0)}^{1,2}(I)$  there exists a sequence  $(w_\varepsilon) \subset W_{(0)}^{2,2}(S)$  such that  $w_\varepsilon \rightharpoonup w$  in  $W^{2,2}(S)$ ,  $\nabla_\varepsilon w_\varepsilon \rightharpoonup (w', \vartheta)$  in  $W^{1,2}(S; \mathbb{R}^2)$ , and

$$\limsup_{\varepsilon \rightarrow 0} J_\varepsilon^{ben}(w_\varepsilon) \leq J^{LvK}(w, \vartheta).$$

**2.2. The von Kármán model.** The statement of our second convergence result needs some further definitions. Let  $Q_0 : \mathbb{R} \rightarrow [0, +\infty)$  be defined by

$$Q_0(\mu) := \min_{z \in \mathbb{R}} Q_1(\mu, z) = \min_{(z_1, z_2) \in \mathbb{R}^2} \left\{ Q_2(M) : M = \begin{pmatrix} \mu & z_1 \\ z_1 & z_2 \end{pmatrix} \right\}.$$

Let  $J^{vK} : BN_{(0)}(S; \mathbb{R}^2) \times W_{(0)}^{2,2}(I) \times W_{(0)}^{1,2}(I) \rightarrow \mathbb{R}$  be defined by

$$J^{vK}(y, w, \vartheta) := \frac{1}{2} \int_S Q_0 \left( \partial_1 y_1 + \frac{|w'|^2}{2} \right) dx + \frac{1}{24} \int_I Q_1(w'', \vartheta') dx_1.$$

**Theorem 2.3.** As  $\varepsilon \rightarrow 0$ , the functionals  $J_\varepsilon^{vK} := J_\varepsilon^{ext} + J_\varepsilon^{ben}$   $\Gamma$ -converge to the functional  $J^{vK}$  in the following sense:

- (i) (liminf inequality) for every pair of sequences  $(y_\varepsilon) \subset W_{(0)}^{1,2}(S; \mathbb{R}^2)$ ,  $(w_\varepsilon) \subset W_{(0)}^{2,2}(S)$ ,  $y \in BN_{(0)}(S; \mathbb{R}^2)$ ,  $w \in W_{(0)}^{2,2}(I)$ , and  $\vartheta \in W_{(0)}^{1,2}(I)$  such that  $y_\varepsilon \rightharpoonup y$  in  $W^{1,2}(S; \mathbb{R}^2)$ ,  $w_\varepsilon \rightharpoonup w$  in  $W^{2,2}(S)$ , and  $\nabla_\varepsilon w_\varepsilon \rightharpoonup (w', \vartheta)$  in  $W^{1,2}(S; \mathbb{R}^2)$ , we have that

$$\liminf_{\varepsilon \rightarrow 0} J_\varepsilon^{vK}(y_\varepsilon, w_\varepsilon) \geq J^{vK}(y, w, \vartheta);$$

- (ii) (recovery sequence) for every  $y \in BN_{(0)}(S; \mathbb{R}^2)$ ,  $w \in W_{(0)}^{2,2}(I)$  and  $\vartheta \in W_{(0)}^{1,2}(I)$  there exists a pair of sequences  $(y_\varepsilon) \subset W_{(0)}^{1,2}(S; \mathbb{R}^2)$ ,  $(w_\varepsilon) \subset W_{(0)}^{2,2}(S)$  such that  $y_\varepsilon \rightharpoonup y$  in  $W^{1,2}(S; \mathbb{R}^2)$ ,  $w_\varepsilon \rightharpoonup w$  in  $W^{2,2}(S)$ ,  $\nabla_\varepsilon w_\varepsilon \rightharpoonup (w', \vartheta)$  in  $W^{1,2}(S; \mathbb{R}^2)$ , and

$$\limsup_{\varepsilon \rightarrow 0} J_\varepsilon^{vK}(y_\varepsilon, w_\varepsilon) \leq J^{vK}(y, w, \vartheta).$$

**2.3. The constrained von Kármán model.** The constrained von Kármán energy of a displacement  $v \in W_{(0)}^{2,2}(S_\varepsilon)$  such that  $\det \nabla^2 v = 0$  a.e. in  $S_\varepsilon$  is  $J_\varepsilon^{ben}(v)$ . We observe that the map  $w$ , defined over the rescaled domain, belongs to the space

$$W_{\det, \varepsilon}^{2,2}(S) := \{w \in W_{(0)}^{2,2}(S) : \det \nabla_\varepsilon^2 w = 0 \text{ a.e. in } S\}.$$

We set  $J_\varepsilon^{CvK} : W_{\det, \varepsilon}^{2,2}(S) \rightarrow \mathbb{R}$  the functional  $J_\varepsilon^{CvK}(w) = J_\varepsilon^{ben}(w)$ .

Let  $\overline{Q} : \mathbb{R} \times \mathbb{R} \rightarrow [0, +\infty)$  be defined by

$$\overline{Q}(\kappa, \tau) := \min_{\gamma \in \mathbb{R}} \left\{ Q_2(M) + \alpha^+ (\det M)^+ + \alpha^- (\det M)^- : M = \begin{pmatrix} \kappa & \tau \\ \tau & \gamma \end{pmatrix} \right\},$$

where

$$\alpha^+ := \sup \{ \alpha > 0 : Q_2(M) + \alpha \det M \geq 0 \text{ for every } M \in \mathbb{R}_{\text{sym}}^{2 \times 2} \}$$

and

$$\alpha^- := \sup \{ \alpha > 0 : Q_2(M) - \alpha \det M \geq 0 \text{ for every } M \in \mathbb{R}_{\text{sym}}^{2 \times 2} \}.$$

Let  $J^{CvK} : W_{(0)}^{2,2}(I) \times W_{(0)}^{1,2}(I) \rightarrow \mathbb{R}$  be defined by

$$J^{CvK}(w, \vartheta) := \frac{1}{24} \int_I \overline{Q}(w'', \vartheta') dx_1.$$

**Theorem 2.4.** As  $\varepsilon \rightarrow 0$ , the functionals  $J_\varepsilon^{CvK}$   $\Gamma$ -converge to the functional  $J^{CvK}$  in the following sense:

(i) (liminf inequality) for every sequence  $(w_\varepsilon)$  with  $w_\varepsilon \in W_{\det, \varepsilon}^{2,2}(S)$ ,  $w \in W_{\langle 0 \rangle}^{2,2}(I)$ , and  $\vartheta \in W_{\langle 0 \rangle}^{1,2}(I)$  such that  $w_\varepsilon \rightharpoonup w$  in  $W^{2,2}(S)$ , and  $\nabla_\varepsilon w_\varepsilon \rightharpoonup (w', \vartheta)$  in  $W^{1,2}(S; \mathbb{R}^2)$ , we have that

$$\liminf_{\varepsilon \rightarrow 0} J_\varepsilon^{CvK}(w_\varepsilon) \geq J^{CvK}(w, \vartheta);$$

(ii) (recovery sequence) for every  $w \in W_{\langle 0 \rangle}^{2,2}(I)$  and  $\vartheta \in W_{\langle 0 \rangle}^{1,2}(I)$  there exists a sequence  $(w_\varepsilon)$  with  $w_\varepsilon \in W_{\det, \varepsilon}^{2,2}(S)$  such that  $w_\varepsilon \rightharpoonup w$  in  $W^{2,2}(S)$ ,  $\nabla_\varepsilon w_\varepsilon \rightharpoonup (w', \vartheta)$  in  $W^{1,2}(S; \mathbb{R}^2)$ , and

$$\limsup_{\varepsilon \rightarrow 0} J_\varepsilon^{CvK}(w_\varepsilon) \leq J^{CvK}(w, \vartheta).$$

**Remark 2.5.** The quadratic energy density  $Q_2$  can be computed from the non-linear energy density  $W$  of the material, also mentioned in the introduction, by first computing the quadratic energy density  $Q_3$ , see [13],

$$Q_3(F) := \frac{\partial^2 W}{\partial F^2}(I)(F, F) = \sum_{i,j,k,l=1}^3 \frac{\partial^2 W}{\partial F_{ij} \partial F_{kl}}(I) F_{ij} F_{kl}, \quad F \in \mathbb{R}^{3 \times 3},$$

and then by minimizing over the third column and row:

$$Q_2(A) := \min\{Q_3(F) : F_{\alpha\beta} = A_{\alpha\beta} \quad \alpha, \beta = 1, 2\}, \quad A \in \mathbb{R}_{\text{sym}}^{2 \times 2}.$$

If the energy density  $W$  is isotropic, the quadratic energy density  $Q_3$  has the following representation:

$$Q_3(F) = 2\mu|F_{\text{sym}}|^2 + \lambda(F_{\text{sym}} \cdot I)^2, \quad F_{\text{sym}} := \frac{F + F^T}{2} \in \mathbb{R}^{3 \times 3},$$

where  $\mu$  and  $\lambda$  are the so-called Lamé coefficients. A simple computation then leads to

$$Q_2(A) = 2\mu|A|^2 + \frac{2\mu\lambda}{2\mu + \lambda}(A \cdot I)^2, \quad A \in \mathbb{R}_{\text{sym}}^{2 \times 2}.$$

The energy densities  $Q_1, Q_0$ , and  $\overline{Q}$ , may be found to have the following representation

$$Q_1(\kappa, \tau) = E_Y \kappa^2 + 4\mu\tau^2,$$

where  $E_Y := \mu \frac{2\mu+3\lambda}{\mu+\lambda}$  is the Young modulus of the material,

$$Q_0(\kappa) = E_Y \kappa^2,$$

and

$$\frac{1}{12} \overline{Q}(\kappa, \tau) = \begin{cases} \mathcal{D} \frac{(\kappa^2 + \tau^2)^2}{\kappa^2} & \text{if } |\kappa| > |\tau|, \\ 4\mathcal{D}\tau^2 & \text{if } |\kappa| \leq |\tau|, \end{cases}$$

where  $\mathcal{D} := \frac{\mu(\lambda+\mu)}{3(2\mu+\lambda)}$  is the bending stiffness.

### 3. PROOFS

This section is devoted to prove the theorems stated in the previous section. For a given function  $u \in L^1(S)$ , we shall denote by  $\langle u \rangle$  the integral mean value of  $u$  on  $S$ , that is,

$$\langle u \rangle := \frac{1}{\ell} \int_S u(x) dx.$$

We use the same notation to denote the average over  $I$  of functions defined on  $I$ .

*Proof of Lemma 2.1.* Let  $(w_\varepsilon) \subset W_{\langle 0 \rangle}^{2,2}(S)$  be a sequence of vertical displacements of  $S$  satisfying (2.2). This bound and the fact that  $Q_2$  is positive definite imply that

$$\|\partial_{11}^2 w_\varepsilon\|_{L^2(S)} + \|\varepsilon^{-1} \partial_{12}^2 w_\varepsilon\|_{L^2(S)} + \|\varepsilon^{-2} \partial_{22}^2 w_\varepsilon\|_{L^2(S)} \leq C \quad (3.1)$$

for any  $\varepsilon$ . Since

$$\int_S w_\varepsilon(x) dx = 0, \quad \int_S \nabla w_\varepsilon(x) dx = 0$$

for every  $\varepsilon > 0$ , by Poincaré-Wirtinger inequality the sequence  $(w_\varepsilon)$  is uniformly bounded in  $W^{2,2}(S)$ . Therefore, there exists  $w \in W_{(0)}^{2,2}(S)$  such that  $w_\varepsilon \rightharpoonup w$  weakly in  $W^{2,2}(S)$ , up to a subsequence.

By the previous bound,  $\nabla(\varepsilon^{-1}\partial_2 w_\varepsilon)$  is a bounded sequence in  $L^2(S; \mathbb{R}^2)$  and, by Poincaré-Wirtinger inequality, also  $(\varepsilon^{-1}\partial_2 w_\varepsilon)$  is bounded in  $L^2(S)$ . It follows that  $w$  is independent of  $x_2$  and there exists  $\vartheta \in W_{(0)}^{1,2}(S)$  such that  $\varepsilon^{-1}\partial_2 w_\varepsilon \rightharpoonup \vartheta$  weakly in  $W^{1,2}(S)$ , up to a subsequence. Moreover, also  $\vartheta$  is independent of  $x_2$ .

By (3.1), up to subsequences, we have that  $\nabla_\varepsilon^2 w_\varepsilon$  converges to a matrix field  $A$  weakly in  $L^2(S; \mathbb{R}_{\text{sym}}^{2 \times 2})$ . By using the convergences established above, it follows that  $A_{11} = w''$  and  $A_{12} = \vartheta'$ . The entry  $A_{22}$ , that cannot be identified in terms of  $w$  and  $\vartheta$ , is denoted by  $\gamma$  in the statement. This proves (2.4).

We now prove the second part of the statement. The bound (2.5) implies that

$$\left\| E^\varepsilon y_\varepsilon + \frac{1}{2} \nabla_\varepsilon w_\varepsilon \otimes \nabla_\varepsilon w_\varepsilon \right\|_{L^2} \leq C \quad (3.2)$$

for any  $\varepsilon$ . Since  $(\nabla_\varepsilon^2 w_\varepsilon)$  is bounded in  $L^2$ , we have

$$\begin{aligned} \|\nabla_\varepsilon w_\varepsilon \otimes \nabla_\varepsilon w_\varepsilon\|_{L^2} &\leq C \|\nabla_\varepsilon w_\varepsilon\|_{L^2}^2 = C \|\nabla_\varepsilon w_\varepsilon\|_{L^4}^2 \leq C(\|\partial_1 w_\varepsilon\|_{L^4}^2 + \|\varepsilon^{-1}\partial_2 w_\varepsilon\|_{L^4}^2) \\ &\leq C(\|\nabla \partial_1 w_\varepsilon\|_{L^2}^2 + \|\nabla \varepsilon^{-1}\partial_2 w_\varepsilon\|_{L^2}^2) \leq C\|\nabla_\varepsilon^2 w_\varepsilon\|_{L^2}^2 \leq C \end{aligned}$$

for any  $\varepsilon$ , and the third to last inequality follows by the imbedding  $W^{1,2}(S) \subset L^q \forall q \in [2, +\infty)$  and Poincaré-Wirtinger inequality. Together with (3.2), this implies that the sequence  $(E^\varepsilon y_\varepsilon)$  is bounded in  $L^2$ .

By the definition of  $E^\varepsilon$  and Korn-Poincaré inequality we have that

$$\|y_\varepsilon\|_{W^{1,2}} \leq C\|E y_\varepsilon\|_{L^2} \leq C\|E^\varepsilon y_\varepsilon\|_{L^2} \leq C. \quad (3.3)$$

Hence, up to subsequences, there exist  $E \in L^2(S; \mathbb{R}_{\text{sym}}^{2 \times 2})$  and  $y \in W_{(0)}^{1,2}(S; \mathbb{R}^2)$  such that

$$\begin{aligned} E^\varepsilon y_\varepsilon &\rightharpoonup E \quad \text{in } L^2(S; \mathbb{R}_{\text{sym}}^{2 \times 2}), \\ y_\varepsilon &\rightharpoonup y \quad \text{in } W^{1,2}(S; \mathbb{R}^2). \end{aligned}$$

By the definition of  $E^\varepsilon$  and (3.3) we have that

$$(E y_\varepsilon)_{12} \rightharpoonup 0 = (E y)_{12}, \quad (E y_\varepsilon)_{22} \rightharpoonup 0 = (E y)_{22};$$

hence,  $y \in BN_{(0)}(S; \mathbb{R}^2)$ . Finally, the observation that  $(E^\varepsilon y_\varepsilon)_{11} = \partial_1(y_\varepsilon)_1 \rightharpoonup \partial_1 y_1$  in  $L^2(S)$  concludes the proof.  $\square$

*Proof of Theorem 2.2-(i).* Let  $(w_\varepsilon) \subset W_{(0)}^{2,2}(S)$  be such that  $w_\varepsilon \rightharpoonup w$  in  $W^{2,2}(S)$ , and  $\nabla_\varepsilon w_\varepsilon \rightharpoonup (w', \vartheta)$  in  $W^{1,2}(S; \mathbb{R}^2)$ , for some  $w \in W_{(0)}^{2,2}(I)$  and  $\vartheta \in W_{(0)}^{1,2}(I)$ . Without loss of generality, we can assume that  $\liminf_{\varepsilon \rightarrow 0} J_\varepsilon^{\text{ben}}(w_\varepsilon) < +\infty$ . By Lemma 2.1 we infer that, up to subsequences,

$$\nabla_\varepsilon^2 w_\varepsilon \rightharpoonup \begin{pmatrix} w'' & \vartheta' \\ \vartheta' & \gamma \end{pmatrix} =: M_\gamma \text{ in } L^2(S; \mathbb{R}_{\text{sym}}^{2 \times 2})$$

for some  $\gamma \in L^2(S)$ . By weak lower semicontinuity and the definition of  $Q_1$  we have

$$\begin{aligned} \liminf_{\varepsilon \rightarrow 0} J_\varepsilon^{\text{ben}}(w_\varepsilon) &= \liminf_{\varepsilon \rightarrow 0} \frac{1}{24} \int_S Q_2(\nabla_\varepsilon^2 w_\varepsilon) dx \\ &\geq \frac{1}{24} \int_S Q_2(M_\gamma) dx \geq \frac{1}{24} \int_I Q_1(w'', \vartheta') dx_1 = J^{L^v K}(w, \vartheta). \end{aligned}$$

$\square$

*Proof of Theorem 2.2-(ii).* Let  $w \in W_{(0)}^{2,2}(I)$  and  $\vartheta \in W_{(0)}^{1,2}(I)$ . We set

$$M_\gamma := \begin{pmatrix} w'' & \vartheta' \\ \vartheta' & \gamma \end{pmatrix},$$

where  $\gamma \in L^2(I)$  is such that

$$Q_1(w'', \vartheta') = Q_2(M_\gamma).$$



The fact that  $\gamma$  belongs to  $L^2(I)$  follows immediately by choosing  $M_0 = w''e_1 \otimes e_1 + \vartheta'(e_1 \otimes e_2 + e_2 \otimes e_1)$  as a competitor in the definition of  $Q_1$  and by using the positive definiteness of  $Q_2$ .

Let  $\vartheta_\varepsilon \in C^\infty(\bar{I})$  be such that  $\int_I \vartheta_\varepsilon(x_1) dx_1 = 0$ ,  $\vartheta_\varepsilon \rightarrow \vartheta$  in  $W^{1,2}(I)$ , and  $\varepsilon \vartheta_\varepsilon'' \rightarrow 0$  in  $L^2(I)$ . Let  $\gamma_\varepsilon \in C^\infty(\bar{I})$  be such that  $\gamma_\varepsilon \rightarrow \gamma$ ,  $\varepsilon \gamma_\varepsilon' \rightarrow 0$  and  $\varepsilon^2 \gamma_\varepsilon'' \rightarrow 0$  in  $L^2(I)$ . Let

$$w_\varepsilon(x) = w(x_1) + \varepsilon x_2 \vartheta_\varepsilon(x_1) + \frac{\varepsilon^2}{2} (x_2^2 \gamma_\varepsilon(x_1) - \langle x_2^2 \gamma_\varepsilon \rangle - x_1 \langle x_2^2 \gamma_\varepsilon' \rangle). \quad (3.4)$$

It turns out that  $w_\varepsilon \in W_{(0)}^{2,2}(S)$  and, by the convergences above, we have  $w_\varepsilon \rightarrow w$  in  $W^{2,2}(S)$ ,  $\nabla_\varepsilon w_\varepsilon \rightarrow (w', \vartheta)$  in  $W^{1,2}(S; \mathbb{R}^2)$ , and  $\nabla_\varepsilon^2 w_\varepsilon \rightarrow M_\gamma$  in  $L^2(S)$ . Moreover, by strong continuity we have

$$\lim_{\varepsilon \rightarrow 0} J_\varepsilon^{ben}(w_\varepsilon) = \lim_{\varepsilon \rightarrow 0} \frac{1}{24} \int_S Q_2(\nabla_\varepsilon^2 w_\varepsilon) dx = \frac{1}{24} \int_I Q_1(w'', \vartheta') dx_1 = J^{LvK}(w, \vartheta).$$

□

*Proof of Theorem 2.3-(i).* Let  $(y_\varepsilon) \subset W_{(0)}^{1,2}(S; \mathbb{R}^2)$ ,  $(w_\varepsilon) \subset W_{(0)}^{2,2}(S)$  be such that  $y_\varepsilon \rightharpoonup y$  in  $W^{1,2}(S; \mathbb{R}^2)$ ,  $w_\varepsilon \rightharpoonup w$  in  $W^{2,2}(S)$ , and  $\nabla_\varepsilon w_\varepsilon \rightharpoonup (w', \vartheta)$  in  $W^{1,2}(S; \mathbb{R}^2)$ , for some  $y \in BN_{(0)}(S; \mathbb{R}^2)$ ,  $w \in W_{(0)}^{2,2}(I)$  and  $\vartheta \in W_{(0)}^{1,2}(I)$ . As usual, we can assume that  $\liminf_{\varepsilon \rightarrow 0} J_\varepsilon^{vK}(y_\varepsilon, w_\varepsilon) < +\infty$  and by Lemma 2.1 we deduce that, up to subsequences,

$$E^\varepsilon y_\varepsilon \rightharpoonup E \quad \text{and} \quad \nabla_\varepsilon^2 w_\varepsilon \rightharpoonup \begin{pmatrix} w'' & \vartheta' \\ \vartheta' & \gamma \end{pmatrix} =: M_\gamma \quad \text{in } L^2(S; \mathbb{R}^{2 \times 2}_{\text{sym}})$$

for some  $E \in L^2(S; \mathbb{R}^{2 \times 2}_{\text{sym}})$  with  $E_{11} = \partial_1 y_1$ , and  $\gamma \in L^2(S)$ . Moreover, by the convergences above,

$$E^\varepsilon y_\varepsilon + \frac{1}{2} \nabla_\varepsilon w_\varepsilon \otimes \nabla_\varepsilon w_\varepsilon \rightharpoonup E + \frac{1}{2} (w', \vartheta) \otimes (w', \vartheta) \quad \text{in } L^2(S; \mathbb{R}^{2 \times 2}_{\text{sym}}).$$

Then, by lower semicontinuity, we have

$$\begin{aligned} \liminf_{\varepsilon \rightarrow 0} J_\varepsilon^{vK}(y_\varepsilon, w_\varepsilon) &\geq \liminf_{\varepsilon \rightarrow 0} \frac{1}{2} J_\varepsilon(y_\varepsilon, w_\varepsilon) + \liminf_{\varepsilon \rightarrow 0} \frac{1}{24} J_\varepsilon^{\text{lin}}(w_\varepsilon) \\ &= \liminf_{\varepsilon \rightarrow 0} \frac{1}{2} \int_S Q_2 \left( E^\varepsilon y_\varepsilon + \frac{1}{2} \nabla_\varepsilon w_\varepsilon \otimes \nabla_\varepsilon w_\varepsilon \right) dx + \liminf_{\varepsilon \rightarrow 0} \frac{1}{24} \int_S Q_2(\nabla_\varepsilon^2 w_\varepsilon) dx \\ &\geq \frac{1}{2} \int_S Q_2 \left( E + \frac{1}{2} (w', \vartheta) \otimes (w', \vartheta) \right) dx + \frac{1}{24} \int_S Q_2(M_\gamma) dx \\ &\geq \frac{1}{2} \int_S Q_0 \left( \partial_1 y_1 + \frac{1}{2} |w'|^2 \right) dx + \frac{1}{24} \int_I Q_1(w'', \vartheta') dx_1 \\ &= J^{vK}(y, w, \vartheta). \end{aligned}$$

□

*Proof of Theorem 2.3-(ii).* Let  $y \in BN_{(0)}(S; \mathbb{R}^2)$ ,  $w \in W_{(0)}^{2,2}(I)$ , and  $\vartheta \in W_{(0)}^{1,2}(I)$ . As before, there exists  $\gamma \in L^2(I)$  such that the matrix

$$M_\gamma := \begin{pmatrix} w'' & \vartheta' \\ \vartheta' & \gamma \end{pmatrix}$$

satisfies

$$Q_1(w'', \vartheta') = Q_2(M_\gamma).$$

There exist  $\xi_1 \in W_{(0)}^{1,2}(I)$  and  $\xi_2 \in W_{(0)}^{1,2}(I) \cap W^{2,2}(I)$  such that  $y_1(x) = \xi_1(x_1) - x_2 \xi_2'(x_1)$  and  $y_2(x) = \xi_2(x_1)$ . Moreover, there exists  $z \in L^2(S; \mathbb{R}^2)$  such that the matrix

$$M_z := \begin{pmatrix} \partial_1 y_1 + \frac{1}{2} |w'|^2 & z_1 \\ z_1 & z_2 \end{pmatrix} = \begin{pmatrix} \xi_1'(x_1) - x_2 \xi_2''(x_1) + \frac{1}{2} |w'(x_1)|^2 & z_1 \\ z_1 & z_2 \end{pmatrix}$$

satisfies

$$Q_0 \left( \partial_1 y_1 + \frac{1}{2} |w'|^2 \right) = Q_2(M_z).$$

It is easily seen that  $z_1$  and  $z_2$  depend linearly on  $\partial_1 y_1 + \frac{1}{2}|w'|^2$ . Since  $\partial_1 y_1(x_1, x_2) + \frac{1}{2}|w'(x_1)|^2 = \xi'_1(x_1) - x_2 \xi''_2(x_1) + \frac{1}{2}|w'(x_1)|^2$ , there exist  $\zeta_\alpha \in L^2(I)$  and  $\eta_\alpha \in L^2(I)$  such that

$$z_\alpha(x_1, x_2) = \zeta_\alpha(x_1) + x_2 \eta_\alpha(x_1), \quad \alpha = 1, 2.$$

Let  $w_\varepsilon$  be as in the proof of Theorem 2.2-(ii) (see (3.4)), and let  $\zeta_\alpha^\varepsilon, \eta_\alpha^\varepsilon \in C^\infty(\bar{I})$  be such that  $\zeta_\alpha^\varepsilon \rightarrow \zeta_\alpha$  and  $\eta_\alpha^\varepsilon \rightarrow \eta_\alpha$  in  $L^2(I)$  and  $\varepsilon \zeta_\alpha^{\varepsilon'} \rightarrow 0$  and  $\varepsilon \eta_\alpha^{\varepsilon'} \rightarrow 0$  in  $L^2(I)$ . Let us define

$$\begin{aligned} (y_\varepsilon)_1(x_1, x_2) &:= \xi_1(x_1) - x_2 \xi'_2(x_1) + \varepsilon(x_2^2 \eta_1^\varepsilon(x_1) - \langle x_2^2 \eta_1^\varepsilon \rangle), \\ (y_\varepsilon)_2(x_1, x_2) &:= \xi_2(x_1) + \varepsilon \left( \int_0^{x_1} (2\zeta_1^\varepsilon(s) - w'(s)\vartheta(s)) ds - \langle \int_0^{x_1} (2\zeta_1^\varepsilon(s) - w'(s)\vartheta(s)) ds \rangle \right) \\ &\quad + \frac{\varepsilon^2}{2} x_2 (2\zeta_2^\varepsilon(x_1) - \vartheta^2(x_1)) + \frac{\varepsilon^2}{2} (x_2^2 \eta_2^\varepsilon(x_1) - \langle x_2^2 \eta_2^\varepsilon \rangle). \end{aligned}$$

Then it is easy to check that

$$E^\varepsilon y_\varepsilon + \frac{1}{2} \nabla_\varepsilon w_\varepsilon \otimes \nabla_\varepsilon w_\varepsilon \rightarrow M_z \text{ in } L^2(S; \mathbb{R}_{\text{sym}}^{2 \times 2}).$$

Thus, by strong continuity,

$$\begin{aligned} \lim_{\varepsilon \rightarrow 0} J_\varepsilon^{vK}(y_\varepsilon, w_\varepsilon) &= \lim_{\varepsilon \rightarrow 0} \left( \frac{1}{2} J_\varepsilon^{ext}(y_\varepsilon, w_\varepsilon) + \frac{1}{24} J_\varepsilon^{ben}(w_\varepsilon) \right) \\ &= \lim_{\varepsilon \rightarrow 0} \left( \frac{1}{2} \int_S Q_2(E^\varepsilon y_\varepsilon + \frac{1}{2} \nabla_\varepsilon w_\varepsilon \otimes \nabla_\varepsilon w_\varepsilon) dx + \frac{1}{24} \int_S Q_2(\nabla_\varepsilon^2 w_\varepsilon) dx \right) \\ &= \frac{1}{2} \int_S Q_2(M_z) dx + \frac{1}{24} \int_S Q_2(M_\gamma) dx \\ &= \frac{1}{2} \int_S Q_0(\partial_1 y_1 + \frac{1}{2}|w'|^2) dx + \frac{1}{24} \int_I Q_1(w'', \vartheta') dx_1 \\ &= J^{vK}(y, w, \vartheta). \end{aligned}$$

□

The proof of the  $\Gamma$ -convergence theorem 2.4 is based on a relaxation result for a quadratic integral functional with a constraint on the determinant, that has been proved in [10, Proposition 9] and recalled here for reader's convenience.

Let  $\mathcal{B}$  be a bounded open subset of  $\mathbb{R}^n$ . Let  $Q : \mathcal{B} \times \mathbb{R}_{\text{sym}}^{2 \times 2} \rightarrow [0, +\infty)$  be measurable in the first variable and quadratic in the second. Define the functional

$$\mathcal{F} : L^2(\mathcal{B}; \mathbb{R}_{\text{sym}}^{2 \times 2}) \rightarrow [0, +\infty]$$

by

$$\mathcal{F}(M) := \begin{cases} \int_{\mathcal{B}} Q(x, M(x)) dx & \text{if } \det M = 0 \text{ a.e. in } \mathcal{B}, \\ +\infty & \text{otherwise.} \end{cases}$$

**Theorem 3.1** ([10]). *The weak- $L^2$  lower semicontinuous envelope of  $\mathcal{F}$  is the functional*

$$\overline{\mathcal{F}} : L^2(\mathcal{B}; \mathbb{R}_{\text{sym}}^{2 \times 2}) \rightarrow [0, +\infty)$$

given by

$$\overline{\mathcal{F}}(M) = \int_{\mathcal{B}} (Q(x, M(x)) + \alpha^+(x)(\det M(x))^+ + \alpha^-(x)(\det M(x))^-) dx,$$

where for every  $x \in \mathcal{B}$

$$\alpha^+(x) := \sup\{\alpha > 0 : Q(x, M) + \alpha \det M \geq 0 \text{ for every } M \in \mathbb{R}_{\text{sym}}^{2 \times 2}\}$$

and

$$\alpha^-(x) := \sup\{\alpha > 0 : Q(x, M) - \alpha \det M \geq 0 \text{ for every } M \in \mathbb{R}_{\text{sym}}^{2 \times 2}\}.$$

*Proof of Theorem 2.4–(i).* Let  $(w_\varepsilon)$  be such that  $w_\varepsilon \in W_{\det, \varepsilon}^{2,2}(S)$ ,  $w_\varepsilon \rightharpoonup w$  in  $W^{2,2}(S)$ , and  $\nabla_\varepsilon w_\varepsilon \rightharpoonup (w', \vartheta)$  in  $W^{1,2}(S; \mathbb{R}^2)$ , for some  $w \in W_{(0)}^{2,2}(I)$  and  $\vartheta \in W_{(0)}^{1,2}(I)$ . Under the assumption that  $\liminf_{\varepsilon \rightarrow 0} J_\varepsilon^{vK}(w_\varepsilon) < +\infty$ , by Lemma 2.1 we deduce that, up to subsequences,

$$\nabla_\varepsilon^2 w_\varepsilon \rightharpoonup \begin{pmatrix} w'' & \vartheta' \\ \vartheta' & \gamma \end{pmatrix} \text{ in } L^2(S; \mathbb{R}_{\text{sym}}^{2 \times 2})$$

for some  $\gamma \in L^2(S)$ . Since  $\det \nabla_\varepsilon^2 w_\varepsilon = 0$ , an application of Theorem 3.1 with  $Q(x, M) := Q_2(M)$  and  $\mathcal{B} = S$  shows that

$$\liminf_{\varepsilon \rightarrow 0} J_\varepsilon^{CvK}(w_\varepsilon) = \liminf_{\varepsilon \rightarrow 0} \frac{1}{24} \mathcal{F}(\nabla_\varepsilon^2 w_\varepsilon) \geq \frac{1}{24} \overline{\mathcal{F}} \begin{pmatrix} w'' & \vartheta' \\ \vartheta' & \gamma \end{pmatrix} \geq \frac{1}{24} \int_I \overline{Q}(w'', \vartheta') dx_1.$$

Therefore, we conclude that

$$\liminf_{\varepsilon \rightarrow 0} J_\varepsilon^{CvK}(w_\varepsilon) \geq J(w, \vartheta).$$

□

*Proof of Theorem 2.4–(ii).* Let  $w \in W_{(0)}^{2,2}(I)$  and  $\vartheta \in W_{(0)}^{1,2}(I)$ . We set

$$M := \begin{pmatrix} w'' & \vartheta' \\ \vartheta' & \gamma \end{pmatrix},$$

where  $\gamma \in L^2(I)$  is such that

$$\overline{Q}(w'', \vartheta') = Q_2(M) + \alpha^+(\det M)^+ + \alpha^-(\det M)^-.$$

As before, the fact that  $\gamma$  belongs to  $L^2(I)$  follows immediately by choosing  $M_0 = w''e_1 \otimes e_1 + \vartheta'(e_1 \otimes e_2 + e_2 \otimes e_1)$  as a competitor in the definition of  $\overline{Q}$  and by using the positive definiteness of  $Q_2$ .

By Theorem 3.1 with  $Q(x, M) := Q_2(M)$  and  $\mathcal{B} = I$ , there exist  $M^j \in L^2(I; \mathbb{R}_{\text{sym}}^{2 \times 2})$  with  $\det M^j = 0$  and such that  $M^j \rightharpoonup M$  weakly in  $L^2(I; \mathbb{R}_{\text{sym}}^{2 \times 2})$  and  $\mathcal{F}(M^j) \rightarrow \overline{\mathcal{F}}(M)$ , as  $j \rightarrow \infty$ . We can also assume that  $M^j \in C^\infty(\bar{I}; \mathbb{R}_{\text{sym}}^{2 \times 2})$ . The proof of this fact relies on a construction described in [9, Theorem 2.2–(ii)]. We give here full details for convenience of the reader. Suppose that  $(M_n)$  be a sequence of matrices with the same properties of  $(M^j)$  apart from the regularity, and denote by  $\lambda_n \in L^2(I)$  the trace of  $M_n$ . Since  $M_n$  is symmetric with  $\det M_n = 0$ , there exists  $\beta_n = \beta_n(x_1) \in (-\pi/2, \pi/2]$  such that

$$M_n = \begin{pmatrix} \cos \beta_n & -\sin \beta_n \\ \sin \beta_n & \cos \beta_n \end{pmatrix} \begin{pmatrix} \lambda_n & 0 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} \cos \beta_n & \sin \beta_n \\ -\sin \beta_n & \cos \beta_n \end{pmatrix},$$

and  $\beta_n$  is uniquely determined if  $\lambda_n \neq 0$ . When  $\lambda_n(x_1) = 0$ , we set  $\beta_n(x_1) = 0$ . We may assume without loss of generality that  $\lambda_n \in L^\infty(I)$ , possibly after truncating  $\lambda_n$  in modulus by  $n$ , while  $M_n$  still enjoys the same properties as before. We can find  $\lambda_{n,k} \in C^\infty(\bar{I})$  and  $\beta_{n,k} \in C^\infty(\bar{I}; (-\pi/2, \pi/2))$  such that, as  $k \rightarrow \infty$ ,  $\lambda_{n,k} \rightarrow \lambda_n$  and  $\beta_{n,k} \rightarrow \beta_n$  in  $L^p(I)$  for every  $p < +\infty$ . Set

$$M_{n,k} := \begin{pmatrix} \cos \beta_{n,k} & -\sin \beta_{n,k} \\ \sin \beta_{n,k} & \cos \beta_{n,k} \end{pmatrix} \begin{pmatrix} \lambda_{n,k} & 0 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} \cos \beta_{n,k} & \sin \beta_{n,k} \\ -\sin \beta_{n,k} & \cos \beta_{n,k} \end{pmatrix}.$$

Then,  $\det M_{n,k} = 0$  for every  $n, k$  and  $M_{n,k} \rightarrow M_n$  in  $L^2(I; \mathbb{R}_{\text{sym}}^{2 \times 2})$ , as  $k \rightarrow \infty$ .

Thus, by a diagonal argument, we may assume that there exist  $\lambda^j \in C^\infty(\bar{I})$  and  $\beta^j \in C^\infty(\bar{I})$  such that  $|\beta^j| < \pi/2$  on  $\bar{I}$ , and with

$$\begin{aligned} M^j &:= \begin{pmatrix} \cos \beta^j & -\sin \beta^j \\ \sin \beta^j & \cos \beta^j \end{pmatrix} \begin{pmatrix} \lambda^j & 0 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} \cos \beta^j & \sin \beta^j \\ -\sin \beta^j & \cos \beta^j \end{pmatrix} \\ &= \lambda^j \begin{pmatrix} \cos^2 \beta^j & \sin \beta^j \cos \beta^j \\ \sin \beta^j \cos \beta^j & \sin^2 \beta^j \end{pmatrix} \end{aligned}$$

we have that  $M^j \in C^\infty(\bar{I}; \mathbb{R}_{\text{sym}}^{2 \times 2})$ ,  $\det M^j = 0$  for every  $j$ ,  $M^j \rightharpoonup M$  in  $L^2(I; \mathbb{R}_{\text{sym}}^{2 \times 2})$ , and  $\mathcal{F}(M_j) \rightarrow \overline{\mathcal{F}}(M)$ , as  $j \rightarrow \infty$ .

For all  $j = 1, 2, \dots$  and all  $k, l \in \{1, 2\}$  we define  $\overline{M}_{kl}^j(x_1) := \int_0^{x_1} M_{kl}^j(s) ds$ ,

$$w^j(x_1) := \int_0^{x_1} (x_1 - s) M_{11}^j(s) ds - \frac{1}{\ell} \int_I \left( \int_0^t (t - s) M_{11}^j(s) ds \right) dt - x_1 \langle \overline{M}_{11}^j \rangle,$$

and

$$\vartheta^j(x_1) := \overline{M}_{12}^j(x_1) - \langle \overline{M}_{12}^j \rangle.$$

It is clear that  $w^j \rightharpoonup w$  weakly in  $W^{2,2}(I)$  and  $\vartheta^j \rightharpoonup \vartheta$  weakly in  $W^{1,2}(I)$ , as  $j \rightarrow \infty$ . Moreover,  $w^j \in W_{(0)}^{2,2}(I)$  and  $\vartheta^j \in W_{(0)}^{1,2}(I)$ .

After extending  $\beta^j$  smoothly to all of  $\mathbb{R}$ , still satisfying  $|\beta^j| < \pi/2$ , we define  $\alpha^j := \frac{\pi}{2} + \beta^j$ ,

$$\tilde{b}^j(\xi_1) := \cos \alpha^j(\xi_1) e_1 + \sin \alpha^j(\xi_1) e_2 \quad \text{and} \quad \Phi^j(\xi_1, \xi_2) := \xi_1 e_1 + \xi_2 \tilde{b}^j(\xi_1).$$

Observe that, by the definition of  $\tilde{b}^j$  and since  $(w^j)'' = M_{11}^j$  and  $(\vartheta^j)' = M_{12}^j$ ,

$$\begin{pmatrix} (w^j)'' \\ (\vartheta^j)' \end{pmatrix} \cdot \tilde{b}^j = 0. \quad (3.5)$$

Arguing as in [10, Lemma 12], we see that for every  $\varepsilon \leq \varepsilon^j$  the matrix  $\nabla \Phi^j(\xi_1, \xi_2)$  is invertible for  $|\xi_2| \leq \varepsilon$ , and the map  $(\Phi^j)^{-1} : S_\varepsilon \rightarrow \mathbb{R}^2$  is well defined. For such  $\varepsilon$  define  $z^j : S_\varepsilon \rightarrow \mathbb{R}$  by setting

$$z^j(\Phi^j(\xi_1, \xi_2)) = w^j(\xi_1) + \xi_2 \tilde{b}^j(\xi_1) \cdot \begin{pmatrix} (w^j)'(\xi_1) \\ \vartheta^j(\xi_1) \end{pmatrix}. \quad (3.6)$$

We clearly have

$$z^j(\cdot, 0) = w^j. \quad (3.7)$$

Moreover, taking derivatives in (3.6) and using (3.5), we obtain

$$\begin{aligned} \nabla z^j(\Phi^j)^T \nabla \Phi^j &= \left( (w^j)'(\xi_1) + \xi_2 (\tilde{b}^j)'(\xi_1) \cdot \begin{pmatrix} (w^j)'(\xi_1) \\ \vartheta^j(\xi_1) \end{pmatrix}, \tilde{b}^j(\xi_1) \cdot \begin{pmatrix} (w^j)'(\xi_1) \\ \vartheta^j(\xi_1) \end{pmatrix} \right) \\ &= \begin{pmatrix} (w^j)'(\xi_1) \\ \vartheta^j(\xi_1) \end{pmatrix}^T \nabla \Phi^j. \end{aligned}$$

Since  $\nabla \Phi^j$  is invertible for small  $|\xi_2|$ , we conclude that for small  $|\xi_2|$  and all  $\xi_1$

$$\nabla z^j(\Phi^j(\xi_1, \xi_2)) = \begin{pmatrix} (w^j)'(\xi_1) \\ \vartheta^j(\xi_1) \end{pmatrix}. \quad (3.8)$$

Taking the derivative with respect to  $\xi_2$ , we conclude that

$$\nabla^2 z^j(\Phi^j(\xi_1, \xi_2)) \tilde{b}^j(\xi_1) = 0.$$

In particular, the kernel is nontrivial, so  $\det \nabla^2 z^j = 0$  on  $S_\varepsilon$ . Since  $M^j \tilde{b}^j = 0$ , in particular we have that  $(\nabla^2 z^j(\cdot, 0) - M^j) \tilde{b}^j = 0$ . But from (3.7) we see that

$$e_1 \cdot (\nabla^2 z^j(\cdot, 0) - M^j) e_1 = \partial_{11}^2 z^j(\cdot, 0) - (w^j)'' = 0.$$

Since  $e_2 \cdot \tilde{b}^j \neq 0$  and since  $(\nabla^2 z^j(\cdot, 0) - M^j)$  is symmetric, we conclude that

$$\nabla^2 z^j(\cdot, 0) = M^j. \quad (3.9)$$

Finally, for  $\varepsilon$  small enough we define  $\tilde{w}_\varepsilon^j : S \rightarrow \mathbb{R}$  by  $\tilde{w}_\varepsilon^j(x_1, x_2) = z^j(x_1, \varepsilon x_2)$ . From (3.7) it follows immediately that  $\tilde{w}_\varepsilon^j \rightarrow w^j$  strongly in  $L^2(S)$ , as  $\varepsilon \rightarrow 0$ . Moreover, since

$$\nabla_\varepsilon \tilde{w}_\varepsilon^j(x) = \nabla z^j(x_1, \varepsilon x_2),$$

equation (3.8) implies that  $\nabla_\varepsilon \tilde{w}_\varepsilon^j \rightarrow ((w^j)', \vartheta^j)$  strongly in  $W^{1,2}(S; \mathbb{R}^2)$ , as  $\varepsilon \rightarrow 0$ . In particular, denoting by  $F_\varepsilon \in \mathbb{R}^2$  the average of  $\nabla_\varepsilon \tilde{w}_\varepsilon^j$  over  $S$ , we have

$$\lim_{\varepsilon \rightarrow 0} F_\varepsilon = \frac{1}{\ell} \int_I \begin{pmatrix} (w^j)' \\ \vartheta^j \end{pmatrix}(x_1) dx_1 = 0,$$

by definition of  $w^j$  and  $\vartheta^j$ . Similarly, denoting by  $c_\varepsilon$  the average of  $\tilde{w}_\varepsilon^j$  over  $S$ , we have  $c_\varepsilon \rightarrow 0$ . Hence the functions  $w_\varepsilon^j : S \rightarrow \mathbb{R}$  defined by

$$w_\varepsilon^j(x) := \tilde{w}_\varepsilon^j(x) - F_\varepsilon \cdot \begin{pmatrix} x_1 \\ \varepsilon x_2 \end{pmatrix} - c_\varepsilon,$$

still satisfy  $\tilde{w}_\varepsilon^j \rightarrow w^j$  strongly in  $L^2(S)$  and  $\nabla_\varepsilon \tilde{w}_\varepsilon^j \rightarrow ((w^j)', \vartheta^j)$  strongly in  $W^{1,2}(S; \mathbb{R}^2)$ . Moreover,  $w_\varepsilon^j \in W_{(0)}^{2,2}(S)$  by definition of  $F_\varepsilon$  and  $c_\varepsilon$ .

Finally, since

$$\nabla_\varepsilon^2 w_\varepsilon^j(x) = \nabla_\varepsilon^2 \tilde{w}_\varepsilon^j(x) = \nabla^2 z^j(x_1, \varepsilon x_2),$$

we have that  $w_\varepsilon^j \in W_{\det, \varepsilon}^{2,2}(S)$ . By (3.9) we deduce that  $\nabla_\varepsilon^2 w_\varepsilon^j \rightarrow M^j$  strongly in  $L^2(S; \mathbb{R}_{\text{sym}}^{2 \times 2})$ , as  $\varepsilon \rightarrow 0$ . Hence,

$$\lim_{\varepsilon \rightarrow 0} J_\varepsilon^{CvK}(w_\varepsilon^j) = \lim_{\varepsilon \rightarrow 0} \frac{1}{24} \int_S Q_2(\nabla_\varepsilon^2 w_\varepsilon^j(x)) dx = \frac{1}{24} \int_S Q_2(M^j(x)) dx = \frac{1}{24} \mathcal{F}(M^j).$$

Therefore, by taking diagonal sequences we obtain the desired maps.  $\square$

**Acknowledgements.** MGM acknowledges support by GNAMPA-INDAM under Project 2016 “Multiscale analysis of complex systems with variational methods” and by the ERC under Grant No. 290888 “Quasistatic and Dynamic Evolution Problems in Plasticity and Fracture”. PH acknowledges support by the DFG.

## REFERENCES

- [1] V. Agostiniani, A. DeSimone, K. Koumatos: Shape programming for narrow ribbons of nematic elastomers. *J. Elasticity*, doi:10.1007/s10659-016-9594-1.
- [2] S. Armon, E. Efrati, R. Kupferman, E. Sharon: Geometry and Mechanics in the Opening of Chiral Seed Pods. *Science* **333** (2011), 1726–1730.
- [3] S. Bartels, P. Hornung: Bending paper and the Möbius strip. *J. Elasticity* **119** (2015), 113–136.
- [4] Z. Chen: Geometric nonlinearity and mechanical anisotropy in strained helical nanoribbons. *Nanoscale* **6** (2014), 9443–9447.
- [5] J. Chopin, V. Démery, B. Davidovitch: Roadmap to the morphological instabilities of a stretched twisted ribbon. *J. Elasticity* **119** (2015), 137–189.
- [6] D.S. Chung, G.B. Benedek, F.M. Konikoff, J.M. Donovan: Elastic free energy of anisotropic helical ribbons as metastable intermediates in the crystallization of cholesterol. *Proc. Natl. Acad. Sci. USA* **90** (1993), 11341–11345.
- [7] M.A. Dias, B. Audoly: “Wunderlich, Meet Kirchhoff”: a general and unified description of elastic ribbons and thin rods. *J. Elasticity* **119** (2015), 49–66.
- [8] E. Efrati: Non-Euclidean ribbons: generalized Sadowsky functional for residually-stressed thin and narrow bodies. *J. Elasticity* **119** (2015), 251–261.
- [9] L. Freddi, P. Hornung, M.G. Mora, R. Paroni: A corrected Sadowsky functional for inextensible elastic ribbons. *J. Elasticity* **123** (2016), 125–136.
- [10] L. Freddi, P. Hornung, M.G. Mora, R. Paroni: A variational model for anisotropic and naturally twisted ribbons. *SIAM J. Math. Anal.* **48** (2016), 3883–3906.
- [11] L. Freddi, M.G. Mora, R. Paroni: Nonlinear thin-walled beams with a rectangular cross-section – Part I. *Math. Models Methods Appl. Sci.* **22** (2012), 1150016 (34 pp).
- [12] L. Freddi, M.G. Mora, R. Paroni: Nonlinear thin-walled beams with a rectangular cross-section – Part II. *Math. Models Methods Appl. Sci.* **23** (2013), 743–775.
- [13] G. Friesecke, R.D. James, S. Müller: A Hierarchy of Plate Models Derived from Nonlinear Elasticity by Gamma-Convergence. *Arch. Rational Mech. Anal.* **180** (2006), 183–236.
- [14] R. Fosdick and E. Fried: *The Mechanics of Ribbons and Möbius Bands*. Springer Netherlands, 2016.
- [15] D.F. Hinz, E. Fried: Translation of Michael Sadowsky’s paper “An elementary proof for the existence of a developable Möbius band and the attribution of the geometric problem to a variational problem”. *J. Elasticity* **119** (2015), 3–6.
- [16] P. Hornung: A remark on constrained von Kármán theories. *Proc. R. Soc. Lond. Ser. A Math. Phys. Eng. Sci.* **470** (2014), no. 2170, 20140346, 8 pp.
- [17] N.O. Kirby, E. Fried: Gamma-limit of a model for the elastic energy of an inextensible ribbon. *J. Elasticity* **119** (2015), 35–47.
- [18] O.O. Kit, T. Tallinen, L. Mahadevan, J. Timonen, P. Koskinen: Twisting graphene nanoribbons into carbon nanotubes. *Phys. Rev. B* **85** (2012), 085428.
- [19] H. Le Dret: *Problèmes variationnels dans les multi-domaines. Modélisation des jonctions et applications*. Recherches en Mathématiques Appliquées, **19**. Masson, Paris, 1991.

- [20] M. Sadowsky: Ein elementarer Beweis für die Existenz eines abwickelbaren Möbiuschen Bandes und die Zurückführung des geometrischen Problems auf ein Variationsproblem. *Sitzungsber. Preuss. Akad. Wiss.* 1930 – Mitteilung vom 26. Juni, 412–415.
- [21] Y. Sawa, F. Yeb, K. Urayamaa, T. Takigawaa, T. Gimenez-Pintob, R.L.B. Selingerb, J.V. Selingerb: Shape selection of twist-nematic-elastomer ribbons. *Proc. Natl. Acad. Sci. USA.* **108** 2011, 6364–6368.
- [22] E.L. Starostin, G.H.M. van der Heijden: Equilibrium shapes with stress localisation for inextensible elastic Möbius and other strips. *J. Elasticity* **119** (2015), 67–112.
- [23] R.E. Todres: Translation of W. Wunderlich’s “On a developable Möbius band”. *J. Elasticity* **119** (2015), 23–34.
- [24] T. von Kármán: Festigkeitsproblem im Maschinenbau, *Encyk. D. Math. Wiss. IV/4* (1910), 311–385.
- [25] K. Washizu: *Variational Methods in Elasticity and Plasticity*, Pergamon Press, New York, 2nd edition, 1975.
- [26] W. Wunderlich: Über ein abwickelbares Möbiusband. *Monatshefte für Mathematik* **66** (1962), 276–289.

(L. Freddi) DIPARTIMENTO DI MATEMATICA E INFORMATICA, VIA DELLE SCIENZE 206, 33100 UDINE, ITALY  
E-mail address: [lorenzo.freddi@uniud.it](mailto:lorenzo.freddi@uniud.it)

(P. Hornung) FACHRICHTUNG MATHEMATIK, TU DRESDEN, 01062 DRESDEN, GERMANY  
E-mail address: [peter.hornung@tu-dresden.de](mailto:peter.hornung@tu-dresden.de)

(M.G. Mora) DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI PAVIA, VIA FERRATA 1, 27100 PAVIA, ITALY  
E-mail address: [mariagiovanna.mora@unipv.it](mailto:mariagiovanna.mora@unipv.it)

(R. Paroni) DADU, UNIVERSITÀ DEGLI STUDI DI SASSARI, PALAZZO DEL POU SALIT, 07041 ALGHERO (SS), ITALY  
E-mail address: [paroni@uniss.it](mailto:paroni@uniss.it)