



UNIVERSITÀ
DEGLI STUDI
DI UDINE

Università degli studi di Udine

The open and clopen Ramsey theorems in the Weihrauch lattice

Original

Availability:

This version is available <http://hdl.handle.net/11390/1195039> since 2021-10-15T13:44:46Z

Publisher:

Published

DOI:10.1017/jsl.2021.10

Terms of use:

The institutional repository of the University of Udine (<http://air.uniud.it>) is provided by ARIC services. The aim is to enable open access to all the world.

Publisher copyright

(Article begins on next page)

THE OPEN AND CLOPEN RAMSEY THEOREMS IN THE WEIHRAUCH LATTICE

ALBERTO MARCONE AND MANLIO VALENTI

Abstract. We investigate the uniform computational content of the open and clopen Ramsey theorems in the Weihrauch lattice. While they are known to be equivalent to ATR_0 from the point of view of reverse mathematics, there is not a canonical way to phrase them as multivalued functions. We identify 8 different multivalued functions (5 corresponding to the open Ramsey theorem and 3 corresponding to the clopen Ramsey theorem) and study their degree from the point of view of Weihrauch, strong Weihrauch and arithmetic Weihrauch reducibility. In particular one of our functions turns out to be strictly stronger than any previously studied multivalued functions arising from statements around ATR_0 .

CONTENTS

1. Introduction	2
1.1. Structure of the paper	3
2. Background	4
2.1. Represented spaces	5
2.2. Weihrauch reducibility	6
3. Ramsey theorems	9
3.1. Some useful tools	9
4. Ramsey theorems in the Weihrauch lattice	14
4.1. Definitions	15
4.2. Problems reducible to $\text{UC}_{\mathbb{N}^{\mathbb{N}}}$	16
4.3. Problems reducible to $\text{C}_{\mathbb{N}^{\mathbb{N}}}$	18
4.4. Problems not reducible to $\text{C}_{\mathbb{N}^{\mathbb{N}}}$	21
4.5. A 0 – 1 law for strong Weihrauch reducibility	28
5. Arithmetic Weihrauch reducibility	29

1991 *Mathematics Subject Classification*. Primary: 03D78; Secondary: 03D30, 03B30, 05C55.

Key words and phrases. Computable analysis, Weihrauch degrees, Homogeneous sets, Open and clopen Ramsey theorem.

We thank Arno Pauly, Vasco Brattka, Jun Le Goh and Eleonora Pippia for useful discussions about the topic of the paper.

We also thank the anonymous referee for his/her careful reading of the paper and many valuable suggestions that improved the presentation. In particular (s)he noticed that the proof of Theorem 4.28 yielded a stronger result than the one we originally stated.

Both author’s research was partially supported by the departmental PRID funding “HiWei — The higher levels of the Weihrauch hierarchy” and by the Italian PRIN 2017 Grant “Mathematical Logic: models, sets, computability”.

§1. Introduction. This work explores the uniform computational strength of some infinite-dimensional generalizations of Ramsey’s theorem. The classical finite-dimensional Ramsey theorem can be stated as follows: for every $A \subset \mathbb{N}$ let $[A]^n := \{B \subset A : |B| = n\}$ be the set of subsets of A with cardinality n . A map $c: [\mathbb{N}]^n \rightarrow k$ is called a k -coloring of $[\mathbb{N}]^n$. A set H s.t. $c([H]^n) = \{i\}$ for some $i < k$ is called *homogeneous* for c .

THEOREM 1.1 (Ramsey’s theorem). *For every $n, k \geq 1$ and every coloring $c: [\mathbb{N}]^n \rightarrow k$ there is an infinite subset $H \subset \mathbb{N}$ that is homogeneous for c .*

There is a vast literature exploring the computational aspects of Ramsey’s theorem. Our focus will be on the infinite generalization of the above result, i.e. we will consider $n = \infty$. In particular we will focus on Nash-Williams’ theorem, also called the open Ramsey theorem:

THEOREM 1.2 (Nash-Williams [20]). *The open subsets of $[\mathbb{N}]^{\mathbb{N}}$ admit infinite homogeneous sets.*

We will also consider the restriction of Nash-Williams’ theorem to clopen subsets of $[\mathbb{N}]^{\mathbb{N}}$.

The notion of Weihrauch reducibility provides a useful tool to explore the uniform computational content of theorems: indeed many statements in mathematics can be written in the form $(\forall x \in X)(\varphi(x) \rightarrow (\exists y \in Y)(\psi(x, y)))$, which can be thought of as a computational problem where x is an instance of a problem and the goal is to find a solution y (which is, in general, not unique). This can be naturally written in the language of partial multivalued functions by considering $f: \subseteq X \rightrightarrows Y$ s.t. $f(x) = \{y \in Y : \psi(x, y)\}$, for every $x \in X$ s.t. $\varphi(x)$.

Starting with the work of Gherardi and Marcone [11], it has been shown that Weihrauch reducibility provides a bridge between computable analysis and reverse mathematics. In particular reverse mathematics focuses on proving the equivalence of statements over a weak base theory. Historically, a large number of theorems turned to be equivalent to one of the so-called *big five*: RCA_0 , WKL_0 , ACA_0 , ATR_0 , $\Pi_1^1\text{-CA}_0$. Oftentimes, the close connection between reverse mathematics and Weihrauch reducibility has been exploited to translate results from one setting into the other.

Following this connection, there are a number of established “analogues” of the big-five in the Weihrauch lattice: RCA_0 corresponds to computable problems, WKL_0 corresponds to $\text{C}_{2^{\mathbb{N}}}$ (i.e. choosing an element from a non-empty closed subset of $2^{\mathbb{N}}$) and ACA_0 corresponds to iterations of lim .

Recently Marcone [8] raised the question “What do the Weihrauch hierarchies look like once we go to very high levels of reverse mathematics strength?”. There have been several works in this direction: Kihara, Marcone and Pauly [18] have studied several principles, like the (strong) comparability of well-orders, the perfect tree theorem and the open determinacy theorem; Goh [13, 12] analyzed

the weak comparability of well-orders and the König duality theorem; Anglès D'Auriac and Kihara [1] dealt with the Σ_1^1 choice on \mathbb{N} and variants thereof.

Our work explores the formalization of the open and clopen Ramsey theorems as multivalued functions and analyzes their position in the Weihrauch lattice (both the open and the clopen Ramsey theorems are known to be equivalent to ATR_0 over RCA_0 , see [26, Sec. V.9]). Notice that, as already occurred to other principles equivalent to ATR_0 ([18, 12]), there is not a single multivalued function corresponding to the open Ramsey theorem. Actually, in our case, the situation is even more complex than for the open determinacy or the perfect tree theorem, as the two alternatives (homogeneous solution on the open side or homogeneous solution on the closed side) given by the open Ramsey theorem are not mutually exclusive. Therefore given an open set we can ask for a homogeneous solution on the open side, a homogeneous solution on the closed side or a homogeneous solution on either side. Altogether we will define five different multivalued functions corresponding to the open Ramsey theorem and three different functions corresponding to the clopen Ramsey theorem.

While it is hard to point out a single analogue of ATR_0 in the Weihrauch lattice, there are a number of degrees that play a central role in calibrating the strength of principles that, from the point of view of reverse mathematics, lie at the level of ATR_0 . Prominent among these are $\text{UC}_{\mathbb{N}^{\mathbb{N}}}$, $\text{C}_{\mathbb{N}^{\mathbb{N}}}$ and $\text{TC}_{\mathbb{N}^{\mathbb{N}}}$ (see Section 2.2 for their precise definitions). It is known that $\text{UC}_{\mathbb{N}^{\mathbb{N}}} <_{\text{W}} \text{C}_{\mathbb{N}^{\mathbb{N}}} <_{\text{W}} \text{TC}_{\mathbb{N}^{\mathbb{N}}}$ (see [18]).

When dealing with multivalued functions that are very high in the Weihrauch lattice it is often appropriate to use arithmetic Weihrauch reducibility rather than (computable) Weihrauch reducibility.

In Figure 1 we summarize the results we obtain both with respect to Weihrauch reducibility and arithmetic Weihrauch reducibility. Notice that the multivalued function $\text{FindHS}_{\Sigma_1^0}$ is stronger than any multivalued function related to ATR_0 considered so far. In fact all these functions are strictly Weihrauch reducible to $\text{TC}_{\mathbb{N}^{\mathbb{N}}}^*$, which, by Corollary 4.27, is strictly below $\text{FindHS}_{\Sigma_1^0}$. Notice also that, since $\text{FindHS}_{\Sigma_1^0}$ is closed under parallel product (Proposition 4.24), it computes $\text{sTC}_{\mathbb{N}^{\mathbb{N}}}^* \equiv_{\text{W}} \text{TC}_{\mathbb{N}^{\mathbb{N}}}^* \times \chi_{\Pi_1^1}^*$, which was suggested as an ATR_0 analogue in [18, Sec. 9].

1.1. Structure of the paper. In Section 2 we briefly introduce the notation and recall the preliminary notions on represented spaces (Section 2.1) and Weihrauch reducibility (Section 2.2). In Section 3 we will recall the precise statement for the open and clopen Ramsey theorems and prove some lemmas that will be useful in proving the results on the Weihrauch degrees. The reader may skip these lemmas on the first read, and return to it as needed. In Section 4 we define the multivalued functions corresponding to the open and clopen Ramsey theorems and study their degrees. In particular we divide the analysis into: functions that are reducible to $\text{UC}_{\mathbb{N}^{\mathbb{N}}}$ (Section 4.2), functions that are reducible to $\text{C}_{\mathbb{N}^{\mathbb{N}}}$ (but not to $\text{UC}_{\mathbb{N}^{\mathbb{N}}}$, Section 4.3) and functions that are not reducible to $\text{C}_{\mathbb{N}^{\mathbb{N}}}$ (Section 4.4). Moreover, in Section 4.5 we characterize the strength of these functions from the point of view of strong Weihrauch reducibility. Finally, in Section 5 we




FIGURE 1. Multivalued functions related to the open and clopen Ramsey theorems in the Weihrauch lattice. Dashed arrows represent Weihrauch reducibility in the direction of the arrow, solid arrows represent strict Weihrauch reducibility. The large rectangles indicate arithmetic Weihrauch equivalence classes. In particular, every function strictly arithmetically reduces to all the functions in rectangles above its own.

focus on the behavior of these functions under arithmetic Weihrauch reducibility and in Section 6 we draw some conclusions and list some open problems.

§2. Background. We refer the reader to [26] for a detailed introduction to reverse mathematics.

Here we introduce the notation we will use in the rest of the paper.

Let $\mathbb{N}^{\mathbb{N}}$ be the Baire space endowed with the usual product topology and let $\mathbb{N}^{<\mathbb{N}}$ be the set of finite sequences of natural numbers. Let also $[\mathbb{N}]^{\mathbb{N}}$ be the space of total functions $f: \mathbb{N} \rightarrow \mathbb{N}$ that are strictly increasing (i.e. $n < m \Rightarrow f(n) < f(m)$). This is sometimes called *Ramsey space* (see e.g. [26, Sec. V.9]). We will use the symbol \sqsubseteq for the prefix relation. For $\sigma, \tau \in \mathbb{N}^{<\mathbb{N}}$ we write: $|\sigma|$ for the length of σ , $\sigma \frown \tau$ for the concatenation of σ and τ , and $\sigma \leq \tau$ for the domination relation, i.e. $|\sigma| = |\tau|$ and $(\forall i < |\sigma|)(\sigma(i) \leq \tau(i))$. The domination

relation applies also to infinite strings. Also if $x \in \mathbb{N}^{<\mathbb{N}} \cup \mathbb{N}^{\mathbb{N}}$ we write $x[m]$ for the prefix of x of length m . If $f, g \in \mathbb{N}^{\mathbb{N}}$ we denote the composition $f \circ g$ by fg . Moreover, we write $f \preceq g$ if f is a subsequence of g , i.e. if there exists $h \in [\mathbb{N}]^{\mathbb{N}}$ s.t. $f = gh$. In particular for every $h \in [\mathbb{N}]^{\mathbb{N}}$ we have $gh \preceq g$. Similarly, we write $\sigma \preceq^* f$ if σ is a finite subsequence of f .

2.1. Represented spaces. We briefly introduce the main concepts on the theory of represented spaces. For a more thorough presentation we refer to [21, 28].

A *represented space* is a pair (X, δ_X) where X is a set and $\delta_X : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow X$ is a partial surjection. For every $x \in X$ a *name* or *code* for x is any element of $\delta_X^{-1}(x)$. When it is clear from the context, we omit to explicitly mention the representation map δ_X .

Given two representation maps δ and δ' of the same set X , we say that δ is *reducible* to δ' (and we write $\delta \leq_c \delta'$) if there is a computable map $F : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ s.t. $\delta(p) = \delta'(F(p))$ for every $p \in \text{dom}(\delta)$. The maps δ and δ' are called *equivalent* if $\delta \leq_c \delta'$ and $\delta' \leq_c \delta$.

Let (X, δ_X) and (Y, δ_Y) be represented spaces and $f : \subseteq X \rightrightarrows Y$ be a partial multivalued function. We say that a partial function $F : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ is a *realizer*¹ for f (and we write $F \vdash f$) if, for every $p \in \text{dom}(f \circ \delta_X)$ we have that $\delta_Y(F(p)) \in f(\delta_X(p))$. The notion of realizer allows us to transfer properties of functions on the Baire space (such as computability or continuity) to multivalued functions on represented spaces. In particular we say that a multivalued function between represented spaces is computable if it has a computable realizer.

Let (X, d) be a separable metric space and $\alpha : \mathbb{N} \rightarrow X$ be a dense sequence in X . The tuple (X, d, α) is called a *computable metric space* if $d \circ (\alpha \times \alpha) : \mathbb{N}^2 \rightarrow \mathbb{R}$ is a computable double sequence in \mathbb{R} .

Fix a computable enumeration $(q_n)_{n \in \mathbb{N}}$ of \mathbb{Q} . If (X, d, α) is a separable metric space, for every $k \geq 1$ we can define the represented spaces $(\Sigma_k^0(X), \delta_{\Sigma_k^0(X)})$, $(\Pi_k^0(X), \delta_{\Pi_k^0(X)})$, $(\Delta_k^0(X), \delta_{\Delta_k^0(X)})$ inductively by:

- $\delta_{\Sigma_k^0(X)}(p) := \bigcup_{\langle i, j \rangle \in \text{ran}(p)} B(\alpha(i), q_j)$;
- $\delta_{\Pi_k^0(X)}(p) := X \setminus \delta_{\Sigma_k^0(X)}(p)$;
- $\delta_{\Sigma_{k+1}^0(X)}(\langle p_0, p_1, \dots \rangle) := \bigcup_{i \in \mathbb{N}} \delta_{\Pi_k^0(X)}(p_i)$;
- $\delta_{\Delta_k^0(X)}(\langle p, q \rangle) := \delta_{\Sigma_k^0(X)}(p)$, iff $p \in \text{dom}(\delta_{\Sigma_k^0(X)})$, $q \in \text{dom}(\delta_{\Pi_k^0(X)})$ and $\delta_{\Sigma_k^0(X)}(p) = X \setminus \delta_{\Pi_k^0(X)}(q)$.

The represented spaces $(\Sigma_k^0(X), \delta_{\Sigma_k^0(X)})$, $(\Pi_k^0(X), \delta_{\Pi_k^0(X)})$ can be defined for any represented space (X, δ_X) , using the fact that the category of represented spaces is cartesian-closed and letting a name for an open set be the name of its characteristic function, where the codomain $\{0, 1\}$ is equipped with the Sierpiński topology. For separable metric spaces, the two representations are equivalent (see [21, 3]).

LEMMA 2.1. *The following maps are computable:*

1. $\Delta_1^0([\mathbb{N}]^{\mathbb{N}}) \hookrightarrow \Sigma_1^0([\mathbb{N}]^{\mathbb{N}}) := D \mapsto D$;
2. $\Delta_1^0([\mathbb{N}]^{\mathbb{N}}) \rightarrow \Delta_1^0([\mathbb{N}]^{\mathbb{N}}) := D \mapsto [\mathbb{N}]^{\mathbb{N}} \setminus D$;

¹The very existence of realizers for every f depends on some form of the axiom of choice.

$$3. \cup: \Sigma_1^0(\mathbb{N}^{\mathbb{N}}) \times \Sigma_1^0(\mathbb{N}^{\mathbb{N}}) \rightarrow \Sigma_1^0(\mathbb{N}^{\mathbb{N}}) := (P, Q) \mapsto P \cup Q.$$

PROOF.

- 1, 2:** follow from the fact that a name for a clopen set is the join $\langle p, q \rangle$ of two names for open sets (one for the set and one for its complement);
3: see [3, Prop. 3.2(5)].

□

The set $\Sigma_1^1(X)$ of analytic subsets of X can be seen as a represented space by defining a name for S to be a name for a closed set $A \subset X \times \mathbb{N}^{\mathbb{N}}$ s.t. $S = \pi_X(A)$ (where π denotes projection). Moreover, we can define a name for a coanalytic set $R \in \Pi_1^1(X)$ to be a name for its complement.

We denote with \mathbf{Tr} the space of trees on \mathbb{N} represented via their characteristic function. Similarly we denote with \mathbf{Ti} the space of trees with strictly increasing strings, represented analogously. The function $[\cdot]: \mathbf{Tr} \rightarrow \Pi_1^0(\mathbb{N}^{\mathbb{N}})$ that maps a tree to the set of its paths is computable with multivalued computable inverse. This implies that a closed set A of $\mathbb{N}^{\mathbb{N}}$ or $[\mathbb{N}]^{\mathbb{N}}$ can be equivalently represented via the characteristic function of a tree T s.t. $[T] = A$.

Similarly, an open set P of $\mathbb{N}^{\mathbb{N}}$ can be equivalently represented via an enumeration p of a prefix-free subset of $\mathbb{N}^{<\mathbb{N}}$ s.t. $P = \{f \in \mathbb{N}^{\mathbb{N}} : (\exists i)(p(i) \sqsubset f)\}$. With a small abuse of notation we write $\tau \in p$ in place of $\tau \in \text{ran}(p)$. The same considerations can be made for the space $[\mathbb{N}]^{\mathbb{N}}$.

We denote by $\text{LO} = (\text{LO}, \delta_{\text{LO}})$ the represented space of countable linear orders, where an order L is represented by the characteristic function of the relation $\{\langle a, b \rangle : a \leq_L b\}$. We also denote by $\text{WO} = (\text{WO}, \delta_{\text{WO}})$ the represented space of countable well-orders, where the representation map δ_{WO} is the restriction of δ_{LO} to codes of well-orders.

For every tree $T \subset \mathbb{N}^{<\mathbb{N}}$ we denote by $\text{KB}(T)$ the Kleene-Brouwer order on T , defined as $\sigma \leq_{\text{KB}(T)} \tau$ iff $\sigma, \tau \in T$ and $\tau \sqsubseteq \sigma$ or $\sigma \leq_{\text{lex}} \tau$. The map $T \mapsto \text{KB}(T)$ from \mathbf{Tr} to LO is computable. It is known that $\text{KB}(T)$ is a well-order iff $[T] = \emptyset$ (see e.g. [26, Lem. V.1.3]).

2.2. Weihrauch reducibility. Let $f : \subseteq X \rightrightarrows Y$ and $g : \subseteq Z \rightrightarrows W$ be multivalued functions between represented spaces. We say that f is *Weihrauch reducible* to g and we write $f \leq_W g$ iff there are two computable maps $\Phi, \Psi : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ s.t., for every realizer $G \vdash g$ we have $p \mapsto \Psi(\langle p, G\Phi(p) \rangle)$ is a realizer for f . We say that f is *strongly Weihrauch reducible* to g , and we write $f \leq_{\text{sw}} g$, if there are two computable maps $\Phi, \Psi : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ s.t., for every realizer $G \vdash g$ we have $\Psi G \Phi \vdash f$. We often say that Φ is the *forward* functional of the (strong) Weihrauch reduction, while Ψ is the *backward* functional.

The relations \leq_W and \leq_{sw} are reflexive and transitive, and therefore induce two degree structures on the family of multivalued functions on represented spaces. For more details on the algebraic properties of the Weihrauch and strong Weihrauch degrees we refer the reader to [16, 9, 7].

In the following we briefly introduce the operations on multivalued functions that we will need in this work:

parallel product: $f \times g : \subseteq X \times Z \rightrightarrows Y \times W := (x, z) \mapsto f(x) \times g(z)$ with $\text{dom}(f \times g) := \text{dom}(f) \times \text{dom}(g)$;

finite parallelization: $f^* : \subseteq X^{<\mathbb{N}} \rightrightarrows Y^{<\mathbb{N}}$ with domain $\text{dom}(f^*) := \{(x_i)_{i < n} : (\forall i < n)(x_i \in \text{dom}(f))\}$ defined as $f^*((x_i)_{i < n}) := \{(y_i)_{i < n} : (\forall i < n)(y_i \in f(x_i))\}$;

parallelization: $\hat{f} : \subseteq X^{\mathbb{N}} \rightrightarrows Y^{\mathbb{N}}$ with domain $\text{dom}(\hat{f}) := \text{dom}(f)^{\mathbb{N}}$ defined as $\hat{f}((x_n)_{n \in \mathbb{N}}) := (f(x_n))_{n \in \mathbb{N}}$;

jump: we define the *jump* of the represented space (X, δ_X) as the represented space $X' = (X, \delta_{X'})$, where a $\delta_{X'}$ -name for x is a string $\langle p_0, p_1, \dots \rangle$ s.t. $\delta_X(\lim_{n \rightarrow \infty} p_n) = x$. The jump of f is defined as $f' : \subseteq X' \rightrightarrows Y := x \mapsto f(x)$. We write $f^{(n)}$ to denote the result of applying the jump operation n times.

We also define the *compositional product* as

$$f * g := \max_{\leq_W} \{f_0 \circ g_0 : f_0 \leq_W f \text{ and } g_0 \leq_W g\}.$$

Intuitively the compositional product captures the idea of applying first g , then using some computable operation on the output of g and then feeding the result as an input for f . The fact that the compositional product is well-defined has been shown in [9, Cor. 3.7]. Notice that, by definition, $f * g$ is a Weihrauch degree and not a specific multivalued function. Nonetheless, with a small abuse of notation, we will write formulas such as $h \leq_W f * g$ with the obvious meaning. We also write $f^{[n]}$ to denote the n -fold compositional product of f with itself, where $f^{[0]} := \text{id}$ and $f^{[1]} := f$.

We say that f is a *cylinder* if $f \equiv_{\text{sw}} \text{id} \times f$, where id denotes the identity on the Baire space. The notion of cylinder is very useful because of the following

PROPOSITION 2.2 ([5, Cor. 3.6]). *For every multivalued functions f and g , if g is a cylinder then $f \leq_W g \iff f \leq_{\text{sw}} g$.*

Cylinders are also useful to deal with the compositional product, thanks to the so-called *cylindrical decomposition*:

PROPOSITION 2.3 ([9, Lem. 3.10]). *For all f, g and all cylinders F, G with $F \equiv_W f$ and $G \equiv_W g$ there exists a computable K such that $f * g \equiv_W F \circ K \circ G$.*

In particular, knowing that for every function f we have that $f \equiv_W \text{id} \times f$ and that the latter is a cylinder, we can always take a representative of $f * g$ of the form $(\text{id} \times f) \circ \Phi_e \circ (\text{id} \times g)$ for some computable function Φ_e .

Finally we define the *total continuation* or *totalization* of f , written $\mathbb{T}f$, as the total multivalued function $\mathbb{T}f(x) := f(x)$ if $x \in \text{dom}(f)$ and $\mathbb{T}f(x) = Y$ otherwise. Clearly $\mathbb{T}f = f$ iff f is total. Notice that the definition of $\mathbb{T}f$ is sensitive to the particular definition of f as a multivalued function between represented spaces. For a more detailed exposition we refer to [6].

Let us now define some well-known multivalued functions that will be useful in the development of the work:

- **LPO:** $\mathbb{N}^{\mathbb{N}} \rightarrow \{0, 1\}$ is defined as $\text{LPO}(p) := 1$ iff $(\exists n)(p(n) = 0)$. It is often convenient to think of LPO as the problem of finding a yes/no answer to a $\Sigma_1^{0,p}$ or $\Pi_1^{0,p}$ question.
- **lim:** $\subseteq (\mathbb{N}^{\mathbb{N}})^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}} := (p_n)_{n \in \mathbb{N}} \mapsto \lim_{n \rightarrow \infty} p_n$, where the domain of lim is the set of Cauchy sequences.

- $\mathbf{NHA} : \mathbb{N}^{\mathbb{N}} \rightrightarrows \mathbb{N}^{\mathbb{N}}$ is defined as $\mathbf{NHA}(p) = \{q : q \text{ is not hyperarithmic in } p\}$.
- $\chi_{\Pi_1^1} : \mathbb{N}^{\mathbb{N}} \rightarrow \{0, 1\}$ is the characteristic function of a Π_1^1 -complete set. It is often convenient to think of $\chi_{\Pi_1^1}$ as the function that takes in input a tree on \mathbb{N} and checks whether it is well-founded.

A central role is played by the choice problems: given a represented space X we define $\mathbf{\Gamma}\text{-}\mathbf{C}_X : \subseteq \mathbf{\Gamma}(X) \rightrightarrows X$ as the multivalued function that chooses an element from a non-empty set $A \in \mathbf{\Gamma}(X)$. If $\mathbf{\Gamma} = \mathbf{\Pi}_1^0$ we simply write \mathbf{C}_X . We also write $\mathbf{\Gamma}\text{-}\mathbf{UC}_X$ if the choice is restricted to singletons.

Three choice problems that turned out to be very relevant to calibrate the strength of multivalued functions that corresponds to theorems around ATR_0 (from the point of view of reverse mathematics) are $\mathbf{UC}_{\mathbb{N}^{\mathbb{N}}}$, $\mathbf{C}_{\mathbb{N}^{\mathbb{N}}}$ and $\mathbf{TC}_{\mathbb{N}^{\mathbb{N}}}$. They have been explored in great detail in [18]. It is known that $\lim^{(n)} <_{\mathbf{W}} \mathbf{UC}_{\mathbb{N}^{\mathbb{N}}}$ for every n (see [4, Sec. 6]) and indeed $\lim^{\dagger} \equiv_{\mathbf{W}} \mathbf{UC}_{\mathbb{N}^{\mathbb{N}}}$, where \lim^{\dagger} corresponds to the iteration of \lim over a countable ordinal ([22]). Moreover $\mathbf{UC}_{\mathbb{N}^{\mathbb{N}}}$ and $\mathbf{C}_{\mathbb{N}^{\mathbb{N}}}$ are closed under compositional product ([4, Thm. 7.3]). In [18] it is proved that $\Sigma_1^1\text{-}\mathbf{UC}_{\mathbb{N}^{\mathbb{N}}} \equiv_{\mathbf{W}} \mathbf{UC}_{\mathbb{N}^{\mathbb{N}}}$ and $\Sigma_1^1\text{-}\mathbf{C}_{\mathbb{N}^{\mathbb{N}}} \equiv_{\mathbf{W}} \mathbf{C}_{\mathbb{N}^{\mathbb{N}}}$. The fact that $\mathbf{UC}_{\mathbb{N}^{\mathbb{N}}} <_{\mathbf{W}} \mathbf{C}_{\mathbb{N}^{\mathbb{N}}}$ follows from the fact that the element of a Σ_1^1 singleton is hyperarithmic, but the hyperarithmic functions are not a basis for the Π_1^0 predicates (see [24, Thm. I.1.6 and thm. III.1.1]). In particular we have

THEOREM 2.4 ([18, Cor. 3.4]). *Let $f : \subseteq \mathbb{N}^{\mathbb{N}} \rightrightarrows X$ be a (partial) multivalued function, for some represented space X . If $f \leq_{\mathbf{W}} \mathbf{UC}_{\mathbb{N}^{\mathbb{N}}}$ then, for every $x \in \text{dom}(f)$, $f(x)$ contains some y hyperarithmic relative to x .*

The Weihrauch degree of $\mathbf{TC}_{\mathbb{N}^{\mathbb{N}}}$ has been explored in [18, Sec. 8]. It is known that $\mathbf{C}_{\mathbb{N}^{\mathbb{N}}} <_{\mathbf{W}} \mathbf{TC}_{\mathbb{N}^{\mathbb{N}}}$ ([18, Prop. 8.2(1)]) and $\mathbf{TC}_{\mathbb{N}^{\mathbb{N}}}^*$ is one of the strongest problem studied so far that is still considered among the “ ATR_0 analogues”.

We conclude this section with the following proposition.

PROPOSITION 2.5. *Let $f : \subseteq X \rightrightarrows Y$ and $g : \subseteq Z \rightrightarrows W$ be multivalued functions between represented spaces and let $A \subset \text{dom}(g)$ be s.t.*

$$\{z \in \text{dom}(g) : (\forall w \in g(z))(w \text{ is not hyperarithmic in } z)\} \subset A.$$

If $f \times \mathbf{NHA} \leq_{\mathbf{W}} g$ then $f \leq_{\mathbf{W}} g|_A$.

PROOF. Assume $f \times \mathbf{NHA} \leq_{\mathbf{W}} g$ and let the reduction be witnessed by the computable functions Φ, Ψ . For every p_x which is the name of some $x \in \text{dom}(f)$, the pair (p_x, p_x) is mapped via Φ to a name p_z for some element $z \in \text{dom}(g)$.

It suffices to show that for every name p_x of an element in the domain of f , the pair (p_x, p_x) is mapped via Φ to a name p_z of an element $z \in A$.

If this were not the case then, for some $x \in \text{dom}(f)$, p_z is the name of some $z \notin A$. By hypothesis, there is a $w \in g(z)$ s.t. w has a name p_w which is hyperarithmic in p_z . Let G be a realizer of g s.t. $p_w = G\Phi(p_x, p_x)$. Since p_w is hyperarithmic in p_z , and hence in p_x , we have reached a contradiction with the fact that $\Psi(p_w, p_x, p_x)$ computes a solution for $\mathbf{NHA}(p_x)$. \square

This result will often be used in combination with Theorem 2.4. In fact if there is a computable $x \in \text{dom}(f)$ s.t. $f(x)$ does not contain any hyperarithmic element, then $f \not\leq_{\mathbf{W}} \mathbf{UC}_{\mathbb{N}^{\mathbb{N}}}$.

§3. Ramsey theorems. The space $[\mathbb{N}]^{\mathbb{N}}$, endowed with the induced topology from the Baire space $\mathbb{N}^{\mathbb{N}}$ is computably isometric to $\mathbb{N}^{\mathbb{N}}$. There is actually a canonical choice for a computable bijection $\mathbb{N}^{\mathbb{N}} \rightarrow [\mathbb{N}]^{\mathbb{N}}$.

Let $P \subset [\mathbb{N}]^{\mathbb{N}}$. We say that f is a *homogeneous solution for P* iff

$$(\forall g \in [\mathbb{N}]^{\mathbb{N}})(fg \in P) \vee (\forall g \in [\mathbb{N}]^{\mathbb{N}})(fg \notin P).$$

If f is homogeneous for P we say that f *lands in P* if the first disjunct holds, i.e. if $(\forall g \in [\mathbb{N}]^{\mathbb{N}})(fg \in P)$. Vice versa, if $(\forall g \in [\mathbb{N}]^{\mathbb{N}})(fg \notin P)$ then we say that f *avoids P* . A set $P \subset [\mathbb{N}]^{\mathbb{N}}$ is called *Ramsey* (or we say that it has the *Ramsey property*) iff it has a homogeneous solution. We will denote the set of homogeneous solutions for P (which may either land in it or avoid it) with $\text{HS}(P)$. Notice that, in general, a set can have both solutions that land in the set and solutions that avoid the set.

In the literature, the symbol $[\mathbb{N}]^{\mathbb{N}}$ is sometimes used to denote the family of all infinite subsets of \mathbb{N} . Also, if X is an infinite subset of \mathbb{N} , $[X]^{\mathbb{N}}$ denotes the family of all infinite subsets of X . It is easy to identify the Ramsey space $[\mathbb{N}]^{\mathbb{N}}$ with the space of infinite subsets of \mathbb{N} (by identifying a function f with its range). With this in mind, we may write $[f]^{\mathbb{N}} := [\text{ran}(f)]^{\mathbb{N}}$ to denote the set of all infinite subsequences of f . The definition of homogeneous solution can now be written as

$$[f]^{\mathbb{N}} \subset P \vee [f]^{\mathbb{N}} \cap P = \emptyset.$$

It is natural to ask which classes of subsets of $[\mathbb{N}]^{\mathbb{N}}$ have the Ramsey property. The problem is well studied and has an extensive literature. The Galvin-Prikry theorem ([10]) states that all Borel subsets of $[\mathbb{N}]^{\mathbb{N}}$ have the Ramsey property. This result can actually be extended to analytic sets ([25]). To go beyond the analytic sets we need axioms above ZFC (see e.g. [26, Rem. VI.7.6, p. 240], [17, pp. 1036–1037]). We will focus on Nash-Williams' theorem ([20]), which states that open sets have the Ramsey property. This is also known as the open Ramsey theorem. It, in turn, implies the clopen Ramsey theorem (which is the restriction of Nash-Williams' theorem to clopen sets). As already mentioned, the open and clopen Ramsey theorems are known to be equivalent to ATR_0 over RCA_0 (see [26, Thm. V.9.7]).

3.1. Some useful tools. Before formalizing the open and clopen Ramsey theorems in the context of Weihrauch reducibility as multivalued functions, let us explicitly state some properties of the set of homogeneous solutions that will turn out to be useful in the rest of the paper. As a notational convenience we will use the letters P, Q, \dots to denote open sets and D, E, \dots to denote clopen sets.

PROPOSITION 3.1. *Let Γ be a definable (boldface) pointclass that is downward closed with respect to Wadge reducibility (i.e. it is a downward closed family of Wadge degrees), such as the families of open and of clopen sets. Assume that every $P \in \Gamma([\mathbb{N}]^{\mathbb{N}})$ is Ramsey and that for every $f \in [\mathbb{N}]^{\mathbb{N}}$,*

$$\Gamma([f]^{\mathbb{N}}) = \{P \cap [f]^{\mathbb{N}} : P \in \Gamma([\mathbb{N}]^{\mathbb{N}})\}.$$

Then for every $f \in [\mathbb{N}]^{\mathbb{N}}$, every $Q \in \mathbf{\Gamma}([f]^{\mathbb{N}})$ is Ramsey. Moreover if $P \in \mathbf{\Gamma}([\mathbb{N}]^{\mathbb{N}})$ and $f \in [\mathbb{N}]^{\mathbb{N}}$ there exists $h \in \text{HS}(P)$ s.t. $h \preceq f$.

PROOF. It is easy to see that every $f \in [\mathbb{N}]^{\mathbb{N}}$ induces a f -computable homeomorphism $\varphi_f: [\mathbb{N}]^{\mathbb{N}} \rightarrow [f]^{\mathbb{N}}$ defined as

$$\varphi_f(p) := n \mapsto f(p(n)).$$

Notice also that

$$(\star) \quad \varphi_f(h)g = fhg = \varphi_f(hg).$$

In particular this homeomorphism preserves subsequences, i.e. for every $q \preceq p$ we have $\varphi_f(q) \preceq \varphi_f(p)$. Fix $f \in [\mathbb{N}]^{\mathbb{N}}$ and let $Q \in \mathbf{\Gamma}([f]^{\mathbb{N}})$. Since $\mathbf{\Gamma}$ is closed under Wadge reducibility we have that

$$P := \varphi_f^{-1}(Q) \in \mathbf{\Gamma}([\mathbb{N}]^{\mathbb{N}}).$$

Moreover, since every pointset in $\mathbf{\Gamma}([\mathbb{N}]^{\mathbb{N}})$ has the Ramsey property, there is $h \in \text{HS}(P)$. Using (\star) , it is straightforward to conclude that $\varphi_f(h) \in \text{HS}(Q)$.

For the second part it suffices to apply the first part to $Q := [f]^{\mathbb{N}} \cap P$, which is in $\mathbf{\Gamma}([f]^{\mathbb{N}})$. \square

PROPOSITION 3.2. Let $A, B \subset \mathbb{N}$ be disjoint. Let $P, Q \in \Sigma_1^0([\mathbb{N}]^{\mathbb{N}})$ be s.t.

1. $(\forall f \in P)(f(0) \in A \text{ and } f(1) \in A)$;
2. $(\forall g \in Q)(g(0) \in B \text{ and } g(1) \in B)$.

If $R := P \cup Q$ then

$$\text{HS}(R) \cap R = (\text{HS}(P) \cap P) \cup (\text{HS}(Q) \cap Q).$$

PROOF. The inclusion $(\text{HS}(P) \cap P) \cup (\text{HS}(Q) \cap Q) \subset \text{HS}(R) \cap R$ is trivial and always holds, so we only need to prove the converse direction. Let $h \in \text{HS}(R) \cap R$ and assume that $h \in P$. By induction we can easily show that $\text{ran}(h) \subset A$. Indeed, by point 1, $h(0) \in A$ and $h(1) \in A$. Moreover, if $h(i) \in A$ then $h(i+1) \in A$ because h lands in R : indeed if not then the substring $\langle h(i), h(i+1), \dots \rangle$ of h can neither be in P nor in Q (by the disjointness of A and B), hence it cannot be in R , contradicting the fact that h lands in R . This shows that $h \in P$ implies $h \in \text{HS}(P) \cap P$. Notice also that, by the disjointness of A and B , $\text{ran}(h) \subset A$ implies $\text{ran}(h) \cap B = \emptyset$ and therefore no subsequence of h is in Q . Similarly we can show that $h \in \text{HS}(R) \cap Q$ implies $h \in (\text{HS}(Q) \cap Q) \setminus (\text{HS}(P) \cap P)$ and therefore the claim follows. \square

The following construction was used by Avigad [2] in his proof of the open Ramsey theorem in ATR_0 .

DEFINITION 3.3. Let $P \in \Sigma_1^0([\mathbb{N}]^{\mathbb{N}})$ and let $\langle P \rangle$ be a name for P . We can define the tree

$$T_{\langle P \rangle} := \{\sigma \in [\mathbb{N}]^{<\mathbb{N}} : (\forall \tau \preceq^* \sigma)(\tau \notin \langle P \rangle)\}.$$

LEMMA 3.4. Let $P \in \Sigma_1^0([\mathbb{N}]^{\mathbb{N}})$. For every name $\langle P \rangle$ of P and every $f \in [\mathbb{N}]^{\mathbb{N}}$ we have

$$f \in \text{HS}(P) \setminus P \iff f \in [T_{\langle P \rangle}].$$

PROOF. If $f \notin [T_{\langle P \rangle}]$ then $(\exists n)(f[n] \notin T_{\langle P \rangle})$, i.e. $(\exists n)(\exists \tau \preceq^* f[n])(\tau \in \langle P \rangle)$. This implies that there exists a $g \preceq f$ s.t. $\tau \sqsubset g$. This shows that $g \in P$ and hence $f \notin \text{HS}(P) \setminus P$.

Let $f \in [T_{\langle P \rangle}]$ and let $g \preceq f$. If $g \in P$ then $(\exists n)(g[n] \in \langle P \rangle)$, contradicting the fact that $f \in [T_{\langle P \rangle}]$ (by definition of $T_{\langle P \rangle}$). Therefore we have that $f \in \text{HS}(P) \setminus P$. \square

Notice that the above lemma shows that $\text{HS}(P) \setminus P$ is closed whenever P is open. In particular, if D is clopen then $\text{HS}(D)$ is closed: indeed, letting $E := [\mathbb{N}]^{\mathbb{N}} \setminus D$, we have $\text{HS}(D) = \text{HS}(E)$ and

$$\text{HS}(D) = (\text{HS}(D) \cap D) \cup (\text{HS}(D) \setminus D) = (\text{HS}(E) \setminus E) \cup (\text{HS}(D) \setminus D)$$

is the union of two closed sets.

On the other hand, the set of solutions for an open set P that lands in P can be Π_1^1 -complete: let $(q_i)_{i \in \mathbb{N}}$ be an enumeration of the rationals. We can define

$$T := \{\sigma \in [\mathbb{N}]^{<\mathbb{N}} : (\forall i < |\sigma| - 1)(q_{\sigma(i+1)} <_{\mathbb{Q}} q_{\sigma(i)})\}.$$

A path through T is an infinite descending sequence in \mathbb{Q} . If we define $P := [\mathbb{N}]^{\mathbb{N}} \setminus [T]$ we have that $\text{HS}(P) \cap P$ is the set of well-suborders of \mathbb{Q} (every suborder of \mathbb{Q} that is not a well-order contains an infinite descending sequence with increasing indexes, and therefore, a subsequence that lands in $[T]$) and hence is Π_1^1 -complete.

This underlines a critical difference between the problem of finding a homogeneous solution that lands in P and finding one that avoids P .

The following construction will be used in the following to move open sets around while “preserving” homogeneous solutions.

DEFINITION 3.5. For every $n > 1$ let $\text{pow}_n := i \mapsto n^{i+1}$. We can define the map $\eta_n : ([\mathbb{N}]^{<\mathbb{N}} \cup [\mathbb{N}]^{\mathbb{N}}) \rightarrow ([\mathbb{N}]^{<\mathbb{N}} \cup [\mathbb{N}]^{\mathbb{N}})$ as

$$\eta_n(f) := \text{pow}_n \circ f = i \mapsto n^{f(i)+1}.$$

It is clear that η_n is a computable injection with computable inverse.

Let $P \in \Sigma_1^0([\mathbb{N}]^{\mathbb{N}})$ and let $\langle P \rangle$ be a name for P . We can define (with a small abuse of notation)

$$\eta_n(\langle P \rangle) := \bigcup_{\sigma \in \langle P \rangle} \{f \in [\mathbb{N}]^{\mathbb{N}} : \eta_n(\sigma) \sqsubset f\}.$$

We can naturally extend the definition to a multivalued map $\Sigma_1^0([\mathbb{N}]^{\mathbb{N}}) \rightrightarrows \Sigma_1^0([\mathbb{N}]^{\mathbb{N}})$ defining

$$\eta_n(P) := \{\eta_n(\langle P \rangle) : \langle P \rangle \text{ is a name of } P\}.$$

LEMMA 3.6. Let $P \in \Sigma_1^0([\mathbb{N}]^{\mathbb{N}})$. Fix $n > 1$ and let $Q := \eta_n(\langle P \rangle)$ for some name $\langle P \rangle$ of P . Then

$$f \in \text{HS}(P) \cap P \Leftrightarrow \eta_n(f) \in \text{HS}(Q) \cap Q.$$

PROOF. It is straightforward to see that, for every $f \in [\mathbb{N}]^{\mathbb{N}}$, $f \in P$ iff $\eta_n(f) \in Q$. Moreover, if $g \in [\mathbb{N}]^{\mathbb{N}}$, then

$$\eta_n(f)g = i \mapsto n^{fg(i)+1} = \eta_n(fg).$$

If f is a homogeneous solution that lands in P then $\eta_n(f)g \in Q$ for every $g \in [\mathbb{N}]^{\mathbb{N}}$, i.e. $\eta_n(f) \in \text{HS}(Q) \cap Q$. Vice versa, if $\eta_n(f) \in \text{HS}(Q) \cap Q$ then for every $g \in [\mathbb{N}]^{\mathbb{N}}$ we have $\eta_n(f)g = \eta_n(fg) \in Q$, which implies $fg \in P$. \square

DEFINITION 3.7. Let $\sigma, \tau \in [\mathbb{N}]^{<\mathbb{N}}$. We define $\sigma \boxtimes \tau$ to be the set of all strings of the form $\langle (\rho(i), \theta(i)) : i < N \rangle$ where $\rho, \theta \in [\mathbb{N}]^{<\mathbb{N}}$ and s.t.

$$\sigma \sqsubseteq \rho \wedge \tau \sqsubseteq \theta \wedge N = \max\{|\sigma|, |\tau|\}.$$

Clearly the map \boxtimes can be extended to infinite strings by defining

$$f \boxtimes g := \langle (f(0), g(0)), (f(1), g(1)), \dots \rangle.$$

For the sake of readability, it is convenient to introduce the following notation: for $i = 1, 2$, we define

$$\pi_i : (\mathbb{N} \times \mathbb{N})^{<\mathbb{N}} \cup (\mathbb{N} \times \mathbb{N})^{\mathbb{N}} \rightarrow \mathbb{N}^{<\mathbb{N}} \cup \mathbb{N}^{\mathbb{N}}$$

as the map that, given in input a finite (resp. infinite) string of pairs, returns the finite (resp. infinite) string of the i -th elements of the pairs.

Let $\langle P \rangle, \langle Q \rangle$ be two names for two open subsets of $[\mathbb{N}]^{\mathbb{N}}$. We can define

$$\langle P \rangle \boxtimes \langle Q \rangle := \bigcup_{\sigma \in \langle P \rangle} \bigcup_{\tau \in \langle Q \rangle} \sigma \boxtimes \tau,$$

which is a name for a new open set.

This leads to a map $\Sigma_1^0([\mathbb{N}]^{\mathbb{N}}) \times \Sigma_1^0([\mathbb{N}]^{\mathbb{N}}) \rightrightarrows \Sigma_1^0([\mathbb{N}]^{\mathbb{N}})$ defined by

$$P \boxtimes Q := \{R \in \Sigma_1^0([\mathbb{N}]^{\mathbb{N}}) : \langle P \rangle \boxtimes \langle Q \rangle \text{ is a name for } R\}.$$

Notice that, in general, it is not true that if f lies in the open set with name $\langle P \rangle \boxtimes \langle Q \rangle$ then $\pi_i f \in [\mathbb{N}]^{\mathbb{N}}$.

LEMMA 3.8. Let $P_1, P_2 \in \Sigma_1^0([\mathbb{N}]^{\mathbb{N}})$ and let $\langle P_1 \rangle, \langle P_2 \rangle$ be names for P_1, P_2 respectively s.t. every string in $\langle P_1 \rangle$ has length at least 2. Let $P \in \Sigma_1^0([\mathbb{N}]^{\mathbb{N}})$ be the open set with name $\langle P_1 \rangle \boxtimes \langle P_2 \rangle$. Then

$$\text{HS}(P) \cap P = \{f \boxtimes g : f \in \text{HS}(P_1) \cap P_1 \text{ and } g \in \text{HS}(P_2) \cap P_2\}.$$

PROOF. Notice first of all that $f \in \text{HS}(P) \cap P$ implies that, for $i = 1, 2$,

$$\pi_i f \in [\mathbb{N}]^{\mathbb{N}}.$$

Indeed, fix $n \in \mathbb{N}$ and consider the substring $g := \langle f(n), f(n+1), \dots \rangle$ of f . Since f is homogeneous we have $g \in P$. In particular, there is $\tau = \langle (\tau_1(i), \tau_2(i)) : i < N \rangle \in \langle P_1 \rangle \boxtimes \langle P_2 \rangle$ s.t. $\tau \sqsubset g$. Fix $\sigma_1 \in \langle P_1 \rangle$ and $\sigma_2 \in \langle P_2 \rangle$ s.t. $\tau \in \sigma_1 \boxtimes \sigma_2$. Since $|\sigma_1| \geq 2$ we have $|\tau| \geq 2$. Moreover

$$(\pi_i f)(n) = \tau_i(0) < \tau_i(1) = (\pi_i f)(n+1).$$

Let now $f \in \text{HS}(P) \cap P$. For every $g \preceq f$ we have that $g \in \text{HS}(P) \cap P$ and $\pi_1 g \in P_1, \pi_2 g \in P_2$ (indeed if $\pi_1 g \notin P_1$ or $\pi_2 g \notin P_2$ then $g \notin P$). Hence $\pi_i f \in \text{HS}(P_i) \cap P_i$ for $i = 1, 2$. The reverse inclusion is straightforward as if $f_i \in \text{HS}(P_i) \cap P_i$ then $f := f_1 \boxtimes f_2$ is a homogeneous solution for P . \square

The following generalizes the tree used by Solovay [27].

DEFINITION 3.9. Let $\phi: \mathbb{N} \rightarrow \mathbb{N}$ be injective. For every tree $T \in \mathbf{Ti}$ we define the *Solovay open set* $W_\phi(T)$ as

$$W_\phi(T) := \{f \in [\mathbb{N}]^{\mathbb{N}} : (\exists k)(\forall \tau \preceq f[k])(\tau \notin T) \text{ and } (\exists n, m \in \mathbb{N})(f(0) = \phi(n) \text{ and } f(1) = \phi(m))\}.$$

If ϕ is the identity function we drop the subscript.

It is easy to see that $W_\phi(T)$ is an open set.

LEMMA 3.10. *Let $T \in \mathbf{Ti}$ and let $W := W_\phi(T)$. Then*

1. *If $[T] = \emptyset$ then $\text{HS}(W) \cap W \neq \emptyset$. Moreover, if ϕ is surjective then $W = [\mathbb{N}]^{\mathbb{N}}$ and therefore $\text{HS}(W) = \text{HS}(W) \cap W = [\mathbb{N}]^{\mathbb{N}}$.*
2. *If $[T] \neq \emptyset$ then $\text{HS}(W) = \text{HS}(W) \setminus W$. Moreover every $f \in \text{HS}(W)$ dominates a path through T .*

PROOF. For each $f \in [\mathbb{N}]^{\mathbb{N}}$ define the tree

$$T_f := \{\sigma \in T : (\forall i < |\sigma|)(\sigma(i) \leq f(i))\}.$$

- 1: Notice that, for every $f \in [\mathbb{N}]^{\mathbb{N}}$, the set T_f is a finitely-branching well-founded subtree of T . By König's lemma, T_f must be finite and therefore there is a k s.t. every string in T_f has length $< k$. This implies that every $\tau \preceq f[k]$ is not in T . If $\text{ran}(f) \subset \text{ran}(\phi)$ (as is always the case if ϕ is surjective) then $f \in \text{HS}(W) \cap W$.
- 2: Notice that $\text{HS}(W) \cap W = \emptyset$ because for every path $x \in [T]$ and every $f \in [\mathbb{N}]^{\mathbb{N}}$, there exists $g \in [\mathbb{N}]^{\mathbb{N}}$ that grows sufficiently quickly s.t. $x \preceq fg$ (as proved in [27, p. 108]).

Moreover, if $f \in \text{HS}(W) \setminus W$ then $(\forall k)(\exists \tau \preceq f[k])(\tau \in T)$. This implies that T_f is infinite and therefore, by König's lemma, $[T_f] \neq \emptyset$. This concludes the proof as $[T_f] \subset [T]$.

□

DEFINITION 3.11. Let $\phi: \mathbb{N} \rightarrow \mathbb{N}$. For every tree $T \in \mathbf{Ti}$ we define the clopen set $D_\phi(T)$ as

$$D_\phi(T) := \{f \in [\mathbb{N}]^{\mathbb{N}} : (\exists \sigma_0, \sigma_1 \in T)(f(0) = \phi(\langle \sigma_0 \rangle) \text{ and } f(1) = \phi(\langle \sigma_1 \rangle) \text{ and } \sigma_0 \sqsubset \sigma_1)\}.$$

If ϕ is the identity function we just drop the subscript.

LEMMA 3.12. *Let $T \in \mathbf{Ti}$ and let $D := D_\phi(T)$ for some computable strictly increasing function $\phi: \mathbb{N} \rightarrow \mathbb{N}$. If $[T] \neq \emptyset$ then $\text{HS}(D) \cap D \neq \emptyset$ and there is a uniform computable surjection $\text{HS}(D) \cap D \rightarrow [T]$. On the other hand, if $[T] = \emptyset$ then $\text{HS}(D) \cap D = \emptyset$.*

PROOF. Let $y \in [T]$ and define $h \in D$ s.t. $h(i) = \phi(\langle y[i] \rangle)$. It is easy to see that h is an homogeneous solution for D landing in D , therefore $\text{HS}(D) \cap D \neq \emptyset$. Moreover, given any f that lands in D we can compute a path x through T as

$$x := \bigcup_{k \in \mathbb{N}} \phi^{-1} f(k).$$

Indeed, since f lands in D we have that $\phi^{-1}f(k) \in T$ for every $k \in \mathbb{N}$. Moreover, for every k , $\langle f(k), f(k+1), \dots \rangle \in D$ so that $\phi^{-1}f(k+1)$ is a finite string that properly extends $\phi^{-1}f(k)$. Therefore x is a well-defined function $\mathbb{N} \rightarrow \mathbb{N}$. Finally it is easy to see that $x \in [T]$: for every i , let $j > i$ s.t. there is a k s.t. $x[j] = \phi^{-1}f(k)$. By definition of D we have that $\phi^{-1}f(k) \in T$. Since T is a tree, we can conclude that $x[i] \in T$. Notice that ϕ^{-1} is computable, therefore x is computable from f . Notice also that, if y and h are as in the beginning of this proof, then $y = \bigcup_{k \in \mathbb{N}} \phi^{-1}h(k)$, which proves that the mapping is a surjection.

On the other hand, if $[T] = \emptyset$ then, for every $f \in D$, there is an i s.t. $\phi^{-1}(f(i)) \notin T$ or $\phi^{-1}(f(i)) \not\sqsubset \phi^{-1}(f(i+1))$, otherwise we could compute a path through T . In any case if $f \in D$ then it is not a homogeneous solution for D . \square

LEMMA 3.13. *The following maps are computable:*

1. $\Sigma_1^0([\mathbb{N}]^{\mathbb{N}}) \rightarrow \Pi_1^0([\mathbb{N}]^{\mathbb{N}}) := P \mapsto \text{HS}(P) \setminus P$;
2. $\eta_n : \Sigma_1^0([\mathbb{N}]^{\mathbb{N}}) \rightrightarrows \Sigma_1^0([\mathbb{N}]^{\mathbb{N}}) := P \mapsto \eta_n(P)$, for every $n \in \mathbb{N}$;
3. $\boxtimes : \Sigma_1^0([\mathbb{N}]^{\mathbb{N}}) \times \Sigma_1^0([\mathbb{N}]^{\mathbb{N}}) \rightrightarrows \Sigma_1^0([\mathbb{N}]^{\mathbb{N}}) := (P, Q) \mapsto P \boxtimes Q$;
4. $W_\phi : \mathbf{Ti} \rightarrow \Sigma_1^0([\mathbb{N}]^{\mathbb{N}}) := T \mapsto W_\phi(T)$, for every injective map $\phi : \mathbb{N} \rightarrow \mathbb{N}$ with computable range;
5. $D_\phi : \mathbf{Ti} \rightarrow \Delta_1^0([\mathbb{N}]^{\mathbb{N}}) := T \mapsto D_\phi(T)$, for every invertible map $\phi : \mathbb{N} \rightarrow \mathbb{N}$ with computable inverse.

PROOF.

- 1: Let $P \subset [\mathbb{N}]^{\mathbb{N}}$ be open and let $\langle P \rangle$ be a name for P . The definition of $T_{\langle P \rangle}$ is computable in $\langle P \rangle$. Moreover, $x \in [T_{\langle P \rangle}]$ iff $x \in \text{HS}(P) \setminus P$ (see Lemma 3.4). Since a name for $A \in \Pi_1^0([\mathbb{N}]^{\mathbb{N}})$ can be a tree T s.t. $A = [T]$, the claim follows.
- 2, 3: Straightforward from the definition.
- 4: Let $T \in \mathbf{Ti}$ be represented by its characteristic function χ_T . We can define

$$\langle W_\phi(T) \rangle := \{\sigma \in [\mathbb{N}]^{<\mathbb{N}} : (\forall \tau \leq \sigma)(\tau \notin T) \text{ and } (\exists n, m \in \mathbb{N})(\sigma(0) = \phi(n) \text{ and } \sigma(1) = \phi(m))\}.$$

Notice that the universal quantifier is bounded, while the formula in the scope of the existential quantifier is equivalent to requiring that $\sigma(0)$ and $\sigma(1)$ are in $\text{ran}(\phi)$, which is computable by hypothesis. Therefore $\langle W_\phi(T) \rangle$ is computable in T .

- 5: Let $T \in \mathbf{Ti}$. By definition of $D_\phi(T)$, the basic clopen cone $\{f \in [\mathbb{N}]^{\mathbb{N}} : \tau \sqsubset f\}$ is a subset of $D_\phi(T)$ iff

$$\phi^{-1}\tau(0) \in T \text{ and } \phi^{-1}\tau(1) \in T \text{ and } \phi^{-1}\tau(0) \sqsubset \phi^{-1}\tau(1).$$

In particular, this shows that we can T -computably obtain open names for $D_\phi(T)$ and its complement. \square

§4. Ramsey theorems in the Weihrauch lattice.

4.1. Definitions. There are several ways to formalize the open Ramsey theorem as a multivalued function.

DEFINITION 4.1 (Open Ramsey Theorem). We define the *full version* of the open Ramsey theorem as the (total) multivalued function

$$\Sigma_1^0\text{-RT} : \Sigma_1^0([\mathbb{N}]^{\mathbb{N}}) \rightrightarrows [\mathbb{N}]^{\mathbb{N}} := P \mapsto \text{HS}(P).$$

We may modify the full version by adding the requirement on “which side” we want the solution to be in. In this case, however, we need to restrict the domain to the family of open sets that admit a solution. We can define the *strong versions* of the open Ramsey theorem as the multivalued functions $\text{FindHS}_{\Sigma_1^0}, \text{FindHS}_{\Pi_1^0} : \subseteq \Sigma_1^0([\mathbb{N}]^{\mathbb{N}}) \rightrightarrows [\mathbb{N}]^{\mathbb{N}}$ with domain respectively

$$\text{dom}(\text{FindHS}_{\Sigma_1^0}) := \{P \in \Sigma_1^0([\mathbb{N}]^{\mathbb{N}}) : \text{HS}(P) \cap P \neq \emptyset\},$$

$$\text{dom}(\text{FindHS}_{\Pi_1^0}) := \{P \in \Sigma_1^0([\mathbb{N}]^{\mathbb{N}}) : \text{HS}(P) \setminus P \neq \emptyset\}$$

and defined as $\text{FindHS}_{\Sigma_1^0}(P) := \text{HS}(P) \cap P$ and $\text{FindHS}_{\Pi_1^0}(P) := \text{HS}(P) \setminus P$. We may strengthen further the requirements, defining the *weak versions* of the open Ramsey theorem: namely we define $\text{wFindHS}_{\Sigma_1^0}$ as the restriction of $\text{FindHS}_{\Sigma_1^0}$ to

$$\text{dom}(\text{wFindHS}_{\Sigma_1^0}) := \{P \in \Sigma_1^0([\mathbb{N}]^{\mathbb{N}}) : \text{HS}(P) \subset P\}.$$

Similarly we can define the weak version of $\text{FindHS}_{\Pi_1^0}$ as the multivalued function $\text{wFindHS}_{\Pi_1^0}$ obtained by restricting $\text{FindHS}_{\Pi_1^0}$ to

$$\text{dom}(\text{wFindHS}_{\Pi_1^0}) := \{P \in \Sigma_1^0([\mathbb{N}]^{\mathbb{N}}) : \text{HS}(P) \cap P = \emptyset\}.$$

Recall that, in general, an open set can have both solutions that land in the set and solutions that avoid the set. The domain of $\text{wFindHS}_{\Sigma_1^0}$ (resp. $\text{wFindHS}_{\Pi_1^0}$) is therefore strictly smaller than the domain of $\text{FindHS}_{\Sigma_1^0}$ (resp. $\text{FindHS}_{\Pi_1^0}$). As we will see the two versions exhibit very different behaviors. Notice also that the weak versions are restrictions of $\Sigma_1^0\text{-RT}$, while the strong versions are not (the set of solutions can be strictly smaller).

As in the case of the open Ramsey theorem, we can consider different multivalued functions corresponding to the clopen Ramsey theorem.

DEFINITION 4.2 (Clopen Ramsey Theorem). We define the full version of the clopen Ramsey theorem as the multivalued function $\Delta_1^0\text{-RT} : \Delta_1^0([\mathbb{N}]^{\mathbb{N}}) \rightrightarrows [\mathbb{N}]^{\mathbb{N}} := D \mapsto \text{HS}(D)$.

The strong version of the clopen Ramsey theorem is the multivalued function

$$\text{FindHS}_{\Delta_1^0} : \subseteq \Delta_1^0([\mathbb{N}]^{\mathbb{N}}) \rightrightarrows [\mathbb{N}]^{\mathbb{N}} := D \mapsto \text{HS}(D) \cap D$$

with domain

$$\text{dom}(\text{FindHS}_{\Delta_1^0}) := \{D \in \Delta_1^0([\mathbb{N}]^{\mathbb{N}}) : \text{HS}(D) \cap D \neq \emptyset\}.$$

The weak version of the clopen Ramsey theorem is the multivalued function $\text{wFindHS}_{\Delta_1^0} : \subseteq \Delta_1^0([\mathbb{N}]^{\mathbb{N}}) \rightrightarrows [\mathbb{N}]^{\mathbb{N}}$ defined as the restriction of $\text{FindHS}_{\Delta_1^0}$ to

$$\text{dom}(\text{wFindHS}_{\Delta_1^0}) := \{D \in \Delta_1^0([\mathbb{N}]^{\mathbb{N}}) : \text{HS}(D) \subset D\}.$$

Notice that we defined only one strong and one weak version of the clopen Ramsey theorem. This is because, using Lemma 2.1.2, it is straightforward to see that the other two are (strongly) Weihrauch equivalent to the ones we defined.

4.2. Problems reducible to $\text{UC}_{\mathbb{N}^{\mathbb{N}}}$. We show that $\text{wFindHS}_{\Sigma_1^0}$, $\text{wFindHS}_{\Delta_1^0}$ and $\Delta_1^0\text{-RT}$ are all Weihrauch equivalent to $\text{UC}_{\mathbb{N}^{\mathbb{N}}}$. None of these principles are strongly Weihrauch equivalent to $\text{UC}_{\mathbb{N}^{\mathbb{N}}}$, as we will show in Proposition 4.29.

LEMMA 4.3. $\text{wFindHS}_{\Sigma_1^0} \leq_W \text{UC}_{\mathbb{N}^{\mathbb{N}}}$.

PROOF. Since $\text{ATR} \equiv_W \text{UC}_{\mathbb{N}^{\mathbb{N}}}$ [18] it suffices to prove that $\text{wFindHS}_{\Sigma_1^0} \leq_W \text{ATR}$. Here we use the formulation of ATR given in [13], namely ATR is defined as the function $\text{ATR} : \text{WO} \times 2^{\mathbb{N}} \times \mathbb{N} \rightarrow 2^{\mathbb{N}}$ mapping a countable well-order L , a set $A \subset \mathbb{N}$ and (a code for) an arithmetic formula $\theta(n, W, V)$ to the unique set $Y \subset \mathbb{N}$ satisfying $H_{\theta}(L, Y, A)$, where $H_{\theta}(X, S, T)$ is the arithmetic formula defined in [26, Def. V.2.2], asserting that X is a linear order and S is the result of iterating θ along X with parameter T . The proof is the direct translation in the context of Weihrauch reducibility of the proof presented in [26, Lem. V.9.4]. More details on the construction can be found in the paper where the proof was first presented, i.e. [2, Sec. 3]. Let $P \in \text{dom}(\text{wFindHS}_{\Sigma_1^0})$. The proof consists of four steps:

1. build the tree $T := T_{\langle P \rangle}$ of homogeneous solutions that avoid P ;
2. build $\text{KB}(T)$;
3. via arithmetic transfinite recursion along $\text{KB}(T)$, obtain a sequence of infinite sets $(U_{\sigma})_{\sigma \in T}$ s.t. $U_{\sigma} \setminus U_{\tau}$ is finite whenever $\tau \leq_{\text{KB}(T)} \sigma$, and classify each $\sigma \in [\mathbb{N}]^{<\mathbb{N}}$ as “good” or “bad”;
4. use this classification to build a solution f .

Notice that steps 1 and 2 are computable (using Lemma 3.13.1). For $\sigma \notin T$, we classify σ as good if the shortest prefix of σ which is not in T belongs to $\langle P \rangle$, and bad otherwise. For $\sigma \in T$, to define U_{σ} and classify σ as good or bad, we first define an infinite set V_{σ} as follows:

- if σ is the minimum of $\text{KB}(T)$ then $V_{\sigma} := \mathbb{N}$;
- if σ is the successor of τ in $\text{KB}(T)$ then $V_{\sigma} := U_{\tau}$;
- if σ is a limit in $\text{KB}(T)$ then we define V_{σ} by diagonal intersection: we computably and uniformly find a sequence τ_j cofinal in σ . Define

$$\begin{aligned} u_0 &:= \min U_{\tau_0}; \\ u_{i+1} &:= \min \left\{ \bigcap_{j \leq i} U_{\tau_j} \setminus \{u_j\} \right\}; \\ V_{\sigma} &:= \{u_i : i \in \mathbb{N}\}. \end{aligned}$$

It is easy to verify that V_{σ} is defined by an arithmetic formula. Let

$$V_{\sigma}^1 := \{m \in V_{\sigma} : \sigma \frown m \text{ is good}\},$$

and similarly $V_{\sigma}^0 := \{m \in V_{\sigma} : \sigma \frown m \text{ is bad}\} = V_{\sigma} \setminus V_{\sigma}^1$. Set

$$n \in U_{\sigma} :\Leftrightarrow (|V_{\sigma}^1| = \infty \text{ and } n \in V_{\sigma}^1) \text{ or } (|V_{\sigma}^1| < \infty \text{ and } n \in V_{\sigma}^0).$$

We now classify σ as good if V_{σ}^1 is infinite, and bad otherwise.

We can obtain the information about $(U_\sigma)_{\sigma \in T}$ and the goodness (or badness) for each $\sigma \in T$ as a name for $Y \in \text{ATR}(\text{KB}(T), P, \theta)$, for an appropriate arithmetic formula θ .

As in [26, 2], one can show that $\langle \rangle$ is good and compute a solution $f \in \text{wFindHS}_{\Sigma_1^0}(P)$ from Y . \square

LEMMA 4.4. $\text{UC}_{\mathbb{N}^{\mathbb{N}}} \leq_W \text{wFindHS}_{\Delta_1^0}$.

PROOF. We follow the proof of the fact that the clopen Ramsey theorem implies ATR_0 over RCA_0 presented in [26, Lem. V.9.6]. We actually prove the reduction $\Sigma_1^1\text{-Sep} \leq_W \text{wFindHS}_{\Delta_1^0}$, as the equivalence $\Sigma_1^1\text{-Sep} \equiv_W \text{UC}_{\mathbb{N}^{\mathbb{N}}}$ has been proved in [18].

Let $((T_k^0, T_k^1))_{k \in \mathbb{N}}$ be a sequence of pairs of trees s.t. for all k at most one of T_k^0 and T_k^1 has a path (i.e. the sequence is a valid input for $\Sigma_1^1\text{-Sep}$). Our goal is to find a set Z s.t. if T_k^0 has a path then $k \in Z$ and if T_k^1 has a path then $k \notin Z$.

Following [26], we can uniformly compute from $((T_k^0, T_k^1))_{k \in \mathbb{N}}$ a name for a clopen $D \in \text{dom}(\text{wFindHS}_{\Delta_1^0})$ s.t. for every $f \in \text{wFindHS}_{\Delta_1^0}(D)$, f and $((T_k^0, T_k^1))_{k \in \mathbb{N}}$ uniformly compute some $Z \in \Sigma_1^1\text{-Sep}(((T_k^0, T_k^1))_{k \in \mathbb{N}})$. \square

THEOREM 4.5. $\text{UC}_{\mathbb{N}^{\mathbb{N}}} \equiv_W \text{wFindHS}_{\Sigma_1^0} \equiv_W \text{wFindHS}_{\Delta_1^0}$.

PROOF. This follows from Lemma 4.3, Lemma 4.4 and the fact that $\text{wFindHS}_{\Delta_1^0}$ is the restriction of $\text{wFindHS}_{\Sigma_1^0}$ to clopen sets. \square

THEOREM 4.6. $\text{UC}_{\mathbb{N}^{\mathbb{N}}} \equiv_W \Delta_1^0\text{-RT}$.

PROOF. By Theorem 4.5 it suffices to prove that $\text{wFindHS}_{\Delta_1^0} \equiv_W \Delta_1^0\text{-RT}$. The reduction $\text{wFindHS}_{\Delta_1^0} \leq_W \Delta_1^0\text{-RT}$ is trivial (the former is a restriction of the latter).

Let us prove the reverse reduction. By Proposition 3.1, for every open $P \subset [\mathbb{N}]^{\mathbb{N}}$, if $f \notin \text{HS}(P)$ then there is a $g \preceq f$ s.t. $g \in \text{HS}(P)$. This implies that the set $[\mathbb{N}]^{\mathbb{N}} \setminus \text{HS}(P)$ has no homogeneous solution that lies in itself.

Let $D \in \Delta_1^0([\mathbb{N}]^{\mathbb{N}})$ and fix a name $\langle p, q \rangle$ for D . Consider the set

$$E := \{f \in [\mathbb{N}]^{\mathbb{N}} : (\exists \sigma, \tau \in p)(\sigma \frown \tau \sqsubset f) \vee (\exists \sigma, \tau \in q)(\sigma \frown \tau \sqsubset f)\}.$$

It is clear that E is a clopen set and a name for E is computable from $\langle p, q \rangle$.

Notice also that $\text{HS}(D) \subset E$. Indeed, let f be a homogeneous solution for D and assume first that $f \in D$. Since D is open there must be a $\sigma \in p$ s.t. $\sigma \sqsubset f$. Moreover, since f is a homogeneous solution, $g := \langle f(|\sigma|), f(|\sigma| + 1), \dots \rangle$ must again be in D , hence there must be a $\tau \in p$ s.t. $\tau \sqsubset g$. This implies that $\sigma \frown \tau \sqsubset f$, i.e. $f \in E$. The case $f \in [\mathbb{N}]^{\mathbb{N}} \setminus D$ is analogous by replacing D with $[\mathbb{N}]^{\mathbb{N}} \setminus D$.

Moreover we can notice that there are no homogeneous solutions that land in $[\mathbb{N}]^{\mathbb{N}} \setminus E$, i.e. $E \in \text{dom}(\text{wFindHS}_{\Delta_1^0})$. Indeed assume that f avoids E and let $\sigma, \tau \in [\mathbb{N}]^{<\mathbb{N}}$ be s.t. $\sigma \frown \tau \sqsubset f$ and $\sigma \in p$ and $\tau \in q$ (the case in which $\tau \in p$ and $\sigma \in q$ is analogous). Let also $g := \langle f(|\sigma|), f(|\sigma| + 1), \dots \rangle$. Since $g \preceq f$ and f

is homogeneous there must be $\rho \in p$ s.t. $\tau \hat{\sqsubset} \rho \sqsubset g$. We can now notice that the subsequence h of f defined as

$$h := \langle f(0), \dots, f(|\sigma| - 1), f(|\sigma| + |\tau|), f(|\sigma| + |\tau| + 1), \dots \rangle$$

is s.t. $\sigma \hat{\sqsubset} \rho \sqsubset h$, and therefore $h \in E$, contradicting the fact that f avoids E .

We now claim that $\text{HS}(D) = \text{HS}(E)$. Once the claim is proved we can use $\text{wFindHS}_{\Delta_1^0}(E) = \Delta_1^0\text{-RT}(D)$ to finish the reduction.

It is straightforward to notice that $\text{HS}(D) \subset E$ implies $\text{HS}(D) \subset \text{HS}(E)$, hence we only need to prove the inclusion $\text{HS}(E) \subset \text{HS}(D)$. Let $f \in \text{HS}(E)$. Since $E \in \text{dom}(\text{wFindHS}_{\Delta_1^0})$ we must have $f \in E$. Assume w.l.o.g. that $f \in D$ and let $\sigma \sqsubset f$ be s.t. $\sigma \in p$. Fix $g \preceq f$ and let $\rho \sqsubset g$ be s.t. $\rho \in p \cup q$. Let $k \in \mathbb{N}$ be s.t. $\max \rho < f(k)$ and $\max \sigma < f(k)$, and consider $\tau \in p \cup q$ be s.t. $\tau \sqsubset \langle f(k), f(k+1), \dots \rangle$. By the homogeneity of f we have that $\tau \in p$ (otherwise every substring of f that begins with $\sigma \hat{\sqsubset} \tau$ would not be in E). Let $h \preceq f$ be s.t. $\rho \hat{\sqsubset} \tau \sqsubset h$. Again, by the homogeneity of f we have that $\rho \in p$, hence $g \in D$. Since g was arbitrary, we have that $f \in \text{HS}(D)$. \square

4.3. Problems reducible to $C_{\mathbb{N}^{\mathbb{N}}}$. Here we consider $\text{wFindHS}_{\Pi_1^0}$, $\text{FindHS}_{\Pi_1^0}$ and $\text{FindHS}_{\Delta_1^0}$.

THEOREM 4.7. $C_{\mathbb{N}^{\mathbb{N}}} \equiv_{\text{sW}} \text{FindHS}_{\Delta_1^0} \equiv_{\text{sW}} \text{FindHS}_{\Pi_1^0}$.

PROOF. We first show that $\text{FindHS}_{\Pi_1^0} \leq_{\text{sW}} C_{\mathbb{N}^{\mathbb{N}}}$. Given a name $\langle P \rangle$ for some open set $P \in \text{dom}(\text{FindHS}_{\Pi_1^0})$, by Lemma 3.13.1 we can compute a name for the closed set $\text{HS}(P) \setminus P$, which by hypothesis is nonempty. Therefore we can use $C_{\mathbb{N}^{\mathbb{N}}}$ to pick a solution.

Since $\text{FindHS}_{\Delta_1^0} \leq_{\text{sW}} \text{FindHS}_{\Pi_1^0}$ is trivial, it remains to show that $C_{\mathbb{N}^{\mathbb{N}}} \leq_{\text{sW}} \text{FindHS}_{\Delta_1^0}$. Let $T \subset [\mathbb{N}]^{<\mathbb{N}}$ be s.t. $[T] \neq \emptyset$ and let $D := D(T)$, i.e.

$$D = \{f \in [\mathbb{N}^{<\mathbb{N}}]^{\mathbb{N}} : f(0) \in T \text{ and } f(1) \in T \text{ and } f(0) \sqsubset f(1)\}.$$

Recall that D is computable from T (see Lemma 3.13.5). Moreover, by Lemma 3.12, we have that $D \in \text{dom}(\text{FindHS}_{\Delta_1^0})$ and that every $f \in \text{FindHS}_{\Delta_1^0}(D)$ uniformly computes a path through T . \square

PROPOSITION 4.8. $\text{UC}_{\mathbb{N}^{\mathbb{N}}} <_{\text{W}} \text{wFindHS}_{\Pi_1^0}$.

PROOF. The reduction $\text{UC}_{\mathbb{N}^{\mathbb{N}}} \leq_{\text{W}} \text{wFindHS}_{\Pi_1^0}$ is straightforward knowing that $\text{UC}_{\mathbb{N}^{\mathbb{N}}} \equiv_{\text{W}} \text{wFindHS}_{\Delta_1^0}$ (Theorem 4.5). The fact that the reduction is strict follows from [27, Sec. 3]. In particular, Solovay showed that there is an open set W with computable code s.t. every homogeneous solution avoids W (hence W is a valid input for $\text{wFindHS}_{\Pi_1^0}$) and is neither Σ_1^1 nor Π_1^1 (in particular it is not hyperarithmetic), while every computable instance of $\text{UC}_{\mathbb{N}^{\mathbb{N}}}$ has an hyperarithmetic solution (Theorem 2.4). \square

PROPOSITION 4.9. $\text{wFindHS}_{\Pi_1^0} \leq_{\text{W}} C_{\mathbb{N}^{\mathbb{N}}} \equiv_{\text{W}} C_{2^{\mathbb{N}}} * \text{wFindHS}_{\Pi_1^0}$.

PROOF. The first reduction follows from Theorem 4.7 as $\text{wFindHS}_{\Pi_1^0}$ is the restriction of $\text{FindHS}_{\Pi_1^0}$ to a smaller domain. The reduction $C_{2^{\mathbb{N}}} * \text{wFindHS}_{\Pi_1^0} \leq_{\text{W}} C_{\mathbb{N}^{\mathbb{N}}}$ is straightforward from $C_{2^{\mathbb{N}}} \leq_{\text{W}} C_{\mathbb{N}^{\mathbb{N}}}$, $\text{wFindHS}_{\Pi_1^0} \leq_{\text{W}} C_{\mathbb{N}^{\mathbb{N}}}$ and $C_{\mathbb{N}^{\mathbb{N}}}$ is closed under compositional product.

Finally the reduction $C_{\mathbb{N}^{\mathbb{N}}} \leq_W C_{2^{\mathbb{N}}} * \mathbf{wFindHS}_{\Pi_1^0}$ is suggested by the proof of the corollary in [27, Sec. 3]. In particular, given an ill-founded tree $T \subset [\mathbb{N}]^{\mathbb{N}}$ we can computably define the open set $W := W(T)$ (Definition 3.9). By Lemma 3.10, $W \in \text{dom}(\mathbf{wFindHS}_{\Pi_1^0})$ and every solution $f \in \mathbf{wFindHS}_{\Pi_1^0}(W)$ dominates a path through T . Let X be the subtree of $\mathbb{N}^{<\mathbb{N}}$ of the strings that are dominated by f and let $T_f := T \cap X$. Since $\emptyset \neq [T_f] \subset [T]$, we can use $C_{[X]}([T_f])$ to compute a path through T . To conclude the proof it is enough to notice that $C_{2^{\mathbb{N}}} \equiv_W C_{[X]}$ ([7, Thm. 7.23]). \square

Notice that, by the choice-elimination principle ([7, Thm. 7.25]), if Y is a computable metric space, $f : \subseteq X \rightarrow Y$ is a single-valued function and $f \leq_W C_{\mathbb{N}^{\mathbb{N}}}$ then $f \leq_W \mathbf{wFindHS}_{\Pi_1^0}$.

Since we are not able to show the equivalence of $\mathbf{wFindHS}_{\Pi_1^0}$ with any known principle, it is worth to study its properties.

PROPOSITION 4.10. $\mathbf{wFindHS}_{\Pi_1^0} \equiv_{sW} \mathbf{wFindHS}_{\Pi_1^0} \times \mathbf{wFindHS}_{\Pi_1^0}$.

PROOF. Notice that if $P, Q \in \text{dom}(\mathbf{wFindHS}_{\Pi_1^0})$ then $P \cup Q \in \text{dom}(\mathbf{wFindHS}_{\Pi_1^0})$. Indeed, for every $f \in [\mathbb{N}]^{\mathbb{N}}$, by the open Ramsey theorem applied to $[f]^{\mathbb{N}} \cap P$ (see Proposition 3.1), there is a $g \preceq f$ s.t. $[g]^{\mathbb{N}} \subset [f]^{\mathbb{N}} \cap P$ or $[g]^{\mathbb{N}} \cap [f]^{\mathbb{N}} \cap P = \emptyset$. In the first case we would have a contradiction as $[g]^{\mathbb{N}} \subset [f]^{\mathbb{N}} \cap P$ implies $[g]^{\mathbb{N}} \subset P$, i.e. $\text{HS}(P) \cap P \neq \emptyset$, against $P \in \text{dom}(\mathbf{wFindHS}_{\Pi_1^0})$. Therefore we have $[g]^{\mathbb{N}} \cap [f]^{\mathbb{N}} \cap P = [g]^{\mathbb{N}} \cap P = \emptyset$, i.e. $g \in \text{HS}(P) \setminus P$. With a similar argument, we can now apply the open Ramsey theorem to $[g]^{\mathbb{N}} \cap Q$ and conclude that there is a $h \preceq g$ s.t. $h \in \text{HS}(Q) \setminus Q$. In particular $h \notin Q$ and $h \notin P$ (as $h \preceq g$ and g avoids P). Therefore h is a subsequence of f that is not in $P \cup Q$. Since f was arbitrary, we have that $\text{HS}(P \cup Q) \cap (P \cup Q) = \emptyset$. This shows that every homogeneous solution $f \in \text{HS}(P \cup Q)$ avoids $P \cup Q$, and, in particular, avoids both P and Q . Since the union is computable (see Lemma 2.1.3) we can compute a solution for $(\mathbf{wFindHS}_{\Pi_1^0} \times \mathbf{wFindHS}_{\Pi_1^0})(P, Q)$ by computing $f \in \mathbf{wFindHS}_{\Pi_1^0}(P \cup Q)$ and returning two copies of f . \square

Let $\Sigma_1^1\text{-}C_{\mathbb{N}}^{\text{cof}} : \subseteq \Sigma_1^1(\mathbb{N}) \rightrightarrows \mathbb{N}$ be the multivalued function that chooses an element from a non-empty Σ_1^1 cofinite subset of \mathbb{N} . It is equivalent to assume that the input is a Σ_1^1 final segment of \mathbb{N} . Indeed, given a cofinite Σ_1^1 set A we can computably define the Σ_1^1 set

$$B := \{n \in \mathbb{N} : (\forall m \geq n)(m \in A)\},$$

which is a (non-empty) final segment of \mathbb{N} . Since $B \subset A$, choosing an element in B yields a solution for $\Sigma_1^1\text{-}C_{\mathbb{N}}^{\text{cof}}(A)$. With this in mind, we can assume that an input for $\Sigma_1^1\text{-}C_{\mathbb{N}}^{\text{cof}}$ is a sequence $(T_m)_{m \in \mathbb{N}}$ of trees s.t. there exists k s.t. $[T_i] = \emptyset$ iff $i < k$.

The problem $\widehat{\Sigma_1^1\text{-}C_{\mathbb{N}}^{\text{cof}}}$ has been studied in [1] under the name $\Sigma_1^1\text{-}AC_{\mathbb{N}^{\mathbb{N}}}^{\text{cof}}$. Moreover, [18] (implicitly) uses $\widehat{\Sigma_1^1\text{-}C_{\mathbb{N}}^{\text{cof}}}$ in the proof of Lemma 4.7 to separate $\Sigma_1^1\text{-}WKL$ from $\widehat{\Sigma_1^1\text{-}C_{\mathbb{N}}}$. It is known that $\widehat{\Sigma_1^1\text{-}C_{\mathbb{N}}^{\text{cof}}} <_W C_{\mathbb{N}^{\mathbb{N}}}$ [1, Thm. 3.34]. Moreover, $UC_{\mathbb{N}^{\mathbb{N}}} <_W \widehat{\Sigma_1^1\text{-}C_{\mathbb{N}}^{\text{cof}}}$ ([15]).

THEOREM 4.11. $\widehat{\Sigma_1^1\text{-}\mathbf{C}_{\mathbb{N}}^{\text{cof}}} \leq_{\text{sW}} \mathbf{wFindHS}_{\Pi_1^0}$.

PROOF. We will use strings σ which are prefixes of an infinite string f obtained by joining countably many strings g_i ; we write $\sigma = \text{dvt}(\tau_0, \dots, \tau_n)$ if τ_i is the prefix of g_i contained in σ . Formally $\sigma = \text{dvt}(\tau_0, \dots, \tau_n)$ iff

- $n = \max\{i : \langle i, 0 \rangle < |\sigma|\}$,
- for each i , $|\tau_i| = \max\{j : \langle i, j \rangle < |\sigma|\} + 1$,
- for each $\langle i, j \rangle < |\sigma|$, $\tau_i(j) = \sigma(\langle i, j \rangle)$.

Let $(T_{n,m})_{n,m \in \mathbb{N}}$ be a double sequence of trees s.t. for every n there is k_n s.t. $[T_{n,m}] = \emptyset$ iff $m < k_n$. For every n we can define

$$T_n := \langle \rangle \cup \bigcup_{m \in \mathbb{N}} \langle m \rangle \frown T_{n,m}.$$

Notice that, by hypothesis, for every n we have $[T_n] \neq \emptyset$. Moreover, if $f \in [T_n]$ then $f(0) \in \Sigma_1^1\text{-}\mathbf{C}_{\mathbb{N}}^{\text{cof}}((T_{n,m})_{m \in \mathbb{N}})$. Define also

$$T := \{\sigma \in \mathbb{N}^{<\mathbb{N}} : \sigma = \text{dvt}(\tau_0, \dots, \tau_k) \wedge (\forall i \leq k)(\tau_i \in T_i)\}.$$

Notice that if $f_n \in [T_n]$ then $\langle f_n \rangle_{n \in \mathbb{N}} \in [T]$, hence $[T] \neq \emptyset$. Moreover, if $f \in [T]$ and $i \leq j$ then, letting $f[i] = \text{dvt}(\tau_0, \dots, \tau_k)$ and $f[j] = \text{dvt}(\rho_0, \dots, \rho_h)$, for every $n \leq k$ we have $\tau_n \sqsubseteq \rho_n$. Therefore

$$f \in [T] \iff f = \langle f_n \rangle_{n \in \mathbb{N}} \text{ and } (\forall n \in \mathbb{N})(f_n \in [T_n]).$$

Let $W := W(T)$ be the Solovay open set for T . By Lemma 3.10, $W \in \text{dom}(\mathbf{wFindHS}_{\Pi_1^0})$ and every $h \in \mathbf{wFindHS}_{\Pi_1^0}(W)$ dominates a path through T .

To conclude the proof we notice that, if $f = \langle f_n \rangle_{n \in \mathbb{N}} \in [T]$ and h dominates f then, for every n ,

$$h(\langle n, 0 \rangle) \geq f_n(0) \in \Sigma_1^1\text{-}\mathbf{C}_{\mathbb{N}}^{\text{cof}}((T_{n,m})_{m \in \mathbb{N}}).$$

In particular $h(\langle n, 0 \rangle) \in \Sigma_1^1\text{-}\mathbf{C}_{\mathbb{N}}^{\text{cof}}((T_{n,m})_{m \in \mathbb{N}})$. \square

Let $\text{ATR}_2 : \text{LO} \times 2^{\mathbb{N}} \times \mathbb{N} \rightrightarrows \{0, 1\} \times \mathbb{N}^{\mathbb{N}}$ be the two sided version of ATR , defined in [13, Def. 8.2 and prop. 8.9] as the multivalued function whose inputs are triples (L, A, θ) and the output is a pair (i, Y) s.t. either $i = 0$ and Y is a $<_L$ -infinite descending chain or $i = 1$ and Y is a (pseudo)hierarchy $\langle Y_a \rangle_{a \in L}$ s.t. for all $b \in L$, $Y_b = \{n : \theta(n, \bigoplus_{a <_L b} Y_a, A)\}$. By a classical argument (see e.g. [26, Lem. V.4.12]), there are ill-founded linear orders which support a pseudo-hierarchy.

It is known that $\text{UC}_{\mathbb{N}^{\mathbb{N}}} <_{\text{W}} \text{ATR}_2 <_{\text{W}} \mathbf{C}_{\mathbb{N}^{\mathbb{N}}}$ ([13, Cor. 8.5 and 8.7]). Jun Le Goh (personal communication) observed the following corollary:

COROLLARY 4.12. $\mathbf{wFindHS}_{\Pi_1^0} \not\leq_{\text{W}} \text{ATR}_2$.

PROOF. The claim follows from the fact that $\mathbf{C}_{\mathbb{N}^{\mathbb{N}}} \leq_{\text{W}} \mathbf{C}_{2^{\mathbb{N}}} * \mathbf{wFindHS}_{\Pi_1^0}$ (Proposition 4.9) while $\mathbf{C}_{\mathbb{N}^{\mathbb{N}}} \not\leq_{\text{W}} \mathbf{C}_{2^{\mathbb{N}}} * \text{ATR}_2$ ([13, Cor. 8.5]). \square

Let us denote with $\text{TwFindHS}_{\Sigma_1^0}$ the total continuation of $\mathbf{wFindHS}_{\Sigma_1^0}$, i.e. the (total) multivalued function with domain $\Sigma_1^0([\mathbb{N}]^{\mathbb{N}})$ defined as

$$\text{TwFindHS}_{\Sigma_1^0}(P) := \begin{cases} \text{HS}(P) & \text{if } P \in \text{dom}(\mathbf{wFindHS}_{\Sigma_1^0}); \\ [\mathbb{N}]^{\mathbb{N}} & \text{otherwise.} \end{cases}$$

The following proposition underlines the gap between $\mathbf{wFindHS}_{\Sigma_1^0}$ and $\mathbf{wFindHS}_{\Pi_1^0}$ ($\mathbf{wFindHS}_{\Sigma_1^0} <_W \mathbf{wFindHS}_{\Pi_1^0}$ by Lemma 4.3 and Proposition 4.8).

PROPOSITION 4.13. $\mathbf{TwFindHS}_{\Sigma_1^0} \leq_W \mathbf{ATR}_2$, and hence $\mathbf{wFindHS}_{\Pi_1^0} \not\leq_W \mathbf{TwFindHS}_{\Sigma_1^0}$.

PROOF. Let $P \in \Sigma_1^0(\mathbb{N}^{\mathbb{N}})$ be an input for $\mathbf{TwFindHS}_{\Sigma_1^0}$ and consider the tree $T_{\langle P \rangle}$. We can computably build the linear order $\mathbf{KB}(T_{\langle P \rangle})$. Let θ be the arithmetic formula defined in the proof of Lemma 4.3. Notice that $\mathbf{KB}(T_{\langle P \rangle})$ is not necessarily a well-order, as we are not assuming $P \in \text{dom}(\mathbf{wFindHS}_{\Sigma_1^0})$ (i.e. there may be solutions that avoid P). We therefore need to modify θ by requiring that for every $\sigma \in T_{\langle P \rangle}$, if either V_σ is finite or $V_\sigma \setminus U_\tau$ is infinite for some $\tau <_{\mathbf{KB}(T_{\langle P \rangle})} \sigma$ (this cannot happen if $\mathbf{KB}(T_{\langle P \rangle})$ is a well-order), then we set $U_\sigma = \mathbb{N}$ and we label σ as bad. Let θ' be the modified formula.

Let $(i, Y) \in \mathbf{ATR}_2(\mathbf{KB}(T_{\langle P \rangle}), P, \theta')$. If $i = 0$ then Y is a $<_{\mathbf{KB}(T_{\langle P \rangle})}$ -infinite descending sequence and $P \notin \text{dom}(\mathbf{wFindHS}_{\Sigma_1^0})$; therefore any $f \in [\mathbb{N}]^{\mathbb{N}}$ is a valid output for $\mathbf{TwFindHS}_{\Sigma_1^0}(P)$. Suppose now that $i = 1$, so that Y is a (pseudo)hierarchy. By construction, Y yields a labeling of each $\sigma \in [\mathbb{N}]^{<\mathbb{N}}$ as “good” or “bad”, and, for each $\sigma \in T_{\langle P \rangle}$, an infinite set U_σ (see the proof of [26, Lem. V.9.4]). The classical proof shows that if $P \in \text{dom}(\mathbf{wFindHS}_{\Sigma_1^0})$ then $\langle \rangle$ is good. In particular if $\langle \rangle$ is bad then we can immediately conclude that $P \notin \text{dom}(\mathbf{wFindHS}_{\Sigma_1^0})$ (and, again, any $f \in [\mathbb{N}]^{\mathbb{N}}$ is a valid solution for the original problem). On the other hand, if $\langle \rangle$ is good then the modifications made to obtain θ' ensure that the intersection of finitely many U_σ is infinite, and therefore we can compute $f \in [\mathbb{N}]^{\mathbb{N}}$ following the construction of the classical proof. Notice that if $P \in \text{dom}(\mathbf{wFindHS}_{\Sigma_1^0})$ then $f \in \mathbf{HS}(P) = \mathbf{TwFindHS}_{\Sigma_1^0}(P)$. On the other hand, if $P \notin \text{dom}(\mathbf{wFindHS}_{\Sigma_1^0})$ then $f \in \mathbf{TwFindHS}_{\Sigma_1^0}(P)$ (trivially).

The second part follows from Corollary 4.12. \square

4.4. Problems not reducible to $\mathbf{C}_{\mathbb{N}^{\mathbb{N}}}$. Let us turn our attention to the last two remaining problems, namely $\Sigma_1^0\text{-RT}$ and $\mathbf{FindHS}_{\Sigma_1^0}$.

PROPOSITION 4.14. $\mathbf{TC}_{\mathbb{N}^{\mathbb{N}}} \leq_W \mathbf{C}_{2^{\mathbb{N}}} * \Sigma_1^0\text{-RT}$ and $\chi_{\Pi_1^1} <_W \mathbf{LPO} * \Sigma_1^0\text{-RT}$.

PROOF. We first show $\mathbf{TC}_{\mathbb{N}^{\mathbb{N}}} \leq_W \mathbf{C}_{2^{\mathbb{N}}} * \Sigma_1^0\text{-RT}$. Let $T \subset [\mathbb{N}]^{\mathbb{N}}$ be a tree and let $W := W(T)$ be the Solovay open set of T . Let also $f \in \Sigma_1^0\text{-RT}(W)$. By Lemma 3.10, if $[T] \neq \emptyset$ then $\mathbf{HS}(W) \cap W = \emptyset$ and f is a bound for some $x \in [T]$. On the other hand, if $[T] = \emptyset$ then $W = [\mathbb{N}]^{\mathbb{N}}$ and f is just an arbitrary infinite string.

Let X be the subtree of $\mathbb{N}^{<\mathbb{N}}$ of the strings that are dominated by f . Notice that $\mathbf{TC}_{[X]} \equiv_W \mathbf{C}_{[X]}$. Indeed, to show that $\mathbf{TC}_{[X]} \leq_W \mathbf{C}_{[X]}$ we can notice that, given a tree $S \subset X$, we can computably define an ill-founded tree R as follows: for each level n we check whether S has no nodes at level n . If this happens for some n , we can (computably) extend S to an ill-founded tree R . If this never happens then $R = S$. It is straightforward to see that $\mathbf{C}_{[X]}([R]) \subset \mathbf{TC}_{[X]}([S])$.

Let $T_f := T \cap X$. By [7, Thm. 7.23], $\mathbf{C}_{2^{\mathbb{N}}} \equiv_W \mathbf{C}_{[X]} \equiv_W \mathbf{TC}_{[X]}$, therefore we can use $\mathbf{C}_{2^{\mathbb{N}}}$ to compute a solution $h \in \mathbf{TC}_{[X]}([T_f])$. Notice that $h \in \mathbf{TC}_{\mathbb{N}^{\mathbb{N}}}([T])$. Indeed, if $[T] \neq \emptyset$ then $[T_f] \neq \emptyset$ and h is a path through T .

A simple modification of the above argument shows that $\chi_{\Pi_1^1} \leq_W \text{LPO} * \Sigma_1^0\text{-RT}$. In fact, we can see the tree T as an input for $\chi_{\Pi_1^1}$. If $f \in \Sigma_1^0\text{-RT}(W(T))$ then T_f is a finitely branching tree. Thus whether T_f is finite is a Σ_1^0 question in T_f . We can therefore use LPO to check if T_f is infinite and hence establish whether it is well-founded or not (by König's lemma, a finitely-branching tree is infinite iff it has a path).

The reduction is trivially strict as $\chi_{\Pi_1^1}$ always has a computable output. \square

It follows from Theorem 4.20 that the reduction $\text{TC}_{\mathbb{N}^{\mathbb{N}}} \leq_W \text{C}_{2^{\mathbb{N}}} * \Sigma_1^0\text{-RT}$ is actually strict.

COROLLARY 4.15. $\Sigma_1^0\text{-RT} \not\leq_W \text{C}_{\mathbb{N}^{\mathbb{N}}}$.

PROOF. If $\Sigma_1^0\text{-RT} \leq_W \text{C}_{\mathbb{N}^{\mathbb{N}}}$ then

$$\chi_{\Pi_1^1} \leq_W \text{LPO} * \Sigma_1^0\text{-RT} \leq_W \text{C}_{\mathbb{N}^{\mathbb{N}}} * \text{C}_{\mathbb{N}^{\mathbb{N}}} \equiv_W \text{C}_{\mathbb{N}^{\mathbb{N}}},$$

contradicting the fact that $\chi_{\Pi_1^1} \not\leq_W \text{C}_{\mathbb{N}^{\mathbb{N}}}$ (see [18, Sec. 7]). \square

COROLLARY 4.16. $\text{wFindHS}_{\Pi_1^0} <_W \Sigma_1^0\text{-RT}$.

PROOF. The fact that $\text{wFindHS}_{\Pi_1^0} \leq_W \Sigma_1^0\text{-RT}$ is trivial since $\text{wFindHS}_{\Pi_1^0}$ is a restriction of $\Sigma_1^0\text{-RT}$ to a smaller domain. The reduction is strict because $\Sigma_1^0\text{-RT} \not\leq_W \text{C}_{\mathbb{N}^{\mathbb{N}}}$ (Corollary 4.15) but $\text{wFindHS}_{\Pi_1^0} \leq_W \text{C}_{\mathbb{N}^{\mathbb{N}}}$ (Proposition 4.9). \square

DEFINITION 4.17. For every represented space X , we define the *strong total continuation* of C_X to be the multivalued function $\text{sTC}_X : \Pi_1^0(X) \rightrightarrows 2 \times X$ defined as

$$\text{sTC}_X(A) := \{(b, x) \in 2 \times X : (b = 0 \rightarrow A = \emptyset) \wedge (b = 1 \rightarrow x \in A)\}.$$

In particular, for $X = \mathbb{N}^{\mathbb{N}}$ (and analogously for $X = 2^{\mathbb{N}}$) we can think of $\text{sTC}_{\mathbb{N}^{\mathbb{N}}}$ as the total multivalued function that, given in input a tree, returns a string $\langle b \rangle \frown x$ s.t. b codes whether the tree is well-founded or not and, if it is ill-founded, then x is a path through T .

It is clear that $\text{TC}_{\mathbb{N}^{\mathbb{N}}} <_W \text{sTC}_{\mathbb{N}^{\mathbb{N}}}$ (the fact that the reduction is strict follows from $\chi_{\Pi_1^1} \not\leq_W \text{TC}_{\mathbb{N}^{\mathbb{N}}}$ [18, Cor. 8.6], while obviously $\chi_{\Pi_1^1} \leq_W \text{sTC}_{\mathbb{N}^{\mathbb{N}}}$). We can also notice the following:

COROLLARY 4.18. $\text{sTC}_{\mathbb{N}^{\mathbb{N}}} \leq_W \text{sTC}_{2^{\mathbb{N}}} * \Sigma_1^0\text{-RT}$.

PROOF. It suffices to repeat the proof of the first statement of Proposition 4.14, using $\text{sTC}_{2^{\mathbb{N}}}$ in place of $\text{C}_{2^{\mathbb{N}}}$. \square

We will prove in Corollary 5.13 that the above reduction is actually strict.

PROPOSITION 4.19. $\widehat{\text{TC}_{\mathbb{N}^{\mathbb{N}}}} \mid_W \text{sTC}_{\mathbb{N}^{\mathbb{N}}}$.

PROOF. The fact that $\text{sTC}_{\mathbb{N}^{\mathbb{N}}} \not\leq_W \widehat{\text{TC}_{\mathbb{N}^{\mathbb{N}}}}$ follows from the obvious observation that $\chi_{\Pi_1^1} \leq_W \text{sTC}_{\mathbb{N}^{\mathbb{N}}}$, while $\chi_{\Pi_1^1} \not\leq_W \widehat{\text{TC}_{\mathbb{N}^{\mathbb{N}}}}$ (see [18, Cor. 8.6]).

On the other hand, if $\widehat{\text{TC}_{\mathbb{N}^{\mathbb{N}}}} \leq_W \text{sTC}_{\mathbb{N}^{\mathbb{N}}}$ then, in particular, $\text{TC}_{\mathbb{N}^{\mathbb{N}}} \times \text{C}_{\mathbb{N}^{\mathbb{N}}} \leq_W \text{sTC}_{\mathbb{N}^{\mathbb{N}}}$. Since $\text{NHA} \leq_W \text{C}_{\mathbb{N}^{\mathbb{N}}}$ (see e.g. [18, Cor. 3.6]), by Proposition 2.5, this implies that $\text{TC}_{\mathbb{N}^{\mathbb{N}}} \leq_W \text{sTC}_{\mathbb{N}^{\mathbb{N}}}|_A$, where A is the set of non-empty closed sets

of $\mathbb{N}^{\mathbb{N}}$ with no hyperarithmetic member (notice that $\text{sTC}_{\mathbb{N}^{\mathbb{N}}}(\emptyset)$ has computable solutions). In particular, this implies that $\text{TC}_{\mathbb{N}^{\mathbb{N}}} \leq_W \text{C}_{\mathbb{N}^{\mathbb{N}}}$, contradicting [18, Prop. 8.2.1]. \square

We will now show that $\Sigma_1^0\text{-RT} \not\leq_W \text{TC}_{\mathbb{N}^{\mathbb{N}}}$. We actually prove a stronger result that will be useful in Section 5.

THEOREM 4.20. *For every $n \in \mathbb{N}$, $\Sigma_1^0\text{-RT} \not\leq_W \text{sTC}_{\mathbb{N}^{\mathbb{N}}} \times \text{lim}^{(n)}$.*

PROOF. Let (X, δ_X) be the represented space of computably open subsets of $[\mathbb{N}]^{\mathbb{N}}$ with no arithmetic homogeneous solution, where δ_X is the restriction of $\delta_{\Sigma_1^0([\mathbb{N}]^{\mathbb{N}})}$ to computable names. Let us define $\Sigma_1^0\text{-RT}_X : X \rightrightarrows [\mathbb{N}]^{\mathbb{N}}$ as $\Sigma_1^0\text{-RT}_X(P) := \text{HS}(P)$.

The reduction $\Sigma_1^0\text{-RT}_X \leq_W \Sigma_1^0\text{-RT}$ holds trivially, hence it is enough to prove that

$$\Sigma_1^0\text{-RT}_X \not\leq_W \text{sTC}_{\mathbb{N}^{\mathbb{N}}} \times \text{lim}^{(n)}.$$

Assume by contradiction that there is a reduction. Since $\text{lim}^{(n)}$ is a cylinder, we can assume that the reduction is a strong Weihrauch reduction. Let Φ_1, Φ_2, Ψ be the maps witnessing the strong reduction, with Φ_1 producing an input for $\text{sTC}_{\mathbb{N}^{\mathbb{N}}}$ and Φ_2 producing an input for $\text{lim}^{(n)}$. Assume that there is a $P \in X$ s.t. $\Phi_1(\langle P \rangle)$ is a name for the empty set, for some name $\langle P \rangle$ of P . By definition, 0^ω is a valid output of $\text{sTC}_{\mathbb{N}^{\mathbb{N}}}(\emptyset)$. Let $q := \text{lim}^{(n)}(\Phi_2(\langle P \rangle))$. Notice that q is arithmetic, as $\langle P \rangle$ is computable by definition of X . We have now reached a contradiction as $\Psi(0^\omega, q)$ is arithmetic, against the fact that P has no arithmetic solution.

This implies that, for every $P \in X$ and every name $\langle P \rangle$ of P , $\Phi_1(\langle P \rangle)$ is a name for a non-empty closed set, hence we have a reduction $\Sigma_1^0\text{-RT}_X \leq_W \text{C}_{\mathbb{N}^{\mathbb{N}}} \times \text{lim}^{(n)} \equiv_W \text{C}_{\mathbb{N}^{\mathbb{N}}}$.

We now claim that $\Sigma_1^0\text{-RT}_X \not\leq_W \text{C}_{\mathbb{N}^{\mathbb{N}}}$, concluding the proof. We will in fact show that $c\chi_{\Pi_1^1} \leq_W \text{LPO} * \Sigma_1^0\text{-RT}_X$, where $c\chi_{\Pi_1^1}$ is the restriction of $\chi_{\Pi_1^1}$ to computable trees. The claim then follows from the fact that $c\chi_{\Pi_1^1} \not\leq_W \text{C}_{\mathbb{N}^{\mathbb{N}}}$ (as $c\chi_{\Pi_1^1}$ is not effectively Borel measurable, see [4, Thm. 7.7]) and the fact that $\text{C}_{\mathbb{N}^{\mathbb{N}}}$ is closed under compositional product.

Let Φ_D be the forward functional witnessing $\Sigma_1^1\text{-UC}_{\mathbb{N}^{\mathbb{N}}} \leq_W \text{wFindHS}_{\Delta_1^0}$ (recall that $\Sigma_1^1\text{-UC}_{\mathbb{N}^{\mathbb{N}}} \equiv_W \text{UC}_{\mathbb{N}^{\mathbb{N}}}$ [18, Thm. 3.11], while $\text{UC}_{\mathbb{N}^{\mathbb{N}}} \leq_W \text{wFindHS}_{\Delta_1^0}$ has been proved in Lemma 4.4). Let also T_{NAR} be a computable input for $\Sigma_1^1\text{-UC}_{\mathbb{N}^{\mathbb{N}}}$ with a single non-arithmetic solution (recall that, by [24, Thm. II.4.2] every H -set is a Π_2^0 singleton).

Let T be an input for $c\chi_{\Pi_1^1}$. We can assume w.l.o.g. that T has no hyperarithmetic path: indeed if S is a computable ill-founded tree with no hyperarithmetic path then

$$T \times S := \{ \langle \langle \sigma(0), \tau(0) \rangle, \dots, \langle \sigma(n-1), \tau(n-1) \rangle \rangle : \sigma \in T \text{ and } \tau \in S \}$$

is ill-founded iff T is, and $T \times S$ has no hyperarithmetic path.

Let $W := W(T)$ be the Solovay open set for T and let Q be the clopen set with name $\Phi_D(T_{NAR})$. Notice that, since $Q \in \text{dom}(\text{wFindHS}_{\Delta_1^0})$, for every f we can computably find a subsequence $g \preceq f$ s.t. $g \in Q$.

We can computably define $P := W \cap Q$ (see [3, Prop. 3.2.4]). Since W and Q are computable then so is P . Let us show that P does not have any arithmetic solution, which implies $P \in X$. We distinguish two cases:

1. $[T] = \emptyset$: by Lemma 3.10 we have that $W = [\mathbb{N}]^{\mathbb{N}}$, hence $P = Q$ and $\text{HS}(P) = \text{HS}(Q)$. Since every solution for Q computes the non-arithmetic solution for T_{NAR} , P does not have arithmetic solutions.
2. $[T] \neq \emptyset$: notice first of all that $P \in \text{dom}(\text{wFindHS}_{\Pi_1^0})$ as $P \subset W$ and $W \in \text{dom}(\text{wFindHS}_{\Pi_1^0})$ (see Lemma 3.10).

Given $f \in \text{HS}(P)$ then, by the above observation, we can computably find a subsolution $g \in \text{HS}(P)$ s.t. $g \in Q$, thus $g \notin W$. By König's lemma such a g is a bound for a path through T (see the proof of Lemma 3.10). This also implies that every $f \in \text{HS}(P)$ is not (hyper)arithmetic (as, by hypothesis, T does not have hyperarithmetic paths).

Given $f \in \Sigma_1^0\text{-RT}_X(P)$ we can computably find $g \preceq f$ s.t. $g \in Q$. Let T_g be the subtree of T bounded by g . Notice that g is a bound for a path through T iff T_g is ill-founded iff T is ill-founded (as shown in case 2 above). Since T_g is a finitely-branching tree, by König's lemma T_g is ill-founded iff it is infinite. Moreover, the problem of checking whether T_g is finite is a $\Sigma_1^{0,g}$ question, hence we can use LPO to solve the problem (as in the proof of Proposition 4.14). \square

PROPOSITION 4.21. *For every (partial) multivalued function f , if $f \times \text{NHA} \leq_W \Sigma_1^0\text{-RT}$ then $f \leq_W \text{C}_{\mathbb{N}^{\mathbb{N}}}$.*

PROOF. Assume $f \times \text{NHA} \leq_W \Sigma_1^0\text{-RT}$ and let $B := \text{dom}(\text{wFindHS}_{\Sigma_1^0})$. Obviously $\Sigma_1^0\text{-RT}|_B = \text{wFindHS}_{\Sigma_1^0}$, hence, since $\text{wFindHS}_{\Sigma_1^0} \equiv_W \text{UC}_{\mathbb{N}^{\mathbb{N}}}$ (Theorem 4.5), the restriction of $\Sigma_1^0\text{-RT}$ to B always has a solution that is hyperarithmetic relative to the input (Theorem 2.4). Since $f \times \text{NHA} \leq_W \Sigma_1^0\text{-RT}$, by Proposition 2.5, we have that f is reducible to the restriction of $\Sigma_1^0\text{-RT}$ to

$$A := \Sigma_1^0([\mathbb{N}]^{\mathbb{N}}) \setminus B = \text{dom}(\text{FindHS}_{\Pi_1^0}).$$

This implies that $f \leq_W \text{FindHS}_{\Pi_1^0}$, as for each $P \in A$ we have $\text{FindHS}_{\Pi_1^0}(P) \subset \Sigma_1^0\text{-RT}(P)$ (and therefore every realizer for $\text{FindHS}_{\Pi_1^0}$ is also a realizer for $\Sigma_1^0\text{-RT}|_A$). The claim follows from the fact that $\text{FindHS}_{\Pi_1^0} \equiv_W \text{C}_{\mathbb{N}^{\mathbb{N}}}$ (Theorem 4.7). \square

COROLLARY 4.22. $\Sigma_1^0\text{-RT} <_W \text{NHA} \times \Sigma_1^0\text{-RT} \leq_W \text{C}_{\mathbb{N}^{\mathbb{N}}} \times \Sigma_1^0\text{-RT}$.

PROOF. The first reduction is straightforward and the second one follows from the fact that $\text{NHA} \leq_W \text{C}_{\mathbb{N}^{\mathbb{N}}}$ (see [18, Cor. 3.6]). The fact that the first reduction is strict follows from Proposition 4.21 and the fact that $\Sigma_1^0\text{-RT} \not\leq_W \text{C}_{\mathbb{N}^{\mathbb{N}}}$ (Corollary 4.15). \square

To have a better understanding of the uniform strength of $\Sigma_1^0\text{-RT}$, we now show that, even with a parallel access to some hyperarithmetic computational power, $\Sigma_1^0\text{-RT}$ does not reach the level of $\text{TC}_{\mathbb{N}^{\mathbb{N}}} \times \text{C}_{\mathbb{N}^{\mathbb{N}}}$. Thus $\Sigma_1^0\text{-RT}$ is not at the level of $\text{TC}_{\mathbb{N}^{\mathbb{N}}}^*$, which is one of the strongest principles considered in [18] to be still at the level of ATR_0 .

PROPOSITION 4.23. *If $f : \subseteq X \rightrightarrows Y$ always has an hyperarithmetic solution relative to the input and $f \leq_W \mathbf{C}_{\mathbb{N}^{\mathbb{N}}}$ then $\mathbf{TC}_{\mathbb{N}^{\mathbb{N}}} \times \mathbf{C}_{\mathbb{N}^{\mathbb{N}}} \not\leq_W f \times \Sigma_1^0\text{-RT}$.*

PROOF. Notice that, if we define $B := \text{dom}(\mathbf{wFindHS}_{\Sigma_1^0})$, then

$$(f \times \Sigma_1^0\text{-RT})|_{X \times B} = f \times \mathbf{wFindHS}_{\Sigma_1^0}.$$

Since $\mathbf{wFindHS}_{\Sigma_1^0} \equiv_W \mathbf{UC}_{\mathbb{N}^{\mathbb{N}}}$ (Theorem 4.5) we have that $(f \times \Sigma_1^0\text{-RT})|_{X \times B}$ always has a solution that is hyperarithmetic relative to the input (Theorem 2.4). Assume by contradiction that the reduction $\mathbf{TC}_{\mathbb{N}^{\mathbb{N}}} \times \mathbf{C}_{\mathbb{N}^{\mathbb{N}}} \leq_W f \times \Sigma_1^0\text{-RT}$ holds. By Proposition 2.5 we have that $\mathbf{TC}_{\mathbb{N}^{\mathbb{N}}}$ is reducible to the restriction of $f \times \Sigma_1^0\text{-RT}$ to

$$A := X \times (\Sigma_1^0([N]^{\mathbb{N}}) \setminus B) = X \times \text{dom}(\mathbf{FindHS}_{\Pi_1^0}).$$

In particular this implies that $\mathbf{TC}_{\mathbb{N}^{\mathbb{N}}} \leq_W f \times \mathbf{FindHS}_{\Pi_1^0}$ (see also the proof of Proposition 4.21). We have therefore reached a contradiction as we would have

$$\mathbf{TC}_{\mathbb{N}^{\mathbb{N}}} \leq_W f \times \mathbf{FindHS}_{\Pi_1^0} \leq_W \mathbf{C}_{\mathbb{N}^{\mathbb{N}}} \times \mathbf{C}_{\mathbb{N}^{\mathbb{N}}} \equiv_W \mathbf{C}_{\mathbb{N}^{\mathbb{N}}}. \quad \square$$

In particular, Proposition 4.23 implies $\mathbf{TC}_{\mathbb{N}^{\mathbb{N}}} \times \mathbf{C}_{\mathbb{N}^{\mathbb{N}}} \not\leq_W \mathbf{UC}_{\mathbb{N}^{\mathbb{N}}} \times \Sigma_1^0\text{-RT}$.

Let us now turn our attention to $\mathbf{FindHS}_{\Sigma_1^0}$. We first notice the following useful property:

PROPOSITION 4.24. $\mathbf{FindHS}_{\Sigma_1^0} \times \mathbf{FindHS}_{\Sigma_1^0} \leq_{sW} \mathbf{FindHS}_{\Sigma_1^0}$.

PROOF. Let $\langle P_1 \rangle, \langle P_2 \rangle$ be names for two open sets $P_1, P_2 \in \text{dom}(\mathbf{FindHS}_{\Sigma_1^0})$. Assume w.l.o.g. that every string $\sigma \in \langle P_1 \rangle$ has length at least 2 (there is no loss of generality as we can computably modify the code of P_1 by replacing a string with length 1 with all its extensions of length 2).

Let P be the open set with name $\langle P_1 \rangle \boxtimes \langle P_2 \rangle$. Recall that P is computable from P_1 and P_2 (see Lemma 3.13). Moreover, by Lemma 3.8

$$\mathbf{HS}(P) \cap P = \{f \boxtimes g : f \in \mathbf{HS}(P_1) \cap P_1 \text{ and } g \in \mathbf{HS}(P_2) \cap P_2\}.$$

Since the projections π_i are computable, it is clear that, from every solution of $\mathbf{FindHS}_{\Sigma_1^0}(P)$, we obtain two homogeneous solutions that land in P_1 and P_2 respectively. \square

COROLLARY 4.25. $\mathbf{FindHS}_{\Sigma_1^0}$ is a cylinder.

PROOF. This follows from Proposition 4.24 and the fact that $\text{id}_{\mathbb{N}^{\mathbb{N}}} \leq_{sW} \mathbf{FindHS}_{\Sigma_1^0}$, as it then follows that

$$\text{id}_{\mathbb{N}^{\mathbb{N}}} \times \mathbf{FindHS}_{\Sigma_1^0} \leq_{sW} \mathbf{FindHS}_{\Sigma_1^0} \times \mathbf{FindHS}_{\Sigma_1^0} \equiv_{sW} \mathbf{FindHS}_{\Sigma_1^0}.$$

To prove that $\text{id}_{\mathbb{N}^{\mathbb{N}}} \leq_{sW} \mathbf{FindHS}_{\Sigma_1^0}$ we proceed as follows: let $p \in \mathbb{N}^{\mathbb{N}}$ and assume w.l.o.g. that $p \in [N]^{\mathbb{N}}$. Consider the tree $T := \{p[k] : k \in \mathbb{N}\}$ of prefixes of p and let $D := D(T)$. By Lemma 3.12 we have that $D \in \text{dom}(\mathbf{FindHS}_{\Sigma_1^0})$ and that every $f \in \mathbf{HS}(D) \cap D$ uniformly computes p . \square

The problem $\mathbf{FindHS}_{\Sigma_1^0}$ is much stronger than all of the other Ramsey-related problems we introduced. We will in fact show that $\Sigma_1^0\text{-RT} <_W \mathbf{FindHS}_{\Sigma_1^0}$ (and this holds even if we consider arithmetic reductions, see Theorem 5.14).

Although we will prove much stronger results it is worth it to sketch a short proof for the reduction $\Sigma_1^0\text{-RT} \leq_W \text{FindHS}_{\Sigma_1^0}$. Given a name $\langle P \rangle$ for an open set P build the open set

$$Q := \eta_2(\langle P \rangle) \cup D_{\psi_3}(T_{\langle P \rangle}),$$

where $\psi_3 := \sigma \mapsto 3^{\langle \sigma \rangle + 1}$ and $\langle \sigma \rangle$ is the code of σ . Using Lemma 3.6 and Lemma 3.12 one can prove that $Q \in \text{dom}(\text{FindHS}_{\Sigma_1^0})$ and that every $f \in \text{HS}(Q) \cap Q$ computes a solution for P .

PROPOSITION 4.26. $\text{sTC}_{\mathbb{N}^{\mathbb{N}}} \leq_W \text{FindHS}_{\Sigma_1^0}$ and hence $\chi_{\Pi_1^1} <_W \text{FindHS}_{\Sigma_1^0}$.

PROOF. Let $\psi_n : \mathbb{N}^{<\mathbb{N}} \rightarrow \mathbb{N} := \sigma \mapsto n^{\langle \sigma \rangle + 1}$, where $\langle \sigma \rangle$ is the code of σ .

Let $T \subset [\mathbb{N}]^{<\mathbb{N}}$ be a tree. We can define the open set $P \subset [\mathbb{N}]^{\mathbb{N}}$ as $P := P_1 \cup P_2$, where $P_1 := D_{\psi_2}(T)$ and $P_2 := W_{\psi_3}(T)$. Notice that, by Lemma 3.12 and Lemma 3.10 we have

$$[T] = \emptyset \iff \text{HS}(P_1) \cap P_1 = \emptyset \iff \text{HS}(P_2) \cap P_2 \neq \emptyset.$$

Moreover, by Proposition 3.2,

$$\text{HS}(P) \cap P = (\text{HS}(P_1) \cap P_1) \cup (\text{HS}(P_2) \cap P_2).$$

This implies that

$$\begin{aligned} [T] \neq \emptyset &\Rightarrow \text{HS}(P) \cap P = \text{HS}(P_1) \cap P_1, \\ [T] = \emptyset &\Rightarrow \text{HS}(P) \cap P = \text{HS}(P_2) \cap P_2. \end{aligned}$$

In particular, given a $f \in \text{HS}(P) \cap P$ we can know whether $f \in \text{HS}(P_1) \cap P_1$ or $f \in \text{HS}(P_2) \cap P_2$ just by checking $f(0)$. If $f(0)$ is a power of 2 then $[T] \neq \emptyset$ and we can compute a path through T by considering the string $x \in [\mathbb{N}]^{\mathbb{N}}$ s.t.

$$x = \bigcup_{i \in \mathbb{N}} \psi_2^{-1}(f(i)).$$

In the other case $[T] = \emptyset$ hence we can just return $\langle 0 \rangle^\frown f$.

The result about $\chi_{\Pi_1^1}$ follows from $\chi_{\Pi_1^1} \leq_W \text{sTC}_{\mathbb{N}^{\mathbb{N}}}$ and $\text{sTC}_{\mathbb{N}^{\mathbb{N}}} \not\leq_W \chi_{\Pi_1^1}$ as $\chi_{\Pi_1^1}$ always has computable output. \square

COROLLARY 4.27. $\text{TC}_{\mathbb{N}^{\mathbb{N}}}^* <_W \text{FindHS}_{\Sigma_1^0}$.

PROOF. The reduction follows from $\text{TC}_{\mathbb{N}^{\mathbb{N}}} \leq_W \text{FindHS}_{\Sigma_1^0}$ (Proposition 4.26) and the fact that $\text{FindHS}_{\Sigma_1^0}$ is closed under product (Proposition 4.24). The fact the reduction is strict follows from the fact that $\text{FindHS}_{\Sigma_1^0}$ computes $\chi_{\Pi_1^1}$ (Proposition 4.26), while $\text{TC}_{\mathbb{N}^{\mathbb{N}}}^*$ does not ([18, Cor. 8.6]). \square

This shows that $\text{FindHS}_{\Sigma_1^0}$ is properly stronger than any multivalued function arising from statements related to ATR_0 studied so far.

THEOREM 4.28. $\text{C}_{\mathbb{N}^{\mathbb{N}}} * \Sigma_1^0\text{-RT} \leq_W \text{FindHS}_{\Sigma_1^0}$.

PROOF. By the cylindrical decomposition we can write

$$\text{C}_{\mathbb{N}^{\mathbb{N}}} * \Sigma_1^0\text{-RT} \equiv_W \text{C}_{\mathbb{N}^{\mathbb{N}}} \circ \Phi_e \circ (\text{id} \times \Sigma_1^0\text{-RT})$$

for some computable function Φ_e . It is enough to show that

$$\text{C}_{\mathbb{N}^{\mathbb{N}}} \circ \Phi_e \circ (\text{id} \times \Sigma_1^0\text{-RT}) \leq_W \text{FindHS}_{\Sigma_1^0} \times \text{FindHS}_{\Sigma_1^0} \times \chi_{\Pi_1^1}$$

and the claim will follow from $\chi_{\Pi_1^1} \leq_W \text{FindHS}_{\Sigma_1^0}$ (Proposition 4.26) and the fact that $\text{FindHS}_{\Sigma_1^0}$ is closed under product (Proposition 4.24).

Let $\langle p_1, p_2 \rangle$ be an input for $\mathbb{C}_{\mathbb{N}^{\mathbb{N}}} \circ \Phi_e \circ (\text{id} \times \Sigma_1^0\text{-RT})$ and let P be the open set with name p_2 . We can consider the tree T_{p_2} of homogeneous solutions for P that avoid P . We can now compute a tree R s.t. for every $x, y \in \mathbb{N}^{\mathbb{N}}$,

$$x \in [T_{p_2}] \text{ and } y \in [\Phi_e(p_1, x)] \iff \langle x, y \rangle \in [R].$$

Using the canonical computable bijection between $\mathbb{N}^{\mathbb{N}}$ and $[\mathbb{N}]^{\mathbb{N}}$ it is easy to transform R into a tree $S \in \mathbf{Ti}$ so that from any path through S we can compute a path through R .

Recall that $\text{TwFindHS}_{\Sigma_1^0} \leq_W \mathbb{C}_{\mathbb{N}^{\mathbb{N}}}$ (see Proposition 4.13). Since $\mathbb{C}_{\mathbb{N}^{\mathbb{N}}}$ is closed under compositional product we have that $\mathbb{C}_{\mathbb{N}^{\mathbb{N}}} \circ \Phi_e \circ (\text{id} \times \text{TwFindHS}_{\Sigma_1^0}) \leq_W \mathbb{C}_{\mathbb{N}^{\mathbb{N}}}$. Let Φ_A, Ψ_A be two computable maps witnessing the reduction. In particular, $\Phi_A(\langle p_1, p_2 \rangle)$ is an ill-founded subtree of $\mathbb{N}^{<\mathbb{N}}$ and every path through $\Phi_A(\langle p_1, p_2 \rangle)$ computes a solution for $\mathbb{C}_{\mathbb{N}^{\mathbb{N}}} \circ \Phi_e \circ (\text{id} \times \text{TwFindHS}_{\Sigma_1^0})$ via Ψ_A . Let also ψ_n be the function that maps σ to $n^{(\sigma)+1}$, where $\langle \sigma \rangle$ is the code of σ .

Let $D := D_{\psi_2}(S)$ and define

$$\begin{aligned} U &:= D \cup D_{\psi_3}(\Phi_A(\langle p_1, p_2 \rangle)); \\ V &:= D \cup W_{\psi_3}(S). \end{aligned}$$

Let us first show that $U, V \in \text{dom}(\text{FindHS}_{\Sigma_1^0})$. Notice that if $P \in \text{dom}(\text{FindHS}_{\Pi_1^0})$ then $[T_{p_2}] \neq \emptyset$ and $[S] \neq \emptyset$. By Lemma 3.12 we have that $\text{HS}(D) \cap D \neq \emptyset$ and therefore $U, V \in \text{dom}(\text{FindHS}_{\Sigma_1^0})$. On the other hand, assume $P \notin \text{dom}(\text{FindHS}_{\Pi_1^0})$. Since $\text{TwFindHS}_{\Sigma_1^0}$ is total we have that $\Phi_A(\langle p_1, p_2 \rangle)$ is ill-founded. This implies that $D_{\psi_3}(\Phi_A(\langle p_1, p_2 \rangle))$ has solutions that land in itself (again by Lemma 3.12), and hence $U \in \text{dom}(\text{FindHS}_{\Sigma_1^0})$. Moreover, since $P \notin \text{dom}(\text{FindHS}_{\Pi_1^0})$ we have that $[S] = \emptyset$ and therefore $\text{HS}(W_{\psi_3}(S)) \cap W_{\psi_3}(S) \neq \emptyset$, which shows that $V \in \text{dom}(\text{FindHS}_{\Sigma_1^0})$.

Let $(f, g, b) \in (\text{FindHS}_{\Sigma_1^0} \times \text{FindHS}_{\Sigma_1^0} \times \chi_{\Pi_1^1})(U, V, S)$. We distinguish 2 cases:

- if $b = 1$ then $[S] = \emptyset$, and hence $P \in \text{dom}(\text{wFindHS}_{\Sigma_1^0})$. By Proposition 3.2 f lands in $D_{\psi_3}(\Phi_A(\langle p_1, p_2 \rangle))$. In particular f computes a path through $\Phi_A(\langle p_1, p_2 \rangle)$ (Lemma 3.12). Moreover $\text{TwFindHS}_{\Sigma_1^0}(P) = \text{HS}(P) = \Sigma_1^0\text{-RT}(P)$, so that f computes also a solution for the compositional product (by applying Ψ_A to the path).
- if $b = 0$ then $[S] \neq \emptyset$ and hence $P \in \text{dom}(\text{FindHS}_{\Pi_1^0})$. Moreover $\text{HS}(V) \cap V = \text{HS}(D) \cap D$. Indeed, by Proposition 3.2,

$$\text{HS}(V) \cap V = (\text{HS}(D) \cap D) \cup (\text{HS}(W_{\psi_3}(S)) \cap W_{\psi_3}(S))$$

and $\text{HS}(W_{\psi_3}(S)) \cap W_{\psi_3}(S) = \emptyset$ by Lemma 3.10. In this case g computes a path through S , hence a path through R , and eventually a solution for the compositional product (by projecting the path through R).

The previous two points describe a way to compute a solution for the compositional product given a solution to $\text{FindHS}_{\Sigma_1^0} \times \text{FindHS}_{\Sigma_1^0} \times \chi_{\Pi_1^1}$, and therefore conclude the proof. \square

Notice that if $P \notin \text{dom}(\text{wFindHS}_{\Sigma_1^0})$ then we cannot (in general) use U to compute a solution for the compositional product. Indeed, it may be that $\text{HS}(P) \cap P \neq \emptyset$ and the solution obtained from $\text{FindHS}_{\Sigma_1^0}(U)$ lands in $D_{\psi_3}(\Phi_A(\langle p_1, p_2 \rangle))$. However, since every string is a valid solution for $\text{TwFindHS}_{\Sigma_1^0}(P)$, the solution we obtain is not guaranteed to have any connection with the original problem.

Notice moreover that $\Sigma_1^0\text{-RT} <_{\text{w}} \text{FindHS}_{\Sigma_1^0}$ as the former is not closed under product with $\mathbb{C}_{\mathbb{N}^{\mathbb{N}}}$ (Corollary 4.22) while the latter is closed under product (Proposition 4.24) and computes $\mathbb{C}_{\mathbb{N}^{\mathbb{N}}}$ (see Proposition 4.26). We will prove a stronger result in Theorem 5.14.

4.5. A 0–1 law for strong Weihrauch reducibility. We now characterize the strength of the Ramsey-related multivalued functions from the point of view of strong Weihrauch reducibility.

PROPOSITION 4.29. *Let Γ be a definable (boldface) pointclass that is downward closed with respect to Wadge reducibility. Assume also that every $P \in \Gamma([\mathbb{N}]^{\mathbb{N}})$ is Ramsey and that, for every $h \in [\mathbb{N}]^{\mathbb{N}}$,*

$$\Gamma([h]^{\mathbb{N}}) = \{P \cap [h]^{\mathbb{N}} : P \in \Gamma([\mathbb{N}]^{\mathbb{N}})\}.$$

If $R : \subseteq \Gamma([\mathbb{N}]^{\mathbb{N}}) \rightrightarrows [\mathbb{N}]^{\mathbb{N}}$ is a multivalued function, s.t. for every $x \in \text{dom}(R)$, $R(x) = \text{HS}(x)$, then

$$\text{id}_2 \not\leq_{\text{sw}} R.$$

In particular id_2 (and, a fortiori, $\text{UC}_{\mathbb{N}^{\mathbb{N}}}$) is not strongly Weihrauch reducible to $\text{wFindHS}_{\Sigma_1^0}$, $\text{wFindHS}_{\Pi_1^0}$, $\text{wFindHS}_{\Delta_1^0}$, $\Sigma_1^0\text{-RT}$, $\Delta_1^0\text{-RT}$.

PROOF. Assume there is a strong Weihrauch reduction witnessed by the computable maps Φ, Ψ . Let $p_i := \Phi(i)$ (with a small abuse of notation we are identifying i with its name) and let $P_i := \delta_{\Gamma([\mathbb{N}]^{\mathbb{N}})}(p_i)$. By definition of strong Weihrauch reducibility, for every $f \in \text{HS}(P_i)$ we have $\Psi(f) = i$. Fix $f \in \text{HS}(P_0)$ and consider the set $[f]^{\mathbb{N}} \cap P_1 \in \Gamma([f]^{\mathbb{N}})$. By Proposition 3.1 we have that every pointset in $\Gamma([f]^{\mathbb{N}})$ has the Ramsey property, therefore there is a $g \preceq f$ s.t. $[g]^{\mathbb{N}} \subset P_1 \cap [f]^{\mathbb{N}}$ or $[g]^{\mathbb{N}} \subset [f]^{\mathbb{N}} \setminus P_1$. In both cases $g \in \text{HS}(P_1)$ and therefore $\Psi(g) = 1$. However $g \preceq f$, hence $g \in \text{HS}(P_0)$ and so $\Psi(g) = 0$, which is a contradiction. \square

On the other hand, Theorem 4.7 shows that $\mathbb{C}_{\mathbb{N}^{\mathbb{N}}} \equiv_{\text{sw}} \text{FindHS}_{\Pi_1^0} \equiv_{\text{sw}} \text{FindHS}_{\Delta_1^0}$, which implies that $\text{FindHS}_{\Pi_1^0}$ and $\text{FindHS}_{\Delta_1^0}$ are cylinders. Since $\text{FindHS}_{\Sigma_1^0}$ is also a cylinder (Corollary 4.25) we have that, for every g and every $f \in \{\text{FindHS}_{\Sigma_1^0}, \text{FindHS}_{\Pi_1^0}, \text{FindHS}_{\Delta_1^0}\}$

$$g \leq_{\text{w}} f \iff g \leq_{\text{sw}} f.$$

This shows that, from the point of view of strong Weihrauch reducibility, the principles related to the open and clopen Ramsey theorems are either very weak (they do not strongly uniformly compute the identity on the 2-element space) or they are as strong as possible (the notions of Weihrauch reducibility and strong Weihrauch reducibility coincide).

§5. Arithmetic Weihrauch reducibility. Let us now define the notion of arithmetic Weihrauch reducibility, which is obtained by relaxing the computability requirements in the definition of Weihrauch reducibility. This was introduced in [12, Def. 1.4] (see also [14, Def. 2.2] and [1, Sec. 2.4]).

DEFINITION 5.1 (Arithmetic Weihrauch reducibility). Let $f : \subseteq X \rightrightarrows Y$, $g : \subseteq Z \rightrightarrows W$ be partial multivalued functions between represented spaces. We say that f is *arithmetically Weihrauch reducible* to g , and we write $f \leq_W^a g$, if

$$(\exists \text{ arithmetic } \Phi, \Psi : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}})(\forall G \vdash g) \Psi(\langle \text{id}_{\mathbb{N}^{\mathbb{N}}}, G\Phi \rangle) \vdash f$$

where a function $F : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ is called *arithmetic* if there is $n \in \mathbb{N}$ s.t. $F \leq_W \lim^{(n)}$.

It is straightforward to see that $f \leq_W g \Rightarrow f \leq_W^a g$. Notice moreover that if $f \leq_W^a g$ then there exists n s.t. $f \leq_W \lim^{(n)} * g * \lim^{(n)}$ (this follows directly from the definition of the compositional product).

PROPOSITION 5.2. *For every multivalued function f*

$$(\exists n)(f \leq_W \lim^{(n)}) \Rightarrow f \leq_W^a \text{id}.$$

PROOF. Assume there is a strong reduction $f \leq_{\text{sw}} \lim^{(n)}$ witnessed by the computable maps Φ_f, Ψ_f . It is easy to see that the maps $\Phi := \Psi_f \circ \lim^{(n)} \circ \Phi_f$ and $\Psi := \text{id}$ witness the reduction $f \leq_W^a \text{id}$. \square

COROLLARY 5.3. $\text{id} \equiv_W^a C_{2^{\mathbb{N}}} \equiv_W^a \text{LPO} \equiv_W^a \lim^{(n)}$.

PROOF. Straightforward from Proposition 5.2 and the fact that id is Weihrauch reducible to $C_{2^{\mathbb{N}}}$, LPO and $\lim^{(n)}$. \square

PROPOSITION 5.4. *For every (partial) multivalued functions f, g , if $f \leq_W^a \text{id}$ then $f * g \equiv_W^a g * f \equiv_W^a g$.*

PROOF. Let us first prove $f * g \equiv_W^a g$, the other equivalence is analogous. We only need to prove that $f * g \leq_W^a g$ as the converse reduction is trivial. We can assume w.l.o.g. that f, g are (partial) multivalued functions $: \subseteq \mathbb{N}^{\mathbb{N}} \rightrightarrows \mathbb{N}^{\mathbb{N}}$ (see e.g. [7, Lem. 3.8]). By the cylindrical decomposition, we can write

$$f * g \equiv_W (\text{id} \times f) \circ \Phi_e \circ (\text{id} \times g)$$

for some computable Φ_e . In particular

$$(\text{id} \times f) \circ \Phi_e \circ (\text{id} \times g)(\langle p_1, p_2 \rangle) = \langle \Phi_1(p_1, g(p_2)), f \circ \Phi_2(p_1, g(p_2)) \rangle$$

where Φ_1, Φ_2 are the computable functions s.t. $\Phi_e(p) = \langle \Phi_1(p), \Phi_2(p) \rangle$.

Let Φ_f, Ψ_f be two arithmetic maps witnessing the reduction $f \leq_W^a \text{id}$. It is straightforward to see that the maps

$$\Phi := \langle p_1, p_2 \rangle \mapsto p_2,$$

$$\Psi := (\langle p_1, p_2 \rangle, q) \mapsto \langle \Phi_1(p_1, q), \Psi_f(\Phi_2(p_1, q), \Phi_f \Phi_2(p_1, q)) \rangle$$

witness the reduction $(\text{id} \times f) \circ \Phi_e \circ (\text{id} \times g) \leq_W^a g$. \square

COROLLARY 5.5. $\text{wFindHS}_{\Pi_1^0} \equiv_W^a C_{\mathbb{N}^{\mathbb{N}}}$.

PROOF. This follows from $C_{\mathbb{N}^{\mathbb{N}}} \equiv_W C_{2^{\mathbb{N}}} * \mathbf{wFindHS}_{\Pi_1^0}$ (Proposition 4.9), using Corollary 5.3 and Proposition 5.4. \square

LEMMA 5.6. *Let g be a (partial multivalued) function that computes every arithmetic function and is closed under compositional product. For every (partial) multivalued function f*

$$f \leq_W^a g \Rightarrow f \leq_W g.$$

PROOF. It is enough to notice that $f \leq_W^a g$ implies that there exists n s.t.

$$f \leq_W \lim^{(n)} * g * \lim^{(n)}.$$

The hypotheses on g immediately yield the claim. \square

COROLLARY 5.7. $C_{\mathbb{N}^{\mathbb{N}}} <_W^a \mathbf{TC}_{\mathbb{N}^{\mathbb{N}}}$.

PROOF. The fact that $C_{\mathbb{N}^{\mathbb{N}}} \leq_W^a \mathbf{TC}_{\mathbb{N}^{\mathbb{N}}}$ is trivial as $C_{\mathbb{N}^{\mathbb{N}}} \leq_W \mathbf{TC}_{\mathbb{N}^{\mathbb{N}}}$. The separation follows from Lemma 5.6 (recall that, for every n , $\lim^{(n)} \leq_W \mathbf{UC}_{\mathbb{N}^{\mathbb{N}}}$, see [7, Prop. 7.50]) and the fact that $\mathbf{TC}_{\mathbb{N}^{\mathbb{N}}} \not\leq_W C_{\mathbb{N}^{\mathbb{N}}}$. \square

THEOREM 5.8. $\mathbf{TC}_{\mathbb{N}^{\mathbb{N}}} \equiv_W^a \mathbf{sTC}_{\mathbb{N}^{\mathbb{N}}}$.

PROOF. This follows from Proposition 5.2, Proposition 5.4 and the fact that

$$\mathbf{TC}_{\mathbb{N}^{\mathbb{N}}} \leq_W \mathbf{sTC}_{\mathbb{N}^{\mathbb{N}}} \leq_W \mathbf{LPO} * \mathbf{TC}_{\mathbb{N}^{\mathbb{N}}}. \quad \square$$

We will now prove the fact that $\mathbf{TC}_{\mathbb{N}^{\mathbb{N}}} <_W^a \Sigma_1^0\text{-RT}$. To do so we will first need some additional results about compositional products of iterations of \lim and $\mathbf{TC}_{\mathbb{N}^{\mathbb{N}}}$.

LEMMA 5.9. *Let $D(X, Y, Z)$ be an arithmetic predicate with free variables among X, Y, Z and let $\Phi: \mathbb{N}^{\mathbb{N}} \times \mathbb{N}^{\mathbb{N}} \rightarrow \mathbf{Tr}$ be computable. Define the Π_1^1 predicate $P(X, Y, Z)$ as*

$$D(X, Y, Z) \wedge ([\Phi(X, Y)] \neq \emptyset \rightarrow Z \in [\Phi(X, Y)]).$$

There exists a Π_1^0 predicate $S(X, Y, Z, W)$ s.t. an index for S is computable from indices for D and Φ s.t.

$$\begin{aligned} (\exists W)(S(X, Y, Z, W)) &\Rightarrow D(X, Y, Z), \\ [\Phi(X, Y)] \neq \emptyset &\Rightarrow (P(X, Y, Z) \iff (\exists W)(S(X, Y, Z, W))). \end{aligned}$$

PROOF. By Kleene's normal form theorem (see e.g. [23, Thm. 16.IV]), there is a Π_1^0 predicate T s.t.

$$D(X, Y, Z) \iff (\exists W)(T(X, Y, Z, W)).$$

Define the Π_1^0 predicate $S(X, Y, Z, W) := T(X, Y, Z, W) \wedge Z \in [\Phi(X, Y)]$. It follows from Kleene's normal form theorem that an index for S is computable from indices for D and Φ . The first property of S is immediate. For the second notice that, if $[\Phi(X, Y)] \neq \emptyset$ then

$$\begin{aligned} P(X, Y, Z) &\iff D(X, Y, Z) \wedge Z \in [\Phi(X, Y)] \\ &\iff (\exists W)(T(X, Y, Z, W) \wedge Z \in [\Phi(X, Y)]) \\ &\iff (\exists W)(S(X, Y, Z, W)). \end{aligned} \quad \square$$

The previous lemma can be interpreted as follows: the predicate P describes the compositional product (on both sides) of $\text{TC}_{\mathbb{N}^{\mathbb{N}}}$ with an arithmetic problem f , while D says that Y is a solution for $f(X, Z)$. Notice that, if we are considering the composition $\text{TC}_{\mathbb{N}^{\mathbb{N}}} * f$ then f (and therefore D) will not depend on the output Z of $\text{TC}_{\mathbb{N}^{\mathbb{N}}}$. On the other hand, if we consider $f * \text{TC}_{\mathbb{N}^{\mathbb{N}}}$ then we need to keep track of Z . The lemma proves that there is a uniform way to build a tree (whose body is the set of solutions to S) s.t., by projecting its paths, we can obtain the solutions to the original problem P . Notice however that the lemma does not guarantee such a tree to be ill-founded. In other words, we can recover (some) solutions to the original problem only if the tree is ill-founded.

Obviously, if D depends only on X, Y and not on Z , then a solution for D can be (arithmetically) computed without first finding a path through the tree $\Phi(X, Y)$.

LEMMA 5.10. *For every $n \in \mathbb{N}$, $\text{TC}_{\mathbb{N}^{\mathbb{N}}} * \lim^{(n)} \equiv_W \text{TC}_{\mathbb{N}^{\mathbb{N}}} \times \lim^{(n)}$.*

PROOF. Fix $n \in \mathbb{N}$. The reduction $\text{TC}_{\mathbb{N}^{\mathbb{N}}} \times \lim^{(n)} \leq_W \text{TC}_{\mathbb{N}^{\mathbb{N}}} * \lim^{(n)}$ trivially follows from the algebraic rules of the operations (see [9, Prop. 4.4]).

To prove the converse reduction, by the cylindrical decomposition we can write

$$\text{TC}_{\mathbb{N}^{\mathbb{N}}} * \lim^{(n)} \equiv_W (\text{id} \times \text{TC}_{\mathbb{N}^{\mathbb{N}}}) \circ \Phi_e \circ \lim^{(n)},$$

for some computable function Φ_e . In particular

$$(\text{id} \times \text{TC}_{\mathbb{N}^{\mathbb{N}}}) \circ \Phi_e \circ \lim^{(n)}(p) = \langle \Phi_1(\lim^{(n)}(p)), \text{TC}_{\mathbb{N}^{\mathbb{N}}} \Phi_2(\lim^{(n)}(p)) \rangle$$

where Φ_1, Φ_2 are the computable functions s.t. $\Phi_e(p) = \langle \Phi_1(p), \Phi_2(p) \rangle$.

Let $D(X, Y)$ be the predicate that says

$$Y = \langle \Phi_1(\lim^{(n)}(X)), \Phi_2(\lim^{(n)}(X)) \rangle.$$

Notice that an index for D can be (uniformly) computed from an index of Φ_e . Define also the predicate $P(X, Y, Z)$ as

$$D(X, Y) \wedge ([\pi_2(Y)] \neq \emptyset \rightarrow Z \in [\pi_2(Y)]),$$

where $\pi_2 := \langle Y_1, Y_2 \rangle \mapsto Y_2$. Since $D(X, Y)$ is arithmetic, we can use Lemma 5.9 to define a computable tree S s.t.

$$\begin{aligned} & (\exists W)((X, Y, Z, W) \in [S]) \Rightarrow D(X, Y), \\ & [\pi_2(Y)] \neq \emptyset \Rightarrow (P(X, Y, Z) \iff (\exists W)((X, Y, Z, W) \in [S])). \end{aligned}$$

For every fixed $p \in \text{dom}((\text{id} \times \text{TC}_{\mathbb{N}^{\mathbb{N}}}) \circ \Phi_e \circ \lim^{(n)})$ we define $S_p := \{\sigma : \langle p \upharpoonright |\sigma|, \sigma \rangle \in S\}$. We now claim that, from an answer to $(\text{TC}_{\mathbb{N}^{\mathbb{N}}} \times \lim^{(n)})([S_p], p)$ we can compute a solution to $(\text{id} \times \text{TC}_{\mathbb{N}^{\mathbb{N}}}) \circ \Phi_e \circ \lim^{(n)}(p)$.

Indeed $\Phi_1 \circ \lim^{(n)}(p)$ is trivially uniformly computed from $\lim^{(n)}(p)$. On the other hand, there is a unique q s.t. $D(p, q)$. Assume $[\pi_2(q)] \neq \emptyset$ and let $z_0 \in [\pi_2(q)]$. Since $P(p, q, z_0)$ holds, we have that $(\exists w)(\langle q, z_0, w \rangle \in [S_p])$, in particular $[S_p] \neq \emptyset$. Let $\langle y, z, w \rangle \in \text{TC}_{\mathbb{N}^{\mathbb{N}}}([S_p])$. Notice that, since $\lim^{(n)}$ is single-valued, we have $y = q$. Hence we can conclude that $P(p, q, z)$ holds, and therefore, by projecting $\langle y, z, w \rangle$ on the second component we obtain a path through $[\pi_2(q)]$. If, on the other hand, $[\pi_2(q)] = \emptyset$, then any z belongs to $\text{TC}_{\mathbb{N}^{\mathbb{N}}}([\pi_2(q)])$. In

both cases, by projecting the output of $\text{TC}_{\mathbb{N}^{\mathbb{N}}}([S_p])$ we can compute a solution to $(\text{TC}_{\mathbb{N}^{\mathbb{N}}} \circ \Phi_2 \circ \lim^{(n)})(p)$ and this concludes the proof. \square

LEMMA 5.11. *For every $n \in \mathbb{N}$, $\lim^{(n)} * \text{TC}_{\mathbb{N}^{\mathbb{N}}} \leq_W \text{sTC}_{\mathbb{N}^{\mathbb{N}}} \times \lim^{(3n+5)}$.*

PROOF. Fix $n \in \mathbb{N}$. By the cylindrical decomposition we can write

$$\lim^{(n)} * \text{TC}_{\mathbb{N}^{\mathbb{N}}} \equiv_W \lim^{(n)} \circ \Phi_e \circ (\text{id} \times \text{TC}_{\mathbb{N}^{\mathbb{N}}})$$

for some computable function Φ_e .

Let us define $F : \mathbb{N}^{\mathbb{N}} \times \mathbb{N}^{\mathbb{N}} \rightrightarrows \mathbb{N}^{\mathbb{N}}$ as $F := \text{T}(\lim^{(n)} \circ \Phi_e)$. Recalling that $\lim^{(n)} = \lim^{[n+1]}$, it is immediate that being in the domain of $\lim^{(n)}$ is a Π_{2n+3}^0 property. On the other hand, whether $\Phi_e(p, q)$ is defined is a Π_2^0 property. This implies that $F \leq_W \lim^{[2n+5+n+1]} = \lim^{(3n+5)}$ and hence to prove the lemma it suffices to show that $\lim^{(n)} \circ \Phi_e \circ (\text{id} \times \text{TC}_{\mathbb{N}^{\mathbb{N}}}) \leq_W \text{sTC}_{\mathbb{N}^{\mathbb{N}}} \times F$.

Let $D(X, Y, Z)$ be the arithmetic predicate

$$Y = \lim^{(n)} \circ \Phi_e(\pi_1(X), Z).$$

Clearly an index for D is computable from an index of Φ_e . Let also $P(X, Y, Z)$ be the predicate

$$D(X, Y, Z) \wedge ([\pi_2(X)] \neq \emptyset \rightarrow Z \in [\pi_2(X)]).$$

Since $D(X, Y, Z)$ is arithmetic, we can use Lemma 5.9 to define a computable tree S s.t.

$$\begin{aligned} (\exists W)((X, Y, Z, W) \in [S]) &\Rightarrow D(X, Y, Z), \\ [\pi_2(X)] \neq \emptyset &\Rightarrow (P(X, Y, Z) \iff (\exists W)((X, Y, Z, W) \in [S])). \end{aligned}$$

For every fixed $p = \langle p_1, p_2 \rangle \in \text{dom}(\lim^{(n)} \circ \Phi_e \circ (\text{id} \times \text{TC}_{\mathbb{N}^{\mathbb{N}}}))$ we define $S_p := \{\sigma : \langle p[[\sigma]], \sigma \rangle \in S\}$. We define the forward Weihrauch functional as the map $\Phi := \langle p_1, p_2 \rangle \mapsto ([S_p], (p_1, 0^\omega))$. Notice that, since F is total, $\Phi(\langle p_1, p_2 \rangle)$ is a correct input for $\text{sTC}_{\mathbb{N}^{\mathbb{N}}} \times F$. Let $(\langle b \rangle \wedge \langle y, z, w \rangle, r) \in (\text{sTC}_{\mathbb{N}^{\mathbb{N}}} \times F)([S_p], (p_1, 0^\omega))$. We claim that a solution for $\lim^{(n)} \Phi_e(p_1, \text{TC}_{\mathbb{N}^{\mathbb{N}}}(p_2))$ is y if $b = 1$ or is r if $b = 0$.

Assume that $b = 1$, i.e. that $\langle y, z, w \rangle \in [S_p]$. Then $D(\langle p_1, p_2 \rangle, y, z)$ holds, i.e. $y = \lim^{(n)}(\Phi_e(p_1, z))$. Therefore it is enough to show that $z \in \text{TC}_{\mathbb{N}^{\mathbb{N}}}([p_2])$. Assume that $[p_2] \neq \emptyset$ (the other case is trivial). Since $\langle y, z, w \rangle \in [S_p]$, we have that $P(\langle p_1, p_2 \rangle, y, z)$ holds and therefore $z \in [p_2]$.

Assume now that $b = 0$, i.e. for all y, z there is no w s.t. $\langle y, z, w \rangle \in [S_p]$. If $[p_2] \neq \emptyset$ then choose $z \in [p_2]$ and let $y = \lim^{(n)} \Phi_e(p_1, z)$. We then have that $D(\langle p_1, p_2 \rangle, y, z)$ and $P(\langle p_1, p_2 \rangle, y, z)$ hold. Therefore there exists w s.t. $\langle y, z, w \rangle \in [S_p]$, which is a contradiction. This implies that $[p_2] = \emptyset$ and therefore $0^\omega \in \text{TC}_{\mathbb{N}^{\mathbb{N}}}([p_2])$ and $(p_1, 0^\omega) \in \text{dom}(\lim^{(n)} \Phi_e)$. Therefore $r = F(p_1, 0^\omega) \in \lim^{(n)} \Phi_e(p_1, \text{TC}_{\mathbb{N}^{\mathbb{N}}}([p_2]))$ and this concludes the proof. \square

We are now ready to prove the following characterization of arithmetic reducibility to $\text{TC}_{\mathbb{N}^{\mathbb{N}}}$, conjectured by Arno Pauly during the BIRS-CMO 2019 workshop “Reverse Mathematics of Combinatorial Principles”.

THEOREM 5.12. *For every multivalued function f ,*

$$f \leq_W^a \text{TC}_{\mathbb{N}^{\mathbb{N}}} \iff (\exists n)(f \leq_W \text{sTC}_{\mathbb{N}^{\mathbb{N}}} \times \lim^{(n)}).$$

PROOF. The right-to-left implication follows from Corollary 5.3 and Proposition 5.4, as $\text{sTC}_{\mathbb{N}^{\mathbb{N}}} \leq_W \text{LPO} * \text{TC}_{\mathbb{N}^{\mathbb{N}}}$.

To prove the left-to-right implication, assume that there exists m s.t. $f \leq_W \lim^{(m)} * \text{TC}_{\mathbb{N}^{\mathbb{N}}} * \lim^{(m)}$. Notice that for every single-valued k and every g, h we have

$$(\star) \quad (g \times h) * k \leq_W (g * k) \times (h * k).$$

This fails for multivalued k , as shown in [9, Prop. 4.9(19)]. We can therefore write

$$\begin{aligned} \lim^{(m)} * \text{TC}_{\mathbb{N}^{\mathbb{N}}} * \lim^{(m)} &\leq_W (\text{sTC}_{\mathbb{N}^{\mathbb{N}}} \times \lim^{[3m+6]}) * \lim^{[m+1]} && \text{Lemma 5.11} \\ &\leq_W (\text{sTC}_{\mathbb{N}^{\mathbb{N}}} * \lim^{[m+1]}) \times \lim^{[4m+7]} && (\star) \\ &\leq_W (\text{LPO} * \text{TC}_{\mathbb{N}^{\mathbb{N}}} * \lim^{[m+1]}) \times \lim^{[4m+7]} \\ &\equiv_W (\text{LPO} * (\text{TC}_{\mathbb{N}^{\mathbb{N}}} \times \lim^{[m+1]})) \times \lim^{[4m+7]} && \text{Lemma 5.10} \\ &\leq_W (\text{LPO} * \lim^{[m+1]} * \text{TC}_{\mathbb{N}^{\mathbb{N}}}) \times \lim^{[4m+7]} \\ &\leq_W (\lim^{[m+2]} * \text{TC}_{\mathbb{N}^{\mathbb{N}}}) \times \lim^{[4m+7]} \\ &\leq_W \text{sTC}_{\mathbb{N}^{\mathbb{N}}} \times \lim^{[3m+9]} \times \lim^{[4m+7]} && \text{Lemma 5.11} \\ &\equiv_W \text{sTC}_{\mathbb{N}^{\mathbb{N}}} \times \lim^{[n]}, \end{aligned}$$

where $n = \max\{3m + 9, 4m + 7\}$. \square

COROLLARY 5.13. $\text{TC}_{\mathbb{N}^{\mathbb{N}}} <_W^a \Sigma_1^0\text{-RT}$.

PROOF. The reduction follows from $\text{TC}_{\mathbb{N}^{\mathbb{N}}} \leq_W \text{C}_{2^{\mathbb{N}}} * \Sigma_1^0\text{-RT}$ (Proposition 4.14) using Corollary 5.3 and Proposition 5.4. The fact that the reduction is strict follows from Theorem 5.12 as $\Sigma_1^0\text{-RT} \not\leq_W \text{sTC}_{\mathbb{N}^{\mathbb{N}}} \times \lim^{(k)}$ for any k (Theorem 4.20). \square

THEOREM 5.14. $\Sigma_1^0\text{-RT} <_W^a \text{FindHS}_{\Sigma_1^0}$.

PROOF. It suffices to show that $\Sigma_1^0\text{-RT} <_W^a \text{C}_{\mathbb{N}^{\mathbb{N}}} \times \Sigma_1^0\text{-RT}$, as $\text{C}_{\mathbb{N}^{\mathbb{N}}} \times \Sigma_1^0\text{-RT} \leq_W \text{FindHS}_{\Sigma_1^0}$ follows from $\Sigma_1^0\text{-RT} \leq_W \text{FindHS}_{\Sigma_1^0}$ (see Theorem 4.28), $\text{C}_{\mathbb{N}^{\mathbb{N}}} \leq_W \text{FindHS}_{\Sigma_1^0}$ (see Proposition 4.26) and the fact that $\text{FindHS}_{\Sigma_1^0}$ is closed under parallel product (Proposition 4.24).

The reduction is trivial, so we only need to prove the separation. Notice that the analogue of Proposition 2.5 for arithmetic Weihrauch reduction holds (the same proof works by replacing “computable” with “arithmetic” and \leq_W with \leq_W^a). This allows us to repeat the proof of Proposition 4.21, obtaining that $\text{C}_{\mathbb{N}^{\mathbb{N}}} \times \Sigma_1^0\text{-RT} \leq_W^a \Sigma_1^0\text{-RT}$ implies $\Sigma_1^0\text{-RT} \leq_W^a \text{C}_{\mathbb{N}^{\mathbb{N}}}$. This contradicts Corollary 5.7 and Corollary 5.13. \square

THEOREM 5.15. $\Sigma_1^0\text{-RT}^* \equiv_W^a \text{TC}_{\mathbb{N}^{\mathbb{N}}}^*$.

PROOF. The right-to-left reduction is a trivial consequence of $\text{TC}_{\mathbb{N}^{\mathbb{N}}} <_W^a \Sigma_1^0\text{-RT}$ (Corollary 5.13).

To prove the left-to-right reduction we first notice that

$$\Sigma_1^0\text{-RT} \leq_W \text{sTC}_{\mathbb{N}^{\mathbb{N}}} \times \text{TwFindHS}_{\Sigma_1^0}.$$

Indeed, given an open set P we can consider the input $([T_{\langle P \rangle}], P)$ for $\text{sTC}_{\mathbb{N}^{\mathbb{N}}} \times \text{TwFindHS}_{\Sigma_1^0}$. This is clearly a valid input as both functions are total. Let $((b)^\wedge x, f) \in (\text{sTC}_{\mathbb{N}^{\mathbb{N}}} \times \text{TwFindHS}_{\Sigma_1^0})([T_{\langle P \rangle}], P)$. If $b = 1$ then $x \in \text{HS}(P) \setminus P$ (Lemma 3.4), and therefore $x \in \Sigma_1^0\text{-RT}(P)$. If $b = 0$ then $[T_{\langle P \rangle}] = \emptyset$, which implies that $P \in \text{dom}(\text{wFindHS}_{\Sigma_1^0})$ and hence $f \in \text{wFindHS}_{\Sigma_1^0}(P)$.

We then have

$$\Sigma_1^0\text{-RT} \leq_W \text{sTC}_{\mathbb{N}^{\mathbb{N}}} \times \text{TwFindHS}_{\Sigma_1^0} \leq_W \text{sTC}_{\mathbb{N}^{\mathbb{N}}} \times \text{C}_{\mathbb{N}^{\mathbb{N}}} \leq_W^a \text{TC}_{\mathbb{N}^{\mathbb{N}}} \times \text{TC}_{\mathbb{N}^{\mathbb{N}}},$$

where $\text{TwFindHS}_{\Sigma_1^0} \leq_W \text{C}_{\mathbb{N}^{\mathbb{N}}}$ follows from $\text{TwFindHS}_{\Sigma_1^0} \leq_W \text{ATR}_2$ (Proposition 4.13). From this $\Sigma_1^0\text{-RT}^* \leq_W^a \text{TC}_{\mathbb{N}^{\mathbb{N}}}^*$ follows immediately. \square

§6. Conclusions. Some problems resisted full characterization. In particular two questions remain open:

QUESTION 6.1. $\text{wFindHS}_{\Pi_1^0} \equiv_W \text{C}_{\mathbb{N}^{\mathbb{N}}}$?

QUESTION 6.2. $\text{C}_{\mathbb{N}^{\mathbb{N}}} \leq_W \Sigma_1^0\text{-RT}$?

Observe that a positive answer to the first question automatically yields a positive answer to the second one by Corollary 4.16. We can expect that answering one of the two questions can shed light on the other.

As already observed in [18], there is not a single “analogue” of ATR_0 in the context of Weihrauch reducibility, and theorems that are equivalent from the reverse mathematics point of view can exhibit very different behaviors when phrased as multivalued functions.

Notice in particular that the classical proofs of the equivalences, over RCA_0 , of ATR_0 and the open and clopen Ramsey theorems ([26]) are useful only to establish that $\text{UC}_{\mathbb{N}^{\mathbb{N}}} \equiv_W \text{wFindHS}_{\Sigma_1^0} \equiv_W \text{wFindHS}_{\Delta_1^0}$.

Finding a homogeneous solution that lands in an open set, when there are also solutions that avoid it, is a much harder problem. In particular, notice that a natural candidate for $\Pi_1^1\text{-CA}_0$ in the Weihrauch lattice is $\widehat{\chi_{\Pi_1^1}}$. The fact that $\text{FindHS}_{\Sigma_1^0}$ computes $\chi_{\Pi_1^1}$ and is closed under parallel product implies that $\chi_{\Pi_1^1}^* \leq_W \text{FindHS}_{\Sigma_1^0}$. This naturally leads to the following question:

QUESTION 6.3. $\widehat{\text{FindHS}_{\Sigma_1^0}} \leq_W \text{FindHS}_{\Sigma_1^0}$?

A positive answer to this question would locate $\text{FindHS}_{\Sigma_1^0}$ in the realm of $\Pi_1^1\text{-CA}_0$, in sharp contrast with what happens in reverse mathematics.

Let us now derive a few computability-theoretic corollaries.

COROLLARY 6.4 (Solovay’s theorem [27, Thm. 1.8]). *If $P \subset [\mathbb{N}]^{\mathbb{N}}$ is open with computable code then either there is an hyperarithmetic homogeneous solution landing in P or there is a homogeneous solution avoiding P .*

PROOF. This follows from Theorem 2.4 and the fact that $\text{wFindHS}_{\Sigma_1^0} \equiv_W \text{UC}_{\mathbb{N}^{\mathbb{N}}}$ (Lemma 4.3). \square

The following result is attributed to Solovay [27] (see [19, Thm. 1] for an explicit proof).

COROLLARY 6.5. *The set of homogeneous solutions for a clopen set with computable code always contains an hyperarithmetic element.*

PROOF. This follows from Theorem 2.4 and the fact that $\Delta_1^0\text{-RT} \equiv_W \text{UC}_{\mathbb{N}^{\mathbb{N}}}$ (Theorem 4.6). \square

COROLLARY 6.6. *There is a clopen set $D \subset [\mathbb{N}]^{\mathbb{N}}$ with computable code s.t. every homogeneous solution that lands in D is not hyperarithmetic.*

PROOF. This follows from the fact that $\text{FindHS}_{\Delta_1^0} \equiv_W \text{C}_{\mathbb{N}^{\mathbb{N}}}$ (Theorem 4.7): if every computable clopen set had an hyperarithmetic solution landing in itself then every computable instance of $\text{C}_{\mathbb{N}^{\mathbb{N}}}$ would have a hyperarithmetic solution, contradicting $\text{NHA} \leq_W \text{C}_{\mathbb{N}^{\mathbb{N}}}$ ([18, Cor. 3.6]). \square

COROLLARY 6.7. *Every open set $P \subset [\mathbb{N}]^{\mathbb{N}}$ with computable code has a homogeneous solution f that is strictly Turing reducible to Kleene's \mathcal{O} .*

PROOF. It follows from the proof of Gandy basis theorem (see [24, Chap. III, Thm. 1.4]) that $\{f : f <_T \mathcal{O}\}$ is a basis for the Σ_1^1 predicates. If $P \in \text{dom}(\text{wFindHS}_{\Sigma_1^0})$ then, by Corollary 6.4, it has an hyperarithmetic solution. Otherwise $P \in \text{dom}(\text{FindHS}_{\Pi_1^0})$ hence, by Lemma 3.4, a homogeneous solution for P can be computed from any element of $[T_{\langle P \rangle}]$ (the tree $T_{\langle P \rangle}$ is computable from $\langle P \rangle$, see Lemma 3.13). By the Gandy basis theorem the claim follows. \square

In particular Corollary 6.7 shows that the difference, in the (arithmetic) Weihrauch lattice, between $\Sigma_1^0\text{-RT}$ and $\text{C}_{\mathbb{N}^{\mathbb{N}}}$ cannot be explained in terms of complexity of the solutions but rests entirely on the lack of uniformity.

REFERENCES

- [1] PAUL-ELLIOT ANGLÈS D'AURIAC and TAKAYUKI KIHARA, *A comparison of various analytic choice principles*, Available at <https://arxiv.org/abs/1907.02769>, 2019.
- [2] JEREMY AVIGAD, *An effective proof that open sets are Ramsey*, *Archive for Mathematical Logic*, vol. 37 (1998), no. 4, pp. 235–240.
- [3] VASCO BRATTKA, *Effective Borel measurability and reducibility of functions*, *Mathematical Logic Quarterly*, vol. 51 (2005), no. 1, pp. 19–44.
- [4] VASCO BRATTKA, MATTHEW DE BRECHT, and ARNO PAULY, *Closed choice and a uniform low basis theorem*, *Annals of Pure and Applied Logic*, vol. 163 (2012), no. 8, pp. 986–1008.
- [5] VASCO BRATTKA and GUIDO GHERARDI, *Weihrauch degrees, omniscience principles and weak computability*, *The Journal of Symbolic Logic*, vol. 76 (2011), no. 1, pp. 143–176, Available at <https://arxiv.org/abs/0905.4679>.
- [6] ———, *Completion of choice*, Available at <https://arxiv.org/abs/1910.13186>, 2019.
- [7] VASCO BRATTKA, GUIDO GHERARDI, and ARNO PAULY, *Weihrauch complexity in computable analysis*, (2017), Available at <https://arxiv.org/abs/1707.03202v1>.
- [8] VASCO BRATTKA, AKITOSHI KAWAMURA, ALBERTO MARCONE, and ARNO PAULY, *Measuring the complexity of computational content (Dagstuhl seminar 15392)*, *Dagstuhl Reports*, vol. 5 (2016), no. 9, pp. 77–104.
- [9] VASCO BRATTKA and ARNO PAULY, *On the algebraic structure of Weihrauch degrees*, *Logical Methods in Computer Science*, vol. 14 (2018), no. 4, pp. 1–36, Available at <https://arxiv.org/abs/1604.08348>.
- [10] FRED GALVIN and KAREL PRIKRY, *Borel sets and Ramsey's theorem*, *The Journal of Symbolic Logic*, vol. 38 (1973), no. 2, pp. 193–198.
- [11] GUIDO GHERARDI and ALBERTO MARCONE, *How incomputable is the separable Hahn-Banach theorem?*, *Notre Dame Journal of Formal Logic*, vol. 50 (2009), no. 4, pp. 393–425, Available at <https://arxiv.org/abs/0808.1663>.

- [12] JUN LE GOH, *Measuring the relative complexity of mathematical constructions and theorems*, **Ph.D. thesis**, Cornell University, 2019, pp. vii–196.
- [13] ———, *Some computability-theoretic reductions between principles around ATR_0* , Available at <https://arxiv.org/abs/1905.06868>, 2019.
- [14] ———, *Embeddings between well-orderings: computability-theoretic reductions*, **Annals of Pure and Applied Logic**, vol. 171 (2020), no. 6.
- [15] JUN LE GOH, ARNO PAULY, and MANLIO VALENTI, *Finding descending sequences through ill-founded linear orders*, In preparation, 2020.
- [16] KOJIRO HIGUCHI and ARNO PAULY, *The degree structure of Weihrauch-reducibility*, **Logical Methods in Computer Science**, vol. 9 (2013), no. 2:02, pp. 1–17, Available at <https://arxiv.org/abs/1101.0112>.
- [17] ILIAS G. KASTANAS, *On the Ramsey property for sets of reals*, **The Journal of Symbolic Logic**, vol. 48 (1983), no. 4, pp. 1035–1045.
- [18] TAKAYUKI KIHARA, ALBERTO MARCONE, and ARNO PAULY, *Searching for an analogue of ATR_0 in the Weihrauch lattice*, **The Journal of Symbolic Logic**, (To appear), Available at <https://arxiv.org/abs/1812.01549>.
- [19] RICHARD MANSFIELD, *A footnote to a theorem of Solovay on recursive encodability*, **Logic colloquium '77** (Angus Macintyre, Leszek Pacholski, and Jeff Paris, editors), Studies in Logic and the Foundations of Mathematics, vol. 96, Elsevier, 1978, pp. 195–198.
- [20] C. ST. J. A. NASH-WILLIAMS, *On well-quasi-ordering transfinite sequences*, **Mathematical Proceedings of the Cambridge Philosophical Society**, vol. 61 (1965), no. 1, pp. 33–39.
- [21] ARNO PAULY, *On the topological aspects of the theory of represented spaces*, **Computability**, vol. 5 (2016), no. 2, pp. 159–180, Available at <https://arxiv.org/abs/1204.3763>.
- [22] ———, *Computability on the space of countable ordinals*, Available at <https://arxiv.org/abs/1501.00386>, 2017.
- [23] HARTLEY ROGERS, *Theory of recursive functions and effective computability*, 1 ed., MIT Press, 1967.
- [24] GERALD E. SACKS, *Higher recursion theory*, 1 ed., Springer Verlag, 1990.
- [25] JACK SILVER, *Every analytic set is Ramsey*, **The Journal of Symbolic Logic**, vol. 35 (1970), no. 1, pp. 60–64.
- [26] STEPHEN G. SIMPSON, *Subsystems of second order arithmetic*, 2 ed., Cambridge University Press, 2009.
- [27] ROBERT M. SOLOVAY, *Hyperarithmetically encodable sets*, **Transactions of the American Mathematical Society**, vol. 239 (1978), pp. 99–122.
- [28] KLAUS WEIHRAUCH, *Computable analysis: An introduction*, 1 ed., Springer, November 2000.

DEPARTMENT OF MATHEMATICS, COMPUTER SCIENCE AND PHYSICS
 UNIVERSITY OF UDINE
 UDINE, UD 33100, IT
E-mail: alberto.marcone@uniud.it

DEPARTMENT OF MATHEMATICS, COMPUTER SCIENCE AND PHYSICS
 UNIVERSITY OF UDINE
 UDINE, UD 33100, IT
E-mail: manlio.valenti@uniud.it