



UNIVERSITÀ  
DEGLI STUDI  
DI UDINE

## Università degli studi di Udine

### On higher Gauss maps

*Original*

*Availability:*

This version is available <http://hdl.handle.net/11390/1069264> since 2021-03-15T12:15:27Z

*Publisher:*

*Published*

DOI:10.1016/j.jpaa.2015.05.009

*Terms of use:*

The institutional repository of the University of Udine (<http://air.uniud.it>) is provided by ARIC services. The aim is to enable open access to all the world.

*Publisher copyright*

(Article begins on next page)

## ON HIGHER GAUSS MAPS

PIETRO DE POI AND GIOVANNA ILARDI

ABSTRACT. We prove that the general fibre of the  $i$ -th Gauss map has dimension  $m$  if and only if at the general point the  $(i + 1)$ -th fundamental form consists of cones with vertex a fixed  $\mathbb{P}^{m-1}$ , extending a known theorem for the usual Gauss map. We prove this via a recursive formula for expressing higher fundamental forms. We also show some consequences of these results.

## 1. INTRODUCTION

Let  $V \subset \mathbb{P}^N$  be a projective embedded algebraic variety and  $P \in V$  a point of it; to  $P$  it is associated the  $i$ -th osculating space  $\tilde{T}_P^{(i)}(V) \subset \mathbb{P}^N$  generated by the  $(i + 1)$ -th infinitesimal neighbourhood (see e. g. [9, Example II.3.2.5]); let  $d_i$  be its dimension. We recall that the  $i$ -th Gauss map  $\gamma^i: V \dashrightarrow \mathbb{G}(d_i, N)$  is the rational map which associates to the general point  $P$  its  $i$ -th osculating space  $\tilde{T}_P^{(i)}(V)$ . For  $i = 1$  this reduces to the usual Gauss map.

The study of the Gauss map has been rather intense, both in classical (see for example [5], [2], [13]) and modern algebraic geometry (see for example [8], [1]). Instead, the study of higher order Gauss map has been much more scarce; as far as we know, we can only cite the classical paper of M. Castellani [3], reviewed in modern form in [7], [11] and the very recent paper [6]. Closely related results have been obtained by R. Piene on higher dual varieties, see for example [12]. We note moreover that a different notion of higher Gauss map has been given by F. Zak: see for example [15].

The main basic difference between the first and higher Gauss maps is—as shown in [3], [7] and [12]—that the fibres of the Gauss map are (Zariski open subsets of) linear spaces, while for higher Gauss maps this is not true in general.

In this paper, the study of higher Gauss maps is taken up further: in particular, we analyse the infinitesimal behaviours of these maps. Surprisingly enough, it turns out, see Theorem 3.6, that the general fibre of the  $i$ -th Gauss map has dimension  $m$  if and only if at the general point the  $(i + 1)$ -th fundamental form consists of cones with vertex a fixed  $\mathbb{P}^{m-1}$ , as it happens for the usual Gauss map: see [8, (2.6)]. This result is obtained by virtue of a recursive formula for expressing the  $(i + 1)$ -th fundamental form in terms of the  $i$ -th fundamental one: see Lemma 3.1. This last result is interesting in its own right: for example, we can prove immediately as a corollary the (well-known) fact that Jacobian system of the  $(i + 1)$ -th fundamental form is contained in the  $i$ -th fundamental form. Finally, we show some results

---

*Date:* 18th February 2018.

*2010 Mathematics Subject Classification.* Primary 53A20. Secondary 14N15; 51N35; 53B99.

*Key words and phrases.* Higher Gauss maps, higher fundamental forms, algebraic varieties.

The authors were partially supported by MIUR, project “Geometria delle varietà algebriche e dei loro spazi di moduli”.

that follow from the main result of the paper: in particular, as an example of the consequences of the theorem, we study the varieties whose image of a higher Gauss map has as image a curve, describing completely their varieties of higher osculating spaces.

*Acknowledgments.* We would like to thank the anonymous referee for valuable suggestions and remarks.

## 2. NOTATION AND PRELIMINARIES

We use notation as in [8], [9], and most of the preliminaries are taken from [4]. Let  $V \subset \mathbb{P}^N$  be a projective variety of dimension  $k$  over  $\mathbb{C}$  that will be always irreducible. For any point  $P \in V$  we use the following notation:  $\tilde{T}_P(V) \subset \mathbb{P}^N$  is the embedded tangent projective space to  $V$  in  $P$  and  $T_P(V)$  is the Zariski tangent space.

As in [8], we abuse notation by identifying the embedded tangent space in  $\mathbb{P}^N$  with the affine cone over it in  $\mathbb{C}^{N+1}$ . With this convention  $T_P(V) \cong \frac{\tilde{T}_P(V)}{\mathbb{C}}$ . We denote by  $\mathbb{G}(t, N)$  the Grassmannian of  $t$ -planes of  $\mathbb{P}^N$ .

To study the behaviour of  $V$  in  $P$ , following [8] (and references [2], [6], [7] and [10] therein), we consider the manifold  $\mathcal{F}(V)$  of frames in  $V$ . An element of  $\mathcal{F}(V)$  is a *Darboux frame* centred in  $P$ . This means an  $(N+1)$ -tuple

$$\{A_0; A_1, \dots, A_k; \dots, A_N\}$$

which is a basis of  $\mathbb{C}^{N+1}$  such that, if  $\pi : \mathbb{C}^{N+1} \setminus \{0\} \rightarrow \mathbb{P}^N$  is the canonical projection,

$$\pi(A_0) = P, \quad \text{and} \quad \pi(A_0), \pi(A_1), \dots, \pi(A_k) \text{ span } \tilde{T}_P(V).$$

Let this frame move in  $\mathcal{F}(V)$ ; then we have the following structure equations (in terms of the restrictions to  $V$  of the Maurer-Cartan 1-forms  $\omega_i, \omega_{i,j}$  on  $\mathcal{F}(\mathbb{P}^N)$ ) for the exterior derivatives of this moving frame

$$(2.1) \quad \begin{cases} \omega_\mu = 0 & \forall \mu > k \\ dA_0 = \sum_{i=0}^k \omega_i A_i \\ dA_i = \sum_{j=0}^N \omega_{i,j} A_j & i = 1, \dots, N \\ d\omega_j = \sum_{h=0}^k \omega_h \wedge \omega_{h,j} & j = 0, \dots, k \\ d\omega_{i,j} = \sum_{h=0}^N \omega_{i,h} \wedge \omega_{h,j} & i = 1, \dots, N, j = 0, \dots, N. \end{cases}$$

*Remark 2.1.* Geometrically, the frame  $\{A_i\}$  defines a coordinate simplex in  $\mathbb{P}^N$ . The 1-forms  $\omega_i, \omega_{i,j}$  give the rotation matrix when the coordinate simplex is infinitesimally displaced; in particular, modulo  $A_0$ , as  $dA_0 \in T_P^*(\mathbb{P}^N)$  (the cotangent space), the 1-forms  $\omega_1, \dots, \omega_k$  give a basis for the cotangent space  $T_P^*(V)$ , the corresponding  $\pi(A_i) = v_i \in T_P(V)$  give a basis for  $T_P(V)$  such that  $v_i$  is tangent to the line  $\overline{A_0 A_i}$ , and  $\omega_{k+1} = \dots = \omega_N = 0$  on  $T_P(V)$ .

We can define now the  $t$ -th fundamental form and the  $t$ -th osculating space at  $P \in V$ , for  $t \geq 2$ , see [8, §1.(b) and (d)] and [4, §1]:

**Definition 2.2.** Let  $P \in V$ , let  $t \geq 2$  be an integer and let  $I = (i_1, \dots, i_k)$  be such that  $|I| \leq t$ . The  $t$ -th osculating space to  $V$  at  $P$  is the subspace  $\tilde{T}_P^{(t)}(V) \subset \mathbb{P}^N$  spanned by  $A_0$  and by all the derivatives  $\frac{d^{|I|} A_0}{dv_1^{i_1} \dots dv_k^{i_k}}$ , where  $v_1, \dots, v_k$  span  $T_P(V)$ .

If the point  $P \in V$  is general, we will put

$$d_t := \dim(\tilde{T}_P^{(t)}(V)).$$

*Remark 2.3.* Obviously, the “expected” dimension of the  $t$ -th osculating space at a general point  $P$  for our  $k$ -dimensional variety  $V \in \mathbb{P}^N$  is the minimum among  $N$  and  $k_t := \binom{k+t}{k} - 1$ ; indeed, for the “general” variety,  $d_t$  has this expected dimension, but, as the next example shows, there are exceptions.

*Example 2.4.* Let us consider the so called *Togliatti surface*  $S \subset \mathbb{P}^5$ , which is a particular projection of the del Pezzo sextic surface embedded in  $\mathbb{P}^6$ , see [14] and [10]; it is the closure of the image of the rational map

$$\begin{aligned} \psi: \mathbb{P}^2 &\dashrightarrow \mathbb{P}^5 \\ (x : y : z) &\mapsto (x^2 y : x y^2 : x^2 z : x z^2 : y^2 z : y z^2). \end{aligned}$$

For the Togliatti surface we have  $d_2 = 4$ : this can be easily seen from the fact that the parametrisation of  $S$  given above, restricted to the open affine set  $x \neq 0$ , satisfies the following second order partial differential equation (classically called *Laplace equation*):

$$u^2 \frac{\partial^2 \mathbf{x}}{\partial u^2} + uv \frac{\partial^2 \mathbf{x}}{\partial u \partial v} + v^2 \frac{\partial^2 \mathbf{x}}{\partial v^2} - 2u \frac{\partial \mathbf{x}}{\partial u} - 2v \frac{\partial \mathbf{x}}{\partial v} + 2\mathbf{x} = 0,$$

where  $u = \frac{y}{x}$ ,  $v = \frac{z}{x}$ , are the affine coordinates, etc.

*Example 2.5.* It is immediate to see that for the Veronese surface  $V_2 \subset \mathbb{P}^5$  we have  $\dim \tilde{T}_P^{(2)}(V_2) = 5$  for every point  $P \in V_2$ : there is not a hyperplane section of  $V_2$  with a triple point.

**Definition 2.6.** Let  $t \geq 2$  and  $V_0 \subseteq V$  be the quasi projective variety of points where  $\tilde{T}_P^{(t)}(V)$  has maximal dimension. The variety

$$\text{Tan}^t(V) := \overline{\bigcup_{P \in V_0} \tilde{T}_P^{(t)}(V)}$$

is called the *variety of osculating  $t$ -spaces to  $V$* .

*Remark 2.7.* Obviously we have

$$d_t \leq d_{t-1} + \binom{k-1+t}{t} \leq \dots \leq \sum_{i=1}^t \binom{k+i-1}{i} = k_t.$$

Following [8] we give

**Definition 2.8.** The  $t$ -th *fundamental form* of  $V$  in  $P$  is the linear system  $|I^t|$  in the projective space  $\mathbb{P}(T_P(V)) \cong \mathbb{P}^{k-1}$  of hypersurfaces of degree  $t$  defined symbolically by the equations:

$$d^t A_0 = 0.$$

More intrinsically, we write  $I^t$  as the map

$$I^t: \text{Sym}^{(t)} T(V) \rightarrow N^t(V)$$

where  $N^t(V)$  is the bundle defined locally as  $N_P^t(V) := \frac{\mathbb{C}^{N+1}}{\tilde{T}_P^{(t-1)}(V)}$  and the map  $I^t$  is defined locally on each  $v \in T_P(V)$  as

$$v^t \mapsto \frac{d^t A_0}{dv^t} \quad \text{mod } \tilde{T}_P^{(t-1)}(V).$$

*Example 2.9.* With an explicit calculation (or from Example 2.11) we can show that the third fundamental form at the general point  $P$  of the Togliatti surface  $S \subset \mathbb{P}^5$  is given by a single cubic in  $\mathbb{P}(T_P(V)) \cong \mathbb{P}^1$ , while for a “general” surface (i.e. if  $d_2 = 5$ ) of  $\mathbb{P}^5$  the third fundamental form is the empty set; this is obviously true for the Veronese surface  $V_2 \subset \mathbb{P}^5$ .

We will denote the dimension of the  $t$ -th fundamental form by  $\Delta_t$ :

$$\Delta_t := \dim(|I^t|),$$

i.e.  $|I^t| \cong \mathbb{P}^{\Delta_t}$ .

We recall that (see [4, Corollary 1.15])

**Proposition 2.10.** *We have that*

$$d_t = d_{t-1} + \Delta_t + 1,$$

and vice versa: if  $d_t = d_{t-1} + \Delta + 1$ , then the  $t$ -th fundamental form has dimension  $\Delta$ .

*Example 2.11.* For the Togliatti surface  $S \subset \mathbb{P}^5$  we have that  $d_3 = 5$  and  $d_2 = 4$ , and therefore  $\Delta_3 = 0$ .

Analogously, for a “general” surface of  $\mathbb{P}^5$ ,  $d_3 = d_2 = 5$ , and  $\Delta_3 = -1$ , i.e.  $|III| = \emptyset$ ; this is true for example for the Veronese surface  $V_2 \subset \mathbb{P}^5$ .

From now on, we will suppose that our Darboux frame

$$(2.2) \quad \{A_0; A_1, \dots, A_k; A_{k+1}, \dots, A_{d_2}; A_{d_2+1}, \dots, A_{d_s}; \dots, A_{d_t}; \dots, A_N\}$$

is such that  $A_0, A_1, \dots, A_{d_s}$  span  $\tilde{T}_P^{(s)}(V)$  for all  $s = 1, \dots, t$ , with  $d_1 := k$ .

**Definition 2.12.** Let  $t \geq 1$ . The  $t$ -th (projective) Gauss map is the rational map

$$\begin{aligned} \gamma^t: V &\dashrightarrow \mathbb{G}(d_t, \mathbb{P}^N) \\ P &\mapsto \tilde{T}_P^{(t)}(V). \end{aligned}$$

We recall that (see for example [4, Theorem 1.18])

**Theorem 2.13.** *The first differential of  $\gamma^t$  at  $P$  is the  $(t+1)$ -th fundamental form at  $P$ .*

The idea of the proof is the following: we have, by the definition of  $\gamma^t$ , that

$$d\gamma_P^t: T_P V \dashrightarrow T_{\tilde{T}_P^{(t)}(V)} \mathbb{G}(d_t, \mathbb{P}^N),$$

and we recall that  $T_{\tilde{T}_P^{(t)}(V)} \mathbb{G}(d_t, \mathbb{P}^N) \cong \text{Hom}(\tilde{T}_P^{(t)}(V), N_P^{t+1}(V))$ ; moreover if we choose a Darboux frame as in (2.2), we have that  $dA_0 \in \tilde{T}_P V \subset \tilde{T}_P^{(t)}(V)$  and

$$\frac{\tilde{T}_P^{(t)}(V)}{\mathbb{C}A_0} = T_P^{(t)}(V)$$

and therefore  $d\gamma_P^t \in \text{Hom}(T_P V \otimes T_P^{(t)}(V), N_P^{t+1}(V))$ .

Now, we remark that, in our Darboux frame, we can interpret  $\gamma^t$  as

$$\gamma^t(P) = A_0 \wedge \cdots \wedge A_{d_t},$$

and therefore by (2.1),

$$d\gamma_P^t \equiv \sum_{\substack{1 \leq i \leq d_t \\ d_t+1 \leq j \leq N}} (-1)^{d_t-i+1} \omega_{i,j} A_0 \wedge \cdots \wedge \hat{A}_i \wedge \cdots \wedge A_{d_t} \wedge A_j, \quad \text{mod } \tilde{T}_P^{(t)}V;$$

now, a basis for  $T_P V \otimes T_P^{(t)}V$  can be expressed by  $(A_\alpha \otimes A_\mu)_{\substack{\alpha=1,\dots,k \\ \mu=1,\dots,d_t}}$ , and

$$d\gamma_P^t(A_\alpha \otimes A_\mu) = \sum_{d_t+1 \leq j \leq N} \omega_{\mu,j}(A_\alpha) A_j \in N_P^{t+1}(V)$$

on the other hand, for the  $(t+1)$ -th fundamental form we have

$$\frac{dA_\mu}{dv_\alpha} \equiv \sum_{d_t+1 \leq j \leq N} \omega_{\mu,j}(A_\alpha) A_j \quad \text{mod } \tilde{T}_P^{(t)}(V).$$

### 3. FIBRES OF HIGHER GAUSS MAPS

We start by writing higher fundamental forms explicitly: choose a Darboux frame

$$\{A_0; A_1, \dots, A_k; A_{k+1}, \dots, A_{d_2}; A_{d_2+1}, \dots, A_{d_s}; \dots, A_{d_t}; \dots, A_N\}$$

such that  $A_0, A_1, \dots, A_{d_s}$  span  $\tilde{T}_P^{(s)}(V)$  for all  $s = 1, \dots, t$ , with  $d_1 := k$ ,  $\tilde{T}_P^{(1)}(V) = \tilde{T}_P(V)$  (and  $d_0 := 0$ ). We use indices  $1 \leq i^{(1)}, i_1^{(1)}, \dots, i_h^{(1)} \leq k = d_1$ ,  $d_{s-1} + 1 \leq i^{(s)} \leq d_s$  for  $s = 2, \dots, t$  and  $d_t + 1 \leq i^{(t+1)} \leq N$ .

With these notations, the second fundamental form is given by the quadrics

$$(3.1) \quad V_{i^{(\ell)}}^{(2)} = \sum_{i_1^{(1)}, i_2^{(1)}} q_{i_1^{(1)}, i_2^{(1)}; i^{(\ell)}} \omega_{i_1^{(1)}} \omega_{i_2^{(1)}},$$

with  $\ell > k$ , where  $q_{i_1^{(1)}, i_2^{(1)}; i^{(\ell)}} (= q_{i_2^{(1)}, i_1^{(1)}; i^{(\ell)}})$  are defined by

$$(3.2) \quad \omega_{i_1^{(1)}, i^{(\ell)}} = \sum_{i_2^{(1)}} q_{i_1^{(1)}, i_2^{(1)}; i^{(\ell)}} \omega_{i_2^{(1)}},$$

which are obtained via the Cartan Lemma from

$$0 = d\omega_{i^{(\ell)}} = \sum_{i_1^{(1)}} \omega_{i_1^{(1)}} \wedge \omega_{i_1^{(1)}, i^{(\ell)}},$$

since  $\omega_{i^{(\ell)}} = 0$  on  $\tilde{T}_P(V)$ .

The higher fundamental forms can be expressed as in the following

**Lemma 3.1.** *The  $(s+1)$ -th fundamental form is given by the polynomials*

$$(3.3) \quad V_{i^{(s+1)}}^{(s+1)} = \sum_{i_1^{(1)}, \dots, i_{s+1}^{(1)}=1}^k q_{i_1^{(1)}, \dots, i_{s+1}^{(1)}; i^{(s+1)}} \omega_{i_1^{(1)}} \cdots \omega_{i_{s+1}^{(1)}},$$

with  $\ell \geq s+1$ , which inductively satisfy the relations

$$(3.4) \quad \sum_{i^{(s)}=d_{s-1}+1}^{d_s} q_{i_1^{(1)}, \dots, i_s^{(1)}; i^{(s)}} \omega_{i^{(s)}, i^{(\ell)}} = \sum_{i_{s+1}^{(1)}=1}^k q_{i_1^{(1)}, \dots, i_{s+1}^{(1)}; i^{(\ell)}} \omega_{i_{s+1}^{(1)}},$$

where  $q_{i_1^{(1)}, \dots, i_s^{(1)}; i^{(s)}}$  are the coefficients of the  $s$ -th fundamental form (and with the natural symmetries of the indices:  $q_{\dots, i_j^{(1)}, \dots, i_k^{(1)}, \dots; i^{(\ell)}} = q_{\dots, i_k^{(1)}, \dots, i_j^{(1)}, \dots; i^{(\ell)}}$ ), and the basis of the induction is the second fundamental form (i.e.  $s = 1$ ) for which relations (3.3) and (3.4) are to be read as, respectively, (3.1) and (3.2).

*Proof.* By the definition of  $\tilde{T}_P^{(s)}(V)$ , where  $s \in \{1, \dots, t-1\}$ , we have that

$$(3.5) \quad dA_{i^{(s-1)}} \equiv 0 \pmod{\tilde{T}_P^{(s)}(V)}$$

and from (2.1) we can write

$$dA_{i^{(s-1)}} = \sum_{j=d_{s-1}+1}^N \omega_{i^{(s-1)}, j} A_j$$

from which we deduce, by (3.5)

$$(3.6) \quad \omega_{i^{(s-1)}, i^{(s+1)}} = \omega_{i^{(s-1)}, i^{(s+2)}} = \dots = \omega_{i^{(s-1)}, i^{(t)}} = \omega_{i^{(s-1)}, i^{(t+1)}} = 0.$$

These equations imply that

$$0 = d\omega_{i^{(s-1)}, i^{(\ell)}} = \sum_{i^{(s)}} \omega_{i^{(s-1)}, i^{(s)}} \wedge \omega_{i^{(s)}, i^{(\ell)}} \quad \ell \geq s+1;$$

now, by the inductive hypothesis, we have

$$(3.7) \quad \sum_{i^{(s-1)}} q_{i_1^{(1)}, \dots, i_{s-1}^{(1)}; i^{(s-1)}} \omega_{i^{(s-1)}, i^{(\ell)}} = \sum_{i_s^{(1)}} q_{i_1^{(1)}, \dots, i_s^{(1)}; i^{(\ell)}} \omega_{i_s^{(1)}}$$

where  $q_{i_1^{(1)}, \dots, i_{s-1}^{(1)}; i^{(s-1)}}$  are the coefficients of the  $(s-1)$ -th fundamental form and  $q_{i_1^{(1)}, \dots, i_s^{(1)}; i^{(\ell)}}$  are those of the  $s$ -th fundamental form (with the convention that  $q_{i_1^{(1)}, \dots, i_s^{(1)}; i^{(\ell)}} = 1$  if  $s \leq 1$ , and if  $s = 2$ , (3.7) is just (3.2) i.e. Cartan lemma used to obtain the second fundamental form (3.1)) and then, applying relations (3.7) to (a linear combination of) (3.6), we deduce

$$\sum_{i^{(s-1)}, i^{(s)}} q_{i_1^{(1)}, \dots, i_{s-1}^{(1)}; i^{(s-1)}} \omega_{i^{(s-1)}, i^{(s)}} \wedge \omega_{i^{(s)}, i^{(\ell)}} = \sum_{i_s^{(1)}, i^{(s)}} q_{i_1^{(1)}, \dots, i_s^{(1)}; i^{(s)}} \omega_{i_s^{(1)}} \wedge \omega_{i^{(s)}, i^{(\ell)}}$$

then, by Cartan lemma

$$\sum_{i^{(s)}} q_{i_1^{(1)}, \dots, i_s^{(1)}; i^{(s)}} \omega_{i^{(s)}, i^{(\ell)}} = \sum_{i_{s+1}^{(1)}} q_{i_1^{(1)}, \dots, i_{s+1}^{(1)}; i^{(\ell)}} \omega_{i_{s+1}^{(1)}};$$

and the  $(s+1)$ -th fundamental form is

$$V_{i^{(\ell)}}^{(s+1)} = \sum_{i_1^{(1)}, \dots, i_{s+1}^{(1)}} q_{i_1^{(1)}, \dots, i_{s+1}^{(1)}; i^{(\ell)}} \omega_{i_1^{(1)}} \cdots \omega_{i_{s+1}^{(1)}}.$$

□

*Remark 3.2.* Alternatively, we can write—for example writing back relations (3.4) in (3.3) or by a direct computation

$$V_{i^{(\ell)}}^{(s+1)} = \sum_{i^{(1)}, \dots, i^{(s)}} \omega_{i^{(1)}, i^{(2)}} \omega_{i^{(2)}, i^{(3)}} \cdots \omega_{i^{(s-1)}, i^{(s)}} \omega_{i^{(s)}, i^{(\ell)}},$$

with  $\ell \geq s+1$  and the sum varies, as above, for  $1 \leq i^{(1)} \leq k = d_1, \dots, d_{s-1} + 1 \leq i^{(s)} \leq d_s$ .

We recall

**Definition 3.3.** Let  $\Sigma$  be the linear system of dimension  $d$  of hypersurfaces of degree  $n > 1$  in  $\mathbb{P}^N$  ( $N > 1$ ), generated by the  $d+1$  hypersurfaces  $f_0 = 0, \dots, f_d = 0$ . The *Jacobian system*  $J(\Sigma)$  of  $\Sigma$  is the linear system defined by the partial derivatives of the forms  $f_0, \dots, f_d$ :

$$J(\Sigma) := (\partial f_i / \partial x_j) \quad i = 0, \dots, d; \quad j = 0, \dots, r.$$

*Remark 3.4.* Obviously, the Jacobian system  $J(\Sigma)$  does not depend on the choice of  $f_0, \dots, f_d$ , but only on  $\Sigma$ .

**Corollary 3.5.** *The  $(s+1)$ -th fundamental form is a linear system of polynomials of degree  $s+1$  whose Jacobian system is contained in the  $s$ -th fundamental form.*

*Proof.* We have, by Lemma 3.1

$$(3.8) \quad \frac{\partial V_{i^{(\ell)}}^{(s+1)}}{\partial \omega_{i_j^{(1)}}} = \sum_{i_1^{(1)}, \dots, i_s^{(1)}, i^{(s)}} q_{i_1^{(1)}, \dots, i_s^{(1)}, i^{(\ell)}} \omega_{i_1^{(1)}} \cdots \omega_{i_s^{(1)}} \frac{\partial \omega_{i^{(s)}, i^{(\ell)}}}{\partial \omega_{i_j^{(1)}}}.$$

□

We are now able to prove the following fact (see [8, (2.6)])

**Theorem 3.6.** *The  $s$ -th Gauss map  $\gamma^s$  ( $s \geq 1$ ) has fibres of dimension  $m$  if and only if at a general point  $P \in V$  the forms of degree  $s+1$  of the  $(s+1)$ -th fundamental form  $|I^{s+1}|$  are cones over a  $\mathbb{P}^{m-1} \subset \mathbb{P}T_P(V)$ .*

*Proof.* We choose a Darboux frame as above, i.e.

$$\{A_0; A_1, \dots, A_k; A_{k+1}, \dots, A_{d_2}; A_{d_2+1}, \dots, A_{d_s}; \dots, A_{d_t}; \dots, A_{d_{t+1}}; \dots, A_N\}$$

such that  $A_0, A_1, \dots, A_{d_s}$  span  $\tilde{T}_P^{(s)}(V)$  and indices  $d_{s-1} + 1 \leq i^{(s)} \leq d_s$  for all  $s = 1, \dots, t+1$ , with  $d_1 := k$ , etc.

In coordinates, the  $s$ -th Gauss map  $\gamma^s$  is

$$\gamma^s(P) = A_0 \wedge \cdots \wedge A_{d_s},$$

and therefore by (2.1), its differential, which is the  $(s+1)$ -th fundamental form, is given by

$$d\gamma_P^s \equiv \sum_{\substack{1 \leq i \leq d_s \\ d_s+1 \leq j \leq N}} (-1)^{d_s-i+1} \omega_{i,j} A_0 \wedge \cdots \wedge \hat{A}_i \wedge \cdots \wedge A_{d_s} \wedge A_j, \quad \text{mod } \tilde{T}_P^{(s)}V;$$

therefore, since  $\gamma^s$  has fibres of dimension  $m$  if and only if  $d\gamma_P^s$  has rank  $k-m$  (recall that  $P$  is a general point)  $\gamma^s$  has fibres of dimension  $m$  if and only if the space

$$U^* := \langle \omega_{i,j} \rangle_{\substack{1 \leq i \leq d_s \\ d_s+1 \leq j \leq N}} \subset T_P^*(V)$$

has dimension  $k-m$ . Dually, this space defines a subspace  $U \subset T_P(V)$  of dimension  $m$ , defined by the equations

$$\omega_{i^{(h)}, i^{(\ell)}} = 0, \quad h = 1, \dots, s; \quad \ell \geq s+1.$$

Now, we prove that  $V_{i^{(\ell)}}^{(s+1)}$  are cones with vertex  $\mathbb{P}(U)$ : let us suppose to choose a frame such that  $\omega_{m+1}, \dots, \omega_k$  form a basis for  $U^*$ , that is

$$(3.9) \quad \langle \omega_{i,j} \rangle_{\substack{1 \leq i \leq d_s \\ d_s+1 \leq j \leq N}} = \langle \omega_{m+1}, \dots, \omega_k \rangle.$$



Moreover, we choose indices  $1 \leq j_1^{(1)}, \dots, j_h^{(1)} \leq m$ , and  $m+1 \leq k_1^{(1)}, \dots, k_h^{(1)} \leq k$ .

From (3.4), the points of  $U$  satisfy also

$$(3.10) \quad 0 = \sum_{i^{(h)}} q_{i_1^{(1)}, \dots, i_h^{(1)}; i^{(h)}} \omega_{i^{(h)}, i^{(\ell)}} = \sum_{i_{h+1}^{(1)}} q_{i_1^{(1)}, \dots, i_{h+1}^{(1)}; i^{(\ell)}} \omega_{i_{h+1}^{(1)}},$$

which implies, for every choice of the indices as above, that

$$q_{i_1^{(1)}, \dots, i_{h+1}^{(1)}; i^{(\ell)}} = 0.$$

From this we infer, from (3.3), that

$$(3.11) \quad V_{i^{(\ell)}}^{(s+1)} = \sum_{k_1^{(1)}, \dots, k_{s+1}^{(1)}} q_{k_1^{(1)}, \dots, k_{s+1}^{(1)}; i^{(\ell)}} \omega_{k_1^{(1)}} \cdots \omega_{k_{s+1}^{(1)}},$$

that is,  $V_{i^{(\ell)}}^{(s+1)}$  is a cone with vertex  $\mathbb{P}^{m-1} = \mathbb{P}(U)$ .

Vice versa, let us suppose that the polynomials  $V_{i^{(\ell)}}^{(s+1)}$  are singular along a  $\mathbb{P}^{m-1} = \mathbb{P}(U)$ ; by choosing our frame, we can suppose that  $U$  is defined by

$$\omega_j = 0, \quad j = m+1, \dots, k,$$

or that (3.9) holds. Then, we have

$$(3.12) \quad 0 = d\omega_{i^{(h)}, i^{(\ell)}} = \sum_{i=1}^{d_s} \omega_{i^{(h)}, i} \wedge \omega_{i, i^{(\ell)}} + \sum_{j=d_s+1}^N \omega_{i^{(h)}, j} \wedge \omega_{j, i^{(\ell)}}$$

and  $\omega_{i, i^{(\ell)}}, \omega_{i^{(h)}, j} \in U^*$  and therefore the Frobenius integrability conditions for  $\omega_{m+1}, \dots, \omega_k$  are satisfied; then, from

$$dA_{i^{(s)}} = \sum_{j=d_s+1}^N \omega_{i^{(s)}, j} A_j$$

we conclude that the  $s$ -th osculating space remains constant along the leaves of the foliation defined by  $\omega_{m+1}, \dots, \omega_k$ .  $\square$

*Remark 3.7.* As we recalled in the introduction, one can deduce, from Theorem 3.6, that, for  $s = 1$  (i.e. the usual Gauss map), the fibres of  $\gamma^1$  are (Zariski open subsets of) linear spaces: see [8, (2.10)]. Moreover, in [8, §3] they give a general description of these varieties; we will pursue further this kind of description for higher  $s$  in Subsection 3.1.

*Example 3.8.* The first nontrivial example of application of the preceding theorem with  $s \geq 2$  is given by surfaces whose second Gauss map  $\gamma^2$  has as image a curve; this case has been studied in [3] and in [7]: they proved that the curves which are fibres of  $\gamma^2$  are indeed contained in 3-dimensional projective spaces; by Theorem 3.6, we deduce that the third fundamental form at the general point is given by a perfect cube.

Let us see the following consequence of the preceding result.

**Corollary 3.9.** *If the  $s$ -th Gauss map  $\gamma^s$  ( $s \geq 1$ ) for a  $k$ -dimensional variety  $V$  has fibres of dimension  $m$ , then*

$$(3.13) \quad d_{s+1} \leq d_s + \binom{k-m+s}{s+1},$$

and the equality is reached if and only if the  $(s+1)$ -th fundamental form  $|I^{s+1}|$  is given by the linear system of all the cones over a  $\mathbb{P}^{m-1} \subset \mathbb{P}T_P(V)$ .

*Proof.* By Proposition 2.10 we have to estimate the dimension  $\Delta_{s+1}$  of the linear system given by the  $(s+1)$ -th fundamental form, but this, by Theorem 3.6, is formed by degree  $(s+1)$  hypersurfaces which are cones over a  $\mathbb{P}^{m-1}$  in a  $\mathbb{P}^{k-1}$  ( $= \mathbb{P}(T_P(V))$ ). Since the linear systems of these cones has (projective) dimension  $\binom{k-m+s}{s+1} - 1$  (is equal to the complete linear system of degree  $(s+1)$  hypersurfaces in a  $\mathbb{P}^{k-m-2}$ ), we have the assertion.  $\square$

*Example 3.10.* Clearly, if  $m = k$ , then  $\gamma^s$  is constant,  $|I^{s+1}| = \emptyset$  and the variety is contained in a  $\mathbb{P}^{d_s}$ .

**3.1. Varieties with a curve as the image of the  $s$ -th Gauss map.** Let us start with the following

**Lemma 3.11.** *If  $d_{s+1} = d_s + 1$ , then  $d_{s+2} \leq d_s + 2$ . Moreover, if  $d_{s+2} = d_s + 2$ , then  $|I^{s+1}|$  and  $|I^{s+2}|$  are generated by powers of the same linear form, and if  $d_s + 1 = d_{s+2}$ , then our variety  $V$  is contained in  $\mathbb{P}^{d_s+2}$ .*

*Proof.* We apply Proposition 2.10 to deduce that the  $(s+1)$ -th fundamental form  $|I^{s+1}|$  is generated by only one form. Call this form  $f$ ; by Corollary 3.5, if  $g$  is a form in  $|I^{s+2}|$ , then all its partial derivatives are in  $|I^{s+1}|$ ; therefore, by Euler's formula

$$g = \frac{1}{s+2} \sum_{i=1}^k \omega_i \frac{\partial g}{\partial \omega_i} = \frac{1}{s+2} \sum_{i=1}^k \omega_i a_i f$$

$a_i \in \mathbb{C}$ ,  $i = 1, \dots, k$ , and we deduce that either  $g$  is zero or  $g = f\ell$ , where  $\ell$  is a linear form. If  $g = f\ell$ , by a change of coordinates we can suppose that  $\ell = \omega_1$ ; then, we have

$$\begin{aligned} \frac{\partial g}{\partial \omega_1} &= f + \omega_1 \frac{\partial f}{\partial \omega_1} \in |I^{s+1}| = (f) \\ \frac{\partial g}{\partial \omega_j} &= \omega_1 \frac{\partial f}{\partial \omega_j} \in |I^{s+1}| = (f) \quad j = 2, \dots, k \end{aligned}$$

which imply that either  $f$  does not depend on  $\omega_1$ , but then  $g$  is the zero form, or  $f$  is reducible of the form  $f = \omega_1 h$ . We can now proceed by induction to prove that  $f = \omega_1^{s+1}$ : if  $f = \omega_1^\ell \phi$ , then

$$\begin{aligned} \frac{\partial g}{\partial \omega_1} &= (\ell + 1)\omega_1^\ell \phi + \omega_1^{\ell+1} \frac{\partial \phi}{\partial \omega_1} = n\omega_1^\ell \phi \\ \frac{\partial g}{\partial \omega_j} &= \omega_1^{\ell+1} \frac{\partial \phi}{\partial \omega_j} = n_j \omega_1^\ell \phi \quad j = 2, \dots, k \end{aligned}$$

$n, n_j \in \mathbb{C}$ ,  $j = 1, \dots, k$ , and, as above, either  $\phi$  does not depend on  $\omega_1$ , but then  $g$  is the zero form, or  $\phi$  is reducible of the form  $\phi = \omega_1 \phi'$ .

If  $g = 0$ , then we can conclude from Theorem 3.6 (or, better, from Example 3.10).  $\square$

**Corollary 3.12.** *If  $d_{s+2} = d_{s+1} + 1 = d_s + 2$ , then the  $s$ -th and  $(s+1)$ -th Gauss maps have fibres of dimension  $k-1$ .*

*Proof.* By Lemma 3.11 we have that the  $|I^{s+1}|$  and  $|I^{s+2}|$  are cones over a  $\mathbb{P}^{k-2}$ , therefore, by Theorem 3.6, the thesis follows.  $\square$

The case we are interested in now is the one with  $m = k - 1$  (i. e. the next after Example 3.10), therefore, by (3.13),  $(d_s \leq) d_{s+1} \leq d_s + 1$ ; if  $d_{s+1} < d_s + 1$ , then  $d_{s+1} = d_s$ , but in this case we would have no elements in  $|I^{s+1}|$ , and therefore we would be in Example 3.10. We deduce that  $d_{s+1} = d_s + 1$ , and we are in the case of Lemma 3.11, and by Theorem 3.6 the image of the  $s$ -th Gauss map has dimension 1; the complete description of such cases will follow from [8, (2.4)]. To use this result, we need some notations and result: we start by recalling the notation introduced in [8, §2(a)]; let  $B$  be an  $r$ -dimensional variety and let

$$f: B \rightarrow \mathbb{G}(d, N)$$

be a morphism; let  $y \in B$  be a general point, and let  $S_y \subset V$  (where  $\mathbb{P}^N = \mathbb{P}(V)$ ) be the  $(d + 1)$ -dimensional vector space which corresponds to  $f(y)$ . Then, the differential of  $f$  in  $y$  can be thought of

$$d_y f: T_y B \rightarrow \text{Hom}(S_y, N_y),$$

where  $N_y := \frac{V}{S_y}$ . More explicitly, if  $y_1, \dots, y_r$  are local coordinates of  $B$  near  $y$ , and if  $e_0(y), \dots, e_d(y)$  is a basis for the  $(d + 1)$ -dimensional vector spaces near  $S_y$ , we have

$$d_y f \left( \frac{\partial}{\partial y_i} \right) (e_j(y)) \equiv \frac{\partial(e_j(y))}{\partial y_i} \pmod{S_y}$$

for  $i = 1, \dots, r$  and  $j = 0, \dots, d$ . Then, fixed  $w \in T_y B$ , one can define the “infinitely near” space

$$\frac{S_y}{dw} := \text{Im } d_y f(w) \subset N_y,$$

and if we denote by  $[v]$  the class of  $v \in V$  in  $N_y$ , we consider the following subspaces of  $V$ :

$$\begin{aligned} S_y + \frac{S_y}{dw} &:= \left\{ v \in V \mid v = s + t, s \in S_y, [t] \in \frac{S_y}{dw} \right\} \\ S_y \cap \frac{S_y}{dw} &:= \left\{ s \in S_y \mid ([0] =) [s] \in \frac{S_y}{dw} \right\} = \ker(d_y f(w)). \end{aligned}$$

Obviously, we have

$$\text{rk}(d_y f) = \rho \iff \dim(S_y + \frac{S_y}{dw}) = d + 1 + \rho \iff \dim(S_y \cap \frac{S_y}{dw}) = d + 1 - \rho.$$

We also denote the projective subspaces of  $\mathbb{P}^N$  associated to the vector spaces just defined by

$$\begin{aligned} \mathbb{P}_y^d + \frac{\mathbb{P}_y^d}{dw} &:= \mathbb{P}(S_y + \frac{S_y}{dw}) \\ \mathbb{P}_y^d \cap \frac{\mathbb{P}_y^d}{dw} &:= \mathbb{P}(S_y \cap \frac{S_y}{dw}). \end{aligned}$$

The case of  $r = 1$  is completely solved in [8]; as in the cited article, with abuse of notation we denote by  $y$  the local coordinate of  $B$  near our general point  $y$ :

**Proposition 3.13.** [8, (2.4)] *With notations as above, if  $r = 1$  and  $\dim(\mathbb{P}_y^d \cap \frac{\mathbb{P}_y^d}{d\omega}) = d - \rho$ , then there exist  $\rho$  curves in  $\mathbb{P}^N$ ,  $C_1, \dots, C_\rho$  parametrised by  $y$ ,  $a_1, \dots, a_\rho \in \mathbb{N}$ , and a fixed  $(d - \sum_{i=1}^\rho a_i)$ -dimensional subspace  $H \subset \mathbb{P}^N$  such that  $\mathbb{P}_y^d$  is the span of  $H$  together with the  $(a_i - 1)$ -osculating  $(a_i - 1)$ -planes  $\tilde{T}_P^{(a_i-1)}(C_i) \subset \mathbb{P}^N$ , that is*

$$\mathbb{P}_y^d = \langle \tilde{T}_P^{(a_1-1)}(C_1), \dots, \tilde{T}_P^{(a_\rho-1)}(C_\rho), H \rangle.$$

Let us now use this result in our case: we take  $B$  as the image of the  $s$ -th Gauss map:

$$\gamma^s: V \rightarrow \text{Im}(\gamma^s) =: B \xrightarrow{i} \mathbb{G}(d_s, N),$$

$\dim B = 1$  and  $S_y$  denotes just the  $d_s$ -dimensional vector subspace  $\tilde{T}_P^{(s)}(V) \subset \mathbb{C}^{N+1}$ , where  $y = \gamma^s(P)$ , defined by—with notations as in the proof of Theorem 3.6— $S_y := \langle A_0, \dots, A_{d_s} \rangle$ . Moreover, by (3.12), we have that  $B$  defines also a ruled variety, which is indeed the variety of osculating  $s$ -spaces to  $V$ :  $\text{Tan}^s(V) := \cup_{y \in B} \mathbb{P}_y^{d_s}$  of dimension  $d_s + 1$ . By Corollary 3.12, if  $\tilde{T}_P^{(s)}(V) \neq \tilde{T}_P^{(s+1)}(V)$ , the same happens with  $\tilde{T}_P^{(s+1)}(V)$ : we have, on  $B$ , the  $S'_y$ 's, which are the  $d_{s+1}$ -dimensional vector subspaces  $\tilde{T}_P^{(s+1)}(V) \subset \mathbb{C}^{N+1}$ , they define the (ruled) variety of osculating  $(s + 1)$ -spaces to  $V$   $\text{Tan}^{s+1}(V) := \cup_{y \in B} \mathbb{P}_y^{d_{s+1}}$  of dimension  $d_s + 2$  such that  $V \subset \text{Tan}^s(V) \subset \text{Tan}^{s+1}(V)$ .

Since, by definition of the  $(s + 1)$  osculating space, we have that, for  $y \in B$ ,

$$\mathbb{P}_y^{d_s} + \frac{\mathbb{P}_y^{d_s}}{dy} \subset \mathbb{P}_y^{d_{s+1}},$$

which implies

$$\dim(\mathbb{P}_y^{d_s} \cap \frac{\mathbb{P}_y^{d_s}}{dy}) = d_s - 1.$$

Then, by Proposition 3.13

**Proposition 3.14.** *If the  $s$ -th Gauss map has dimension one, then there exists a curve  $C \subset \mathbb{P}^N$  having  $(a - 1)$ -osculating  $(a - 1)$  planes  $\mathbb{P}_y^{a-1} := \tilde{T}_y^{(a-1)}(C)$  and a fixed  $\mathbb{P}^{N-a-1}$  such that*

$$\mathbb{P}_y^{d_s} = \mathbb{P}_y^{a-1} + \mathbb{P}^{N-a-1}.$$

*In other words,  $\text{Tan}^s(V)$ , is the cone over  $\mathbb{P}^{N-a-1}$  of  $\text{Tan}^{a-1}(C)$  (and the same holds true for higher Gauss maps).*

Clearly, one can deduce more general results on the varieties of osculating  $s$ -spaces in the same vein as at the end of [8, §3].

## REFERENCES

- [1] M. A. Akivis and V. V. Goldberg, *Differential Geometry of Varieties with Degenerate Gauss Maps*, CMS Books in Mathematics/Ouvrages de Mathématiques de la SMC, 18. Springer-Verlag, New York, 2004.
- [2] E. Bertini, *Introduzione alla geometria proiettiva degli iperspazi con appendice sulle curve algebriche e loro singolarità*, E. Spoerri, Pisa, 1907.
- [3] M. Castellani, *Sulle superficie i cui spazi osculatori sono biosculatori*, Rom. Acc. L. Rend. (5) **31** (1922), no. 1, 347–350.
- [4] P. De Poi, R. Di Gennaro and G. Ilardi, *On varieties with higher osculating spaces*, Rev. Mat. Iberoam. **29** (2013), no. 4, 191–1210.

- [5] P. del Pezzo, *Sugli spazii tangenti ad una superficie o ad una varietà immersa in uno spazio di più dimensioni*, Nap. Rend. **XXV** (1886), 176–180.
- [6] S. Di Rocco, K. Jabbusch and A. Lundman, *A note on higher order Gauss maps*, arXiv:1410.4811[math.AG]
- [7] D. Franco and G. Ilardi, *On multiosculating spaces*, Commun. Algebra, **29** (2001), no. 7, 2961–2976 .
- [8] P. Griffiths and J. Harris *Algebraic geometry and local differential geometry*, Ann. Sci. École Norm. Sup. (4), **12** (1979), no. 3, 355–452.
- [9] R. Hartshorne, *Algebraic Geometry*, Number 52 in Graduate Text in Mathematics. Springer-Verlag, New York, 1977.
- [10] G. Ilardi, *Togliatti systems*, Osaka J. Math. **43** (2006), no. 1, 1–12.
- [11] J. M. Landsberg, *On second fundamental forms of projective varieties*, Invent. Math. **117** (1994), no. 2, 303–315.
- [12] R. Piene, *A note on higher order dual varieties, with an application to scrolls*, Singularities, Part 2 (Arcata, Calif., 1981), 1983, pp. 335–342.
- [13] C. Segre, *Preliminari di una teoria delle varietà luoghi di spazii*, Palermo Rend. **30** (1910), 87–121.
- [14] E. Togliatti, *Alcuni esempi di superficie algebriche degli iperspazii che rappresentano un'equazione di Laplace*, Comm. Math. Helvetici **1** (1929), 255–272.
- [15] F. L. Zak, *Tangents and secants of algebraic varieties*, Translations of Mathematical Monographs, vol. 127, American Mathematical Society, Providence, RI, 1993. Translated from the Russian manuscript by the author.

DIPARTIMENTO DI MATEMATICA E INFORMATICA, UNIVERSITÀ DEGLI STUDI DI UDINE, VIA DELLE SCIENZE, 206, 33100 UDINE, ITALY

*E-mail address:* `pietro.depoi@uniud.it`

DIPARTIMENTO DI MATEMATICA E APPLICAZIONI, UNIVERSITÀ DEGLI STUDI DI NAPOLI “FEDERICO II”, 80126 NAPOLI, ITALY

*E-mail address:* `giovanna.ilardi@unina.it`