



UNIVERSITÀ
DEGLI STUDI
DI UDINE

Università degli studi di Udine

A general tool for consistency results related to I1

Original

Availability:

This version is available <http://hdl.handle.net/11390/1109744> since 2021-03-16T16:52:33Z

Publisher:

Published

DOI:10.1007/s40879-015-0092-y

Terms of use:

The institutional repository of the University of Udine (<http://air.uniud.it>) is provided by ARIC services. The aim is to enable open access to all the world.

Publisher copyright

(Article begins on next page)

A general tool for consistency results related to $I1$

Vincenzo Dimonte*, Liuzhen Wu[†]

December 15, 2014

Abstract

In this paper we provide a general tool to prove the consistency of $I1(\lambda)$ with various combinatorial properties at λ typical at settings with $2^\lambda > \lambda^+$, that does not need a profound knowledge of the forcing notions involved. Examples of such properties are the first failure of GCH, a very good scale and the negation of the approachability property, or the tree property at λ^+ and λ^{++} .

Keywords: Infinite combinatorics, rank-into-rank, Prikry forcing, singular cardinal hypothesis.

2010 Mathematics Subject Classifications: 03E55, 03E05, 03E35(03E45)

1 Introduction

While Cantor gave us the means to conceive infinite cardinals, it is clear that to use them in a fruitful way a thorough study of their structure is needed, and this is the aim of the field usually called “infinite combinatorics”. The key turning point for this study, as old as Set Theory, was the introduction of forcing [1]: it was clear then that much of the structural properties are independent from ZFC, therefore shifting the focus of the study from what is the structure of infinite cardinals, to what it could be. In the few years after Cohen’s seminal results, the analysis of regular cardinals was pretty much complete, with the definition of many forcings that can change effortlessly the

*Technische Universität Wien, Wiedner Hauptstraße 8–10, 1040 Wien, Austria *E-mail address:* vincenzo.dimonte@gmail.com, *URL:* <http://www.logic.univie.ac.at/~dimonte/>

[†]Institute of Mathematics, Chinese Academy of Sciences, East Zhong Guan Cun Road No 55, Beijing 100190 China *E-mail address:* lzwu@math.ac.cn

combinatorial properties not only of a single cardinal, but, using a method introduced by Easton [6], to all regular cardinals at once.

Changing combinatorial properties of singular cardinals, especially those of cofinality ω , proved to be much harder than in the regular cardinal case. As a result, the research on such properties is rich and varied, it provided and still provides many challenges. Some situations are even impossible: Silver [16] proved that SCH cannot fail first at a singular cardinal of uncountable cofinality, and Solovay [17] proved that above a strongly compact cardinal SCH must hold. The typical way to make SCH fail at a singular cardinal (i.e. blowing up the cardinality of its powerset) is to start with κ measurable, blowing up its power and then adding an ω -sequence cofinal to κ with Prikry forcing, to make it of cofinality ω . But $2^\kappa > \kappa^+$ permits a multitude of properties to hold, and it is an ongoing research to find more and more sophisticated variations of the Prikry forcing that permit different combinations of specific combinatorics on a singular cardinal.

While this research heavily involves large cardinals, their role has almost always been giving consistency strength to a certain scenery, but they rarely appear directly with the desired combinatorial property, for the simple reason that the great majority of large cardinals are regular cardinals, therefore unrelated to the problem. Moreover some large cardinal simply do not accept a lot of variety on the structure of singular cardinals, as noted above. Going up the hierarchy, however, one can find an exception. The strongest large cardinal axioms, called rank-into-rank axioms, do involve a singular cardinal of countable cofinality, and it is very natural to question the position of them in this field: as they imply all known large cardinal hypothesis, knowing their structure is crucial in the large cardinal field, as it trickle down to all the hierarchy.

Woodin in [18] introduced what he called “Generic Absoluteness for I_0 ”, and this proved to be key for this study: starting with I_0 , adding a Prikry sequence to its critical point κ , an action that we noted typical for proving consistency results for singular cardinals, adds in fact I_1 . In [4] this was exploited to prove that it was possible to have $j : V_{\lambda+1} \prec V_{\lambda+1}$ and $2^\lambda > \lambda^+$ at the same time, in the same way it was proved just with a measurable.

If the proof of $2^\lambda > \lambda^+$ uses Prikry forcing, could we use the sophisticated variations of the Prikry forcing that appear in literature to prove the consistency of different combinations of specific combinatorics on a singular cardinal with I_1 ? In this paper, we extend such theorem, describing a general procedure to be applied to many of the refined Prikry forcings, therefore automatically transferring the already known results about the combinatorics of singular cardinal of countable cofinality to cardinals that moreover satisfy very large cardinal properties, providing therefore a number of new results

and a tool that any researcher can use without going into the original details of the forcing notions involved.

In Section 2 all the preliminary facts are collected. In Section 3, the general procedure is described: the notion of κ -geometric forcing is introduced, and this is the key notion that will permit the proof to work; the procedure is tested with Prikry forcing and tree Prikry forcing. In Section 4, the procedure is applied to the extender-based Prikry forcing, to provide $I1(\lambda)$ and the first failure of GCH at λ (this answers a question in [4]). In Section 5, the procedure is applied to two different flavors of diagonal supercompact Prikry forcing, to achieve results on pcf combinatorics and the Tree Property. In Section 6, we see another application to the Tree Property. In Section 7 we note some possible directions for future research on the subject.

The first author would like to thank the FWF (Austrian Science Fund) for its generous support through project M 1514-N25, and the kind hospitality of the Kurt Gödel Research Center, Beijing Normal University and the Chinese Academy of Sciences. The second author would like to acknowledge the support through the funding projects NCSF 11321101 and 11401567.

2 Preliminaries

To avoid confusion or misunderstandings, all notation and standard basic results are collected here.

Elementary embeddings have a key role in the definitions of all large cardinals from measurable to above.

If M and N are sets or classes, $j : M \prec N$ denotes that j is an elementary embedding from M to N . We write the case in which the elementary embedding is the identity, i.e., M is an elementary submodel of N , simply as $M \prec N$, while when j is indicated we always suppose that it is not the identity.

If $j : M \prec N$ and either $M \models \text{AC}$ or $N \subseteq M$ then it moves at least one ordinal. The *critical point*, $\text{crt}(j)$, is the least ordinal moved by j .

If $j : M \prec N$ and $N \subseteq M$, we define j^n as the composition of n copies of j , i.e., $j^1 = j$ and $j^{n+1} = j \circ j^n$.

Let j be an elementary embedding and $\kappa = \text{crt}(j)$. Define $\kappa_0 = \kappa$ and $\kappa_{n+1} = j(\kappa_n)$. Then $\langle \kappa_n : n \in \omega \rangle$ is the *critical sequence* of j .

Kunen [11] proved under AC that if $M = N = V_\eta$ for some ordinal $\eta \leq \text{Ord}$, and λ is the supremum of the critical sequence, then η cannot be bigger than $\lambda + 1$ (and of course cannot be smaller than λ).

This at the time was considered a stop to the large cardinal study, as a $j : V \prec V$ would have been the largest possible cardinal, but Kunen's result

leaves room for a new breed of large cardinal hypotheses, sometimes referred to in the literature as rank-into-rank hypotheses:

- I3** iff there exists λ s.t. $\exists j : V_\lambda \prec V_\lambda$;
- I2** iff there exists λ s.t. $\exists j : V \prec M$, with $V_\lambda \subseteq M$ and λ is the supremum of the critical sequence;
- I1** iff there exists λ s.t. $\exists j : V_{\lambda+1} \prec V_{\lambda+1}$.

The consistency order of the above hypotheses is reversed with respect to their numbering: I1 is strictly stronger than I2, which in turn is strictly stronger than I3 (see [12]). All of these hypotheses are strictly stronger than all of the large cardinal hypotheses outside the rank-into-rank umbrella (see [10], 24.9 for n -huge cardinals, or [2] for the Wholeness Axiom). I3 enjoyed a particularly rich literature, as it has an interesting algebraic content [3].

Note that if j witnesses a rank-into-rank hypothesis, then λ is uniquely determined by j , so in the following λ always denotes the first nontrivial fixed point of the embedding j under consideration. We write $I1(\lambda)$ for $\exists j : V_{\lambda+1} \prec V_{\lambda+1}$, and $I1(\kappa, \lambda)$ for $\exists j : V_{\lambda+1} \prec V_{\lambda+1}$ and $\text{crt}(j) = \kappa$.

In the early 1980's Woodin proposed an axiom even stronger than all the previous ones:

- I0** For some λ there exists a $j : L(V_{\lambda+1}) \prec L(V_{\lambda+1})$, with $\text{crt}(j) < \lambda$.

Again, $I0(\lambda)$ and $I0(\kappa, \lambda)$ express what is expected.

Note that if λ witnesses I0, then $L(V_{\lambda+1}) \not\models \text{AC}$, because otherwise $L(V_{\lambda+1}) \models \text{ZFC}$, and we would contradict the proof of Kunen's Theorem [11], which shows that one cannot have $j : V \prec V$ with critical point less than λ and a well-order of $V_{\lambda+1}$ in V . The fact that I0 is strictly stronger than I1 was proved by Laver [12].

I0 is probably the most interesting of the rank-into-rank axioms: it is the only very large cardinal that induces a structure on an inner model, therefore creating a new field of research and new tools, and moreover the structure is reminiscent of the one induced by the Axiom of Determinacy, for reasons that are still not completely understood [18].

An embedding that witnesses I0 has an ultrapower structure:

Lemma 2.1. *Let $j : L(V_{\lambda+1}) \prec L(V_{\lambda+1})$ be such that $\text{crt}(j) < \lambda$. Let*

$$U = U_j = \{X \in L(V_{\lambda+1}) \cap V_{\lambda+2} : j \restriction V_\lambda \in j(X)\}.$$

Then U is an $L(V_{\lambda+1})$ -ultrafilter such that $\text{Ult}(L(V_{\lambda+1}), U)$ is well-founded. By condensation the collapse of $\text{Ult}(L(V_{\lambda+1}), U)$ is $L(V_{\lambda+1})$, and $j_U : L(V_{\lambda+1}) \prec$

$L(V_{\lambda+1})$, the inverse of the collapse, is an elementary embedding. Moreover, there is an elementary embedding $k_U : L(V_{\lambda+1}) \prec L(V_{\lambda+1})$ with $\text{crt}(k_U) > \Theta^{L(V_{\lambda+1})}$ such that $j = k_U \circ j_U$.

We can use the ultrapower structure to define iterability for j :

Definition 2.2. Let $j : L(V_{\lambda+1}) \prec L(V_{\lambda+1})$ with $\text{crt}(j) < \lambda$ be an elementary embedding, and suppose j is generated by $U = U_j$. Define

$$j(U) = \bigcup \{j(\text{ran}(\pi)) : \pi \in L(V_{\lambda+1}), \pi : V_{\lambda+1} \rightarrow U\}$$

and then define j_2 as the map associated to $j(U)$.

Define the successive iterates in the usual way: let α be an ordinal. Then

- if $\alpha = \beta + 1$, M_β is well-founded and $j_\beta : M_\beta \prec M_\beta$ is the ultrapower via W , then $M_\alpha = \text{Ult}(M_\beta, j_\beta(W))$ and $j_\alpha = j_\beta(j_\beta)$.
- if α is a limit, let (M_α, j_α) be the direct limit of (M_β, j_β) with $\beta < \alpha$.

We say that j is iterable, if for every $\alpha \in \text{Ord}$, M_α is well-founded and $j_\alpha : M_\alpha \prec M_\alpha$. In this case, we call $j_{\alpha,\beta}$ the natural embeddings between M_α and M_β .

The following is a conjunction of Lemma 16 and Lemma 21 in [18]:

Theorem 2.3. Let $j : L(V_{\lambda+1}) \prec L(V_{\lambda+1})$ with $\text{crt}(j) < \lambda$ be a proper elementary embedding. Then j is iterable. Moreover, for any $n \in \omega$, $j_n : L(V_{\lambda+1}) \prec L(V_{\lambda+1})$.

Theorem 2.3 states that $M_n = L(V_{\lambda+1})$ for $n < \omega$, but M_ω is definitively different. The key point is that $j_{0,\omega}(\text{crt}(j)) = \lambda$, so many characteristics of the critical point of j are transferred by elementarity to λ in M_ω . For example, in $L(V_{\lambda+1})$, $\text{crt}(j)$ is measurable and there is a well-ordering of V_λ , therefore λ is measurable in M_ω and there is a well-ordering of $V_{j_{0,\omega}(\lambda)}^{M_\omega} = V_{j_{0,\omega}(\lambda)} \cap M_\omega$ in M_ω .

Trees are a typical structure that is investigated in combinatorics. Let α be an ordinal. For any $s \in [\alpha]^n$, $\text{lh}(s) = n$. A tree on α is a subset of $[\alpha]^{<\omega}$ closed under initial segments. If T is a tree, for any $s \in T$, denote $T_s = \{t \in T : t \subseteq s \wedge s \subseteq t\}$, $\text{Suc}_T(s) = \{\beta \in \alpha : t \frown \langle \beta \rangle \in T\}$ and finally for any $n \in \omega$, $\text{Lev}_n(T) = \{s \in T : \text{lh}(s) = n\}$.

3 General procedure

In [18] Woodin introduced Generic Absoluteness for $I0$, while in [4] one of the authors and Sy Friedman used it to prove a single result about the power function and rank-into-rank embeddings. Here we introduce a general procedure that extends the scope of [4] to many more kinds of forcing, and in the next sections we will give some important examples.

One of the most important forcing in dealing with the combinatorics of singular cardinals of cofinality ω is Prikry forcing. It adds a cofinal sequence to a measurable cardinal.

Definition 3.1. *A cardinal κ is measurable iff there exists a κ -complete ultrafilter on κ .*

Definition 3.2. *Let κ be a measurable cardinal. Fix U a normal measure on κ . Define $p \in \mathbb{P}$ iff $p = (s, A)$, where $s \in [\kappa]^{<\omega}$, $A \in U$ and $\bigcup s < \bigcap A$. For $p = (s, A)$, $q = (t, B) \in \mathbb{P}$, we say $q \leq p$ iff $s \subseteq t$, $B \subseteq A$ and $t \setminus s \subseteq A$.*

Prikry forcing is useful because it is a very ”‘delicate’” forcing [8]: it does not add bounded subsets of κ , and is κ^+ -cc, so it does not change the cardinal structure above κ . In other words, it makes κ singular while changing the universe at least as possible.

The following is instead the tree Prikry forcing:

Definition 3.3. *Let κ be a measurable cardinal. Fix U an ultrafilter on κ . The tree Prikry forcing \mathbb{P} is the set of conditions $p = (s_p, T^p)$, where s_p is a finite sequence of ordinals in κ , and T^p is a tree of increasing sequences in κ with stem s_p , such that for any $t \in T^p$, $\text{Suc}_{T^p}(t) \in U$. We say that $p \leq q$ if $s_p \supseteq s_q$ and $T^p \subseteq T^q$. We say that $p \leq^* q$ if $p \leq q$ and $s_p = s_q$. For any $p \in \mathbb{P}$ and $t \in T^p$, we write $p \oplus t$ for $(t, (T^p)_t)$.*

The difference between the two forcings is minimal: the only difference is that standard Prikry forcing uses a normal ultrafilter, while for tree Prikry forcing normality is not needed. As for the majority of times the ultrafilters are normal, the two forcing are interchangeable, and using one or the other is a matter of better clarity of the proof.

The general technique, in short, will be to start with $I0(\kappa, \lambda)$, and then the following Theorem 3.11 expresses the fact that if one forces with an ”Easton-like” forcing and then with a ”Prikry-like” forcing at κ , by Generic Absoluteness one can have $I1(\kappa)$. While we have already a formal definition for ”Easton-like” (reverse Easton iteration), we need a definition of ”Prikry-like” suitable to our wants.

Definition 3.4. Let \mathbb{P} be a forcing notion and κ a cardinal. We say that \mathbb{P} is κ -geometric if

- there exists a length measure of the conditions of \mathbb{P} , i.e. $l : \mathbb{P} \rightarrow \omega$ such that $l(1_{\mathbb{P}}) = 0$ and for any $p, q \in \mathbb{P}$, if $p \leq q$ then $l(p) \geq l(q)$.
- for any $\alpha < \kappa$, if $\langle D_\beta : \beta < \alpha \rangle$ is a collection of open dense sets, then there is a condition q such that whenever a filter contains q and meets all the dense open sets $E_n = \{p : l(p) > n\}$, it also meets all the D_β 's.

This notion implies the notion of κ -goodness that was first introduced by Woodin in [18], and then perfected by Shi in [15]. The change is due to the fact that κ -geometric is more natural in working with tree Prikry-like forcings, even if it works in general:

Lemma 3.5. *Prikry forcing on κ is κ -geometric.*

Proof. It is a well known fact that for any D dense set and any $p \in \mathbb{P}$, there exists $q = (s, A) \leq^* p$ and $n \in \omega$ such that for any $t \in [A]^n$, $(q \cup t, A \setminus (\max(t) + 1)) \in D$, see for example Lemma 1.13 in [8]. Now, suppose $\alpha < \kappa$ and $\langle D_\beta : \beta < \alpha \rangle$ is a collection of open dense sets. Let $\langle q_\beta : \beta \leq \alpha \rangle$ be the sequence built with an iteration of the first sentence, i.e., for any $\beta \leq \alpha$, $q_{\beta+1} \leq^* q_\beta$ is such that there exists an $n_{\beta+1}$ such that for any t of length n , $(q \cup t, A \setminus (\max(t) + 1)) \in D_\beta$, and if $\beta \leq \alpha$ is limit, then by κ -closeness of U let q_β be just $(s, \bigcap_{\gamma < \beta} A_{q_\gamma})$.

Therefore q_α is as wanted: let $q' \leq q_\alpha = (s_{q_\alpha}, A_{q_\alpha})$ such that $q' \in E_{n_\beta}$. Then

$$q' < (q_\alpha \cup t, A_{q_\alpha} \setminus (\max(t) + 1)) \leq (q_\beta \cup t, A_{q_\beta} \setminus (\max(t) + 1)) \in D_\beta$$

for some t with length bigger than n_β . \square

The following is the most basic non-immediate example, and the method used in the proof is also the base for the more sophisticated methods in the next sections:

Lemma 3.6. *Let κ be a measurable cardinal. Then the tree Prikry forcing \mathbb{P} on κ is κ -geometric.*

Proof. Fix U a measure on κ . For any $p \in \mathbb{P}$, we define $l(p) = \text{lh}(s_p)$. The proof is in four steps:

- in the first claim, we modify the tree T^p so that if some condition $r \leq p$ is in D , then all the conditions $t \leq p$ such that $s_r = s_t$ are in D ;

- in the second claim, we modify the previous tree, so that if some condition $r \leq p$ with $T^s = (T^p)_s$ is in D , then all the conditions $t \leq p$ with $T^t = (T^p)_t$ and $\text{lh}(s_r) = \text{lh}(s_t)$ are in D ;
- in the third claim, we put the two claims together, so that there always exists a fixed length so that all the conditions that extend p of that length are in D ;
- finally, with the third claim we prove the lemma.

Claim 3.7 (First claim). *For any D open dense set and for any $p \in \mathbb{P}$, there exists $q \leq^* p$ such that if there exists $r = (s_r, T^r) \leq q$ (i.e. $r \leq^* q \oplus s_r$) such that $r \in D$, then $q \oplus s_r \in D$.*

Proof of claim. We can suppose $p = 1_{\mathbb{P}}$. It is done by induction. Informally, we consider $T^{1_{\mathbb{P}}}$, and we restrict it asking at each level whether there is a possible way to shrink it to reach D : if there is, then we just shrink it; otherwise we do nothing. More formally:


- if there exists a $r \leq^* 1_{\mathbb{P}}$ such that $r \in D$, then let S_0 be T^r ; otherwise $S_0 = T^{1_{\mathbb{P}}}$ (note that in the first case $(\langle \rangle, S_0) = r \in D$);
- if there exists a $r \leq (\langle \rangle, S_n)$ such that $l(s_r) = n + 1$ and $r \in D$, then let $(S_{n+1})_{s_r} = (T^r)_{s_r}$; otherwise $(S_{n+1})_{s_r} = (S_n)_{s_r}$ (note that in the first case $(s_r, (S_{n+1})_{s_r}) = r \in D$); to complete the definition, let $S_{n+1} \upharpoonright [\kappa]^{n+1} = S_n \upharpoonright [\kappa]^{n+1}$.

Let $S = \bigcap_{n \in \omega} S_n$ and $q = (\langle \rangle, S)$. Then q is as desired: since the n -th level is changed just in the first n steps, we have that for any $s \in S$ of length $n \in \omega$, $\text{Suc}_S(s) = \bigcap_{m < n} \text{Suc}_{S_m}(s) \in U$, therefore $q \in \mathbb{P}$. Let $t \leq^* q \oplus s_t$, i.e. $s_t \in S$ and $T^t \subseteq S$, with $t \in D$. Suppose $\text{lh}(s_t) = n$. Then also $s_t \in S_n$, as $S_{n+1} \upharpoonright [\kappa]^n = S_n \upharpoonright [\kappa]^n$, so $t \leq (\langle \rangle, S_n)$. This means that in the construction the first case was true, therefore $q \oplus s_t = (s_t, S_{s_t}) \leq (s_t, (S_{n+1})_{s_t}) \in D$. \square

Claim 3.8 (Second claim). *For any D open dense set and for any $p \in \mathbb{P}$, there exists $q \leq^* p$ and $n \in \omega$ such that for any $s_1, s_2 \in T^q$ such that $l(s_1) = l(s_2) = n$, $(s_1, (T^q)_{s_1}) \in D$ iff $(s_2, (T^q)_{s_2}) \in D$.*

Proof. We can still assume $p = 1_{\mathbb{P}}$. Let $R = \{s \in T^{1_{\mathbb{P}}} : p \oplus s \in D\}$.

Informally, we are climbing up level by level, deleting at each level either the branches that are in R or the ones that are not, so that the sets of successors are still in U . The first step, then, will be simple: let $B_{\langle \rangle}^0 = \{\delta \in \text{Suc}_{T^{1_{\mathbb{P}}}}(\langle \rangle) : \langle \delta \rangle \in R\}$. Then either $B_{\langle \rangle}^0$ or $\text{Suc}_{T^{1_{\mathbb{P}}}}(\langle \rangle) \setminus B_{\langle \rangle}^0$ are in U . Call such $A_{\langle \rangle}^0$. Then let $\langle \mu_0, \dots, \mu_l \rangle \in S^0$ iff $\mu_0 \in A_{\langle \rangle}^0$.




Note that in S^0 , $\text{Suc}_{S^0}(\langle \rangle) = \text{Suc}_{T^{1\mathbb{P}}}(\langle \rangle) \cap A_{\langle \rangle}^0$, while for all $s \in S^0$ of length ≥ 1 , $\text{Suc}_{S^0}(s) = \text{Suc}_{T^{1\mathbb{P}}}(s)$, and the sequences in S^0 of length 1 are either all in R or all outside.


The second step shows more complexity. First, for any $\langle \mu \rangle \in S^0$ we restrict its successors so that they are either all in R or all outside R . Therefore let

$$B_{\langle \mu \rangle}^1 = \{\delta \in \text{Suc}_{S^0}(\langle \mu \rangle) : \langle \mu, \delta \rangle \in R\}$$

for any $\langle \mu \rangle \in \text{Suc}_{S^0}(\langle \rangle)$. Then either $B_{\langle \mu \rangle}^1$ or $\text{Suc}_{S^0}(\langle \mu \rangle) \setminus B_{\langle \mu \rangle}^1$ is in U . Call it $A_{\langle \mu \rangle}^1$. Now define $S^{1,0}$ so that $\langle \mu_0, \dots, \mu_l \rangle \in S^{1,0}$ iff $\mu_0 \in A_{\langle \rangle}^0$ and $\mu_1 \in A_{\langle \mu_0 \rangle}^1$. Note that for all $s \in S^{1,0}$ with $\text{lh}(s) = 1$, $\text{Suc}_{S_0^1}(s) = \text{Suc}_{S^0}(s) \cap A_{\langle s(0) \rangle}^1$, while if $\text{lh}(s) \neq 1$ then $\text{Suc}_{S_0^1}(s) = \text{Suc}_{S^0}(s)$.

This is not enough. Singularly, all the 2-sequences that share the same root are either all in R or all outside, but it can be that all the 2-sequences that start with μ_1 are in R , and all the 2-sequences that start with μ_2 are not in R . Therefore we must choose only the μ 's that give a consistent result.





Let

$$B_{\langle \rangle}^1 = \{\mu \in \text{Suc}_{S^0}(\langle \rangle) : \text{Suc}_{S^{1,0}}(\langle \mu \rangle) = B_{\langle \mu \rangle}^1\},$$

i.e. the set of μ 's such that for any $\delta \in \text{Suc}_{S^{1,0}}(\langle \mu \rangle)$, $\langle \mu, \delta \rangle \in R$. Then either $B_{\langle \rangle}^1$ or $\text{Suc}_{S^0}(\langle \rangle) \setminus B_{\langle \rangle}^1$ is in U . Let $A_{\langle \rangle}^1$ be it. Now define $S^{1,1} = S^1$ as $\langle \mu_0, \dots, \mu_l \rangle \in S^1$ iff $\langle \mu_0, \dots, \mu_l \rangle \in S^{1,0}$ and $\mu_0 \in A_{\langle \rangle}^1$. Note that for all $s \in S^1$, if $\text{lh}(s) = 0$ then $\text{Suc}_{S^1}(s) = \text{Suc}_{S^{1,0}}(s) \cap A_{\langle \rangle}^1$, otherwise $\text{Suc}_{S^1}(s) = \text{Suc}_{S^{1,0}}(s)$. The sequences in S^1 of length 2 are either all in R or all outside it.

By induction the construction continues level-by-level, each time starting with $S^{n+1,0} \subseteq S^n$, and then going down to S^n , a tree such that all the $n+1$ -branches are either all in R or all outside it. More technically, suppose S^n is defined. For all $t \in S^n$, $\text{lh}(t) = n+1$, define $B_t^{n+1} = \{\delta \in \text{Suc}_{S^n}(t) : t \frown \langle \delta \rangle \in R\}$. Then either B_t^{n+1} or $\text{Suc}_{S^n}(t) \setminus B_t^{n+1}$ is in U . Let A_t^{n+1} be it. Define $\langle \mu_0, \dots, \mu_l \rangle \in S^{n+1,0}$ iff $\langle \mu_0, \dots, \mu_l \rangle \in S^n$ and $\mu_{n+1} \in A_{\langle \mu_0, \dots, \mu_n \rangle}^{n+1}$. Note that for all $s \in S^{n+1,0}$, $\text{lh}(s) = n+1$,

$$\text{Suc}_{S^{n+1,0}}(s) = \text{Suc}_{S^n}(s) \cap A_s^{n+1},$$

otherwise $\text{Suc}_{S^{n+1,0}}(s) = \text{Suc}_{S^n}(s)$.

Let $t \in \text{Lev}_m S^{n+1}$ and suppose that $S^{n+1,n-m}$, B_s^{n+1} and A_s^{n+1} are defined for all $s \in S^{n+1}$ with $\text{lh}(s) = m+1$. Let

$$B_t^{n+1} = \{\delta \in \text{Suc}_{S^{n+1}}(t) : \text{Suc}_{S^{n+1,n-m}}(t \frown \langle \delta \rangle) = B_{t \frown \langle \delta \rangle}^{n+1}\}.$$

Then either B_t^{n+1} or $\text{Suc}_{S^{n+1}}(t) \setminus B_t^{n+1}$ is in U . Let A_t^{n+1} be it.

Suppose A_t^{n+1} is defined for all $t \in S^{n+1}$ of length m . Then $\langle \mu_0, \dots, \mu_l \rangle \in S^{n+1,n+1-m}$ iff $\langle \mu_0, \dots, \mu_l \rangle \in S^{n+1,n-m}$ and $\mu_i \in A_{\langle \mu_0, \dots, \mu_m \rangle}^{n+1}$. Note that for all $s \in S^{n+1,n+1-m}$ of length $n+1-m$,

$$\text{Suc}_{S^{n+1,n+1-m}}(s) = \text{Suc}_{S^{n+1,n-m}}(s) \cap A_s^{n+1},$$

otherwise $\text{Suc}_{S^{n+1,n+1-m}}(s) = \text{Suc}_{S^{n+1,n-m}}(s)$. Call $S^{n+1,n+1} = S^{n+1}$. Then all the sequences in S^{n+1} of length $n+1$ either are all in R or all outside it.

Now, let $S = \bigcap_{n \in \omega} S^n$. The last remark is sufficient to prove the claim. We prove that $(\langle \rangle, S) \in \mathbb{P}$. It suffices to prove that for any $t \in S$, $\text{Suc}_S(t) \in U$.

So let $t \in S$, $\text{lh}(t) = n$. Then $\text{Suc}_S(t)$ will be modified in the construction of S only in the stages $S^{n+1, n}$ with $i \in \omega$, therefore

$$\text{Suc}_S(t) = \text{Suc}_{T^1\mathbb{P}}(t) \cap \bigcap_{i \in \omega} A_t^{n+i},$$

that is a countable intersection of elements of U , and therefore in U . \square

Claim 3.9 (Third claim). *For any $p \in \mathbb{P}$ and for any D open dense there exists a $p \leq^* q$ and an $n \in \omega$ such that for any $t \leq q$ with $\text{lh}(t) = n$, $t \in D$.*

Proof of claim. Putting the first and second claims together, we have that for any D dense set in \mathbb{P} and for any $p \in \mathbb{P}$ there exists a $q \leq^* p$ and a $n \in \omega$ such that for all $s \in T^q$ with $\text{lh}(s) = n$, $q \oplus s \in D$. Pick a $q \leq^* p$ as the first claim and a $q' \leq^* q$ as the second claim. By density, there exists a $r \leq q$, $r \in D$. Let $n = \text{lh}(r)$. Then by the first claim $q \oplus s_r \in D$, and by the second claim we proved that all the extensions of q of the same length of r are in D . \square

Now the proof is exactly as in Lemma 3.5. Let $\alpha < \kappa$ and $\langle D_\beta : \beta < \alpha \rangle$ be a collection of open dense sets. Let $\langle q_\beta : \beta \leq \alpha \rangle$ be the sequence built with an iteration of the third claim, and let q_α be the \leq^* -infimum of the q_β 's. Then q_α is as wanted: let $q' \leq q_\alpha$ such that $q' \in E_{n_\beta}$. Then $q' < q_\alpha \oplus t \leq q_\beta \oplus t \in D_\beta$ for some t with length bigger than n_β . \square

Definition 3.10. *Let \mathbb{P}_λ be a forcing iteration of length λ , where λ is either a strong limit cardinal or is equal to ∞ , the class of all ordinals. We say that \mathbb{P}_λ is*

- reverse Easton if nontrivial forcing is done only at infinite cardinal stages, direct limits are taken at all inaccessible cardinal limit stages, and inverse limits are taken at all other limit stages; moreover, \mathbb{P}_λ is the direct limit of the $\langle \mathbb{P}_\delta, \delta < \lambda \rangle$ if λ is regular or ∞ , the inverse limit of the $\langle \mathbb{P}_\delta, \delta < \lambda \rangle$, otherwise;
- directed closed if for all $\delta < \lambda$, \mathbb{Q}_δ is $< \delta$ -directed closed, i.e., for any $D \subseteq \mathbb{Q}_\delta$, $|D| < \delta$ such that for any $d_1, d_2 \in D$ there is an $e \in D$ with $e \leq d_1, e \leq d_2$, there exists $p \in \mathbb{Q}_\delta$ such that $p \leq d$ for any $d \in D$;
- λ -bounded if for all $\delta < \lambda$, \mathbb{Q}_δ has size $< \lambda$. Note that in the case $\lambda = \infty$, this just means that each \mathbb{Q}_δ is a set-forcing;

Moreover, if j is any elementary embedding such that $j''\lambda \subset \lambda$ and $\mathbb{P}_\lambda \subset \text{dom}(j)$, we say that \mathbb{P}_λ is j -coherent if for any $\delta < \lambda$, $j(\mathbb{P}_\delta) = \mathbb{P}_{j(\delta)}$.

The following theorem summarizes the general procedure:

Theorem 3.11. *Let $j : L(V_{\lambda+1}) \prec L(V_{\lambda+1})$ with $\text{crt}(j) = \kappa < \lambda$. Let \mathbb{P} be a directed closed, λ -bounded, j -coherent reverse Easton iteration. Let \mathbb{Q} be a κ -geometric forcing in $(V_\lambda)^{V^\mathbb{P}}$ that adds a Prikry sequence to κ . Then there exist G generic for \mathbb{P} and H $V[G]$ -generic for \mathbb{Q} such that $V[G][H] \models \exists k : V_{\kappa+1} \prec V_{\kappa+1}$.*

The relevant point of the proof is the forcing \mathbb{Q} , as by Lemma 3.6 and Lemma 3.7 in [4] combined, for any G generic for \mathbb{P} , $V[G] \models \exists j : L(V_{\lambda+1}) \prec L(V_{\lambda+1})$, $\text{crt}(j) = \kappa$. So, for better readability, from now on we call the generic extension of \mathbb{P} just V .

Let $j_{0,\omega} : L(V_{\lambda+1}) \prec M_\omega$ the ω -th iterate of j . Then $j_{0,\omega}(\mathbb{Q})$ is a λ -geometric forcing that adds a Prikry sequence to λ in M_ω .

Lemma 3.12. *In V there are only λ open dense sets of $j_{0,\omega}(\mathbb{Q})$.*

Proof. As $\mathbb{Q} \in V_\lambda$, there exists $n \in \omega$ such that $\mathbb{Q} \in V_{\kappa_n}$. In particular $j_{0,\omega}(\mathbb{Q}) \in M_\omega \cap V_{j_{0,\omega}(\kappa_n)}$ and its dense sets are in $j_{0,\omega}(\mathbb{Q}) \in M_\omega \cap V_{j_{0,\omega}(\kappa_{n+1})}$. But $|M_\omega \cap V_{j_{0,\omega}(\lambda)}| = \lambda$. \square

Proposition 3.13. *There exists a generic ultrafilter $G \in V$ of $j_{0,\omega}(\mathbb{Q})$.*

Proof. Let $\langle D_\alpha : \alpha < \lambda \rangle$ be an enumeration of the dense sets of $j_{0,\omega}(\mathbb{Q})$ in V . For every $n \in \omega$, fix q_n that witnesses λ -geometricity for $\langle D_\alpha : \alpha < \kappa_n \rangle$. Then for every m there exists a $q'_{n,m} < q_n$ such that $q'_{n,m} \in E_m$. Let G be the filter $\bigcup_{n,m \in \omega} \mathcal{F}_{q'_{n,m}}$, with \mathcal{F}_q the filter generated by q . Then G is generic. \square

The following appeared in [18].

Theorem 3.14 (Generic Absoluteness). *Let $j : L(V_{\lambda+1}) \prec L(V_{\lambda+1})$ with $\text{crt}(j) < \lambda$ be a proper elementary embedding. Let (M_ω, j_ω) be the ω -th iterate of j and let $\langle \eta_i : i < \omega \rangle \in V$ be a sequence generic for the Prikry forcing on λ in M_ω .*

Then for all $\alpha < \lambda$ there exists an elementary embedding

$$\pi : L_\alpha(M_\omega[\langle \eta_n : n \in \omega \rangle] \cap V_{\lambda+1}) \prec L_\alpha(V_{\lambda+1})$$

such that $\pi \upharpoonright \lambda$ is the identity.

In particular $M_\omega[\langle \eta_n : n \in \omega \rangle] \models I1(\lambda)$.

Proof of Theorem 3.11. Let $H \in V$ be $j_{0,\omega}(\mathbb{P})$ -generic. Fix g , one of the Prikry sequences added by $j_{0,\omega}(\mathbb{Q})$. Then $M_\omega \subseteq M_\omega[g] \subseteq M_\omega[H] \subseteq V$. But

$M_\omega[g]$ satisfies Generic absoluteness' conditions, therefore $M_\omega[g] \models I1(\lambda)$. But also $V \models I1(\lambda)$, therefore it must be $M_\omega[H] \models I1(\lambda)$.

We just proved that

$$M_\omega \models \exists p \in j_{0,\omega}(\mathbb{Q}) \ p \Vdash_{j_{0,\omega}(\mathbb{Q})} \exists i : (V_{\lambda+1}) \prec (V_{\tilde{\lambda}+1}),$$

Applying j^{-1} , we have that

$$L(V_{\lambda+1}) \models \exists p \in \mathbb{Q} \ p \Vdash_{\mathbb{Q}} \exists i : (V_{\tilde{\kappa}+1}) \prec (V_{\tilde{\kappa}+1}),$$

as we wanted to prove. □

4 Extender-based Prikry forcing

The first application of κ -geometricity will be on the extender-based Prikry forcing. It was introduced by Gitik and Magidor, and the reader can find an exhaustive description in [8]. The aim of the forcing is to add many Prikry sequences to a strong enough cardinal, blowing up its power while not changing the power function below it. This is more difficult than just having λ singular and $2^\lambda > \lambda^+$: the proof for this is to take λ measurable, forcing $2^\lambda > \lambda^+$ and then adding a Prikry sequence to λ . But Dana Scott [14] proved that if λ is measurable and $2^\lambda > \lambda^+$, then for a measure one set below λ , $2^\kappa > \kappa^+$, therefore this method would not give the first failure of GCH on λ . The solution is to exploit the extender structure of the cardinal to add many Prikry sequences, at the same time blowing up the power and changing the cofinality.

Definition 4.1. *Let κ and γ be cardinal. Then κ is γ -strong iff there is a $j : V \prec M$ such that $\text{crt}(j) = \kappa$, $\gamma < j(\kappa)$ and $V_{\kappa+\gamma} \subseteq M$.*

We write the definition as it is in [8].

Suppose GCH, and let λ be a 2-strong cardinal.

For any $\alpha < \lambda^{++}$, define a λ -complete normal ultrafilter on λ as $X \in U_\alpha$ iff $\alpha \in j(X)$. For any $\alpha, \beta < \lambda^{++}$, define $\alpha \leq_E \beta$ iff $\alpha \leq \beta$ and for some $f \in {}^\lambda \lambda$, $j(f)(\beta) = \alpha$. Then by a result in [8], $\langle \lambda^{++}, \leq \rangle$ is a λ^{++} -directed order, and there exists $\langle \pi_{\alpha\beta} : \alpha, \beta \in \lambda^{++}, \alpha \leq_E \beta \rangle$ such that $\langle \lambda^{++}, \langle U_\alpha : \alpha < \lambda^{++} \rangle, \leq_E \rangle$ is a nice system. There is no need to define a nice system here, the term is introduced only because the extender-based Prikry forcing is built on a nice system, the full definition can be found in [8].

Fix a nice system $\langle \lambda^{++}, \langle U_\alpha : \alpha < \lambda^{++} \rangle, \leq_E \rangle$. For any $\nu < \lambda$ and $\lambda < \alpha < \lambda^{++}$, let us denote $\pi_{\alpha,0}(\nu)$ by $\nu^{\alpha,0}$. We will write just ν^0 when α is obvious. By a $^\circ$ -increasing sequence of ordinals we mean a sequence $\langle \nu_0, \dots, \nu_n \rangle$ of

ordinals below λ such that $\nu_0^0 < \dots < \nu_n^0$. We say that μ is permitted for $\langle \nu_0, \dots, \nu_n \rangle$ iff $\mu^0 > \nu_i^0$ for all $i = 0 \dots n$.

Also, choose the system so that if $A \in U_\alpha$, $\mu_0, \mu_1 \in A$ and $\mu_0^0 < \mu_1^0$, then $|\{\mu \in A : \mu^0 = \mu_0^0\}| < \mu_1^0$.

Definition 4.2. *The set of forcing conditions \mathbb{P} consists of all the elements p of the form*

$$\{\langle \gamma, p^\gamma \rangle \mid \gamma \in g \setminus \{\max(g)\}\} \cup \{\langle \max(g), p^{\max(g)}, T \rangle\},$$

where

1. $g \subseteq \lambda^{++}$ of cardinality $\leq \lambda$ which has a maximal element according to \leq_E and $0 \in g$.
2. for $\gamma \in g$, p^γ is a finite $^\circ$ -increasing sequence of ordinals $< \lambda$.
3. T is a tree, with a trunk $p^{\max(g)}$, consisting of $^\circ$ -increasing sequences. All the splittings in T are required to be on sets in $U_{\max(g)}$, i.e., for every $\eta \in T$, if $\eta \geq p^{\max(g)}$ then the set

$$\text{Suc}_T(\eta) = \{\mu < \lambda : \eta^\frown \langle \mu \rangle \in T\} \in U_{\max(g)}.$$

Also require that for $\eta_1 \geq_T \eta_2 \geq_T p^{mc}$, $\text{Suc}_T(\eta_1) \subseteq \text{Suc}_T(\eta_2)$.

4. For every $\mu \in \text{Suc}_T(p^{\max(g)})$, $|\{\gamma \in g : \mu \text{ is permitted for } p^\gamma\}| \leq \mu^0$.
5. For every $\gamma \in g$, $\pi_{\max(g), \gamma}(\max(p^{\max(g)}))$ is not permitted for p^γ .
6. $\pi_{\max(g), 0}$ projects $p^{\max(g)}$ onto p^0 (so $p^{\max(g)}$ and p^0 are of the same length).

Let us denote g by $\text{supp}(p)$, $\max(g)$ by $\text{mc}(p)$, T by T^p , $p^{\max(g)}$ by p^{mc} and $\text{bas}(p) = p \upharpoonright (\text{supp}(p) \setminus \text{mc}(p))$.

Definition 4.3. *Let $p, q \in \mathbb{P}$. We say that p extends q and denote this by $p < q$ iff*

1. $\text{supp}(p) \supseteq \text{supp}(q)$.
2. For every $\gamma \in \text{supp}(q)$, p^γ is an end-extension of q^γ .
3. $p^{\text{mc}(q)} \in T^q$.

4. For every $\gamma \in \text{supp}(q)$,

$$p^\gamma \setminus q^\gamma = \pi_{\text{mc}(q), \gamma}[(p^{\text{mc}(q)} \setminus q^{\text{mc}(q)}) \upharpoonright (\text{lh}(p^{\text{mc}}) \setminus (i+1))],$$

where $i \in \text{dom}(p^{\text{mc}(q)})$ is the largest such that $p^{\text{mc}(q)}(i)$ is not permitted for q^γ .

5. $\pi_{\text{mc}(p), \text{mc}(q)}$ projects $T_{p^{\text{mc}}}^p$ into $T_{q^{\text{mc}}}^q$.

6. For every $\gamma \in \text{supp}(q)$ and $\mu \in \text{Suc}_{T^p}(p^{\text{mc}})$, if μ is permitted for p^γ , then $\pi_{\text{mc}(p), \gamma}(\mu) = \pi_{\text{mc}(q), \gamma}(\pi_{\text{mc}(p), \text{mc}(q)}(\mu))$.

Definition 4.4. Let $p, q \in \mathbb{P}$. We say that p is a direct extension of q and denot this by $p <^* q$ iff

1. $p < q$
2. for every $\gamma \in \text{supp}(q)$, $p^\gamma = q^\gamma$.

Definition 4.5. Let $p \in \mathbb{P}$ and $t \in T_{p^{\text{mc}}}^p$. Then $p \oplus t$ is defined as follows:

1. $\text{supp}(p \oplus t) = \text{supp}(p)$;
2. $(p \oplus t)^{\text{mc}} = p^{\text{mc}} \hat{\wedge} t$;
3. $T^{p \oplus t} = \{s \in T^p : s \subseteq (p \oplus t)^{\text{mc}} \vee (p \oplus t)^{\text{mc}} \subseteq s\}$;
4. if $\gamma \in \text{supp}(p)$,

$$(p \oplus t)^\gamma = p^\gamma \hat{\wedge} \pi_{\text{mc}(p), \gamma}[t \upharpoonright (\text{lh}(t) \setminus (i_\gamma + 1))],$$

where i_γ is the largest such that $t(i)$ is not permitted by p^γ .

If $s = \text{bas}(p)$ for some $p \in \mathbb{P}$, $\alpha \geq_E \gamma$ for all $\gamma \in \text{supp}(p) \setminus \text{mc}(p)$ and t is a finite $^\circ$ -increasing sequence of ordinals $< \lambda$, then $s \oplus (\alpha, t)$ is defined as follows:

1. $\text{supp}(s \oplus t) = \text{supp}(s)$;
2. if $\gamma \in \text{supp}(s)$,

$$(s \oplus t)(\gamma) = p^\gamma \hat{\wedge} \pi_{\alpha, \gamma}[t \upharpoonright (\text{lh}(t) \setminus (i_\gamma + 1))],$$

where i_γ is the largest such that $t(i)$ is not permitted by $s(\gamma)$.

Note that the previous definition is independent from p , and $\text{bas}(p \oplus t) = \text{bas}(p) \oplus (\text{mc}(p), p^{\text{mc}} \hat{\wedge} t)$.

A condition in \mathbb{P} is therefore a set of finite sequences and T indicating the possible extensions not only of the last one, but, via projection, of all of them. Morally speaking, $p \oplus t$ is the largest extension of p that we can have choosing t (and its projections) as extension.

Theorem 4.6 (Gitik, Magidor). *Let \mathbb{P} as above. Then*

$$V^{\mathbb{P}} \models 2^\lambda = \lambda^{++} \quad \wedge \quad \forall \kappa < \lambda \quad 2^\kappa = \kappa^+$$

Proposition 4.7. *Let \mathbb{P} as above. Then \mathbb{P} is λ -geometric.*

Proof. The proof goes through the same three claims as the proof for Lemma 3.6. Suppose that D is a dense open subset of \mathbb{P} and $p \in \mathbb{P}$. Without loss of generality, we can assume $p = \{\langle 0, \langle \rangle, TC \rangle\}$, where TC is the complete tree of the increasing finite sequences in λ . Note that in this case any $q \in \mathbb{P}$ with $q^{\text{mc}} = \langle \rangle$ is a direct extension of p . Fix an elementary submodel N of $H(\nu)$ with ν sufficiently large to contain all the relevant information of cardinality λ^+ and closed under λ -sequences of its elements. Pick $\alpha < \lambda^{++}$ above all the elements of $N \cap \lambda^{++}$.

Let T be a tree such that $\{\langle 0, \langle \rangle\} \cup \{\langle \alpha, \langle \rangle, T \rangle\}$ is in \mathbb{P} .

Claim 4.8 (First claim). *There exists $r \cup \{\langle \alpha, \langle \rangle, S \rangle\}$, with $S \subseteq T$, such that for every $t \in S$, if for some $q, R \in N$, $q \cup \{\langle \alpha, t, R \rangle\} \leq^* (r \cup \{\langle \alpha, \langle \rangle, S \rangle\}) \oplus t$ and $q \cup \{\langle \alpha, t, R \rangle\} \in D$, then $(r \cup \{\langle \alpha, \langle \rangle, S \rangle\}) \oplus t \in D$;*

Proof. If there is a $r \in N$ and a $T' \subseteq T$ such that $r \cup \{\langle \alpha, \langle \rangle, T' \rangle\} \in D$, then we put $q = r \cup \{\langle \alpha, \langle \rangle, T' \rangle\}$, $k = 0$ and we're done.

Let $A = \text{Suc}_T(\langle \rangle)$. We shall define by recursion the sequences $\langle r_\mu : \mu \in A \rangle$ and $\langle T^\mu : \mu \in A \rangle$, the first one increasing.

Let $\mu = \min(A)$. If there are an $s \in N$ and a $T' \subseteq T$ with trunk $\langle \mu \rangle$ such that $s \cup \{\langle \alpha, \langle \mu \rangle, T' \rangle\} \in D$, then set $r_\mu = s$ and $T^\mu = T'$. Otherwise do nothing, i.e., $r_\mu = \{\langle 0, \langle \rangle\}$ and $T^\mu = T$.

Suppose now that r_ξ and T^ξ are defined for any $\xi < \mu$ in A . Let $r''_\mu = \bigcup_{\xi \in \mu \cap A} r_\xi$ and consider $r'_\mu = r''_\mu \oplus (\alpha, \langle \mu \rangle)$. There are two cases:

1. If there are an $s \in N$ and a $T' \subseteq T$ such that

$$D \ni s \cup \{\langle \alpha, \langle \mu \rangle, T' \rangle\} <^* r'_\mu \cup \{\langle \alpha, \langle \mu \rangle, T \rangle\},$$

then set $r_\mu = r''_\mu \cup ((s \oplus (\alpha, \langle \mu \rangle)) \setminus r'_\mu)$ and $T^\mu = T'$.

2. Otherwise do nothing, i.e., $r_\mu = r''_\mu$ and $T^\mu = T$.

Subclaim 4.9. *For any $\gamma \in \text{supp}(r) \setminus \text{supp}(r''_\mu)$, μ is not permitted for $(r_\mu)^\gamma$.*

Proof of Subclaim. By definition, as if γ is not in $\text{supp}(r''_\mu)$, it must be in $\text{supp}(s \oplus (\alpha, \langle \mu \rangle))$, and μ is not permitted for $s \oplus (\alpha, \langle \mu \rangle)(\gamma)$. \square

Let $s_1 = \bigcup_{\mu \in A} r_\mu$. We need to trim T to some S^1 so that $s_1 \cup \{\langle \alpha, \langle \rangle, S^1 \rangle\}$ is an element of \mathbb{P} .

For $i < \lambda$ let

$$C_i = \begin{cases} A & \text{if there is no } \mu \in A \text{ such that } \mu^0 = i; \\ \bigcap_{\mu \in A, \mu^0 = i} \text{Suc}_{T^\mu}(\langle \mu \rangle) & \text{otherwise.} \end{cases}$$

Note that $A \in U_\alpha$, and therefore by our choice of the nice system we have that for any $i \in \lambda$, if there is a $\mu_1 \in A$ such that $\mu_1^0 = i$, for any $\mu_1 < \mu_2 \in A$, $|\{\mu \in A : \mu^0 = i\}| < \mu_2$, so by λ -completeness $C_i \in U_\alpha$. Set $A^* = A \cap \Delta_{i < \lambda}^* C_i$. Then for every $\delta \in A^*$ and for every $\mu \in A$ if $\delta^0 < \mu^0$ then $\mu \in \text{Suc}_{T^\mu}(\langle \delta \rangle)$. S^1 will be the tree obtained from T by eliminating all the branches that do not start with $\mu \in A^*$, replacing $T_{\langle \mu \rangle}$ with $T_{\langle \mu \rangle}^\mu$ and intersecting all the levels with A^* , i.e., $\langle \delta_0, \dots, \delta_n \rangle \in S^1$ iff $\langle \delta_0, \dots, \delta_n \rangle \in T^{\delta_0}$ and $\forall i \leq n, \delta_i \in A^*$.

Subclaim 4.10. $s_1 \cup \{\alpha, \langle \rangle, S^1\} \in \mathbb{P}$.

Proof of Subclaim. The only non-trivial point is to show condition (4) of the definition of \mathbb{P} , i.e., that for any $\delta \in \text{Suc}_{S^1}(\langle \rangle) = A^*$,

$$|\{\gamma \in \text{supp}(s_1) : \delta \text{ is permitted for } r^\gamma\}| \leq \delta^0.$$

Let

$$B_\delta = \{\gamma \in \text{supp}(s_1) : \delta \text{ is permitted for } r^\gamma\}.$$

Since $\text{supp}(s_1) = \bigcup_{\mu \in A} \text{supp}(r_\mu)$, we can divide B_δ in

$$B_{\delta, \mu} = \{\gamma \in \text{supp}(r_\mu) : \delta \text{ is permitted for } r^\gamma\}.$$

We can also suppose that μ is such that $r_\mu \neq r''_\mu$, i.e., μ is a stage that follows step (1). By Subclaim 4.9 if $\gamma \in \text{supp}(r_\mu) \setminus \bigcup_{\xi \in A} \text{supp}(r_\xi)$, then μ is not permitted for p^γ , so we can restrict the division to $B_\delta = \bigcup_{\mu \in A, \mu^0 < \delta^0} B_{\delta, \mu}$. Again, by our choice of the nice system, if $\mu^0 < \delta^0$ then there are less than δ^0 other elements $\xi \in A$ such that $\xi^0 = \mu^0$, therefore the former is a union of $\leq \delta^0$ elements.

Now fix a $B_{\mu, \delta}$. Since $\delta \in A^*$, by definition of A^* $\delta \in \text{Suc}_{T^\mu}(\langle \mu \rangle)$. Since $s \cup \{\langle \alpha, \langle \mu \rangle, T^\mu \rangle\} \in \mathbb{P}$, by point (4) of the definition of \mathbb{P} we have

$$|\{\gamma \in \text{supp}(s) : \delta \text{ is permitted for } s^\gamma\}| \leq \delta^0.$$

But $\text{supp}(s) = \text{supp}(r_\mu)$, and δ is permitted for s^γ iff δ is permitted for $(r_\mu)^\gamma$, as either $s^\gamma = (r_\mu)^\gamma$, or $s^\gamma = (r''_\mu \oplus \langle \mu \rangle)^\gamma$, $(r_\mu)^\gamma = (r''_\mu)^\gamma$ and μ is permitted for $(r''_\mu)^\gamma$, but in the second case δ is trivially permitted both for s^γ and $(r_\mu)^\gamma$. Therefore $|B_\delta| \leq \delta^0$. \square

Subclaim 4.11. For every $\delta \in \text{Suc}_{S^1}(\langle \rangle)$, if for some $q, R \in N$,

$$q \cup \{\alpha, \langle \delta \rangle, R\} \leq^* (r \cup \{\alpha, \langle \rangle, S^1\}) \oplus \langle \delta \rangle$$

and $q \cup \{\alpha, \langle \delta \rangle, R\} \in D$, then $(r \cup \{\alpha, \langle \rangle, S^1\}) \oplus \langle \delta \rangle \in D$.

Proof of Subclaim. Recall the construction of r at the δ -th stage. Since $r \oplus \langle \delta \rangle \upharpoonright \text{supp}(s) = s$, we have also that $q \cup \{\alpha, \langle \delta \rangle, R\} \leq^* s \cup \{\langle \alpha, \langle \rangle, T\}\}$ and $q \cup \{\alpha, \langle \delta \rangle, R\} \in D$, therefore the construction followed step (1). This implies that there exists a s such that $s \cup \{\langle \alpha, \langle \delta \rangle, T^\delta\}\} \in D$, with

$$(r_\delta)^\gamma = \begin{cases} (r_\xi)^\gamma & \text{if there exists } \xi \in \delta \cap A, \gamma \in \text{supp}(r_\xi); \\ s^\gamma & \text{otherwise.} \end{cases}$$

By the fact that $S_{\langle \delta \rangle}^1 \subseteq T_{\langle \delta \rangle}^\delta$ and by Subclaim 4.9, this implies that

$$r \cup \{\langle \alpha, \langle \rangle, S^1\}\} \oplus \langle \delta \rangle \leq^* s \cup \{\langle \alpha, \langle \delta \rangle, T^\delta\}\},$$

and by density we proved the claim. \square

Now we climb up the tree, by induction. Suppose that the first n levels are already defined.

We define r_t and T^t for any $t \in S^n$ of length $n+1$, by induction on the lexicographical order.

Let $r_t'' = s_n \cup \bigcup_{s < t} r_t$ and $r_t' = r_t'' \oplus (\alpha, t)$. There are two cases:

1. If there are an $s \in N$ and a $T' \subseteq S^n$ such that

$$D \ni s \cup \{\langle \alpha, t, T'\}\} <^* r_t' \cup \{\langle \alpha, \langle \rangle, S^n\}\} \oplus t,$$

then set $r_t = r_t'' \cup ((s \oplus (\alpha, t)) \setminus r_t')$ and $T^t = T'$.

2. Otherwise do nothing, i.e., $r_t = r_t''$ and $T^t = S^n$.

Let $s_{n+1} = \bigcup_{t \in \text{Lev}_n(S^n)} r_t$.

Subclaim 4.12. For any $\gamma \in \text{supp}(s_{n+1}) \setminus \text{supp}(r_\mu'')$, μ is not permitted for $(r_\mu)^\gamma$.

Proof. As before. \square

For $i < \lambda$ let

$$C_i = \begin{cases} \{t(n) : t \in \text{Lev}_n(S^n)\} & \text{if there is no } t \in \text{Lev}_n(S^n) \text{ such that } t(n)^0 = i; \\ \bigcap_{t \in \text{Lev}_n(S^n), t(n)^0 = i} \text{Suc}_{T^t}(t) & \text{otherwise.} \end{cases}$$

As before, $C_i \in U_\alpha$, we define $A^* = \text{Suc}_{S^n}(\langle \rangle) \cap \Delta_{i < \lambda}^* C_i$. Let $\langle \mu_0, \dots, \mu_i \rangle \in S^{n+1}$ iff $\langle \mu_0, \dots, \mu_i \rangle \in S^n$, $\forall l \leq i$, $\mu_l \in A^*$ and if $i \geq n$, $\langle \mu_0, \dots, \mu_i \rangle \in T^{\langle \mu_0, \dots, \mu_{n-1} \rangle}$.

Subclaim 4.13. $s_{n+1} \cup \{\alpha, \langle \rangle, S^{n+1}\} \in \mathbb{P}$.

Proof of Subclaim. The proof is similar to the previous one. In this case, we split B_δ in the union of

$$\{\gamma \in \text{supp}(s_n) : \delta \text{ is permitted for } (s_{n+1})^\gamma = (s_n)^\gamma\}$$

and

$$B_{t,\delta} = \{\gamma \in \text{supp}(r_t) : \delta \text{ is permitted for } (s_{n+1})^\gamma = (r_t)^\gamma\},$$

with $t(n)^0 < \delta^0$, thanks to Subclaim 4.12. By induction the first one has $\leq \delta_0$ elements, and the rest is as Subclaim 4.10, \square

Subclaim 4.14. For every $t \in S^{n+1}$, if for some $q, R \in N$,

$$q \cup \{\alpha, t, R\} \leq^* (s_{n+1} \cup \{\alpha, \langle \rangle, S^{n+1}\}) \oplus t$$

and $q \cup \{\alpha, t, R\} \in D$, then $(s_{n+1} \cup \{\alpha, \langle \rangle, S^{n+1}\}) \oplus t \in D$.

Proof of Subclaim. As before. \square

Finally, let $r = \bigcup_{n \in \omega} s_n$ and $S = \bigcap_{n \in \omega} S^n$. It is in \mathbb{P} and satisfies the first claim. \square

Claim 4.15 (Second claim). *There exists $r \cup \{\langle \alpha, \langle \rangle, S^*\}\}$, with $S^* \subseteq S$, if $t_1, t_2 \in S$ are of the same length, then $(r \cup \{\langle \alpha, \langle \rangle, S^*\}) \oplus t_1 \in D$ iff $(r \cup \{\langle \alpha, \langle \rangle, S^*\}) \oplus t_2 \in D$.*

Proof. The proof follows closely the proof of the second claim in Lemma 3.6, but it needs more care because now we require for $\eta_1 \geq_T \eta_2 \geq_T p^{mc}$, $\text{Suc}_T(\eta_1) \subseteq \text{Suc}_T(\eta_2)$.

Let

$$R = \{t \in S : r \cup \{\langle \alpha, \langle \rangle, S\} \oplus t \in D\}.$$

Therefore we have to find $S^* \subseteq S$ such that for any $t_1, t_2 \in S^*$ of the same length, $t_1 \in R$ iff $t_2 \in R$.

Let $B_\emptyset^0 = \{\delta \in \text{Suc}_S(\langle \rangle) : t \in R\}$. Then either B_\emptyset^0 or $\text{Suc}_S(\langle \rangle) \setminus B_\emptyset^0$ are in U_α . Call such A_\emptyset^0 . Then let $\langle \mu_0, \dots, \mu_l \rangle \in S^0$ iff $\forall i \mu_i \in A_\emptyset^0$. We are intersecting all the levels of S to A_\emptyset^0 so that for any $\eta_1 \leq_{S^0} \eta_2$, $\text{Suc}_{S^0}(\eta_2) \subseteq \text{Suc}_{S^0}(\eta_1)$, and we are going to this this repeatedly without further comment. Note that for all $s \in S^0$, $\text{Suc}_{S^0}(s) = \text{Suc}_S(s) \cap A_\emptyset^0$, and the sequences in S^0 of length 1 are either all in R or all outside.

By induction the construction continues level-by-level, each time starting with $S^{n+1,0} \subseteq S^n$, and then going down to S^{n+1} , a tree such that all the $n+1$ -branches are either all in R or all outside it. More technically, Suppose

S^n is defined. For all $t \in S^n$, $\text{lh}(t) = n + 1$, define $B_t^{n+1} = \{\delta \in \text{Suc}_{S^n}(t) : t \smallfrown \langle \delta \rangle \in R\}$. Then either B_t^{n+1} or $\text{Suc}_{S^n}(t) \setminus B_t^{n+1}$ is in U_α . Let A_t^{n+1} be it. Define $\langle \mu_0, \dots, \mu_l \rangle \in S^{n+1,0}$ iff $\langle \mu_0, \dots, \mu_l \rangle \in S^n$ and $\forall i > n \mu_i \in A_{\langle \mu_0, \dots, \mu_n \rangle}^{n+1}$. Note that for all $s \in S^{n+1,0}$, $\text{lh}(s) \geq n + 1$,

$$\text{Suc}_{S^{n+1,0}}(s) = \text{Suc}_{S^n}(s) \cap A_{\langle s(0), \dots, s(n) \rangle}^{n+1}.$$

Let $t \in \text{Lev}_m S^{n+1}$ and suppose that $S^{n+1,n-m}$, B_s^{n+1} and A_s^{n+1} are defined for all $s \in S^{n+1}$ with $\text{lh}(s) = m + 1$. Let

$$B_t^{n+1} = \{\delta \in \text{Suc}_{S^{n+1}}(t) : \text{Suc}_{S^{n+1,n-m}}(t \smallfrown \langle \delta \rangle) = B_{t \smallfrown \langle \delta \rangle}^{n+1}\}.$$

Then either B_t^{n+1} or $\text{Suc}_{S^{n+1}}(t) \setminus B_t^{n+1}$ is in U_α . Let A_t^{n+1} be it.

Suppose A_t^{n+1} is defined for all $t \in S^{n+1}$ of length m . Then $\langle \mu_0, \dots, \mu_l \rangle \in S^{n+1,n+1-m}$ iff $\forall i \geq m \mu_i \in A_{\langle \mu_0, \dots, \mu_m \rangle}^{n+1}$. Call $S^{n+1} = S^{n+1,n+1}$. Note that for all $s \in S^{n+1,n+1-m}$ of length bigger than m ,

$$\text{Suc}_{S^{n+1,n+1-m}}(s) = \text{Suc}_{S^{n+1,n-m}}(s) \cap A_{\langle s(0), \dots, s(m) \rangle}^{n+1}$$

and all the sequences in S^{n+1} of length $n + 1$ either are all in R or all outside it.

Now, let $S^* = \bigcap_{n \in \omega} S^n$. The last remark is sufficient to prove the claim. We prove that $r \cup \{\langle \alpha, \langle \rangle, S^* \rangle\} \in \mathbb{P}$. Since $S^* \subseteq S$, and we were careful to build it so that $\text{Suc}_{S^*}(t) \subseteq \text{Suc}_{S^*}(s)$ when $t \supseteq s$, it suffices to prove that for any $t \in S^*$, $\text{Suc}_{S^*}(t) \in U_\alpha$. So let $t \in S^*$, $\text{lh}(t) = m$. Then $\text{Suc}_S(t)$ will be modified in the construction of S only in the stages $S^{n,n-i}$ where $i < m$, therefore

$$\text{Suc}_{S^*}(t) = \text{Suc}_S(t) \cap \bigcap_{i \leq m, n \in \omega} A_{\langle t(0), \dots, t(i) \rangle}^n,$$

that is a countable intersection of elements of U_α , and therefore in U_α . \square

Claim 4.16 (Third claim). *Suppose that D is a dense open subset of \mathbb{P} and $p \in \mathbb{P}$, then there is a $q <^* p$ and $k \in \omega$ such that for any $t \in T_{q^{\text{mc}}}^q$ with $\text{lh}(t) = k$, $q \oplus t \in D$.*

Proof. In the proof of Lemma 3.6, the first and second claims were enough to prove that if D is a dense open subset of \mathbb{P} and $p \in \mathbb{P}$, then there is a $q <^* p$ and $k \in \omega$ such that for any $t \in T_{q^{\text{mc}}}^q$ with $\text{lh}(t) = k$, $q \oplus t \in D$. This needs more work.

For ease of notation, let us call the previous condition of \mathbb{P} , $r \cup \{\langle \alpha, \langle \rangle, S \rangle\}$. This is our q . We just need to prove that there are $s, R \in N$ such that

$$s \cup \{\langle \alpha, t, R \rangle\} \leq^* (r \cup \{\langle \alpha, \langle \rangle, S \rangle\}) \oplus t = q \oplus t$$

and $s \cup \{\langle \alpha, t, R \rangle\} \in D$, and then by the two properties the Lemma is proved.

Pick some $\beta \in N \cap \lambda$ which is \leq_E above every element of $\text{supp}(r)$. This is possible since $\text{supp}(r) \in N$. Shrink S to a tree S^* to insure that for every $\mu \in \text{Suc}_{S^*}(\langle \rangle)$ and $\gamma \in \text{supp}(r)$, if μ is permitted for r^γ , then $\pi_{\alpha, \gamma}(\mu) = \pi_{\beta, \gamma}(\pi_{\alpha, \beta}(\mu))$.

Subclaim 4.17. *The former is possible.*

Proof. For any $\mu \in \text{Suc}_S(\langle \rangle)$, let

$$B_\mu = \{\gamma \in \text{supp}(r) : \mu \text{ is permitted for } p^\gamma\}.$$

Then we have $|B_\mu| \leq \mu^0$. Let $\langle \xi_i : i < \lambda \rangle$ an enumeration of $\text{supp}(r)$ such that for any $\mu \in \text{Suc}_S(\langle \rangle)$, $B_\mu \subseteq \{\xi_i : i < \mu^0\}$. For any $i < \lambda$, let

$$C_i = \{\mu \in \text{Suc}_S(\langle \rangle) : \pi_{\alpha, \xi_i}(\mu) = \pi_{\beta, \xi_i}(\pi_{\alpha, \beta}(\mu))\}.$$

Let $A^* = \Delta_{i < \lambda}^* C_i$ and let S^* be the intersection of S with A^* . □

Let S^{**} be the projection of S^* to β via $\pi_{\alpha, \beta}$. Let $r^* = r \cup \{\langle \beta, \langle \rangle, S^{**} \rangle\}$. Then $r^* \in N$, and since N is an elementary submodel there exists $s \in N$, $s < r^*$ and $s \in D$. By definition of extension, $s(\beta) \in S^{**}$, therefore there exists a $t \in S^*$ such that $\pi_{\alpha, \beta}(t) = s(\beta)$. Note also that $\text{mc}(s) <_E \alpha$ by the choice of N . Let R be the tree with stem t , derived intersecting S_t^* with $(\pi_{\alpha, \text{mc}(s)}^{-1})''T^s$ and shrinking, if necessary, in order to insure the equality of projections $\pi_{\alpha, \gamma}$ and $\pi_{\text{mc}(s), \gamma} \circ \pi_{\alpha, \text{mc}(s)}$ for the relevant γ 's in $\text{supp}(s)$. Then $\text{bas}(s) \cup \{\langle \beta, s(\beta) \rangle\} \cup \{\langle \alpha, t, R \rangle\} < s$, therefore it is in D . But we also have

$$\text{bas}(s) \cup \{\langle \beta, s(\beta) \rangle\} \cup \{\langle \alpha, t, R \rangle\} \leq^* (r \cup \{\langle \alpha, \langle \rangle, S \rangle\}) \oplus t,$$

and this proves that there is a $q <^* p$ and $k \in \omega$ such that for any $t \in T_{q^{\text{mc}}}^q$ with $\text{lh}(t) = k$, $q \oplus t \in D$. □

Let $\langle D_\beta : \beta < \alpha \rangle$ a sequence of open dense sets, with $\alpha < \lambda$. Let $\langle q_\beta : \beta \leq \alpha \rangle$ be the sequence built with an iteration of the third claim, i.e., for any $\beta \leq \alpha$, $q_{\beta+1} \leq^* q_\beta$ is such that there exists an $n_{\beta+1}$ such that for any $t \in T^{q_\beta}$ of length $n_{\beta+1}$, $q_{\beta+1} \oplus t \in D_\beta$, and if $\beta \leq \alpha$ is limit, then by λ -closeness of \leq^* let q_β be such that $q_\beta \leq^* q_\gamma$ for all $\gamma < \beta$.

Subclaim 4.18. *For any $\beta < \alpha$, for any $t \in T^{q_\alpha}$, $q_\alpha \oplus t \leq^* q_\beta \oplus \pi_{\text{mc}(q_\alpha), \text{mc}(q_\beta)}'' t$.*

Proof. Note that $q_\alpha^{\text{mc}} = q_\beta^{\text{mc}} = \langle \rangle^1$. We prove it point by point.

¹The claim is in fact true in general, but in this case calculation is easier

- $q_\beta \oplus \pi''_{\text{mc}(q_\alpha), \text{mc}(q_\beta)} t$ is well defined, i.e., $\pi''_{\text{mc}(q_\alpha), \text{mc}(q_\beta)} t \in T^{q_\beta}$. This is true for Definition 4.3(5).
- $\text{supp}(q_\alpha \oplus t) \supseteq \text{supp}(q_\beta \oplus \pi''_{\text{mc}(q_\alpha), \text{mc}(q_\beta)} t)$. This is true because $\text{supp}(q_\alpha \oplus t) = \text{supp}(q_\alpha)$,

$$\text{supp}(q_\beta \oplus \pi''_{\text{mc}(q_\alpha), \text{mc}(q_\beta)} t) = \text{supp}(q_\beta)$$

and by Definition 4.3(1).

- for any $\gamma \in \text{supp}(q_\alpha)$, $(q_\alpha \oplus t)^\gamma = (q_\beta \oplus \pi''_{\text{mc}(q_\alpha), \text{mc}(q_\beta)} t)^\gamma$. By definition, and since $(q_\alpha)^\gamma = (q_\beta)^\gamma$, this is true if and only if

$$\pi_{\text{mc}(q_\alpha), \gamma}[t \upharpoonright (\text{lh}(t) \setminus (i_\gamma + 1))] = \pi_{\text{mc}(q_\beta), \gamma}[\pi''_{\text{mc}(q_\alpha), \text{mc}(q_\beta)} t \upharpoonright (\text{lh}(t) \setminus (j_\gamma + 1))],$$

where i_γ is the largest such that $t(i_\gamma)$ is not permitted by $(q_\alpha)^\gamma$ and j_γ is the largest such that $\pi''_{\text{mc}(q_\alpha), \text{mc}(q_\beta)} t(j_\gamma)$ is not permitted by $(q_\beta)^\gamma = (q_\alpha)^\gamma$. By Definition 4.2(6) $(q_\alpha)^0 = (q_\beta)^0 = \langle \rangle$, therefore by Definition 4.3(6)

$$\pi_{\text{mc}(q_\alpha), 0} \upharpoonright t = \pi_{\text{mc}(q_\beta), 0} \circ \pi_{\text{mc}(q_\alpha), \text{mc}(q_\beta)} \upharpoonright T,$$

so $i_\gamma = j_\gamma$. Therefore this point is true by Definition 4.3(6).

- $(q_\alpha \oplus t)^{\text{mc}(q_\beta \oplus \pi''_{\text{mc}(q_\alpha), \text{mc}(q_\beta)} t)} \in T^{q_\beta \oplus \pi''_{\text{mc}(q_\alpha), \text{mc}(q_\beta)} t}$. First of all, $\text{mc}(q_\beta \oplus \pi''_{\text{mc}(q_\alpha), \text{mc}(q_\beta)} t) = \text{mc}(q_\beta)$. By definition,

$$(q_\alpha \oplus t)^{\text{mc}(q_\beta)} = (q_\alpha)^{\text{mc}(q_\beta)} \wedge \pi_{\text{mc}(q_\alpha), \text{mc}(q_\beta)}[t \upharpoonright (\text{lh}(t) \setminus (i_{\text{mc}(q_\beta)} + 1))],$$

with $i_{\text{mc}(q_\beta)}$ as above. By Definition 4.3(2) $(q_\alpha)^{\text{mc}(q_\beta)} = q_\beta^{\text{mc}} = \langle \rangle$, and therefore $i_{\text{mc}(q_\beta)} = 0$, so the point follows simply by definition of $T^{q_\beta \oplus \pi''_{\text{mc}(q_\alpha), \text{mc}(q_\beta)} t}$.

- $\pi_{\text{mc}(q_\alpha), \text{mc}(q_\beta)}$ projects $T_t^{q_\alpha \oplus t}$ into $T_{\pi''_{\text{mc}(q_\alpha), \text{mc}(q_\beta)} t}^{q_\beta \oplus \pi''_{\text{mc}(q_\alpha), \text{mc}(q_\beta)} t}$. By definition, $T_t^{q_\alpha \oplus t} = T_t^{q_\alpha}$ and $T_{\pi''_{\text{mc}(q_\alpha), \text{mc}(q_\beta)} t}^{q_\beta \oplus \pi''_{\text{mc}(q_\alpha), \text{mc}(q_\beta)} t} = T_{\pi''_{\text{mc}(q_\alpha), \text{mc}(q_\beta)} t}^{q_\beta}$, so this is true by Definition 4.3(5).
- for any $\gamma \in \text{supp}(q_\alpha)$ and $\mu \in \text{Suc}_{T^{q_\alpha \oplus t}}(t)$, if μ is permitted for $(q_\alpha \oplus t)^\gamma$ then

$$\pi_{\text{mc}(q_\alpha), \gamma}(\mu) = \pi_{\text{mc}(q_\beta), \gamma}(\pi_{\text{mc}(q_\alpha), \text{mc}(q_\beta)}(\mu)).$$

As $T^{q_\alpha \oplus t}$ is a subtree of T^{q_α} , and μ is permitted for $(q_\alpha \oplus t)^\gamma$ means that it is also permitted for $(q_\alpha)^\gamma$, this is a direct consequence of Definition 4.3(6).

□

Therefore q_α is as wanted: let $q' \leq q_\alpha$ such that $q' \in E_{n_\beta}$. Then $q' < q_\alpha \oplus t$ for some $t \in T^{q_\alpha}$ with length n_β . But by the claim $q_\alpha \oplus t \leq^* q_\beta \oplus \pi''_{\text{mc}(q_\alpha), \text{mc}(q_\beta)} t$, and by the Lemma $q_\beta \oplus \pi''_{\text{mc}(q_\alpha), \text{mc}(q_\beta)} t \in D_\beta$, so also $q' \in D_\beta$. □

Corollary 4.19. *Suppose $I0(\kappa, \lambda)$. Then there exists a generic extension in which $I1(\kappa) + 2^\kappa = \kappa^{++} + \forall \eta < \kappa \ 2^\eta = \eta^+$.*

Proof. We want to apply Theorem 3.11. so first note that the forcing that forces GCH below λ is a directed closed, λ -bounded, j -coherent reverse Easton iteration, while the forcing that forces GCH above λ is λ -closed, and therefore does not touch $I0$. $I0$ clearly implies $I2$, and it is a well-known fact (see e.g. Proposition 24.2 in [10]) that this is equivalent to the existence of $k : V \prec M$ with $V_\lambda \subseteq M$. We can construct k so that $k \restriction V_\lambda = j \restriction V_\lambda$, so κ is 2-strong. Therefore we can apply the extender-based Prikry forcing to κ . The elements of \mathbb{P} on κ are κ -sequences of triples of elements of κ^{++} , finite sequences in κ and functions from κ^ω to $\mathcal{P}(\kappa)$, so we can say that the forcing is in V_{κ_1} . The forcing adds a Prikry sequence to κ and it is κ -geometric, therefore the conditions of Theorem 3.11 are met. □

5 Diagonal Supercompact Prikry forcings

There are many versions of the diagonal supercompact Prikry forcing, we are going to use the one in [9] (and later the one in [13]). First, there is a preparation forcing that forces $2^\alpha = \alpha^{+\omega+2}$ for all α inaccessible. Then, the diagonal supercompact forcing exploits the fine normal ultrafilters that come from enough supercompactness of a cardinal to add Prikry sequences to it, while inducing an interesting pcf structure.

Definition 5.1. *Let κ, γ be cardinals. We say that κ is γ -supercompact iff there exists a fine normal measure on $\mathcal{P}_\kappa(\gamma)$, i.e., a measure U such that for any $f : \mathcal{P}_\kappa(\gamma) \rightarrow \gamma$ such that $f(x) \in x$ for almost every x , f is constant on a set in U .*

One interesting combinatorial principle is \square_κ , that states the existence of a coherent collection of clubs. While many combinatorial principles are consistent with the existence of large cardinals, \square_κ fails above large enough cardinals (Solovay). It is of interest, therefore, investigating weakenings of such principle.

Definition 5.2. *We say that a cardinal κ has the approachability property, AP_λ , iff there exists a sequence $\langle C_\alpha : \alpha < \kappa^+ \rangle$ such that*

- for α limit, C_α is a club in α and $\text{ot}(C_\alpha) = \text{cof}(\alpha)$;
- there is a club $D \subseteq \kappa^+$ such that for any $\alpha \in D$, for any $\beta < \alpha$ there exists $\gamma < \alpha$ such that $C_\alpha \cap \beta = C_\gamma$.

It is not difficult to see that it is a weakening of \square_κ .

Another field of research in infinite combinatorics is pcf theory: given a cardinal κ and $\langle \mu_n : n \in \omega \rangle$ cofinal, it investigates the structure of the functions in $\prod_{n \in \omega} \mu_n$, and it is a standard tool for the analysis of the combinatorics of a cardinal of cofinality ω .

Definition 5.3. Let $\langle \mu_n : n \in \omega \rangle$ be a sequence cofinal in κ . A sequence $\langle f_\alpha : \alpha < \kappa^+ \rangle \subseteq \prod_{n \in \omega} \mu_n$ is a very good scale iff

- $\langle f_\alpha : \alpha < \kappa^+ \rangle$ is a scale, i.e. such that for every $\alpha < \beta < \kappa^+$, $f_\alpha(m) < f_\beta(m)$ for almost every m and for every $f \in \prod_{n \in \omega} \mu_n$ there exists $\beta < \kappa^+$ and $n \in \omega$ with $f(m) < f_\beta(m)$ for every $m > n$.
- for every $\beta < \kappa^+$ such that $\omega < \text{cof}(\beta)$ there exists a club C of β and $n < \omega$ such that $f_{\gamma_1}(m) < f_{\gamma_2}(m)$ for every $\gamma_1 < \gamma_2 \in C$ and $m > n$.

If κ is as above, we say that there exists a very good scale in κ , VGS_κ .

Both these properties don't hold above a supercompact cardinal, and in [9] it is proven that having a very good scale does not imply the approachability property. We will prove that this holds also under rank-into-rank hypotheses.

Let \mathbb{E} be the reverse Easton forcing of length λ that force $2^\alpha = \alpha^{+\omega+2}$ for all α inaccessible. This forcing is:

- directed closed: \mathbb{Q}_α , the forcing that adds $\alpha^{+\omega+2}$ subsets of α , is $< \alpha$ -directed closed;
- λ -bounded: as λ is strong limit $|\mathbb{Q}_\alpha| < \lambda$.
- j -coherent: as $j(\mathbb{Q}_\alpha)$ is the poset consisting of the functions whose domain is a subset of $j(\alpha)^{+\omega+2}$ of size less than $j(\alpha)$, and whose range is a subset of the partial functions between $j(\alpha)$ and $j(\alpha)$, that is, $j(\mathbb{Q}_\alpha) = \mathbb{Q}_{j(\alpha)}$.

Let κ be $\kappa^{+\omega+2}$ -supercompact, with U_ω witnessing it and let U_n be the projection of U_ω on $\mathcal{P}_\kappa(\kappa^{+n})$, i.e.,

$$X \in U_n \text{ iff } \{P \in \mathcal{P}_\kappa(\kappa^{+\omega+2}) : P \cap \kappa^{+n} \in X\} \in U_\omega.$$

Clearly U_n is a normal ultrafilter on $\mathcal{P}_\kappa(\kappa^{+n})$.

Let $a, b \in \mathcal{P}_\kappa(\kappa^{+n})$ and $b \cap \kappa \in \kappa$. Set

$$a \subseteq b \leftrightarrow a \subseteq b \wedge \text{ot}(a) < b \cap \kappa.$$

Definition 5.4. $p \in \mathbb{Q}$ iff $p = \langle a_0^p, a_1^p, \dots, a_{n-1}^p, X_n^p, X_{n+1}^p, \dots \rangle$ where

1. $\forall l < n$ $a_l^p \in \mathcal{P}_\kappa(\kappa^{+n})$ and $a_l^p \cap \kappa$ is an inaccessible cardinal;
2. $\forall m \geq n$, $X_m^p \in U_m$;
3. $\forall m \geq n$ $\forall b \in X_m^p$ $\forall l < n$ $a_l^p \subseteq_\sim b$;
4. $\forall i < j < l$ $a_i^p \subseteq_\sim a_j^p$.

For $p = \langle a_0^p, a_1^p, \dots, a_{n-1}^p, X_n^p, X_{n+1}^p, \dots \rangle$, let us denote n as $l(p)$. Moreover, for any collection of $A_i \subseteq \mathcal{P}_\kappa(\kappa^{+i})$, let

$$\tilde{\Pi}_{n \in m} A_n = \{ \langle a_0, \dots, a_{m-1} \rangle : \forall i < j < m \ a_i \in A_i \wedge a_i \subseteq_\sim a_j \}.$$

For any collection of A_a , $a \in \mathcal{P}_\kappa(\kappa^{+n})$, let

$$\Delta A_a = \{ b \in \mathcal{P}_\kappa(\kappa^{+n}) : \forall a \in \mathcal{P}_\kappa(\kappa^{+n}) \ a \subseteq_\sim b \rightarrow b \in A_a \}.$$

It is a standard result that if each $A_a \in U_n$, then $\Delta A_a \in U_n$.

Definition 5.5. Let $p, q \in \mathbb{Q}$. Then $p \leq^* q$ iff

1. $l(p) = l(q)$;
2. $\forall l < l(p)$ $a_l^p = a_l^q$;
3. $\forall m \geq l(p)$ $X_m^p \subseteq X_m^q$.

Definition 5.6. Let $p \in \mathbb{Q}$ and $\vec{a} \in \tilde{\Pi}_{l(p) \leq n \leq m} X_n^p$. Then we denote by $p \oplus \vec{a}$ the sequence $\langle a_0^p, \dots, a_{l(p)}^p, a(l(p)), \dots, a(m), Y_{m+1}^p, \dots \rangle$, where

$$Y_n = \{ b \in X_n^p : \forall l(p) \leq i \leq m \ a(i) \subseteq_\sim b \}.$$

Then $p \leq q$ iff there exists \vec{a} such that $p \leq^* q \oplus \vec{a}$.

Theorem 5.7 (Gitik, Sharon). Let G generic for \mathbb{P} and H generic for \mathbb{Q} as above. Then $V[H][G] \models 2^\kappa > \kappa^+ \wedge \neg AP_\kappa \wedge VGS_\kappa$.

Proposition 5.8. \mathbb{Q} as above is κ -geometric.

Proof.

Claim 5.9 (First Claim). Let $p \in \mathbb{Q}$. For any dense set D there exists $q \leq^* p$ such that for any $\vec{a} \in \bigcup_{n \in \omega} \tilde{\Pi}_{m \in n} \mathcal{P}_\kappa(\kappa^{+m})$ if there exists \vec{X} such that $\vec{X} \oplus \vec{a} \in D$, then $\vec{Y} \oplus \vec{a} \in D$.

Proof. We can suppose $p = 1_{\mathbb{P}}$. For any $\vec{a} = \langle a_0, \dots, a_{n-1} \rangle \in \tilde{\Pi}_{m \in n} \mathcal{P}_{\kappa}(\kappa^{+m})$, if there exists $\vec{X} = \langle X_n, X_{n+1} \dots \rangle$ such that $\vec{a} \smallfrown \vec{X} \in D$, then let $X_{\vec{a}}$ be such \vec{X} , otherwise let $X_{\vec{a}}$ be just $\langle \mathcal{P}_{\kappa}(\kappa^{+n}), \mathcal{P}_{\kappa}(\kappa^{+n+1}) \dots \rangle$. Code $\langle a_0, \dots, a_n \rangle$ as an element of $\mathcal{P}_{\kappa}(\kappa^{+n})$ and define $Y_m = \Delta X_{\vec{a}}(m)$. Then for any $\vec{a} = \langle a_0, \dots, a_n \rangle$ we have $\{b \in Y_m : a_n \subseteq b\} \subseteq X_{\vec{a}}(m)$. \square

Claim 5.10 (Second Claim). *There exist $Y'_m \subseteq Y_m$ such that for any n either all $\vec{Y}' \oplus \vec{a}$ with \vec{a} of length n are in D , or they all are not in D .*

Proof. It is possible to do this via diagonal intersection, but with some care: the first step is, of course, to intersect Y_m either with all the a_0 such that $\vec{Y} \oplus \langle a_0 \rangle$ is in D or its complement (the one in U_0), so that we still have an element of \mathbb{Q} . Then shrink Y_1 so that for any a_0 , either for all the a_1 , $\vec{Y} \oplus \langle a_0, a_1 \rangle \in D$ (case 1), or the opposite is true (case 2), then shrink again Y_0 so that either case 1 or case 2 is true for all the a_0 , and so on, This is the formal definition (with the assumption that $\langle a_0, \dots, a_n \rangle$ for $n = -1$ is the empty sequence):

- $Y_m^{(0)} = Y_m$;
- let $\vec{a} = \langle a_0, \dots, a_n \rangle \in \tilde{\Pi}_{m \in n} \mathcal{P}_{\kappa}(\kappa^{+m})$. Then $B_{\vec{a}}^{(n+2)} = \{a \in Y_{n+1} : \vec{a} \smallfrown \langle a \rangle \smallfrown \langle Y_{n+2}, Y_{n+3}, \dots \rangle \in D\}$;
- either $B_{\vec{a}}^{(n+2)}$ or $Y_{n+1} \setminus B_{\vec{a}}^{(n+2)}$ is in U_{n+1} . Let it be $A_{\vec{a}}^{(n+2)}$;
- $Y_0^{(m)} = A_{\emptyset}^{(m)}$;
- for $m > n + 2$, $B_{\vec{a}}^{(m)} = \{a \in Y_{n+1}^{(m-1)} : B_{\vec{a} \smallfrown \langle a \rangle}^{(m)} = A_{\vec{a} \smallfrown \langle a \rangle}^{(m)}\}$;
- either $B_{\vec{a}}^{(m)}$ or $Y_{n+1}^{(m-1)} \setminus B_{\vec{a}}^{(m)}$ is in U_{n+1} . Let it be $A_{\vec{a}}^{(m)}$;
- for $m > n > 0$, $Y_n^{(m)} = \Delta \{A_{\langle a_0, \dots, a_{n-1} \rangle}^{(m)} : a_i \in Y_i^{(m)}\}$;
- finally, $Y'_n = \bigcap_{m > n} Y_n^{(m)}$.

The last step can be carried on because the ultrafilters are complete. \square

Claim 5.11 (Third Claim). *For any dense set D there is a $q \leq^* p$ and an n such that for all $\vec{a} \in \tilde{\Pi}_{l(p) \leq i < l(p)+n} X_i^q$, $q \oplus \vec{a} \in D$.*

Proof. Let $q = \vec{Y}'$. By density there exists $r < p$ such that $r \in D$, and by the first paragraph we can assume that $r = q \oplus \vec{a}$ for some \vec{a} . By construction, if there exists \vec{a} such that $q \oplus \vec{a} \in D$, then for all \vec{b} of the same length $q \oplus \vec{b} \in D$. \square

Then it is exactly as in Lemma 3.6. \square

Corollary 5.12. *Suppose $I0(\kappa, \lambda)$. Then there exists a generic extension in which there exists $j : V_{\kappa+1} \prec V_{\kappa+1}$, $2^\kappa > \kappa^+$, there is a very good scale in κ but the approachability property does not hold in κ .*

Proof. Let H be \mathbb{E} -generic for V . By 3.11 (without the Prikry part) $I0(\kappa, \lambda)$ still holds in $V[H]$, say witnessed by k . Then

$$U_\omega = \{X \supseteq \mathcal{P}_\kappa(\kappa^{+\omega+2}) : k^n \restriction \kappa^{+\omega+2} \in k^n(X)\},$$

defined in $V[H]$, witnesses that $\kappa = \text{crt}(k)$ is $\kappa^{+\omega+2}$ -supercompact. Therefore we can force on κ with \mathbb{Q} . $\mathbb{Q} \subseteq \prod_{n \in \omega} \mathcal{P}_\kappa(\kappa^{+n})$, so $\mathbb{Q} \subseteq V_{\kappa_1}$. The hypotheses of Theorem 3.11 are then satisfied, and Theorem 5.7 proves the Corollary. \square

In [13] Neeman introduced a variation on Gitik-Sharon forcing, that has a more structured preparation forcing and needs more large cardinal power. The result will involve the Tree Property:

Definition 5.13. *Let κ be a cardinal. Then the tree property holds at κ , $TP(\kappa)$, if every tree of height κ and such that all levels have size $< \kappa$ has a cofinal branch.*

Suppose j witnesses $I0(\kappa, \lambda)$ and let $\langle \kappa_i : i \in \omega \rangle$ be the critical sequence of j .

Lemma 5.14. $V_\lambda \models \kappa_1$ is limit of supercompact cardinals.

Proof. It is by reflection of rank-into-rank embeddings: for any $\gamma < \kappa$, the sentence " $\exists k : V_\lambda \prec V_\lambda, j(\gamma) < \text{crt}(k) < j(\kappa)$ " is true, witnessed by $j \restriction V_\lambda$ (note that $j(\gamma) = \gamma$). Then, by elementarity, there exists $k : V_\lambda \prec V_\lambda$ with critical point between γ and κ . Such critical point is supercompact in V_λ , and choosing different γ 's we have that the cardinals supercompact in V_λ form an unbounded subset of κ . By elementarity, this is true also for κ_1 . \square

Let $\mu_0 = \kappa$ and μ_{i+1} the smallest cardinal supercompact in V_λ larger than μ_i , and let $\nu = \sup_{i \in \omega} \mu_i$. By the lemma above, $\nu < \kappa_1$.

Suppose GCH.

Proposition 5.15 (Shi [15]). *There is a generic extension of V such that $I0(\kappa, \lambda)$ holds and if μ is a cardinal supercompact in V_λ , μ is indestructible by μ -directed closed forcing.*

So, we can suppose that all the μ_i are closed under μ_i -directed closed forcing.

Let \mathbb{A}_λ the reverse Easton iteration that is not trivial only on V_λ -supercompact cardinals limits of V_λ -supercompact cardinals, and if η is such a cardinal, \mathbb{Q}_η is the forcing that adds $\nu(\eta)^{++}$ subsets to η , with conditions of size $< \eta$, where $\nu(\eta)$ is the sup of the ω V_λ -supercompact cardinals above η . As noted before, if $\eta < \lambda$ then $\nu(\eta) < \lambda$.

- \mathbb{A}_λ is directed closed, because each \mathbb{Q}_η is $< \eta$ -directed closed;
- \mathbb{A}_λ is λ -bounded, because for each η , $|\mathbb{Q}_\eta| = \nu(\eta)^\eta < \lambda$, and λ is strong limit;
- \mathbb{A}_λ is j -coherent, because its definition depends only on λ .

Let E be generic for \mathbb{A}_λ . Then in $V[E]$:

- by Theorem 3.11, $I0(\kappa, \lambda)$ holds;
- by indestructibility, κ is V_λ -supercompact, and since the forcing is trivial from $\kappa + 1$ to ν , and closed enough, $2^\kappa = \nu^{++}$.
- by Gitik-Sharon [9], there exists a ν^+ supercompactness measure on κ .

We say that π is a ν^+ supercompactness measure on κ if $\pi : V[E] \prec M$, $\text{crt}(\pi) = \kappa$ and $M \restriction \pi(\kappa) = \{\pi(f)(\kappa) : f : \mathcal{P}_\kappa(\nu^+) \rightarrow \kappa\}$.

Let U be the ν^+ supercompactness measure on κ , and U_n the μ_n supercompactness measure on κ induced by U , i.e., $X \in U_n$ iff $\pi''\mu_n \in \pi(X)$.

Now the definition of the forcing is the same as 5.4, with μ_n instead of κ^{+n} .

Let $a, b \in \mathcal{P}_\kappa(\mu_n)$ and $b \cap \kappa \in \kappa$. Set

$$a \subseteq_\sim b \leftrightarrow a \subseteq b \wedge \text{ot}(a) < b \cap \kappa.$$

Definition 5.16. $p \in \mathbb{Q}$ iff $p = \langle a_0^p, a_1^p, \dots, a_{n-1}^p, X_n^p, X_{n+1}^p, \dots \rangle$ where

1. $\forall l < n$ $a_l^p \in \mathcal{P}_\kappa(\mu_n)$ and $a_l^p \cap \kappa$ is an inaccessible cardinal;
2. $\forall m \geq n$, $X_m^p \in U_m$;
3. $\forall m \geq n$ $\forall b \in X_m^p$ $\forall l < n$ $a_l^p \subseteq_\sim b$;
4. $\forall i < j < l$ $a_i^p \subseteq_\sim a_j^p$.

For $p = \langle a_0^p, a_1^p, \dots, a_{n-1}^p, X_n^p, X_{n+1}^p, \dots \rangle$, let us denote n as $l(p)$. Moreover, for any collection of $A_i \subseteq \mathcal{P}_\kappa(\kappa^{+i})$, let

$$\tilde{\Pi}_{n \in m} A_n = \{ \langle a_0, \dots, a_{m-1} \rangle : \forall i < j < m \ a_i \in A_i \wedge a_i \subseteq_{\sim} a_j \}.$$

For any collection of A_a , $a \in \mathcal{P}_\kappa(\mu_n)$, let

$$\Delta A_a = \{ b \in \mathcal{P}_\kappa(\mu_n) : \forall a \in \mathcal{P}_\kappa(\mu_n) \ a \subseteq_{\sim} b \rightarrow b \in X_a \}.$$

It is a standard result that if each $A_a \in U_n$, then $\Delta A_a \in U_n$.

Definition 5.17. Let $p, q \in \mathbb{Q}$. Then $p \leq^* q$ iff

1. $l(p) = l(q)$;
2. $\forall l < l(p) \ a_l^p = a_l^q$;
3. $\forall m \geq l(p) \ X_m^p \subseteq X_m^q$.

Note that if G is generic for \mathbb{Q} as above as defined in $V[E]$, $(2^\kappa)^{V[E][G]} = (\nu^{++})$. As ν is collapsed to κ , and no other cardinal is collapsed, $V[E][G] \models 2^\kappa = \kappa^{++}$

Theorem 5.18 (Neeman). *If G is generic for \mathbb{Q} as above as defined in $V[E]$, then $TP(\kappa^+)$.*

Lemma 5.19. \mathbb{Q} as above is κ -geometric.

Proof. The proof is exactly the same as in 5.7 □

Corollary 5.20. *Suppose $I0(\kappa, \lambda)$. Then there exists a generic extension in which $I1(\kappa) + 2^\kappa > \kappa^+ + TP(\kappa^+)$ holds.*

Proof. The remarks above show that the hypotheses for Theorem 3.11 are satisfied: there are three preparation forcing (one for GCH, one for the indestructibility of supercompactness, and one for blowing up the power of κ) and they are all reverse Easton iterations with the properties needed. The forcing \mathbb{Q} is a subset of $\Pi_{n \in \omega} \mathcal{P}_\kappa(\mu_n)$, therefore in $V[E]_{\kappa_1}$ by Lemma 5.14, and it is κ -geometric. So in $V[E][H]$ $I1(\kappa)$ holds, but also (see above) $2^\kappa = \kappa^{++}$ and $TP(\kappa^+)$ □

6 Tree property at the double successor

Some results can be achieved using the general procedure without much further effort. This is the case for the forcing introduced by Dobrinen and Friedman in [5], to prove the tree property at a double successor of a singular cardinal.

Note that for $TP(\kappa^{++})$ to hold, it must be that $2^\kappa > \kappa^{++}$, so it is natural to ask whether $I1(\kappa)$ holds at the same time. This is another property that is implied by \square_κ .

Definition 6.1. *For any κ inaccessible, the forcing $\text{Sacks}(\kappa)$ is the set of subsets of $2^{<\kappa}$ such that:*

- $s \in p, t \subseteq s \rightarrow t \in p$;
- each $s \in p$ has a proper extension in p ;
- for any $\alpha < \kappa$, if $\langle s_\beta : \beta < \alpha \rangle$ is a \subseteq -increasing sequence of elements of p , then $\bigcup_{\beta < \alpha} s_\beta \in p$;
- there exists a club $C(p)$ such that $\{s \in p : s \restriction 0 \in p \wedge s \restriction 1 \in p\} = \{s \in p : \text{lh}(s) \in C(p)\}$.

Extension is simply the inclusion.

Definition 6.2. *For any κ inaccessible and $\gamma(\kappa)$ the first weakly compact above it, $\text{Sacks}^+(\kappa)$ is the $\gamma(\kappa)$ iteration of $\text{Sacks}(\kappa)$ with supports of size $\leq \kappa$.*

Let \mathbb{P} be the reverse Easton forcing of length λ such that $\mathbb{P}_\alpha = \text{Sacks}^+(\alpha)$. This forcing is clearly λ -bounded and j -coherent. Fact 2.7 in [5] states that it is also closed. Moreover, Theorem 3.2 in the same paper shows that in the extension the Tree Property holds in the extension for α^{++} , for any α inaccessible.

Corollary 6.3. *Suppose $I0(\kappa, \lambda)$. There exists a generic extension of V such that $I1(\kappa) + TP(\kappa^{++})$ holds.*

Proof. Let \mathbb{P} be $\text{Sacks}(\kappa)$ and \mathbb{Q} the Prikry forcing on κ . By Theorem 3.11 and the remarks above, it suffices to show that \mathbb{Q} is κ -geometric and it does not kill the tree property in κ^{++} . The first is Lemma 3.5, the second is Theorem 2 in [7], and we're done. \square

7 Open questions

The general procedure introduced in Theorem 3.11 has its own shortcomings. Among the many "Prikry-like" forcings, there are some that exploit the full supercompactness of one or many cardinal. A priori, this is not immediately useful: under $I0(\kappa, \lambda)$, κ is just λ -supercompact. Also, in [4] there is a proof that it is consistent to have κ less than the least supercompact, so it cannot be a consequence of $I0$. Thus, we can ask this:

Question 1. *What is the consistency strength of $I0 + L(V_{\lambda+1}) \models \kappa$ is λ^+ -supercompact?*

Note that we want κ to be supercompact in $L(V_{\lambda+1})$ because the Prikry forcing in the general procedure must be in $L(V_{\lambda+1})$, and as $L(V_{\lambda+1})$ does not satisfy AC , but just DC_λ , asking for more than λ^+ -supercompactness can be improper.

It is also possible that there are ways to make the general procedure, always or just in some cases, obsolete. For now, there is no proof that $I0$ is needed for the consistency of $I1$ and the combinatorial properties above. The usual large cardinals analysis, in fact, many times has results that have the same large cardinal consistency: this case is different because, while usually one starts with a model with a large cardinal, forces the combinatorial property and proves that the large cardinal is intact, in this case the forcing "reflects" the large cardinal to a cardinal that had already the property desired. So we can ask:

Question 2. *Is it possible to have the results in Corollaries 4.19, 5.12, 5.20 and 6.3 with hypotheses weaker than $I0$? Or is it possible to have the consequences with hypotheses stronger than $I1$?*

With generic absoluteness, it is already possible to raise $I1$ to $j : L_\alpha(V_{\lambda+1}) \prec L_\alpha(V_{\lambda+1})$, with $\alpha < \lambda$. An improvement of generic absoluteness could improve also this, up to the so-called "internal $I0$ ", i.e., the existence for any $\alpha < \Theta$ of $j : L_\alpha(V_{\lambda+1}) \prec L_\alpha(V_{\lambda+1})$, but for $I0$ a different approach could be needed.

References

- [1] P. Cohen, *The independence of the Continuum Hypothesis*. Proceedings of the National Academy of Sciences U.S.A. **50** (1963), 1143–1148.
- [2] P. Corazza, *The wholeness axiom and Laver sequences*. Annals of Pure and Applied Logic **105** (2000), 157–260.

- [3] P. Dehornoy, *Elementary Embeddings and Algebra*. Handbook of Set Theory, Vol 2, 752–789. Springer, Dordrecht 2010.
- [4] V. Dimonte, S. Friedman, *Rank-into-rank hypotheses and the failure of GCH*, Archive for Mathematical Logic **53** (2014), 351–366.
- [5] N. Dobrinen, S. Friedman, *The consistency strength of the tree property at the double successor of a measurable cardinal*. Fundamenta mathematicae **208** (2010), 123–153.
- [6] W. B. Easton, *Powers of regular cardinals*. Annals of Mathematical Logic, **1** (1970), 139–178.
- [7] S. Friedman, A. Halivović, *The tree property at $\aleph_{\omega+2}$* . Journal of Symbolic Logic **76** (2011), 477–490.
- [8] M. Gitik, *Prikry-type forcings*. Handbook of set theory, Vols 2, 1351–1447. Springer, Dordrecht 2010.
- [9] M. Gitik, A. Sharon, *On SCH and the approachability property*. Proceedings of the American Mathematical Society **136** (2008), 311–320.
- [10] A. Kanamori, *The Higher Infinite*. Springer, Berlin 1994.
- [11] K. Kunen, *Elementary embeddings and infinite combinatorics*. Journal of Symbolic Logic **36** (1971), 407–413.
- [12] R. Laver, *Implications between strong large cardinal axioms*. Annals of Pure and Applied Logic **90** (1997), 79–90.
- [13] I. Neeman, *Aronszajn trees and failure of the Singular Cardinal Hypothesis*. Journal of Mathematical Logic **9** (2009), 139–157.
- [14] D. Scott, *Measurable cardinals and constructible sets*. Bulletin de l’Académie Polonaise des Sciences, Série des Sciences Mathématiques, Astronomiques et Physiques **9** (1961), 521–524.
- [15] X. Shi, *Axiom I_0 and higher degree theory*, pre-print.
- [16] J. Silver, *On the singular cardinals problem*. Proceedings of the International Congress of Mathematicians, Vancouver 1974, 265–268
- [17] R. Solovay, *Strongly compact cardinals and the GCH*. Proceedings of the Tarski Symposium, Proceedings of Symposia in Pure Mathematics **25**, Providence, American Mathematical Society, 1974, 365–372.

- [18] W. H. Woodin, *Suitable extender models II: Beyond ω -huge*. Journal of Mathematical Logic **11** (2011), 115–436.