



UNIVERSITÀ
DEGLI STUDI
DI UDINE

Università degli studi di Udine

The derived sequence of a pre-Jaffard family

Original

Availability:

This version is available <http://hdl.handle.net/11390/1216604> since 2025-01-10T08:47:24Z

Publisher:

Published

DOI:10.1007/s00009-022-02028-z

Terms of use:

The institutional repository of the University of Udine (<http://air.uniud.it>) is provided by ARIC services. The aim is to enable open access to all the world.

Publisher copyright

(Article begins on next page)

THE DERIVED SEQUENCE OF A PRE-JAFFARD FAMILY

DARIO SPIRITO

ABSTRACT. We introduce the concept of *pre-Jaffard family*, a generalization of Jaffard families obtained by substituting the locally finite hypothesis with a much weaker compactness hypothesis. From any such family, we construct a sequence of overrings of the starting domain that allows to decompose stable semistar operations and singular length functions in more cases than what is allowed by Jaffard families. We also apply the concept to one-dimensional domains, unifying the treatment of sharp and dull degree of a Prüfer domain.

1. INTRODUCTION

A *Jaffard family* of an integral domain D is a family of flat overrings of D that satisfy some strong independence property, while simultaneously respecting the structure of the whole ring (see Definition 3.7 for a precise definition). This notion allows to extend several results of h -local domains to more general rings; in particular, it was used to extend factorization properties from domains of Dedekind type to a wider class of domains [13, Chapter 6]. Jaffard families were subsequently used to factorize the set of star operations [5, 8] and the set of length functions [7] on an integral domain as the product of the analogous sets on the members of a Jaffard family.

In this paper, we introduce two generalizations of Jaffard families, namely *weak Jaffard families* and *pre-Jaffard families*.

Weak Jaffard families (see Section 5) are very similar to Jaffard families, with the exception that we allow for a single member of the family to behave “badly”.

Pre-Jaffard families (see Section 4), on the other hand, need to satisfy weaker hypothesis, but for these reason are much more common; for example, the set of localizations at the maximal ideals of a domain of dimension 1 is always a pre-Jaffard family. We show in Section 6 how every pre-Jaffard family Θ generates a sequence of weak Jaffard families; this sequence is constructed very similarly to the sequence of derived sets of a topological space, and for this reason we call it the

Date: August 15, 2021.

2010 Mathematics Subject Classification. 13A15, 13F05, 13G05.

Key words and phrases. Jaffard families; length functions; stable operations; sharp degree; dull degree; flat overrings.

derived sequence of Θ . Indeed, when the dimension of the base ring D is 1, the members of the derived sequence of Θ correspond naturally to the member of the sequence of derived sets of the maximal space of D , endowed with the inverse topology. In particular, our construction is a generalization of the study of sharp and dull primes tackled in [16, Section 6] for one-dimensional Prüfer domains; moreover, our terminology symmetrizes some of their results by unifying the concept of sharp and dull degree of a one-dimensional domain into the concept of *Jaffard degree* of a pre-Jaffard family. See Section 8 for the discussion.

In Section 7, we apply weak Jaffard families to the study of singular length functions and of star operations: we show that, given a pre-Jaffard family, we can factorize their set through the derived sequence (Theorem 7.5) allowing a wide generalization of the results on Jaffard families and of [7, Example 6.9].

2. PRELIMINARIES

Throughout the paper, all rings will be commutative, unitary and without zero-divisors, i.e., integral domains; we denote such a ring by D , and we will always use K to denote its quotient field.

We use $\text{Spec}(D)$ and $\text{Max}(D)$, respectively, to denote the spectrum and the maximal spectrum of D , and we denote by $\mathcal{D}(I)$ and $\mathcal{V}(I)$, respectively, the open and the closed set of $\text{Spec}(D)$ associated to the ideal I . The *inverse topology* on $\text{Spec}(D)$ is the topology generated by the $\mathcal{V}(I)$, as I ranges among the finitely generated ideals of D . We denote by Δ^{inv} a subset $\Delta \subseteq \text{Spec}(D)$ endowed with the inverse topology.

The *constructible topology* is the topology generated by the $\mathcal{D}(I)$ and the $\mathcal{V}(J)$ (as I ranges among all ideals and J among all finitely generated ideals); the constructible topology is still compact, but it is also Hausdorff. We denote by Δ^{cons} a subset $\Delta \subseteq \text{Spec}(D)$ endowed with the constructible topology. See [14, Chapter 1] for the construction and properties of the inverse and the constructible topology.

2.1. Overrings. An *overring* of D is a ring T such that $D \subseteq T \subseteq K$; the set of all overrings of D is denoted by $\text{Over}(D)$. This set can be endowed with a topology (called the *Zariski topology*) by taking as a subbasis the family of sets

$$\mathcal{B}(x_1, \dots, x_n) := \{T \in \text{Over}(D) \mid x_1, \dots, x_n \in T\},$$

as x_1, \dots, x_n range in K . Under this topology, $\text{Over}(D)$ is a compact space that is not Hausdorff, and furthermore it is a *spectral space* in the sense of Hochster [15], i.e., there is a ring A (in general not determined explicitly) such that $\text{Spec}(A)$ (endowed with the Zariski topology) is homeomorphic to $\text{Over}(D)$ (see e.g. [1, Proposition 3.5]). The name

“Zariski topology” is also due to the fact that the localization map

$$\begin{aligned}\lambda: \operatorname{Spec}(D) &\longrightarrow \operatorname{Over}(D), \\ P &\longmapsto D_P\end{aligned}$$

is continuous and, indeed, a topological embedding [9, Lemma 2.4].

The *closure under generizations* of a set $\Theta \subseteq \operatorname{Over}(D)$ is

$$\Theta^\uparrow := \{T \in \operatorname{Over}(D) \mid T \supseteq S \text{ for some } S \in \Theta\};$$

a family Θ is *closed by generizations* if $\Theta = \Theta^\uparrow$. The family of all sets that are closed by generizations and compact with respect to the Zariski topology is the family of closed sets of a topology, called the *inverse topology* of $\operatorname{Over}(D)$; equivalently, the inverse topology is the topology generated by the complements of the sets $\mathcal{B}(x_1, \dots, x_n)$.

The *constructible topology* on $\operatorname{Over}(D)$ is the topology generated by both the sets $\mathcal{B}(x_1, \dots, x_n)$ and its complements. The space of all overrings, under both the inverse and the constructible topology, is again compact and a spectral space; moreover, under the constructible topology it is Hausdorff. Every set that is closed in the constructible topology is compact with respect to the Zariski topology.

2.2. Isolated points. Let X be a topological space. A point $x \in X$ is *isolated* if $\{x\}$ is an open set; we denote the set of isolated points of X by $\mathcal{I}(X)$. The set of non-isolated (i.e., limit) points is called the *derived set* of X and is denoted by $\mathcal{D}(X)$.

We set $\mathcal{D}^0(X) := X$ and, for every ordinal α , we define:

$$\mathcal{D}^\alpha(X) := \begin{cases} \mathcal{D}(\mathcal{D}^\gamma(X)) & \text{if } \alpha = \gamma + 1 \text{ is a successor ordinal,} \\ \bigcap_{\beta < \alpha} \mathcal{D}^\beta(X) & \text{if } \alpha \text{ is a limit ordinal.} \end{cases}$$

The set $\mathcal{D}^\alpha(X)$ is called the α -th *Cantor-Bendixson derivative* of X , and the smallest ordinal α such that $\mathcal{D}^\alpha(X) = \mathcal{D}^{\alpha+1}(X)$ is called the *Cantor-Bendixson rank* of X . If $\mathcal{D}^\alpha(X) = \emptyset$ for some ordinal α , the space X is said to be *scattered*; equivalently, X is scattered if and only if every nonempty subspace has an isolated point.

2.3. Semistar operations and length functions. Let D be a domain and let $\mathbf{F}_D(K)$ be the set of D -submodules of K . A *semistar operation* on D is a map $\star : \mathbf{F}_D(K) \longrightarrow \mathbf{F}_D(K)$, $I \mapsto I^\star$, such that, for every $I, K \in \mathbf{F}_D(K)$ and every $x \in K$:

- $I \subseteq I^\star$;
- if $I \subseteq J$, then $I^\star \subseteq J^\star$;
- $(I^\star)^\star = I^\star$;
- $x \cdot I^\star = (xI)^\star$.

If $(I \cap J)^\star = I^\star \cap J^\star$ for every I, J , we say that \star is *stable*. We denote the sets of semistar operations and of stable semistar operations, respectively, by $\operatorname{SStar}(D)$ and $\operatorname{SStar}_{\text{st}}(D)$. These two sets have a partial order, given by $\star_1 \leq \star_2$ if and only if $I^{\star_1} \subseteq I^{\star_2}$ for every ideal I ; the

infimum of a family Δ of semistar operations is the map $\sharp : I \mapsto \bigcap \{I^\star \mid \star \in \Delta\}$. If $\Delta \subseteq \text{SStar}_{\text{st}}(D)$, then also \sharp is stable.

Let $\text{Mod}(D)$ be the category of D -modules. A *length function* on D is a function $\ell : \text{Mod}(D) \longrightarrow \mathbb{R}^{\geq 0} \cup \{\infty\}$ such that:

- $\ell(0) = 0$;
- if $0 \longrightarrow N \longrightarrow M \longrightarrow P \longrightarrow 0$ is an exact sequence, then $\ell(M) = \ell(P) + \ell(N)$;
- for every module M , we have $\ell(M) = \sup\{\ell(N) \mid N \text{ is a finitely generated submodule of } M\}$.

The sum of a family Λ of length functions is defined as the map such that

$$\left(\sum_{\ell \in \Lambda} \ell \right) (M) = \sup\{\ell_1(M) + \cdots + \ell_n(M)\},$$

as $\{\ell_1, \dots, \ell_n\}$ ranges among the finite subsets of Λ .

If T is a flat overring of D , then we can associate to any length function ℓ on D a length function ℓ^D on T by restriction of scalars, i.e., setting $\ell^D(M) := \ell(M)$ for all T -modules M . Moreover, we can defined a new length function $\ell \otimes T$ on D by setting

$$(\ell \otimes T)(M) := \ell(M \otimes T).$$

for all D -modules M .

By [7, Theorem 6.5] and the subsequent discussion, there is a bijection between the set $\mathcal{L}_{\text{sing}}(D)$ of length functions such that $\ell(M) \in \{0, +\infty\}$ for all $M \in \text{Mod}(D)$ and the set $\text{SStar}_{\text{st}}(D)$ of stable semistar operations on D , and by [7, Proposition 6.6] the infimum of a family of stable operations correspond to the sum of the corresponding length functions. Moreover, the passage from a length function ℓ on a flat overring T to ℓ^D correspond to the passage from the a stable operation \star on T to the closure $I \mapsto (IT)^\star$ on D .

3. JAFFARD OVERRINGS

In this paper we will mostly use families consisting of *flat overrings*, i.e., overrings of a domain D that are flat when considered as D -modules. However, in many case we will define rings by intersecting localizations; thus we need the following definition.

Definition 3.1. *An overring T of D is a sublocalization of D if there is a set $\Delta \subseteq \text{Spec}(D)$ such that $T = \bigcap \{D_P \mid P \in \Delta\}$.*

If T is a sublocalization of D , we set:

- $\sigma(T) := \{Q \cap D \mid Q \in \text{Spec}(T)\}$;
- $\Sigma(T) := \{P \in \text{Spec}(D) \mid T \subseteq D_P\}$;
- $T^\perp := \bigcap \{D_P \mid P = (0) \text{ or } P \in \text{Spec}(D) \setminus \Sigma(T)\}$.

Note that, by definition, T^\perp is a sublocalization too, and thus it makes sense to consider $\sigma(T^\perp)$ and $\Sigma(T^\perp)$.

Lemma 3.2. *For every sublocalization T of D , we have $\Sigma(T) \cup \Sigma(T^\perp) = \text{Spec}(D)$.*

Proof. If $P \notin \Sigma(T)$, then by definition $T^\perp \subseteq D_P$, and thus $P \in \Sigma(T^\perp)$. \square

Every flat overring is a sublocalization [10, Corollary to Theorem 2], but the converse is not true (see [18] and [6, Example 6.3]). We can characterize when a sublocalization is flat.

Lemma 3.3. *Let T be a sublocalization of D . Then, T is flat over D if and only if $\Sigma(T) = \sigma(T)$.*

Proof. The containment $\Sigma(T) \subseteq \sigma(T)$ holds for every sublocalization.

If T is flat and $P \in \sigma(T)$, then $PT \neq T$, and by [10, Theorem 1] we have $T \subseteq D_P$, i.e., $P \in \Sigma(T)$. Conversely, if $\sigma(T) = \Sigma(T)$ and $PT \neq T$, let Q be a prime ideal of T above PT : then, $P' := Q \cap D \in \sigma(T)$ and thus $T \subseteq D_{P'}$. Hence, $PT \cap D \subseteq PD_{P'} \cap D = P$, and so $P' = P$, and in particular $T \subseteq D_P$. Again by [10, Theorem 1], T is flat. \square

Lemma 3.4. *Let A be a flat overring and B a sublocalization of D . Then, $AB = K$ if and only if $\sigma(A) \cap \sigma(B) = \{(0)\}$.*

Proof. Suppose that $AB = K$, and let $P \in \sigma(A) \cap \sigma(B)$. Since A is flat, $P \in \Sigma(A)$ and so $A \subseteq D_P$; on the other hand, if Q is a prime ideal of B above P , then $D_P \subseteq B_Q$. Hence, $K = ABD_P = (AD_P)(BD_P) \subseteq D_P B_Q = B_Q$. It follows that $Q = (0)$ and so $P = (0)$ too.

Conversely, if $\sigma(A) \cap \sigma(B) = \{(0)\}$ then the claim follows from [13, Lemma 6.2.1]. \square

Definition 3.5. *Let D be an integral domain with quotient field K and let Θ be a family of overrings of D . We say that Θ is:*

- complete if, for every ideal I of D , we have $I = \bigcap \{IT \mid T \in \Theta\}$;
- independent if, for every $A, B \in \Theta$ such that $A \neq B$, we have $\sigma(A) \cap \sigma(B) = \{(0)\}$;
- strongly independent if, for every $A \in \Theta$, we have

$$A \cdot \left(\bigcap_{\substack{B \in \Theta \\ B \neq A}} B \right) = K$$

- locally finite if every $x \in K$, $x \neq 0$, is a nonunit in only finitely many members of Θ .

By Lemma 3.4, if Θ is a family of flat subsets, then Θ is independent if and only if $AB = K$ for every $A \neq B$ in Θ ; in particular, a strongly independent set of flat overrings is always independent. We can prove when the converse happens.

Proposition 3.6. *Let Θ be a complete and independent set of flat overring of D . Then, Θ is strongly independent if and only if Θ is locally finite.*

Proof. If Θ is locally finite, the claim follows from [13, Theorem 6.3.1(4)] (see below for the definitions used in the reference). Suppose that Θ is strongly independent but not locally finite: then, there is a nonzero $x \in D$ such that $xT \neq T$ for an infinite family $\Theta' \subseteq \Theta$. Hence, for each $T \in \Theta'$ there is a prime ideal $P_T \in \text{Spec}(D)$ such that $x \in P_T$ and $P_T T \neq T$; let Λ be the family of such ideals. Then, Λ is an infinite subset of the compact space $\text{Spec}(D)^{\text{cons}}$, and thus it has a limit point Q ; furthermore, Λ is contained in the clopen set $\mathcal{V}(x)$ of $\text{Spec}(D)^{\text{cons}}$ and thus also $Q \in \mathcal{V}(x)$, i.e., $x \in Q$; in particular, $Q \neq (0)$.

Since Θ is complete and independent, there is a unique $S \in \Theta$ such that $QS \neq S$; let $A := \bigcap \{T \in \Theta \mid T \neq S\}$. Then, A is a sublocalization of D ; by [10, Theorem 1], $\sigma(A)$ is the image of $\text{Spec}(A)$ under the restriction map $Z \mapsto Z \cap D$, and thus $\sigma(A)$ is a closed set in the constructible topology. Moreover, $\sigma(A)$ contains all the elements of Λ except one (the ideal P_S), and thus it must contain also the limit point Q of $\Lambda \setminus \{P_S\}$. Therefore, $D_Q A \neq K$. However, $S \subseteq D_Q$ since S is flat; therefore, $AS \subseteq D_Q S \neq K$. This contradicts the fact that Θ is strongly independent: hence Θ must be locally finite. \square

Note that the above result does not hold without the hypothesis that Θ is complete: see Example 6.7 below.

Families satisfying the hypothesis of the previous proposition have their own name.

Definition 3.7. *Let Θ be a family of overrings of D . We say that Θ is a Jaffard family of D if:*

- *either $K \notin \Theta$ or $\Theta = \{K\}$;*
- *every $T \in \Theta$ is flat;*
- *Θ is complete;*
- *Θ is independent;*
- *Θ is locally finite.*

We say that an overring T of D is a Jaffard overring if it belongs to a Jaffard family of D .

Remark 3.8.

- (1) Definition 3.7 is not the original one of a Jaffard family, but it is the one most useful for our purpose; see [13, Section 6.3] and [5, Proposition 4.3].
- (2) By Proposition 3.6 (see also [13, Theorem 6.3.1(4)]) the two conditions “ Θ is independent” and “ Θ is locally finite” can be unified into the single one “ Θ is strongly independent”.

- (3) If P is a nonzero prime of D , then there is exactly one $T \in \Theta$ such that $PT \neq T$: indeed, such a T must exist since Θ is complete, while there cannot be two of them due to independence condition and the fact that a flat extension satisfies going-down. In particular, Θ induces a partition on $\text{Max}(D)$, called a *Matlis partition* [13, Section 6.3].
- (4) If $\Theta \in K$ (i.e., $\Theta = \{K\}$) then since Θ must be complete we must have also $D = K$, i.e., D must be a field.

Proposition 3.9. *Let T be a flat overring of D . Then, the following are equivalent:*

- (i) T is a Jaffard overring of D ;
- (ii) $T \cdot T^\perp = K$;
- (iii) $\{T, T^\perp\}$ is a Jaffard family of D ;
- (iv) $\sigma(T) \cap \sigma(T^\perp) = \{(0)\}$;
- (v) for every nonzero $P \in \sigma(T)$, we have $PT^\perp = T^\perp$;
- (vi) there is a sublocalization A of D such that $\{T, A\}$ is complete and $TA = K$.

Proof. (iii) \implies (i) follows from the definitions.

(i) \implies (ii) Let Θ be a Jaffard family containing T , and let $\Theta^\perp(T) := \bigcap \{S \in \Theta \mid S \neq T\}$. For every nonzero prime ideal P of D out of $\Sigma(T)$, there is a unique $S \in \Theta$ such that $PS \neq S$; since S is flat, we have $S \subseteq D_P$, and so

$$\Theta^\perp(T) \subseteq \bigcap \{P \in \text{Spec}(D) \mid P \notin \Sigma(T)\} = T^\perp.$$

Since Θ is strongly independent, $T \cdot \Theta^\perp(T) = K$, and so $TT^\perp = K$.

(ii) \implies (iii) Let $\Theta := \{T, T^\perp\}$. Clearly, Θ is locally finite and complete, while it is independent by Lemma 3.4 (since T is flat by hypothesis). Since $T \cap T^\perp = D$, by [13, Theorem 6.2.2(1)] T^\perp is also flat; hence, Θ is a Jaffard family.

(ii) \iff (iv) is exactly Lemma 3.4.

(ii) \implies (vi) is obvious. To show (vi) \implies (i), it is enough to show that $\Theta := \{T, A\}$ is a Jaffard family. By hypothesis, Θ is complete and locally finite, while it is independent by Lemma 3.4. In particular, by [13, Theorem 6.2.2(1)] A is also flat, and thus Θ is a Jaffard family.

(iii) \implies (v) If $P \in \sigma(T)$, $P \neq (0)$, then $PT \neq T$; but since $\{T, T^\perp\}$ is a Jaffard family, no nonzero prime can survive in both T and T^\perp , and so $PT^\perp = T^\perp$.

(v) \implies (ii) Suppose that $T \cdot T^\perp \neq K$: then, there is a nonzero $Q \in \text{Spec}(D)$ such that $QTT^\perp \neq TT^\perp$, and so both $QT \neq T$ and $QT^\perp \neq T^\perp$. However, the first condition implies that $Q \in \sigma(T)$, contradicting the hypothesis. \square

Corollary 3.10. *Let Θ be a complete and independent family of flat overrings of D . Then, Θ is a Jaffard family if and only if each $T \in \Theta$ is a Jaffard overring.*

Proof. If Θ is a Jaffard family then every $T \in \Theta$ is a Jaffard overring by definition. Conversely, suppose each $T \in \Theta$ is a Jaffard overring; by Proposition 3.6 we only need to show that Θ is strongly independent.

Fix $T \in \Theta$. If $S \in \Theta \setminus \{T\}$, then $\sigma(S) \cap \sigma(T) = \Sigma(S) \cap \Sigma(T) = \emptyset$, and thus $T^\perp \subseteq S$. Therefore,

$$T \left(\bigcap_{S \in \Theta \setminus \{T\}} S \right) \supseteq TT^\perp = K$$

using Proposition 3.9. Hence, Θ is strongly independent and thus a Jaffard family. \square

We conclude this section with a lemma that will be useful later.

Lemma 3.11. *Let Θ be a complete and independent family of flat overrings of D and let $P \neq (0)$ be a prime ideal of D . For every $S \in \Theta$, either $PS \neq S$ or $D_P S = K$.*

Proof. Suppose that $PS = S$. Since Θ is complete, there is a $S' \in \Theta$ such that $PS' \neq S'$; since S' is flat, by Lemma 3.3 $S' \subseteq D_P$. Hence, $SS' \subseteq SD_P$; however, since Θ is independent $SS' = K$. Hence $K = SD_P$. \square

4. PRE-JAFFARD FAMILIES

The hypothesis that a family is locally finite is usually very strong. To expand our reach beyond Jaffard families, we define a new class of families by weakening this condition.

Definition 4.1. *Let Θ be a family of overrings of D . We say that Θ is a pre-Jaffard family of D if:*

- *either $K \notin \Theta$ or $\Theta = \{K\}$;*
- *every element of Θ is flat over D ;*
- *Θ is independent;*
- *Θ is complete;*
- *Θ is compact in the Zariski topology.*

Remark 4.2. We do not know any example of a family of overring satisfying the first three conditions of Definition 4.1 but that is not compact; it is possible that the compactness condition is actually redundant.

Proposition 4.3. *A pre-Jaffard family is Hausdorff, with respect to the inverse topology.*

Proof. Without loss of generality we can suppose that $K \notin \Theta$. Fix two distinct overrings $T, S \in \Theta$. Let $\mathcal{C} := \{\mathcal{B}(x) \mid x \in T \setminus S\} \cup \{\mathcal{B}(y) \mid y \in S \setminus T\}$; we claim that \mathcal{C} is a cover of Θ .

Since $ST = K$, we have $S \subsetneq T$ and $T \subsetneq S$, and thus $T \setminus S$ and $S \setminus T$ are both nonempty; it follows that S, T belong to some member of \mathcal{C} .

Let $A \in \Theta \setminus \{S, T\}$: then,

$$A \cap (T \setminus S) = A \cap T \cap (K \setminus S),$$

and thus if $A \cap (T \setminus S) = \emptyset$ then $A \cap T \subseteq S$. However, $(A \cap T)S = AS \cap TS = K$; hence, $A \cap (T \setminus S) \neq \emptyset$ and so there is an $x \in A \cap (T \setminus S)$, i.e., $A \in \mathcal{B}(x)$. Thus, any such A belong to some member of \mathcal{C} , and \mathcal{C} is a cover of Θ .

Since Θ is compact in the Zariski topology, we can find $x_1, \dots, x_n, y_1, \dots, y_m$ such that $\{\mathcal{B}(x_1), \dots, \mathcal{B}(x_n), \mathcal{B}(y_1), \dots, \mathcal{B}(y_m)\}$ is a finite subcover. Let

$$\Omega_1 := \bigcap_{i=1}^n \mathcal{B}(x_i)^c \quad \text{and} \quad \Omega_2 := \bigcap_{j=1}^m \mathcal{B}(y_j)^c;$$

then, Ω_1 and Ω_2 are both open in the inverse topology, since they are a finite intersection of subbasic open sets. Moreover, $S \in \Omega_1$ since $x_i \notin S$ for every i , while $T \in \Omega_2$ since $y_j \notin T$ for every j . Finally,

$$\Omega_1 \cap \Omega_2 = \bigcap_{i=1}^n \mathcal{B}(x_i)^c \cap \bigcap_{j=1}^m \mathcal{B}(y_j)^c = \left(\bigcup_{i=1}^n \mathcal{B}(x_i) \cup \bigcup_{j=1}^m \mathcal{B}(y_j) \right)^c \subseteq \Theta^c,$$

that is, $\Omega_1 \cap \Omega_2$ does not intersect Θ . Hence, $\Omega_1 \cap \Theta$ and $\Omega_2 \cap \Theta$ are disjoint neighborhood of S and T in Θ , with respect to the inverse topology. Thus, Θ is Hausdorff in the inverse topology. \square

Proposition 4.4. *A Jaffard family of D is a pre-Jaffard family.*

Proof. By definition, any Jaffard family of D is independent, complete and composed of flat overrings. Furthermore, any locally finite family of overrings is compact (see e.g. the proof of [2, Corollary 8]), and thus a Jaffard family is also pre-Jaffard. \square

Proposition 4.5. *Let Θ be a pre-Jaffard family of D , and let $T \in \Theta$. Then, T is a Jaffard overring of D if and only if $\Theta \setminus \{T\}$ is compact in the Zariski topology. Furthermore, if this happens, then T is isolated in Θ , with respect to the inverse topology.*

Proof. If T is a Jaffard overring, by Proposition 3.9 we have $TT^\perp = K$, and in particular no overring of D different from K contains both T^\perp and T . Then, $\Theta^\uparrow \cap \{T^\perp\}^\uparrow = (\Theta \setminus \{T\})^\uparrow$: however, since Θ^\uparrow and $\{T^\perp\}^\uparrow$ are closed in the inverse topology (the former since Θ is compact by hypothesis), then also $(\Theta \setminus \{T\})^\uparrow$ is inverse-closed. Therefore, $\Theta \setminus \{T\}$, which is the set of minimal elements of $(\Theta \setminus \{T\})^\uparrow$, is compact with respect to the Zariski topology. This also shows that T is isolated in Θ , with respect to the inverse topology.

Suppose $\Theta \setminus \{T\}$ is compact, and let $A := \bigcap \{S \mid S \in \Theta, S \neq T\}$. Then, A is a sublocalization of D ; moreover, since T is flat and $\Theta \setminus \{T\}$ is compact, by [2, Corollary 5] we have

$$TA = T \bigcap_{S \in \Theta} S = \bigcap_{S \in \Theta} TS = K.$$

Hence, T is a Jaffard overring by Proposition 3.9. \square

Remark 4.6. Proposition 4.5 cannot be improved to a full equivalence between being a Jaffard overring and being an isolated point of Θ^{inv} . Consider the ring D defined in [17, Example 2]. Then, D is a two-dimensional domain such that:

- all its finitely generated ideals are principal (i.e., D is a Bézout domain); in particular, D_M is a valuation domain for all maximal ideals M ;
- all its maximal ideals, except for one (say M_∞), have height 1;
- M_∞ is the radical of a principal ideal;
- the unique nonzero, nonmaximal prime ideal P is contained in a unique maximal ideal (M_∞), but also in the union of all maximal ideals distinct from M_∞ .

Let $\Theta := \{D_M \mid M \in \text{Max}(D)\}$. Then, every $T \in \Theta$ is flat and Θ is complete; furthermore, Θ is independent (if $D_M D_N \neq K$, then $M \cap N$ should contain a nonzero prime ideal, a contradiction) and compact in the Zariski topology (since the localization map is continuous by [9, Lemma 2.4] and so Θ is homeomorphic to $\text{Max}(D)$); thus, Θ is a pre-Jaffard family.

Let $V := D_{M_\infty}$: then, T is not a Jaffard overring of D . Indeed, $V^\perp = \bigcap \{D_M \mid M \in \text{Max}(D), M \neq M_\infty\}$ is such that $PV^\perp \neq V^\perp$: otherwise, since D is Bézout, there would be an $a \in P$ such that $aV^\perp = V^\perp$. However, there is also a maximal ideal $M \neq M_\infty$ such that $a \in M$: hence, $MV^\perp = V^\perp$, against the fact that $V^\perp \subseteq D_M$ by construction. Hence, $PV^\perp \neq V^\perp$ and so $V^\perp \subseteq D_P$, so that $VV^\perp \subseteq VD_P = D_P$ (since $V = D_{M_\infty} \subseteq D_P$). By Proposition 3.9, V is not a Jaffard overring of D .

We claim that V is isolated in Θ , with respect to the inverse topology. Indeed, M_∞ is the radical of a principal ideal, say bD ; hence, M_∞ is the unique $T \in \Theta$ such that $b^{-1} \notin T$, i.e., $\mathcal{B}(b^{-1})^c \cap \Theta = \{M_\infty\}$. However, $\mathcal{B}(b^{-1})^c$ is the complement of an open and compact subset of $\text{Over}(D)$, and thus it is open in the inverse topology; hence, V is isolated in Θ^{inv} .

The following two results show how to construct pre-Jaffard families from other such families by taking intersections.

Lemma 4.7. *Let Θ be a family of overrings that is compact in the Zariski topology. Let $\Theta_1, \dots, \Theta_n$ be subsets of Θ , and for each i let $S_i := \bigcap \{T \mid T \in \Theta_i\}$. Then, the family*

$$\Theta' := \left(\Theta \setminus \bigcup_{i=1}^n \Theta_i \right) \cup \{S_1, \dots, S_n\}$$

is compact in the Zariski topology.

Proof. Let $\Omega := \{\Omega_\alpha\}_{\alpha \in A}$ be an open cover of Θ' . If $S_i \in \Omega_\alpha$, then $\Theta_i \subseteq \Omega_\alpha$ (since this holds for every subbasic open set $\mathcal{B}(x)$); hence, Ω

is also an open cover of Θ . Since Θ is compact, we can find a finite subcover $\{\Omega_{\alpha_1}, \dots, \Omega_{\alpha_k}\}$ of Θ ; furthermore, for every i we can find a $\beta_i \in A$ such that $S_i \in \Omega_{\beta_i}$. Then, $\{\Omega_{\alpha_1}, \dots, \Omega_{\alpha_k}, \Omega_{\beta_1}, \dots, \Omega_{\beta_n}\}$ is a finite subcover of Θ' . Thus, Θ' is compact. \square

Proposition 4.8. *Let Θ be a pre-Jaffard family, and let $\Theta_1, \dots, \Theta_n$ be pairwise disjoint subsets of Θ that are compact in the Zariski topology. For every i , let $S_i := \bigcap \{T \mid T \in \Theta_i\}$. Then, the family*

$$\Theta' := \left(\Theta \setminus \bigcup_{i=1}^n \Theta_i \right) \cup \{S_1, \dots, S_n\}$$

is a pre-Jaffard family.

Proof. By induction, it is enough to prove the claim for $n = 1$; let $S := S_1$.

By construction, Θ' is complete, and by Lemma 4.7 it is compact in the Zariski topology. It is independent: indeed, take $T_1, T_2 \in \Theta'$. If $T_1 \neq S \neq T_2$ then $T_1 T_2 = K$ since Θ is independent, while if $S = T_2$ then

$$T_1 S = T_1 \bigcap_{T \in \Theta_1} T = \bigcap_{T \in \Theta_1} T_1 T = K$$

by [2, Corollary 5], since Θ_1 is compact and every $T \in \Theta_1$ is in Θ and is different from T_1 . Thus, Θ' is independent.

We only need to prove that S is flat. By construction, S is a sublocalization. If $P \in \sigma(\text{Spec}(S))$ is a nonzero prime, then $PT = T$ for every $T \in \Theta \setminus \{S\}$ (otherwise $K = TS \subseteq D_P$, a contradiction); on the other hand, there is a $S' \in \Theta_1$ such that $P \in \Sigma(S')$, and thus $P \in \Sigma(S)$. Hence, $\sigma(\text{Spec}(S)) = \Sigma(S)$, and S is flat by Lemma 3.3. \square

5. WEAK JAFFARD FAMILIES

Let Θ be a pre-Jaffard family. In general, we cannot expect the properties of a Jaffard family to hold also for Θ ; however, we want to show that at least some properties hold also under the weaker pre-Jaffard hypothesis. To do so, we want to proceed “step-by-step”, isolating first the Jaffard overring belonging to Θ ; from a technical point of view, we need the following definition.

Definition 5.1. *Let Θ be a family of overrings of D and let $T_\infty \in \Theta$. We say that Θ is a weak Jaffard family of D pointed at T_∞ if:*

- *either $K \notin \Theta$ or $\Theta = \{K\}$;*
- *Θ is complete and independent;*
- *every $T \in \Theta \setminus \{T_\infty\}$ is a Jaffard overring of D ;*
- *T_∞ is flat over D .*

Lemma 5.2. *Let Θ be a family of flat overrings of D , and let B be an overring of D . Let $\Theta_B := \{TB \mid T \in \Theta\}$*

- (a) If Θ is independent, Θ_B is independent.
- (b) If Θ is complete with respect to D , then Θ_B is complete with respect to B .
- (c) If every $T \in \Theta$ is flat as a D -module, every $TB \in \Theta_B$ is flat as a B -module.
- (d) If T is a Jaffard overring of B and $T \neq K$, then TB is a Jaffard overring of B .
- (e) If Θ is a Jaffard family of B , then $\Theta_B \setminus \{K\}$ is a Jaffard family of B .

Proof. (a) If $TB \neq T'B$ with $T \neq T'$ in Θ , then $(TB)(T'B) = (TT')B = K$.

(b) Let I be a B -submodule of the quotient field K . Then, $IB = B$, and thus

$$I = IB = \bigcap_{T \in \Theta} (IB)T = \bigcap_{T \in \Theta} I(BT) = \bigcap_{S \in \Theta_B} IS$$

so that Θ_B is complete with respect to B .

(c) Since B is an overring, the extension $A \subseteq B$ is an epimorphism, and thus $TB \simeq T \otimes B$ [4, Lemma 1.0]. The claim follows.

(d) If T is a Jaffard overring, then by Proposition 3.9 $\Theta := \{T, T^\perp\}$ is a Jaffard family of D . By the previous points, $\Theta_B = \{TB, T^\perp B\}$ is complete, independent and formed by flat overrings; since it is clearly locally finite, it is a Jaffard family, and thus TB is a Jaffard overring.

(e) follows from the previous points and from Corollary 3.10. \square

A Jaffard family is always a weak Jaffard family, pointed to any of its elements. Analogously, a weak Jaffard family is pre-Jaffard, as we show next.

Proposition 5.3. *Let Θ be a weak Jaffard family of D pointed at S .*

- (a) *If J is a proper ideal of D and $JS = S$, then $JT \neq T$ for only finitely many $T \in \Theta$.*
- (b) *Θ is a pre-Jaffard family of D .*

Proof. (a) Let $B := \bigcap \{D_M \mid M \in \mathcal{V}(J)\}$: we claim that $BS = K$. Indeed, $\mathcal{V}(J)$ is a compact subset of $\text{Spec}(D)$, and thus

$$BS = \left(\bigcap_{M \in \mathcal{V}(J)} D_M \right) S = \bigcap_{M \in \mathcal{V}(J)} D_M S = K$$

since if $J \subseteq M$ then $JT \neq T$ for some $T \in \Theta$ and thus $D_M S \supseteq TS = K$.

Consider the family $\Theta_B := \{BT \mid T \in \Theta\} \setminus \{K\}$: by Lemma 5.2, Θ_B is a complete and independent set of flat overrings of B , and all its element except BS are Jaffard overrings of B . By Corollary 3.10, Θ_B is a Jaffard family of B , and thus it is locally finite. For every $T \in \Theta$ such that $JT \neq T$, also $JTB \neq TB$; hence, there are at most finitely many elements of Θ such that $JT \neq T$, as claimed.

(b) We need only to show that Θ is compact, with respect to the Zariski topology. Let $\{\mathcal{B}(x_\alpha)\}_{\alpha \in \mathcal{I}}$ be an open cover of Θ , and suppose $S \in \mathcal{B}(x)$. Let $J := (D :_D x)$: then, $JS = (S :_S x) = S$ (using the flatness of S), and thus there are only finitely many $T \in \Theta$ such that $JT \neq T$; call them T_1, \dots, T_n . Therefore, if $A \in \Theta \setminus \{T_1, \dots, T_n\}$ then $JA = A$ and $(A :_A x) = A$, i.e., $x \in A$; thus $\mathcal{B}(x) \setminus \Theta$ is finite. It follows that we can find a subcover $\{\mathcal{B}(x), \mathcal{B}(x_1), \dots, \mathcal{B}(x_n)\}$ by choosing x_1, \dots, x_n such that $x_i \in T_i$. Since the cover was arbitrary, $\Theta \cup \{S\}$ is compact. \square

Weak Jaffard families are much more ubiquitous than Jaffard families; the main reason is that a weak Jaffard family has a place to “hide the singularities” of D (namely, the ring T_∞ to which the family is pointed), while a Jaffard family does not have such a luxury. A first way in which weak Jaffard families arise is from a set of Jaffard overrings.

Proposition 5.4. *Let Θ be an independent set of Jaffard overrings of D , and let*

$$S := \bigcap \{D_P \mid PT = T \text{ for every } T \in \Theta\}.$$

Then, $\Theta \cup \{S\}$ is a weak Jaffard family of D pointed at S .

Proof. Let $P \in \text{Spec}(D)$. If $PT \neq T$ for some $T \in \Theta$, then $T \subseteq D_P$ since T is flat [10, Theorem 1]; if $PT = T$ for every $T \in \Theta$, then $S \subseteq D_P$. Therefore, every localization D_P of D contains at least one element of $\Theta \cup \{S\}$. Hence, for every ideal I of D ,

$$I = \bigcap_{M \in \text{Max}(D)} ID_M \supseteq \bigcap_{T \in \Theta \cup \{S\}} IT \supseteq I.$$

Thus $\Theta \cup \{S\}$ is complete.

If $T, T' \in \Theta$, $T \neq T'$, then $TT' = K$ by hypothesis. To show that Θ is independent, let $\Sigma := \{P \in \text{Spec}(D) \mid PT = T \text{ for every } T \in \Theta\}$. We claim that $\Sigma = \bigcap \{\text{Spec}(D) \setminus \Sigma(T) \mid T \in \Theta\}$. Indeed, if $P \in \Sigma$ then $PT = T$ for every $T \in \Theta$, and thus P is in the intersection; conversely, if $P \notin \Sigma(T)$ for every $T \in \Theta$, then (since each T is flat) we have by Lemma 3.3 $P \notin \sigma(T)$, and thus $PT = T$, so that $P \in \Sigma$. By Proposition 3.9, $\text{Spec}(D) \setminus \Sigma(T) = \Sigma(T^\perp) \setminus \{(0)\}$; hence,

$$\Sigma \cup \{(0)\} = \bigcap_{T \in \Theta} (\Sigma(T^\perp) \setminus \{(0)\}) \cup \{(0)\} = \bigcap_{T \in \Theta} \Sigma(T^\perp)$$

Again by Proposition 3.9, T^\perp is a Jaffard overring of D , and thus it is flat; hence, $\Sigma(T^\perp) = \sigma(T^\perp)$ is closed in the constructible topology, and thus also $\Sigma \cup \{(0)\}$ is closed in the constructible topology, and in particular it is compact. Fix now a $T \in \Theta$. Using the flatness of T , we

have

$$TS = T \left(\bigcap_{P \in \Sigma \cup \{0\}} D_P \right) = K \cap \bigcap_{P \in \Sigma} TD_P = K,$$

since $D_P \supseteq T^\perp$ for every nonzero $P \in \Sigma \supseteq \Sigma(T^\perp)$. Hence, $\Theta \cup \{S\}$ is independent.

Since every $T \in \Theta$ is a Jaffard overring, we only need to show that S is flat. Suppose that $P \in \sigma(S)$: since $\Theta \cup \{S\}$ is complete and independent, we must have $PT = T$ for every $T \in \Theta$, and thus $P \in \Sigma$, so that, by definition $S \subseteq D_P$. Thus $P \in \Sigma(S)$ and $\sigma(S) = \Sigma(S)$; by Lemma 3.3, S is flat. \square

The previous two propositions provide a way to pass from a pre-Jaffard family to a weak Jaffard family.

Definition 5.5. *If Θ is a pre-Jaffard family of D , we denote by Θ_J the set of Jaffard overrings contained in Θ .*

Proposition 5.6. *Let Θ be a pre-Jaffard family of D , and suppose that $\Theta_J \neq \Theta$. Let $S := \bigcap \{T \mid T \in \Theta \setminus \Theta_J\}$. Then:*

- (a) $\Theta_J \cup \{S\}$ is a weak Jaffard family of D pointed at S ;
- (b) $\Theta \setminus \Theta_J$ is a pre-Jaffard family of S .

Proof. (a) The set Θ_J is a set of Jaffard overrings of D ; we claim that $S = \bigcap \{D_P \mid PT = T \text{ for every } T \in \Theta\}$. Indeed, since every $T \in \Theta$ is flat we have

$$S = \bigcap_{T \in \Theta \setminus \Theta_J} \bigcap_{P \in \Sigma(T)} D_P = \bigcap_{P \in \Sigma} D_P$$

where $\Sigma := \bigcup \{\Sigma(T) \mid T \in \Theta \setminus \Theta_J\}$. Hence, $\Theta_J \cup \{S\}$ is a weak Jaffard family by Proposition 5.4.

(b) Let $\Theta' := \Theta \setminus \Theta_J$. Then, Θ' is an independent family of flat overrings of S , and is complete with respect to S , since if I is an ideal of S then

$$I = IS = \bigcap_{T \in \Theta} IST = \bigcap_{T \in \Theta_J} IST \cap \bigcap_{T \in \Theta'} IST = \bigcap_{T \in \Theta'} IS$$

as $S \subseteq T$ if $T \in \Theta'$ while $ST = K$ if $T \in \Theta_J$. We thus need to show that Θ' is compact.

For every $T \in \Theta_J$, by Proposition 4.5 $\Theta \setminus \{T\}$ is compact in the Zariski topology, and thus $\Lambda_T := (\Theta \setminus \{T\})^\uparrow$ is closed in the inverse topology. Thus, also the intersection $\Lambda := \bigcap \{\Lambda_T \mid T \in \Theta_J\}$ is closed in the inverse topology. However, since $\{S\}^\uparrow \cap \{S'\}^\uparrow = \{K\}$ for every $S \neq S'$ in Θ , the set of minimal elements of Λ is exactly $\Theta \setminus \Theta_J = \Theta'$; hence, Θ' is compact, as claimed. \square

6. THE DERIVED SEQUENCE

Let Θ be a pre-Jaffard family of D , and let Θ_J be the set of Jaffard overrings inside Θ . If $\Theta = \Theta_J$, then by Corollary 3.10 Θ is a Jaffard family; on the other hand, if $\Theta \neq \Theta_J$ then by Proposition 5.6 we can define an overring T_1 of D such that:

- $\Theta_J \cup \{T_1\}$ is a weak Jaffard family of D ;
- $\Theta_1 := \Theta \setminus \Theta_J$ is a pre-Jaffard family of T_1 .

In particular, we can repeat the same construction on Θ_1 : either Θ_1 is a Jaffard family of T_1 or we can find an overring T_2 of T_1 (and so of D) such that $(\Theta_1)_J \cup \{T_2\}$ is a weak Jaffard family of T_1 and $\Theta_2 := \Theta_1 \setminus (\Theta_1)_J$ is a pre-Jaffard family of T_2 ; then we can use the same construction on T_2 , and so on. We now want to define rings T_α and subfamilies Θ_α for every ordinal α .

To start, define $T_0 := D$ and $\Theta_0 := \Theta$.

Suppose that for every ordinal $\beta < \alpha$ we have defined a ring T_β and a subset $\Theta_\beta \subseteq \Theta$ that is a pre-Jaffard family of T_β . Then:

- if $\alpha = \gamma + 1$ is a successor ordinal, we define

$$\Theta_\alpha := \Theta_\gamma \setminus (\Theta_\gamma)_J;$$

- if α is a limit ordinal, we define

$$\Theta_\alpha := \bigcap_{\beta < \alpha} \Theta_\beta.$$

In both cases, we define

$$T_\alpha := \bigcap \{S \mid S \in \Theta_\alpha\}$$

with the convention that $T_\alpha := K$ if $\Theta_\alpha = \emptyset$.

Proposition 6.1. *Preserve the notation above. Then:*

- Θ_α is a pre-Jaffard family of T_α ;
- if $\alpha = \gamma + 1$ is a successor ordinal, then $\Theta_\alpha \cup \{T_\alpha\}$ is a weak Jaffard family of T_γ pointed at T_α .

Proof. We proceed by transfinite induction: the claim is true by hypothesis for $\alpha = 0$. Suppose that it holds for every $\beta < \alpha$. If $\alpha = \gamma + 1$ is a successor ordinal, then the two statements are exactly Proposition 5.6. Suppose thus that α is a limit ordinal.

Each $T \in \Theta_\alpha$ is flat over D and thus over T_α ; furthermore, Θ_α is independent since it is contained in the independent set Θ . Since also every Θ_β (for $\beta < \alpha$) is independent, as in the proof of Proposition 5.6 we have

$$\Theta_\alpha^\uparrow = \left(\bigcap_{\beta < \alpha} \Theta_\beta \right)^\uparrow = \bigcap_{\beta < \alpha} \Theta_\beta^\uparrow$$

which is closed in the inverse topology since each Θ_β is compact with respect to the Zariski topology (being a pre-Jaffard family by inductive

hypothesis), and so also Θ_α , which is the set of minimal elements of Θ_α^\uparrow , is compact in the Zariski topology. Thus, we only need to show that Θ_α is complete. Let P be a nonzero prime ideal of T_α , and suppose that $PS = S$ for some $S \in \Theta_\alpha$: then, by Lemma 3.11, $SD_P = K$. Therefore, if $PS = S$ for all $S \in \Theta_\alpha$ then, by the flatness of D_P and the compactness of Θ_α , by [2, Corollary 5] we have

$$D_P = D_P T_\alpha = D_P \bigcap_{S \in \Theta_\alpha} S = \bigcap_{S \in \Theta_\alpha} D_P S = K,$$

a contradiction. Hence Θ_α is complete and thus it is a pre-Jaffard family. \square

Definition 6.2. *We call the family $\{T_\alpha\}$ defined in this way the derived sequence with respect to Θ .*

By construction, the derived sequence of Θ is an ascending chain of rings:

$$D = T_0 \subseteq T_1 \subseteq T_2 \subseteq \cdots \subseteq T_\omega \subseteq \cdots,$$

which corresponds to a descending chain of sets of overrings:

$$\Theta = \Theta_0 \supseteq \Theta_1 \supseteq \Theta_2 \supseteq \cdots \supseteq \Theta_\omega \supseteq \cdots.$$

Proposition 6.3. *Preserve the notation above. There is an ordinal α such that $T_\alpha = T_{\alpha'}$ for all $\alpha' > \alpha$ (equivalently, such that $\Theta_\alpha = \Theta_{\alpha'}$ for all $\alpha' > \alpha$).*

Proof. Note that, if $T_\alpha = T_{\alpha+1}$, then $T_\alpha = T_{\alpha'}$ for all $\alpha' > \alpha$; thus, suppose by contradiction that $T_\alpha \subsetneq T_{\alpha+1}$ for all α . Then, $T_{\alpha+1} \setminus T_\alpha$ is nonempty for all α ; but since all the T_α are contained inside K , this is impossible if the cardinality of α is larger than the cardinality of K . \square

Definition 6.4. *We call the minimal ordinal α such that $T_\alpha = T_{\alpha'}$ for all $\alpha' > \alpha$ the Jaffard degree of the family Θ , and we call T_α the dull limit of Θ . We say that Θ is:*

- a sharp family if $T_\alpha = K$;
- a dull family if $T_\alpha \neq K$.

Equivalently, Θ is a sharp family if $\Theta_\alpha = \emptyset$ for some α , while it is a dull family otherwise.

The terminology sharp/dull family is chosen in analogy with [12] and [16], where sharp and dull domains (and, in correspondence, sharp and dull degrees) are defined, respectively, for almost Dedekind domains and for one-dimensional Prüfer domains; our definition can be seen as a wide generalization of their concept. However, we do not use the terminology “sharp degree” and “dull degree”, both because the definition of Jaffard degree unifies them and because there is actually a small difference in the sharp case. See Section 8 for a more detailed discussion.

Example 6.5. Let Θ be a Jaffard family of D , with $D \neq K$. Then, $\Theta = \Theta_J$, and so $\Theta_1 = \emptyset$; thus, $T_1 = K = T_\alpha$ for all ordinals $\alpha > 0$. Hence, Θ is sharp with Jaffard degree 1.

Example 6.6. Let Θ be a weak Jaffard family of D pointed at S . Then, $\Theta_1 = \{S\}$, and so $T_1 = S$; on the other hand, $\Theta_2 = \emptyset$ and so $T_2 = K$. Thus Θ is sharp with Jaffard degree 2.

Example 6.7. Let D be an almost Dedekind domain with only finitely many maximal ideals that are not finitely generated, say M_1, \dots, M_n . (Those rings do indeed exist: see [11].) Let $\Theta := \{D_M \mid M \in \text{Max}(D)\}$: then, Θ is a pre-Jaffard family of D (see Proposition 8.1 below).

If P is a maximal ideal of D different from the M_i , then D_P is a Jaffard overring of D , since $\text{Max}(D) \setminus \{P\}$ is compact and thus $D_P D_P^\perp = K$. On the other hand, each D_{M_i} is not a Jaffard overring, since $D_Q^\perp = \bigcap \{D_Q \mid Q \in \text{Max}(D) \setminus \{M_i\}\} = D$; in particular, there is no weak Jaffard family of D that can contain at the same time every D_{M_i} . Furthermore, the family $\Theta' := \{D_M \mid M \in \text{Max}(D), M \neq M_1, \dots, M_n\}$ is strongly independent (since every D_M is a Jaffard overring and D_M^\perp contains the intersection of all $T \in \Theta' \setminus \{D_M\}$), but it is not locally finite, since otherwise the whole $\Theta = \Theta' \cup \{D_{M_1}, \dots, D_{M_n}\}$ would be locally finite and thus a Jaffard family.

The set Θ_1 is equal to $\{M_1, \dots, M_n\}$ and thus it is finite; moreover, $T_1 = D_{M_1} \cap \dots \cap D_{M_n}$ is a semilocal almost Dedekind domain, and thus a PID. Therefore, $\Theta_2 = \emptyset$ and $T_2 = K$, so that Θ is sharp with Jaffard degree 2.

Example 6.8. Let D be the ring of all algebraic integers, i.e., the integral closure of \mathbb{Z} in $\overline{\mathbb{Q}}$. Then, D is a one-dimensional Bézout (in particular, Prüfer) domain such that none of its maximal ideals are finitely generated, nor any nonzero primary ideal is finitely generated.

Therefore, $\Theta := \{D_M \mid M \in \text{Max}(D)\}$ is a pre-Jaffard family of D (as in the previous example), but none of its elements are Jaffard overrings: hence $\Theta_1 = \emptyset$ and $T_1 = D = T_0$. Therefore, Θ is dull with Jaffard degree 0 and its dull limit is D itself.

Let now $\Theta' := \{D_M \mid M \in \text{Max}(D), 2 \in M\} \cup \{D[1/2]\}$. Then, Θ' is obtained from Θ with the construction of Proposition 4.8 applied to $\Theta_1 := \{S \in \Theta \mid 1/2 \in S\} = \mathcal{B}(1/2) \cap \Theta$ (which is compact), and thus is a pre-Jaffard family. The ring $D[1/2]$ is a Jaffard overring of D , since it belongs to the Jaffard family $\{D[1/p] \mid p \text{ is a prime number}\}$, while no other element of Θ' is a Jaffard overring; hence, $(\Theta')_1 = \{D_M \mid M \in \text{Max}(D), 2 \in M\}$, while $(\Theta')_2 = (\Theta')_1$. Hence Θ' is dull with Jaffard degree 1, and its dull limit is

$$T_1 = \bigcap_{\substack{M \in \text{Max}(D) \\ 2 \in M}} D_M = D[1/3, 1/5, \dots] = D[1/p \mid p \neq 2 \text{ is a prime number}].$$

Remark 6.9. Note that, if D is not a field and Θ is sharp, then the Jaffard degree of D cannot be 0.

7. STABLE OPERATIONS

Let T be an overring of D . Then, $\mathbf{F}_D(K) \subseteq \mathbf{F}_T(K)$, and the image of any T -submodule of K by any semistar operation on D is still a T -module. Then, we have two ways to relate the semistar operations on D and T : the first one is the restriction map

$$\begin{aligned} \psi_T: \text{SStar}(D) &\longrightarrow \text{SStar}(T), \\ \star &\longmapsto \star|_{\mathbf{F}_T(K)}, \end{aligned}$$

while the second is the extension map

$$\begin{aligned} \phi_T: \text{SStar}(T) &\longrightarrow \text{SStar}(D), \\ \star &\longmapsto \phi_T(\star): \begin{array}{l} \mathbf{F}_D(K) \longrightarrow \mathbf{F}_D(K), \\ I \longmapsto (IT)^\star. \end{array} \end{aligned}$$

If now we have a family Θ of overrings of D , then we can put together the maps relative to each member of the family: we obtain a restriction map

$$\begin{aligned} \Psi_\Theta: \text{SStar}(D) &\longrightarrow \prod_{T \in \Theta} \text{SStar}(T), \\ \star &\longmapsto (\Psi_T(\star)) \end{aligned}$$

and an extension map

$$\begin{aligned} \Phi_\Theta: \prod_{T \in \Theta} \text{SStar}_{\text{st}}(T) &\longrightarrow \text{SStar}_{\text{st}}(D), \\ (\star^{(T)})_{T \in \Theta} &\longmapsto \inf_{T \in \Theta} \Phi_T(\star^{(T)}). \end{aligned}$$

All these maps are order-preserving when $\text{SStar}(D)$ and $\text{SStar}(T)$ are endowed with the natural order, and when the product is endowed with the product order.

Proposition 7.1. *Let Θ be a complete and independent family of overrings of D . Then, $\Psi_\Theta \circ \Phi_\Theta$ is the identity on $\prod_{T \in \Theta} \text{SStar}(T)$.*

Proof. For every T , let $\star^{(T)} \in \text{SStar}(T)$, and let $\star := \Phi_\Theta((\star^{(T)})_{T \in \Theta})$. Fix $S \in \Theta$ and let $I \in \mathbf{F}_S(D)$. Then, $I = IS \neq (0)$; since I is complete, we have

$$I^{\star^S} = I^\star = \bigcap_{T \in \Theta} (IT)^{\star^{(T)}} = (IS)^{\star^{(S)}} \cap \bigcap_{T \in \Theta \setminus \{S\}} (IST)^{\star^{(T)}} = I^{\star^{(S)}}$$

as $ST = K$ for every $T \in \Theta \setminus \{S\}$ (since Θ is independent). The claim is proved. \square

Definition 7.2. Let Θ be a family of overrings of D . We say that Θ is stable-preserving if, for every $\star \in \text{SStar}_{\text{st}}(D)$ and every $I \in \mathbf{F}_D(K)$, we have

$$I^\star = \bigcap_{T \in \Theta} (IT)^\star.$$

A stable semistar operation is uniquely determined by its action on proper ideals of D . Hence, if \star is a stable semistar operation fixing D , then the notion of extension of a star operation studied in [5] can be used to show that if Θ is a Jaffard family then $I^\star = \bigcap_{T \in \Theta} (IT)^\star$ (see, in particular, [5, Theorems 5.4 and 5.6]); a similar result, without the hypothesis $D = D^\star$, can be shown joining the results in Sections 3 and 6 of [7] (passing through length functions), so that any Jaffard family is stable-preserving. We want to generalize this case, but we first point out why stable-preserving properties are useful.

Proposition 7.3. Let Θ be a stable-preserving family of flat overrings of D . Then, Ψ_Θ and Φ_Θ establish an order-preserving isomorphism between $\text{SStar}_{\text{st}}(D)$ and $\prod \{\text{SStar}_{\text{st}}(T) \mid T \in \Theta\}$.

Proof. If \star is a stable semistar operation, then the restriction $\Psi_T(\star)$ is stable for every overring T ; conversely, the infimum of a family of restriction of stable operations is still stable, since

$$\begin{aligned} (I \cap J)^{\Phi_\Theta(\star^{(T)})} &= \bigcap_{T \in \Theta} ((I \cap J)T)^{\star^{(T)}} = \bigcap_{T \in \Theta} ((IT \cap JT))^{\star^{(T)}} = \\ &= \bigcap_{T \in \Theta} (IT)^{\star^{(T)}} \cap (JT)^{\star^{(T)}} = I^{\Phi_\Theta(\star^{(T)})} \cap J^{\Phi_\Theta(\star^{(T)})}, \end{aligned}$$

using the flatness of the members of Θ . Hence, the maps Φ_Θ and Ψ_Θ restrict to maps from $\text{SStar}_{\text{st}}(D)$ to $\prod_{T \in \Theta} \text{SStar}_{\text{st}}(T)$.

By Proposition 7.1, $\Psi_\Theta \circ \Phi_\Theta$ is the identity. Let now $\star \in \text{SStar}_{\text{st}}(D)$. Then,

$$I^{\Phi_\Theta \circ \Psi_\Theta(\star)} = \bigcap_{T \in \Theta} (IT)^\star = I^\star$$

since Θ is stable-preserving. Hence, $\Phi_\Theta \circ \Psi_\Theta$ is the identity of $\text{SStar}_{\text{st}}(D)$, and thus Φ_Θ and Ψ_Θ are isomorphism. \square

Proposition 7.4. A weak Jaffard family is stable-preserving.

Proof. Let Θ be a weak Jaffard family pointed at T_∞ . Fix any $\star \in \text{SStar}_{\text{st}}(D)$, and let \sharp be the map $I \mapsto \bigcap_{T \in \Theta} (IT)^\star$. Then, \sharp is stable, and $\star \leq \sharp$; in particular, if $1 \in I^\star$ then $1 \in I^\sharp$.

Conversely, let $I \subseteq D$ be such that $1 \in I^\sharp$; without loss of generality, we can suppose that $I = I^\star$. Let $T \in \Theta \setminus \{T_\infty\}$: then, then, T is a Jaffard overring of D , and thus $\{T, T^\perp\}$ is a Jaffard family of D by Proposition 3.9. Hence,

$$IT = I^\star T = (IT \cap IT^\perp)^\star T = ((IT)^\star \cap (IT^\perp)^\star) T = (IT)^\star \cap (IT^\perp)^\star T.$$

By definition, $I^\sharp \subseteq (IT)^\star$, and thus $1 \in (IT)^\star$; on the other hand, $TT^\perp = K$ and thus $(IT^\perp)^\star T = K$. Thus, $1 \in IT$ and $IT = T$.

Since Θ is complete, we thus have $I = (IT_\infty \cap D)$, and so

$$IT_\infty = I^\star T_\infty = (IT_\infty \cap D)^\star T_\infty = (IT_\infty)^\star \cap D^\star T_\infty.$$

Again, by construction 1 belongs to both $(IT_\infty)^\star$ and $D^\star T_\infty$, and thus $1 \in IT_\infty$, so that $IT_\infty = T_\infty$. Hence, $IT = T$ for every $T \in \Theta$, and thus $I = D$. Therefore, for every $I \subseteq D$ we have $1 \in I^\star$ if and only if $1 \in I^\sharp$; since \star and \sharp are stable, it follows that $\star = \sharp$, as claimed. Thus, Θ is stable-preserving, as claimed. \square

Theorem 7.5. *Let Θ be a Jaffard family, α an ordinal, and let $\Theta' := (\Theta \setminus \Theta_\alpha) \cup \{T_\alpha\}$. Then, Θ' is stable-preserving.*

Proof. For every $\beta \leq \alpha$, let $\Lambda_\beta := (\Theta \setminus \Theta_\beta) \cup \{T_\beta\}$: we want to show by induction that Λ_β is stable-preserving. Note that each Λ_β is complete and, by definition, $\Lambda_\alpha = \Theta'$.

If $\beta = 0$ then $\Lambda_0 = (\Theta \setminus \Theta) \cup \{T_0\} = \{T_0\} = \{D\}$ is clearly stable-preserving; suppose thus $\beta > 0$, and suppose that the claim holds for every $\gamma < \beta$; we distinguish two cases.

If $\beta = \gamma + 1$ is a successor ordinal, then $\Theta_\beta = \Theta_\gamma \setminus (\Theta_\gamma)_J$ and thus

$$\Lambda_\beta = (\Theta \setminus (\Theta_\gamma \setminus (\Theta_\gamma)_J)) \cup \{T_\beta\} = (\Theta \setminus \Theta_\gamma) \cup (\Theta_\gamma)_J \cup \{T_\beta\}.$$

Let $\Lambda' := \Theta \setminus \Theta_\gamma$, and take a stable semistar operation on D . By inductive hypothesis, $\Lambda_\gamma = \Lambda' \cup \{T_\gamma\}$ is stable-preserving, and thus

$$I^\star = \bigcap_{A \in \Lambda'} (IA)^\star \cap (IT_\gamma)^\star.$$

Moreover, by construction, $(\Theta_\gamma)_J \cup \{T_\beta\}$ is a weak Jaffard family of T_γ pointed at T_β ; by Proposition 7.4, it is stable-preserving on T_β , and thus

$$(IT_\gamma)^\star = \bigcap_{A \in \Theta_\gamma \cup \{T_\beta\}} (IT_\gamma A)^\star = \bigcap_{A \in \Theta_\gamma \cup \{T_\beta\}} (IA)^\star,$$

so that

$$I^\star = \bigcap_{A \in \Lambda'} (IA)^\star \cap \bigcap_{A \in \Theta_\gamma \cup \{T_\beta\}} (IA)^\star = \bigcap_{A \in \Lambda_\beta} (IA)^\star.$$

Hence, Λ_β is stable-preserving.

Suppose now that β is a limit ordinal: then, $\Theta_\beta = \bigcap_{\gamma < \beta} \Theta_\gamma$, and thus

$$\Lambda_\beta = \left(\Theta \setminus \bigcap_{\gamma < \beta} \Theta_\gamma \right) \cup \{T_\beta\} = \bigcup_{\gamma < \beta} (\Theta \setminus \Theta_\gamma) \cup \{T_\beta\}.$$

Let \star be a stable semistar operation, and let \sharp be the map

$$\sharp : I \mapsto \bigcap_{A \in \Lambda_\beta} (IA)^\star = \left[\bigcap_{\gamma < \beta} \bigcap_{A \in \Theta_\gamma} (IA)^\star \right] \cap (IT_\beta)^\star.$$

Then, \sharp is a stable semistar operation, and $I^\star \subseteq I^\sharp$ for all ideals I (as I^\star is contained in all $(IA)^\star$ and in $(IT_\beta)^\star$). We claim that it is equal to \star , and to do so it is enough to show that if $1 \in I^\sharp$ then also $1 \in I^\star$, where I is a proper ideal of D (this follows from condition (4) of [3, Theorem 2.6]).

Take thus a proper ideal I such that $1 \in I^\sharp$, and let $\Gamma := \{\gamma < \beta \mid IT \neq T \text{ for some } T \in \Theta_\gamma \setminus \Theta_{\gamma+1}\}$.

Suppose that Γ is nonempty: then, it has a minimum γ . Since Λ_γ is complete and $IS = S$ for all $S \in \Theta_\delta$ with $\delta < \gamma$, we must have $I = IT_\gamma \cap D$; as above, it follows that $I^\star = (IT_\gamma)^\star \cap D^\star$. Let $T \in \Theta_\gamma \setminus \Theta_{\gamma+1}$: then, T is a Jaffard overring of T_γ . Let

$$A := \bigcap \{(T_\gamma)_P \mid P \in \text{Spec}(T_\gamma), PT = T\},$$

that is, $A = T^\perp$ with respect to T_γ . Then, $TA = K$ and $J = JT \cap JA$ for all T_γ -submodules J of K . Thus,

$$(IT_\gamma)^\star T = (IT_\gamma T \cap IT_\gamma A)^\star T = (IT)^\star \cap ((IA)^\star) T = (IT)^\star.$$

Therefore,

$$I^\star T = ((IT_\gamma)^\star \cap D^\star) T = (IT)^\star \cap D^\star T$$

contains 1 since it contains I^\sharp . Since T was arbitrary in $\Theta_\gamma \setminus \Theta_{\gamma+1}$, this is a contradiction.

Therefore, Γ must be empty. Since β is a limit ordinal, Λ_β is also equal to $\bigcup_{\gamma < \beta} (\Theta_\gamma \setminus \Theta_{\gamma+1})$; therefore, since Λ_β is complete and I is proper, we must have $I = IT_\beta \cap D$; therefore, $I^\star = (IT_\beta \cap D)^\star = (IT_\beta)^\star \cap D^\star$ since \star is stable. However, $1 \in (IT_\beta)^\star$ since $(IT_\beta)^\star$ contains I^\sharp , while obviously $1 \in D^\star$; hence, $1 \in I^\star$.

By induction, it follows that $\Lambda_\alpha = \Theta'$ is stable-preserving, as claimed. \square

Corollary 7.6. *Let Θ be a Jaffard family, α an ordinal, and let $\Theta' := (\Theta \setminus \Theta_\alpha) \cup \{T_\alpha\}$. Then:*

- (1) *for every stable semistar operation \star on D , we have $I^\star = \bigcap \{(IT)^\star \mid T \in \Theta'\}$;*
- (2) $\text{SStar}_{\text{st}}(D) \simeq \prod \{\text{SStar}_{\text{st}}(T) \mid T \in \Theta'\}$.

Proof. The first part follows by joining Theorem 7.5 with Definition 7.2, the second part from Theorem 7.5 and Proposition 7.3. \square

From the correspondence between stable semistar operations and length functions we have the following.

Proposition 7.7. *Let D be an integral domain and let Θ be a stable-preserving family of flat overrings of D . Then, for every $\ell \in \mathcal{L}_{\text{sing}}(D)$, we have*

$$\ell = \sum_{T \in \Theta} \ell \otimes T.$$

In particular, this holds for Jaffard families, as was proved in [7, Theorem 3.10]; likewise, the analogue of Corollary 7.6 holds.

Corollary 7.8. *Let Θ be a Jaffard family, α an ordinal, and let $\Theta' := (\Theta \setminus \Theta_\alpha) \cup \{T_\alpha\}$. Then:*

- (1) *for every length function ℓ on D , we have $\ell = \sum \{\ell \otimes T \mid T \in \Theta'\}$;*
- (2) $\mathcal{L}_{\text{sing}}(D) \simeq \prod \{\mathcal{L}_{\text{sing}}(T) \mid T \in \Theta'\}$.

Obviously, the previous results are at their strongest when α is the Jaffard degree of Θ , so that T_α is the dull limit of Θ .

8. THE DIMENSION 1 CASE

In this section, we specialize the results of the previous sections to domains of dimension 1. In this case, there is a natural pre-Jaffard family to consider.

Proposition 8.1. *Let D be a domain of dimension 1, and let $\Theta := \{D_M \mid M \in \text{Max}(D)\}$. Then, Θ is a pre-Jaffard family of D .*

Proof. The family Θ is clearly complete, independent (no nonzero prime survives in D_M and in D_N for $M \neq N$) and composed of flat overrings. Furthermore, the localization map $\lambda : \text{Spec}(D) \rightarrow \text{Over}(D)$ is a homeomorphism between $\text{Spec}(D)$ and $\lambda(\text{Spec}(D))$ when the spaces are endowed with the respective Zariski topologies [9, Lemma 2.4]; in particular, $\Theta = \lambda(\text{Max}(D))$ is compact. Hence, Θ is a pre-Jaffard family. \square

Definition 8.2. *Let D be a one-dimensional integral domain, and let $\Theta := \{D_M \mid M \in \text{Max}(D)\}$. We say that D is:*

- ultimately sharp if Θ is sharp;
- ultimately dull if Θ is dull.

The second reason why the dimension 1 hypothesis is powerful is that we can improve Proposition 4.5.

Proposition 8.3. *Let D be a domain of dimension 1, and let $\Theta := \{D_M \mid M \in \text{Max}(D)\}$. Let $M \in \text{Max}(D)$. Then, the following are equivalent:*

- (i) D_M is a Jaffard overring of D ;
- (ii) $\Theta \setminus \{D_M\}$ is compact, with respect to the Zariski topology;
- (iii) $\text{Max}(D) \setminus \{M\}$ is compact, with respect to the Zariski topology;
- (iv) M is isolated in $\text{Max}(D)$, with respect to the inverse topology.

Proof. The equivalence of (i) and (ii) follows from Proposition 4.5 (and Proposition 8.1), while the equivalence of (ii) and (iii) from the fact that $\Theta \setminus \{D_M\} \simeq \text{Max}(D) \setminus \{M\}$ via the localization map. Again by Proposition 4.5, (i) implies (iv).

Suppose (iv) holds. Then, $\text{Max}(D) \setminus \{M\}$ is closed in $\text{Max}(D)$, with respect to the inverse topology. Since D has dimension 1, it follows that $\text{Spec}(D) \setminus \{M\}$ is closed in the inverse topology, and thus that $\text{Max}(D) \setminus \{M\}$ is compact. Hence, (iii) holds and all the conditions are equivalent. \square

Recall that, for a topological space X , $\mathcal{I}(X)$ and $\mathcal{D}(X)$ are, respectively, the set of isolated points and the set of limit points of X . Proposition 8.3 allows to describe the derived sequence in a purely topological way.

Theorem 8.4. *Let D be a one-dimensional domain, let $\Theta := \{D_M \mid M \in \text{Max}(D)\}$ and let $X := \text{Max}(D)^{\text{inv}}$. Let $\{T_\alpha\}$ be the derived sequence of Θ and let $\{\Theta_\alpha\}$ be the corresponding chain of subsets of Θ . Then, for every ordinal α and every $M \in \text{Max}(D)$, we have $MT_\alpha \neq T_\alpha$ if and only if $M \in \mathcal{D}^\alpha(X)$, and $\Theta_\alpha = \{D_M \mid M \in \mathcal{D}^\alpha(X)\}$.*

Proof. Let $\Lambda_\alpha := \{M \in \text{Max}(D) \mid MT_\alpha \neq T_\alpha\}$; then, $\{\Lambda_\alpha\}$ is a descending chain of subsets of $\text{Max}(D)$, and we need to show that $\Lambda_\alpha = \mathcal{D}^\alpha(X)$. By definition, Θ_α is a pre-Jaffard family of T_α ; it follows that $M \in \Lambda_\alpha$ if and only if $D_M \in \Theta_\alpha$.

We proceed by transfinite induction. For $\alpha = 0$, $\Theta_0 = \text{Max}(D)$ and $\mathcal{D}^0(X) = X$, so the claim is proved. Suppose that the claim holds for every $\beta < \alpha$; we distinguish two cases.

Suppose first that $\alpha = \gamma + 1$ is a successor ordinal. Then, by hypothesis, $\{M \in \text{Max}(D) \mid MT_\gamma \neq T_\gamma\} = \mathcal{D}^\gamma(X)$; hence, the restriction map $\rho : \text{Max}(T_\gamma) \rightarrow \text{Max}(D)$, $P \mapsto P \cap D$ establishes a homeomorphism between $\text{Max}(T_\gamma)$ and its image $\mathcal{D}^\gamma(X) = \Lambda_\gamma(X)$ both in the Zariski and in the inverse topology. By definition and by Proposition 8.3, $\Theta_\alpha = \Theta_{\gamma+1} = \{(T_\gamma)_P \mid P \in (\text{Max}(T_\gamma))^{\text{inv}}\}$, i.e., given a maximal ideal M of D , we have $D_M \in \Theta_\alpha$ if and only if MT_γ is a limit point of $(\text{Max}(T_\gamma))^{\text{inv}}$. Since ρ is a homeomorphism in the inverse topology, this is equivalent to saying that $M \in \mathcal{D}^{\gamma+1}(X) = \mathcal{D}^\alpha(X)$; that is, $D_M \in \Theta_\alpha$ if and only if $M \in \mathcal{D}^\alpha(X)$. By the remark at the beginning of the proof, we have our claim.

Suppose now that α is a limit ordinal. If $M \in \Lambda_\alpha$, then $M \in \Lambda_\beta$ for all $\beta < \alpha$, and thus by induction $M \in \mathcal{D}^\beta(X)$ for all $\beta < \alpha$; by definition, this is exactly the condition $M \in \mathcal{D}^\alpha(X)$. Conversely, if $M \in \mathcal{D}^\alpha(X)$ then $M \in \mathcal{D}^\beta(X)$ for all $\beta < \alpha$, and thus by induction $D_M \in \Theta_\beta$ for all $\beta < \alpha$; therefore, $D_M \in \Theta_\alpha$ by definition and $M \in \Lambda_\alpha$. Thus $M \in \Lambda_\alpha$ if and only if $M \in \mathcal{D}^\alpha(X)$, and the claim is proved. \square

Corollary 8.5. *Let D be a one-dimensional integral domain. Then, the Jaffard degree of Θ is equal to the Cantor-Bendixson rank of $\text{Max}(D)^{\text{inv}}$.*

Proof. By definition, the Cantor-Bendixson rank of X is the least ordinal α such that $\mathcal{D}^\alpha(X) = \mathcal{D}^{\alpha+1}(X)$. By Theorem 8.4, when $X =$

$\text{Max}(D)^{\text{inv}}$ the latter condition is equivalent to $\Theta_\alpha = \Theta_{\alpha+1}$, and thus α is also the Jaffard degree of Θ . \square

Corollary 8.6. *Let D be a one-dimensional integral domain, and let $X := \text{Max}(D)^{\text{inv}}$. Then, D is ultimately sharp if and only if X is a scattered space.*

Proof. By Theorem 8.4, $\mathcal{D}^\alpha(X)$ becomes empty if and only if there is an α such that $MT_\alpha = T_\alpha$ for all maximal ideal M , where T_α is the α -th element of the derived sequence of Θ . However, the latter condition is equivalent to $T_\alpha = K$, i.e., to the fact that D is ultimately sharp. \square

When D is a Prüfer domain, a similar construction has been given in [16, Section 6], following ideas introduced in [12]. In this case, a maximal ideal M is said to be *sharp* if $\bigcap \{D_N \mid N \in \text{Max}(D), N \neq M\} \not\subseteq D_M$, while *dull* otherwise; by [16, Lemma 6.3(2)] and Proposition 8.3 (or by direct proof) we have that M is sharp if and only if D_M is a Jaffard overring of D . If $\text{Max}_\sharp(D)$ is the set of sharp maximal ideals of D , they define recursively an ascending sequence of rings by

$$D_\alpha := \begin{cases} \bigcap \{(D_\gamma)_M \mid M \in \text{Max}_\sharp(D_\gamma)\} & \text{if } \alpha = \gamma + 1 \text{ is a successor ordinal,} \\ \bigcup \{D_\gamma \mid \gamma < \alpha\} & \text{if } \alpha \text{ is a limit ordinal,} \end{cases}$$

and they show [16, Lemma 6.5(2)] that $\text{Max}(D_\alpha) = \{MD_\alpha \mid M \in \mathcal{D}^\alpha(\text{Max}(D)^{\text{inv}})\}$, so that D_α actually coincides with our T_α , the α -th element of the derived sequence of $\Theta := \{D_M \mid M \in \text{Max}(D)\}$ (this also, *a fortiori*, for limit ordinals α , for which the definitions of T_α and D_α do not coincide). Then, they say that D has *sharp degree* α if $D_\alpha \neq K$ while $D_{\alpha+1} = K$, and that D has *dull degree* α if $D_\alpha = D_{\alpha+1} \neq K$ and $D_\beta \neq D_\alpha$ for all $\beta < \alpha$.

In the case of dull degree, our definition agrees with theirs: it is straightforward to see (using $D_\alpha = T_\alpha$) that D has a dull degree if and only if D is ultimately dull, and that the dull degree of D coincides with the Jaffard degree of Θ .

On the other hand, for sharp degree, there is a difference: indeed, if D has sharp degree α then Θ has Jaffard degree $\alpha + 1$, and conversely if the Jaffard degree of Θ is a successor ordinal $\alpha + 1$ then D has sharp degree α . However, if the Jaffard degree of Θ is a limit ordinal α , then the sharp degree of D does not exist, because the definition requires that the first β such that $D_\beta = K$ is a successor ordinal. Thus, D has a sharp degree if and only if D is ultimately sharp and the Jaffard degree of Θ is a successor ordinal.

Our approach allows to give a simpler form to some of their results. Indeed, Corollary 8.6 above is a more symmetric version of [16, Theorem 6.6], because it gives a complete equivalence between a topological fact ($\text{Max}(D)^{\text{inv}}$ is scattered) and the sharpness of D , without requiring that the Jacobson radical of D is nonzero (as in part (2) of the

reference) and we do not need to separate the dull and the sharp case (as in parts (1) and (3)).

To conclude the paper, we apply the results of Section 7 to one-dimensional domain.

Proposition 8.7. *Let D be a one-dimensional domain. Let T_α be the dull limit of $\Theta := \{D_M \mid M \in \text{Max}(D)\}$. Then, the family*

$$\Theta' := \{D_M \mid M \in \text{Max}(D), MT_\infty = T_\alpha\} \cup \{T_\alpha\}$$

is stable-preserving.

Proof. The claim is a direct consequence of Theorem 7.5. \square

Proposition 8.8. *Let D be a one-dimensional Prüfer domain. Then, the family $\Theta := \{D_M \mid M \in \text{Max}(D)\}$ is stable-preserving if and only if D is ultimately sharp.*

Proof. If D is ultimately sharp, then its dull limit T_α is equal to K , and thus the family $\Theta' = \{D_M \mid M \in \text{Max}(D), MT_\infty = T_\alpha\} \cup \{T_\alpha\}$ of Θ coincides with $\Theta \cup \{K\}$. Hence, $\Theta \cup \{K\}$ is stable-preserving and so is Θ .

Suppose D is ultimately dull, and let $T := T_\alpha$ be its dull limit. Consider the set $\Lambda := \{N \mid NT \neq T\}$. For every $N \in \Lambda$, $D_N = T_{NT}$ is not a Jaffard overring of T (by construction of T_α), and thus $\text{Max}(T) \setminus \{NT\}$ is not compact; hence, $\bigcap_{N' \neq N} T_{N'} = T$. For every $N \in \Lambda$, let \star_N be the stable semistar operation

$$\star_N : I \mapsto \bigcap_{\substack{N' \in \Lambda \\ N' \neq N}} ID_{N'}.$$

Then, \star_N fixes T , but $I^{\star_N} = K$ for all I that are D_N -modules or D_M -modules for some $M \notin \Lambda$. Let \star be the supremum of all the \star_N ; then, \star fixes T . However, $(TD_M)^{\star_N} = K$ for every $M \in \text{Max}(D)$, and thus

$$K = \bigcap_{M \in \text{Max}(D)} (TD_M)^{\star_N} \neq T^{\star_N}.$$

Hence, Θ is not stable-preserving. \square

Corollary 8.9. *Let D be an almost Dedekind domain that is ultimately sharp. Then, there is a natural bijection between $\text{SStar}_{\text{st}}(D)$ and the power set of $\text{Max}(D)$.*

Proof. By Propositions 7.3 and 8.8, $\text{SStar}_{\text{st}}(D)$ is isomorphic to the product $\prod \{\text{SStar}_{\text{st}}(D_M) \mid M \in \text{Max}(D)\}$. However, each D_M is a discrete valuation ring, and thus $\text{SStar}_{\text{st}}(D_M)$ contains exactly two operations (the identity and the one sending everything to K). The claim follows. \square

REFERENCES

- [1] Carmelo A. Finocchiaro. Spectral spaces and ultrafilters. *Comm. Algebra*, 42(4):1496–1508, 2014.
- [2] Carmelo A. Finocchiaro and Dario Spirito. Topology, intersections and flat modules. *Proc. Amer. Math. Soc.*, 144(10):4125–4133, 2016.
- [3] D. D. Anderson and Sylvia J. Cook. Two star-operations and their induced lattices. *Comm. Algebra*, 28(5):2461–2475, 2000.
- [4] Daniel Lazard. Autour de la platitude. *Bull. Soc. Math. France*, 97:81–128, 1969.
- [5] Dario Spirito. Jaffard families and localizations of star operations. *J. Commut. Algebra*, 11(2):265–300, 2019.
- [6] Dario Spirito. Topological properties of localizations, flat overrings and sublocalizations. *J. Pure Appl. Algebra*, 223(3):1322–1336, 2019.
- [7] Dario Spirito. Decomposition and classification of length functions. *Forum Math.*, 32(5):1109–1129, 2020.
- [8] Dario Spirito. The sets of star and semistar operations on semilocal Prüfer domains. *J. Commut. Algebra*, 12(4), 2020.
- [9] David E. Dobbs, Richard Fedder, and Marco Fontana. Abstract Riemann surfaces of integral domains and spectral spaces. *Ann. Mat. Pura Appl. (4)*, 148:101–115, 1987.
- [10] Fred Richman. Generalized quotient rings. *Proc. Amer. Math. Soc.*, 16:794–799, 1965.
- [11] K. Alan Loper. Sequence domains and integer-valued polynomials. *J. Pure Appl. Algebra*, 119(2):185–210, 1997.
- [12] K. Alan Loper and Thomas G. Lucas. Factoring ideals in almost Dedekind domains. *J. Reine Angew. Math.*, 565:61–78, 2003.
- [13] Marco Fontana, Evan Houston, and Thomas Lucas. *Factoring Ideals in Integral Domains*, volume 14 of *Lecture Notes of the Unione Matematica Italiana*. Springer, Heidelberg; UMI, Bologna, 2013.
- [14] Max Dickmann, Niels Schwartz, and Marcus Tressl. *Spectral spaces*, volume 35 of *New Mathematical Monographs*. Cambridge University Press, Cambridge, 2019.
- [15] Melvin Hochster. Prime ideal structure in commutative rings. *Trans. Amer. Math. Soc.*, 142:43–60, 1969.
- [16] Olivier A. Heubo-Kwegna, Bruce Olberding, and Andreas Reinhart. Group-theoretic and topological invariants of completely integrally closed Prüfer domains. *J. Pure Appl. Algebra*, 220(12):3927–3947, 2016.
- [17] Robert W. Gilmer, Jr. and William J. Heinzer. Overrings of Prüfer domains. II. *J. Algebra*, 7:281–302, 1967.
- [18] William Heinzer and Moshe Roitman. Well-centered overrings of an integral domain. *J. Algebra*, 272(2):435–455, 2004.

DIPARTIMENTO DI MATEMATICA “TULLIO LEVI-CIVITA”, UNIVERSITÀ DEGLI STUDI DI PADOVA, PADOVA, ITALY

Email address: `spirito@math.unipd.it`