

Treatment with Olanzapine is Associated with Modulation of the Default Mode Network in Patients with Schizophrenia

Fabio Sambataro^{1,3}, Giuseppe Blasi¹, Leonardo Fazio¹, Grazia Caforio¹, Paolo Taurisano¹, Raffaella Romano¹, Annabella Di Giorgio¹, Barbara Gelao¹, Luciana Lo Bianco¹, Apostolos Papazacharias¹, Teresa Popolizio², Marcello Nardini¹ and Alessandro Bertolino^{*,1,2}

¹Psychiatric Neuroscience Group, Department of Neurological and Psychiatric Sciences, University of Bari, Bari, Italy; ²Department of Neuroradiology, IRCCS 'Casa Sollievo della Sofferenza', San Giovanni Rotondo, FG, Italy; ³Clinical Brain Disorders Branch, Genes, Cognition and Psychosis, NIMH, NIH, Bethesda, MD, USA

Earlier studies have shown widespread alterations of functional connectivity of various brain networks in schizophrenia, including the default mode network (DMN). The DMN has also an important role in the performance of cognitive tasks. Furthermore, treatment with second-generation antipsychotic drugs may ameliorate to some degree working memory (WM) deficits and related brain activity. The aim of this study was to evaluate the effects of treatment with olanzapine monotherapy on functional connectivity among brain regions of the DMN during WM. Seventeen patients underwent an 8-week prospective study and completed two functional magnetic resonance imaging (fMRI) scans at 4 and 8 weeks of treatment during the performance of the N-back WM task. To control for potential repetition effects, 19 healthy controls also underwent two fMRI scans at a similar time interval. We used spatial group-independent component analysis (ICA) to analyze fMRI data. Relative to controls, patients with schizophrenia had reduced connectivity strength within the DMN in posterior cingulate, whereas it was greater in precuneus and inferior parietal lobule. Treatment with olanzapine was associated with increases in DMN connectivity with ventromedial prefrontal cortex, but not in posterior regions of DMN. These results suggest that treatment with olanzapine is associated with the modulation of DMN connectivity in schizophrenia. In addition, our findings suggest critical functional differences in the regions of DMN.

Neuropsychopharmacology (2010) **35**, 904–912; doi:10.1038/npp.2009.192; published online 2 December 2009

Keywords: default mode network; schizophrenia; antipsychotics; working memory; ventromedial prefrontal cortex; fMRI

INTRODUCTION

Alterations of functional connectivity of the brain have been implicated in the pathophysiology of schizophrenia (Friston and Frith, 1995; Meyer-Lindenberg *et al*, 2001; Weinberger *et al*, 1992). Although initial studies focused on abnormal dynamic interactions among functionally segregated brain regions co-activated during a task (Meyer-Lindenberg *et al*, 2001), a growing number of reports has recently suggested that disruptions in the default mode network (DMN) may have a role in the cognitive deficits found in schizophrenia (see Broyd *et al*, 2009) for a review).

The DMN comprises a set of brain regions whose activity increases during rest and is attenuated during an active task (Mazoyer *et al*, 2001; Raichle *et al*, 2001). This brain

network includes midline areas such as the ventromedial prefrontal cortex (vmPFC), the posterior cingulate cortex (pCC), the precuneus, as well as bilateral inferior parietal lobule (IPL) (Raichle *et al*, 2001). Activity in the DMN has been associated with stimulus-unrelated thoughts, lapses of attention and mind wandering (Mason *et al*, 2007). Different roles have been hypothesized for this network ranging from monitoring external environment ('watchfulness'), to internal mentation, for example, autobiographical memory, theory of mind, moral decision making and prospective thoughts (Gusnard *et al*, 2001). Interestingly, activity of the DMN shows a strong negative correlation, 'anticorrelation', with activity in task-related networks (Fox *et al*, 2005). This finding suggests the existence of two different modes of information processing: one with predominant DMN activity, which is relatively unfocused on external objects and characterized by mental exploration, whereas the other is focused on active processing of external stimuli (see (Buckner *et al*, 2008) for a review). Importantly, integrity of the DMN is also important for allocation of attentional resources needed for cognitive processing and has been associated with working memory

*Correspondence: Professor A Bertolino, Dipartimento di Scienze Neurologiche e Psichiatriche, Università degli Studi di Bari, Piazza Giulio Cesare, Bari 9 - 70124, Italy, Tel: +39 080 5478572, Fax: +39 080 5593204, E-mail: a.bertolino@psichiat.uniba.it

Received 21 July 2009; revised 7 October 2009; accepted 26 October 2009

(WM) performance (Hampson *et al*, 2006; Sambataro *et al*, 2008).

Although brain functions proposed for the DMN are impaired in patients with schizophrenia (Buckner *et al*, 2008), neuroimaging studies have reported inconsistent findings in the DMN of these individuals. Increased (Garrity *et al*, 2007; Harrison *et al*, 2007; Whitfield-Gabrieli *et al*, 2009; Zhou *et al*, 2007) or decreased (Liang *et al*, 2006; Pomarol-Clotet *et al*, 2008) activity and connectivity have been reported in the DMN network at the level of individual regions as well as of the global network. Different issues may have contributed to lack of consistent findings, including the method used for estimation of connectivity (Murphy *et al*, 2009). For example, the use of global signal regression, a standard preprocessing step of univariate seed-based connectivity analyses, has been questioned for the study of DMN as having a potential biasing effect in the estimation of the magnitude of the anticorrelations. Given that the use of this technique is still controversial (Fox *et al*, 2009; Murphy *et al*, 2009; Weissenbacher *et al*, 2009), other statistical approaches that do not rely on this preprocessing method such as independent component analysis (ICA) may yield more reliable results.

Another crucial determinant potentially impacting data consistency and DMN activity may be the effect of pharmacological treatment with antipsychotics. Although some reports are questioning the size of the effect and specificity of the cognitive effects of second-generation antipsychotics (Goldberg *et al*, 2007; Keefe *et al*, 2007), some evidence shows that this treatment is associated with improvement of some of the cognitive deficits of schizophrenia including WM (Bertolino *et al*, 2004; Purdon *et al*, 2000). Furthermore, second-generation antipsychotic treatment appears to modulate brain physiology underlying WM function with reduced activation of dorsolateral prefrontal cortex (DLPFC) and parietal cortex for better levels of performance, suggesting increased neural efficiency (Bertolino *et al*, 2004). Moreover, other studies have indicated that systemic administration of levodopa and apomorphine in healthy humans or in patients with Parkinson's disease can modify the activity of the DMN (Argyelan *et al*, 2008; Nagano-Saito *et al*, 2009) further suggesting the potential relationship between dopamine signaling and DMN modulation. On the other hand, despite these earlier studies, the effect of treatment with antipsychotics on the DMN is still unknown.

In this study, we hypothesized that the effects of treatment with olanzapine, a second-generation antipsychotic drug, on WM brain physiology and behavioral performance can also be associated with modulation of connectivity within the DMN. Thus, we performed a longitudinal study in previously untreated patients with schizophrenia undergoing 8 weeks of treatment with olanzapine monotherapy. Patients were studied using spatial group ICA with fMRI during WM at 4 (t_1) and 8 weeks of treatment (t_2). To control for potential well-known effects of task practice and learning (Goldberg *et al*, 2007), we also scanned normal controls twice (t_1 , t_2) with a time interval similar to that of patients.

MATERIALS AND METHODS

Subjects

A total of 19 patients with schizophrenia and 19 healthy controls, matched for age, gender, race, handedness participated (Oldfield, 1971) and socioeconomic status (Hollingshead and Redlich, 1958) participated in this study (Table 1). All subjects underwent clinical assessment following the Structured Clinical Interview for DSM-IV Axis I Disorders (SCID-IV; First *et al*, 1996). Exclusion criteria included past history or presence of any medical, neurological disorders, drug or alcohol abuse, active drug use in the past year, past head trauma with loss of consciousness. Normal controls were excluded if presented with past history or current diagnosis of psychiatric disorders as documented by SCID-IV.

At the time of enrollment, all patients were experiencing acute exacerbation of psychosis requiring hospitalization. Nine patients were diagnosed with schizophreniform disorder and all of them were diagnosed with schizophrenia at a later follow-up. In all 13 patients were drug-naïve and 6 had been drug-free for at least 2 weeks from oral antipsychotics or from at least two cycles of depot antipsychotic treatment. Clinical symptoms in patients were assessed on the day of study entry, at 4 weeks (t_1), and at 8 weeks (t_2) by a trained psychiatrist using the Positive and Negative Syndrome Scale (PANSS; see Table 2 for clinical data). All patients were treated with oral olanzapine monotherapy. Titration was allowed for the first 2 weeks, and then the dose was kept constant throughout the remaining 6 weeks of treatment (mean dose 20 mg/day; SD = 6.9 mg/day). Given the lack of reliable algorithms to handle missing data points in statistical imaging, only subjects with two longitudinal fMRI scans were selected for this study.

All participants gave written informed consent, approved by the Institutional Review Board of the University of Bari, Bari (Italy), to take part in the experiment.

Experimental Task

All subjects performed the N-back WM task as described elsewhere (Callicott *et al*, 1999). Briefly, subjects saw numbers (1–4) shown in pseudorandom sequence and displayed at the corners of a diamond-shaped box. A non-memory guided control condition (0-back) that required subjects to identify the number currently seen, alternated with the 2-back WM condition, in which subjects were asked to recall the number seen two stimuli before, while continuing to encode additionally incoming stimuli.

Table 1 Demographics of the Sample

	Patients	Normal controls	Difference
N	19	19	
Male/Female ratio	15:4	15:4	NS
Age (M ± SD, years)	24.8 ± 5.7	26.3 ± 6.9	$p = 0.47$
Handedness	0.80 ± 0.4	0.64 ± 0.4	$p = 0.22$
Hollingshead (M ± SD)	28.8 ± 15.6	32.8 ± 16.0	$p = 0.43$

Table 2 Clinical Data

	M	SD
Drug free (months), N = 6	11.2	13.2
Drug-naive, N = 13, 9 of whom First-Episode		
Age of onset	21.4	4.2
Length of illness (months)	44.1	67
Mean dose olanzapine (mg/day)	20	6.9
PANSS baseline		
PANSS total	101.6	25.2
PANSS positive	25.6	6.0
PANSS negative	23.9	10.7
PANSS general	52.2	13.7
PANSS 4 weeks		
PANSS total	68.47	18.88*
PANSS positive	16.95	4.61*
PANSS negative	17.89	6.87*
PANSS general	35.26	8.90*
PANSS 8 weeks		
PANSS total	60.63	15.66* [†]
PANSS positive	14.16	4.65* [†]
PANSS negative	15.58	7.05* [†]
PANSS general	30.37	7.48* [†]

Abbreviations: M, Mean; PANNS, Positive and Negative Syndrome Scale.

* $p < 0.005$ from baseline; [†] $p < 0.005$ from 4 weeks.

Performance was recorded through a fiber optic response box which allowed measurement of behavioral data as the number of correct responses (accuracy) and reaction time (RT). Data recordings of two patients were lost because of a computer glitch. The stimuli were arranged in a block-design, consisting of eight 30-s blocks: four blocks of the control condition alternating with four blocks of the 2-back WM condition.

Image Acquisition

Blood-oxygen level dependent fMRI was performed on a GE Signa (Milwaukee, WI) 3T scanner at t_1 and t_2 . A gradient echo Blood-oxygen level dependent-EPI pulse sequence was used to acquire 120 images per run. Each functional image consisted of 20 6-mm-thick axial slices covering the entire cerebrum and most of the cerebellum (TR = 2000 ms; TE = 30 ms; field of view = 24 cm; flip angle = 90°).

Data Analysis

Demographics, behavioral, and clinical data. Two sample *t*-tests and repeated measures ANOVAs were used to compare demographics and behavioral data, respectively. Chi-square analyses were performed to compare categorical variables. Clinical data were compared using paired *t*-tests.

Image analysis.

Preprocessing: Data were pre-processed and analyzed using Statistical Parametrical Mapping 2 (SPM; <http://www.fil.ion.ucl.ac.uk>). The first four volumes were discarded to allow for T1 equilibration effects. All functional volumes were realigned to the first volume acquired and spatially normalized to the Montreal Neurological Institute (MNI) standard brain in the space of Talairach and Tournoux (1988). Smoothing was carried out with a 10-mm full width half maximum isotropic 3D Gaussian kernel.

Independent component analysis: One group of spatial ICA was performed on preprocessed data using the Group ICA of fMRI Toolbox [GIFT; <http://icatb.sourceforge.net>, (Calhoun *et al*, 2001)] entering the data of all subjects for both time points. The dimensionality of functional data for each subject was reduced using three consecutive steps of Principal Component Analysis alternating with data concatenation across the subjects, resulting in one aggregate mixing matrix for all subjects. An ICA decomposition using the Infomax algorithm was used to extract 21 independent components (IC), consisting of group spatial maps and the related time courses (TC) of the estimated signal. Individual participant's spatial maps for each IC, which are the voxel-wise IC loadings, represent local strength of functional connectivity and reflect the correspondence between the estimated TC in each voxel and the average TC of the network itself. Given that in spatial ICA, the individual subject TC is assumed to be constant across the entire brain, the voxel-wise estimated signal deviations from the average network TC reflects local changes in the strength of functional connectivity within a given network. The minimum description length criteria were used to estimate the order selection that is the number of ICs from the smoothed data sets after taking into account the spatial and temporal correlation of the fMRI data (Li *et al*, 2007). These components were then used to back-reconstruct to individual ICs using the aggregate mixing matrix created during the reduction steps of dimensionality data. Each voxel value of the IC map for each participant was calibrated using z-scores for between participant comparisons.

Eventually, two components that resulted unstable as suggested by a coefficient of stability (Iq) lower than 0.95 calculated by 50 bootstrapped permuted estimations of the ICs (ICASSO; Himberg *et al*, 2004) or showed artifactual patterns defined by those ICs with a spatial correlation greater than $R^2 = 0.02$ with white matter and $R^2 = 0.05$ for CSF (Kim *et al*, 2009) were removed from the analysis. The ICs that had (1) the highest negative Pearson's *r* correlation between its TC and the 2-back condition (a box-car function convolved with a double-Gaussian hemodynamic response function) and (2) the highest spatial correlation with a DMN template provided with the ICA toolbox were selected for further analyses.

Spatial analyses: Each subject's DMN-IC was entered into SPM5 and analyzed using second level random-effects analyses. Repeated measures ANCOVA with the normalized 2-back performance (ratio between 2-back and 0-back accuracy) as a covariate of no interest and time as a within subjects variable was used to compare DMN-IC spatial

maps across diagnostic groups at each time point. Additional planned comparisons were performed to assess differences across diagnostic groups at each time point. To examine differences in positive and negative spatial maps separately, we created a positive and negative mask, respectively, using a one-sample *t*-test across all subjects. Average connectivity strength was extracted in significant clusters and correlated with WM performance computed as a ratio between percent accuracy at 2-back and at 0-back.

A statistical threshold of $p < 0.05$ with False-Discovery-Rate (FDR; Genovese *et al*, 2002) small-volume-correction with $q\text{-FDR} = 0.05$ was used to identify significant differences within regions of interest. Regions of interest were created using WFU pickatlas software version 1.04 (Functional MRI Laboratory at the Wake Forest University School of Medicine, <http://www.rad.wfubmc.edu/fmri>) and comprised the following *a priori* regions as part of the standard default mode: vmPFC, anterior cingulate cortex (BA24/32), and the IPL, as well as the posterior cingulate, and precuneus (Raichle *et al*, 2001). All coordinates are reported in MNI system.

Stimulus-related response: Averaged IC responses to the WM task were calculated for the DMN-IC TC for patients and normal controls separately during the 2-back condition using the deconvolution algorithm by Eichele *et al* (2008). Stimulus-related response depicts the extent of the modulation of the DMN TC during presentation of a WM stimulus of null duration. Averaged TCs were overlaid for both

groups for 17 s following the stimulus onset to estimate the full extent of the TC response to stimuli.

RESULTS

Behavioral Performance

There was a significant effect of diagnosis [$F(1, 34) = 13.596$, $p = 0.0007$] on task accuracy: patients with schizophrenia had reduced performance relative to normal controls. There was also an effect of time [$F(1, 34) = 9.151$, $p = 0.004$] indicating that subjects at t_1 had lower performance relative to t_2 . There was no interaction of diagnosis by time of observation ($p > 0.1$). Even though both groups of subjects tended to improve accuracy over time, a *t*-test for dependent samples demonstrated that the difference was significant in patients ($t_{16} = 2.05$, $p = 0.05$) but not in controls ($t_{18} = 1.61$, $p = 0.12$), suggesting an effect of olanzapine on WM accuracy in patients. There was a significant effect of diagnosis [$F(1, 34) = 5.1299$, $p = 0.03$] on reaction time, with patients with schizophrenia being slower relative to normal controls. There was no effect of time of observation ($p > 0.1$) or interaction of diagnosis by time of observation ($p > 0.1$) on reaction time.

Imaging Results

Default mode network component. One IC spanning across the DMN regions and anticorrelated with the task was identified for all subjects across both time points (Figure 1a).




Figure 1 Default mode network (DMN). One DMN-IC comprising spatial maps and time courses was identified across both time points for all the subjects. (a) Axial maps display the spatial pattern of the independent components identified. (b) Time courses represent the temporal profile of DMN each component (blue continuous line) overlaid on the paradigm 'box-car' design (red dashed line). All images are thresholded at $p = 0.05$ corrected for multiple comparisons. Color bar indicates *t*-scores.

This IC included midline regions vmPFC, pCC, precuneus (BA5/7), as well as bilateral IPL. This IC had high spatial correlation with a DMN template, $R^2 = 0.33$ and high temporal anticorrelation with the timing of the 2-back condition of task, $r = -0.65$ (Figure 1b).

Between group comparisons.

Spatial extent: Effect of Diagnosis Patients with schizophrenia had increased connectivity with the DMN in left IPL (BA39, $x = -56$, $y = -67$, $z = 23$, $Z = 4.03$, $p = 0.00003$ FDR corrected) as well as in precuneus (BA7, $x = 4$, $y = -79$, $z = 45$, $Z = 3.33$, $p = 0.0004$ with q -FDR = 0.09). In addition, patients had decreased connectivity with the DMN in pCC (BA23/31, $x = 4$, $y = -52$, $z = 26$, $Z = 3.34$, $p = 0.0004$ FDR corrected; Figure 2).

Effect of time There was no significant effect of time.

Time by diagnosis There was a time by diagnosis interaction in vmPFC (BA10, $x = 0$, $y = 49$, $z = -7$, $Z = 3.02$, $p = 0.001$ FDR corrected; Figure 3a). In patients with schizophrenia the strength of connectivity increased at t_2 relative to t_1 with respect to normal controls (Figure 3b). **Post hoc** analysis revealed that patients with schizophrenia had significantly greater connectivity strength extracted in the vmPFC cluster at t_2 relative to t_1 ($p = 0.01$). However, this was not true in normal controls ($p > 0.1$). In addition, at t_2 connectivity strength in patients in this region was greater relative to normal controls ($p = 0.01$). Strength of connectivity in vmPFC was positively correlated with WM accuracy in patients, but not in normal controls (controls: t_1 : $r = -0.20$, $p > 0.1$; t_2 : $r = -0.15$, $p > 0.1$; Figure 3c).




Figure 2 Effect of diagnosis on brain connectivity in the DMN. Patients with schizophrenia show decreased connectivity strength within the DMN in pCC (a) as well as increased connectivity in left-Inferior Parietal Lobule (BA39) and precuneus (BA7) relative to normal controls (b). Statistical t -map of DMN connectivity overlaid on the MNI brain template. Color bar indicates t -scores.

Furthermore, in patients the correlation was significant at t_1 but not at t_2 (patients: t_1 : $r = 0.49$, $p = 0.05$; t_2 : $r = 0.16$, $p > 0.1$).

Average stimulus-related responses. Average stimulus responses of DMN IC to the WM condition had a smaller negative modulation of this signal in patients relative to normal controls. This difference in WM-related responses decreased at t_2 relative to t_1 (Figure 4).

Negative DMN component

The negative component of DMN had a pattern similar to the typical WM task described during 2-back task (Supplementary Figure S1). This network of brain regions included bilaterally dorsolateral (DLPFC) and ventrolateral prefrontal cortex, frontal pole, caudate, anterior cingulate, and lateral posterior parietal cortex. This network demonstrated main effects and interaction between diagnosis and treatment in the region of DLPFC previously shown to be responsive to treatment with olanzapine as in earlier studies (Bertolino *et al*, 2004; see Supplementary Materials for further details). In normal controls (t_1 : $r = 0.40$, $p = 0.09$; $t_2 = 0.39$, $p = 0.1$) but not patients (t_1 : $r = -0.03$, $p > 0.1$; $t_2 = -0.14$, $p > 0.1$) there was a statistical trend for correlation between DLPFC connectivity strength and WM accuracy.

DISCUSSION

In this study we investigated the effect of olanzapine on DMN connectivity in patients with schizophrenia. We found that patients have greater strength of connectivity in the IPL and precuneus as well as reduced connectivity in pCC. These effects were not significantly affected by treatment with olanzapine. On the other hand, our data indicate an interaction of time by diagnosis on connectivity strength in vmPFC where patients with schizophrenia had greater connectivity strength with the rest of DMN during olanzapine treatment. In patients with schizophrenia the extent of DMN connectivity strength in vmPFC positively correlated with WM accuracy at t_1 but not at t_2 .

Previous imaging studies have reported widespread alterations of DMN in patients with schizophrenia (see Broyd *et al*, 2009 for a review). Our findings indicate decreased functional connectivity with the DMN in pCC as well as increased connectivity strength in parietal cortex, including precuneus and IPL, in patients with schizophrenia. pCC is an important node of the DMN and a crucial cortical hub for interconnecting segregated brain areas, that is, anterior and posterior midline systems that are mPFC and entorhinal cortex (Greicius *et al*, 2009). As a result of these connections pCC may have an important role in cognitive functions including episodic memory (Greicius *et al*, 2003) and WM (Hampson *et al*, 2006; Sambataro *et al*, 2008). In addition, disruption of functional coupling with pCC is associated with ADHD (Castellanos *et al*, 2008) and cognitive decline in physiological (Sambataro *et al*, 2008) and pathological aging (Greicius *et al*, 2004). The relative lack of disease specificity of these findings suggests that this mechanism reflects basic disruption of the brain systems associated with memory dysfunction across different




Figure 3 Time by diagnosis interaction on brain connectivity in the DMN. In patients with schizophrenia (SCZ) connectivity strength within the DMN increased in ventromedial prefrontal cortex (vmPFC) (a); at t_2 (red) relative to t_1 (blue) when compared with normal controls (NC) (b); connectivity strength in vmPFC was positively correlated with task performance in patients at t_1 ($r=0.45$, $p=0.04$) (c); statistical t-maps of DMN connectivity are overlaid on the MNI brain template. Color bar indicates t-scores. Bar plots report parameter estimates of the time by diagnosis contrast from the ANCOVA extracted from the vmPFC cluster and measured in arbitrary units (a.u.). Dashed lines indicate the trendline. * $P=0.01$.




Figure 4 Time by diagnosis interaction on stimulus-related responses. At t_1 patients had a smaller stimulus-related decrease of the time course amplitude relative to normal controls, and this difference decreased at t_2 . Y axis indicates the difference (patients—normal controls, SCZ—NC) of time course amplitude of the DMN-IC patients with schizophrenia relative to normal controls in arbitrary units (a.u.) as calculated by deconvolving the estimated signal for the 2-back condition, X axis indicates latency from the working memory stimulus onset expressed in seconds. The dashed arrow indicates the effect of time on the group differences.

categorical diagnostic conditions. Thus, decreased connectivity strength between pCC and the other regions of DMN may contribute to the WM deficits observed in patients with schizophrenia. On the other hand, connectivity of the parietal regions, that is, precuneus and IPL, with the DMN has crucial anatomical and functional differences within posterior DMN regions. Cytoarchitecturally, the precuneus is a fully differentiated isocortex as opposed to the limbic cortex in pCC, and it also has structural connections with different thalamic nuclei and with occipital, parietal, and frontal areas (Buckwalter *et al*, 2008). This region is

associated with visual processing, motor planning (Cavada and Goldman-Rakic, 1989), cognitive functions, such as motor imagery, spatial navigation (Sato *et al*, 2006), memory retrieval (Lundstrom *et al*, 2005), and also self-perception and internal mentation (Gusnard *et al*, 2001). On the lateral parietal cortex, IPL which is a heteromodal association area, also has a crucial role in integrating incoming sensory information (Lynch, 1980) and in executive functions, specifically WM, sustained attention and dual task performance (Torrey, 2007). Increased connectivity of parietal regions within the DMN may be compensatory for deficits in other brain regions within the DMN, for example, pCC.

The DMN anti-correlated network comprised a set of brain regions that are recruited during WM tasks (Callicott *et al*, 1999). Patients with schizophrenia had increased connectivity in DLPFC which reflects the well-known prefronto-cortical dysfunction in schizophrenia (Weinberger *et al*, 1986). Consistent with these prefrontal effects during working memory, after 8 weeks of treatment with olanzapine, patients with schizophrenia had greater connectivity in vmPFC. Interestingly, connectivity in vmPFC correlated significantly with behavioral accuracy in patients at the first time point only and not in controls. On the other hand, DLPFC activation in healthy subjects correlated with WM behavioral performance both at t_1 and t_2 . Alterations in vmPFC have been reported in patients with schizophrenia and this finding has been associated with abnormal prefronto-cortical function (Meyer-Lindenberg *et al*, 2001; Pomarol-Clotet *et al*, 2008). vmPFC is involved in monitoring and integrating emotional and cognitive processing through its projections to limbic regions, orbitofrontal cortex, ventral striatum, amygdala, and hypothalamus (Gusnard *et al*, 2001).

This brain region has an important role in decision making as well as in emotion and motivation associated with a cognitive task. Neurons in vmPFC encode the relative current value and the outcome of chosen options during decisions by comparing available options (Boorman *et al*, 2009; Luk and Wallis, 2009). In addition, vmPFC regulates both the emotional context and motivational milieu during a cognitive task (Groenewegen and Uylings, 2000) so that this region inhibits emotional processing that may interfere with cognitive performance in a 'dynamic interplay' with the regions of DMN anticorrelated network and specifically DLPFC (Longe *et al*, 2009). In keeping with this, Uddin *et al* (2009) found that DMN activity in vmPFC 'Granger-causes' the activity in the anticorrelated network and not vice versa. Decreased temporal anticorrelation between the DMN and task-related network may impair task performance either for decreased allocation of attentional resources for the task at hand (Fransson, 2006) or for direct interference of non suppressed DMN ('default mode interference hypothesis' Sonuga-Barke and Castellanos, 2007). Our present data in vmPFC are consistent with this earlier literature. Moreover, they suggest that olanzapine treatment is associated with a more physiological modulation of the relationship between vmPFC connectivity and behavioral accuracy which is found only in patients and only at t_1 . This interpretation is also consistent with the possibility that olanzapine treatment may modulate DMN directly or acting on both anticorrelated networks (Bertolino *et al*, 2004; Blasi *et al*, 2009). As a result of this modulation, olanzapine can improve the temporal characteristics of brain response (Schlagenhauf *et al*, 2008) of these networks so that patients have a faster and deeper hemodynamic response to the task at hand (Figure 4) which may improve WM performances.

Olanzapine may affect DMN connectivity via dopaminergic system either directly on pCC or indirectly via PFC modulation. Evidence from animal studies indicates that olanzapine increases extracellular levels of dopamine in PFC (Gessa *et al*, 2000; Ichikawa *et al*, 2001). Dopamine can regulate brain networks by modulating the synchronization among different brain regions and membrane oscillation frequencies (Seamans and Yang, 2004) thus increasing stability of attentional networks, which ultimately results in decreased trial-by-trial variability and increased signal-to-noise ratio (Rolls *et al*, 2008). Decreased dopamine levels as resulting from carrying val alleles of COMTval¹⁵⁸met polymorphism have been associated with increased brain deactivations in pCC during a prosaccade task (Ettinger *et al*, 2008) as well as a decreased sequence learning-related deactivation in mPFC (Argyelan *et al*, 2008). Thus increased levels of dopamine in the hubs of DMN may affect the connectivity in the whole network. Alternatively, treatment with olanzapine may also indirectly regulate the DMN connectivity via an increase of dopamine levels in the anticorrelated network (Bertolino *et al*, 2004).

A caveat to our study is the lack of fMRI data during WM at baseline that for clinical reasons could not be acquired. It is extremely difficult to obtain co-operation from patients to undergo complicated research experiments during acute exacerbations requiring hospitalization. We cannot exclude that patients may have already had an improvement at the level of brain physiology and of behavioral performance before 4 weeks of treatment with olanzapine relative to the

baseline. Nevertheless, our data acquired within the time-frame of this study suggest that olanzapine modulates not only DMN connectivity, but, more importantly, its relationship with WM performance.

Another limitation of this study is the intrinsic reliability of the fMRI data in longitudinal studies. Manoach *et al* (2001), have reported a low test-retest reliability of the magnitude and extent of brain activations during a Sternberg Item recognition task in patients with schizophrenia across scan sessions. To control this variance, we used a bootstrapped ICA that is a technique well-known to be able to identify and isolate artifactual and noise components (Calhoun *et al*, 2001).

Finally, although estimation of DMN may be more optimal during non complex cognitive tasks (Esposito *et al*, 2006), some studies have reported similar spatial extent of DMN during rest and during variously demanding cognitive tasks (Calhoun *et al*, 2008; Smith *et al*, 2009). Moreover, Calhoun *et al* (2008) found similar results during rest and during an oddball task when comparing results between patients with schizophrenia and normal controls. Furthermore, most of the studies on DMN in patients with schizophrenia have used complex cognitive tasks such as an oddball task (Garrity *et al*, 2007), Multi-Source Interference Task (Harrison *et al*, 2007) and 1-back and 2-back (Pomarol-Clotet *et al*, 2008; Whitfield-Gabrieli *et al*, 2009). These studies have reported results that are similar to those obtained with rest data. The objective of our study was to investigate potential effects of treatment using a within subject design under the same task load conditions. Therefore, this design has allowed estimation and evaluation of our objective under very similar physiological conditions thus contributing to temper potential limitations associated with assessing DMN during cognition.

In conclusion, our data suggest that treatment with olanzapine can modulate brain connectivity in DMN and in its anticorrelated network, and their task-related responses. Furthermore, our findings shed new light on the functional anatomy underlying the DMN network, suggesting also potential different roles of posterior DMN regions in the pathogenesis of schizophrenia.

ACKNOWLEDGEMENTS

We are thankful to Riccarda Lomuscio, BA, for help with acquisition of the data, and to all subjects who have participated in the study.

DISCLOSURE

Dr Bertolino has received consultant fees from Eli-Lilly, Astra-Zeneca, and Janssen. The other authors declare that they do not have any commercial or financial involvements that might present an appearance of a conflict of interest in connection with the submitted article.

REFERENCES

Argyelan M, Carbon M, Ghilardi MF, Feigin A, Mattis P, Tang C *et al* (2008). Dopaminergic suppression of brain deactivation responses during sequence learning. *J Neurosci* 28: 10687–10695.

- Bertolino A, Caforio G, Blasi G, De Candia M, Latorre V, Petruzzella V et al (2004). Interaction of COMT (Val(108/158)Met) genotype and olanzapine treatment on prefrontal cortical function in patients with schizophrenia. *Am J Psychiatry* **161**: 1798–1805.
- Blasi G, Popolizio T, Taurisano P, Caforio G, Romano R, Di Giorgio A et al (2009). Changes in prefrontal and amygdala activity during olanzapine treatment in schizophrenia. *Psychiatry Res* **173**: 31–38.
- Boorman ED, Behrens TE, Woolrich MW, Rushworth MF (2009). How green is the grass on the other side? Frontopolar cortex and the evidence in favor of alternative courses of action. *Neuron* **62**: 733–743.
- Broyd SJ, Demanuele C, Debener S, Helps SK, James CJ, Sonuga-Barke EJ (2009). Default-mode brain dysfunction in mental disorders: a systematic review. *Neurosci Biobehav Rev* **33**: 279–296.
- Buckner RL, Andrews-Hanna JR, Schacter DL (2008). The brain's default network: anatomy, function, and relevance to disease. *Ann NY Acad Sci* **1124**: 1–38.
- Buckwalter JA, Parvizi J, Morecraft RJ, van Hoesen GW (2008). Thalamic projections to the posteromedial cortex in the macaque. *J Comp Neurol* **507**: 1709–1733.
- Calhoun VD, Adali T, Pearlson GD, Pekar JJ (2001). A method for making group inferences from functional MRI data using independent component analysis. *Hum Brain Mapp* **14**: 140–151.
- Calhoun VD, Kiehl KA, Pearlson GD (2008). Modulation of temporally coherent brain networks estimated using ICA at rest and during cognitive tasks. *Hum Brain Mapp* **29**: 828–838.
- Callicott JH, Mattay VS, Bertolino A, Finn K, Coppola R, Frank JA et al (1999). Physiological characteristics of capacity constraints in working memory as revealed by functional MRI. *Cereb Cortex* **9**: 20–26.
- Castellanos FX, Margulies DS, Kelly C, Uddin LQ, Ghaffari M, Kirsch A et al (2008). Cingulate-precuneus interactions: a new locus of dysfunction in adult attention-deficit/hyperactivity disorder. *Biol Psychiatry* **63**: 332–337.
- Cavada C, Goldman-Rakic PS (1989). Posterior parietal cortex in rhesus monkey: I. Parcellation of areas based on distinctive limbic and sensory corticocortical connections. *J Comp Neurol* **287**: 393–421.
- Eichele T, Debener S, Calhoun VD, Specht K, Engel AK, Hugdahl K et al (2008). Prediction of human errors by maladaptive changes in event-related brain networks. *Proc Natl Acad Sci USA* **105**: 6173–6178.
- Esposito F, Bertolino A, Scarabino T, Latorre V, Blasi G, Popolizio T et al (2006). Independent component model of the default-mode brain function: assessing the impact of active thinking. *Brain Res Bull* **70**: 263–269.
- Ettinger U, Kumari V, Collier DA, Powell J, Luzi S, Michel TM et al (2008). Catechol-O-methyltransferase (COMT) val158met genotype is associated with BOLD response as a function of task characteristic. *Neuropsychopharmacology* **33**: 3046–3057.
- First MB, Gibbon M, Spitzer RL, Williams JBW (1996). *User's Guide for the Structured Clinical Interview for DSM IV Axis I Disorders (SCID-I, Research Version)*. Biometric Research Department: New York.
- Fox MD, Snyder AZ, Vincent JL, Corbetta M, Van Essen DC, Raichle ME (2005). The human brain is intrinsically organized into dynamic, anticorrelated functional networks. *Proc Natl Acad Sci USA* **102**: 9673–9678.
- Fox MD, Zhang D, Snyder AZ, Raichle ME (2009). The global signal and observed anticorrelated resting state brain networks. *J Neurophysiol* **101**: 3270–3283.
- Fransson P (2006). How default is the default mode of brain function? Further evidence from intrinsic BOLD signal fluctuations. *Neuropsychologia* **44**: 2836–2845.
- Friston KJ, Frith CD (1995). Schizophrenia: a disconnection syndrome? *Clin Neurosci* **3**: 89–97.
- Garrity AG, Pearlson GD, McKiernan K, Lloyd D, Kiehl KA, Calhoun VD (2007). Aberrant 'default mode' functional connectivity in schizophrenia. *Am J Psychiatry* **164**: 450–457.
- Genovese CR, Lazar NA, Nichols T (2002). Thresholding of statistical maps in functional neuroimaging using the false discovery rate. *Neuroimage* **15**: 870–878.
- Gessa GL, Devoto P, Diana M, Flore G, Melis M, Pistis M (2000). Dissociation of haloperidol, clozapine, and olanzapine effects on electrical activity of mesocortical dopamine neurons and dopamine release in the prefrontal cortex. *Neuropsychopharmacology* **22**: 642–649.
- Goldberg TE, Goldman RS, Burdick KE, Malhotra AK, Lencz T, Patel RC et al (2007). Cognitive improvement after treatment with second-generation antipsychotic medications in first-episode schizophrenia: is it a practice effect? *Arch Gen Psychiatry* **64**: 1115–1122.
- Greicius MD, Krasnow B, Reiss AL, Menon V (2003). Functional connectivity in the resting brain: a network analysis of the default mode hypothesis. *Proc Natl Acad Sci USA* **100**: 253–258.
- Greicius MD, Srivastava G, Reiss AL, Menon V (2004). Default-mode network activity distinguishes Alzheimer's disease from healthy aging: evidence from functional MRI. *Proc Natl Acad Sci USA* **101**: 4637–4642.
- Greicius MD, Supekar K, Menon V, Dougherty RF (2009). Resting-state functional connectivity reflects structural connectivity in the default mode network. *Cereb Cortex* **19**: 72–78.
- Groenewegen HJ, Uylings HB (2000). The prefrontal cortex and the integration of sensory, limbic and autonomic information. *Prog Brain Res* **126**: 3–28.
- Gusnard DA, Akbudak E, Shulman GL, Raichle ME (2001). Medial prefrontal cortex and self-referential mental activity: relation to a default mode of brain function. *Proc Natl Acad Sci USA* **98**: 4259–4264.
- Hampson M, Driesen NR, Skudlarski P, Gore JC, Constable RT (2006). Brain connectivity related to working memory performance. *J Neurosci* **26**: 13338–13343.
- Harrison BJ, Yucel M, Pujol J, Pantelis C (2007). Task-induced deactivation of midline cortical regions in schizophrenia assessed with fMRI. *Schizophr Res* **91**: 82–86.
- Himberg J, Hyvarinen A, Esposito F (2004). Validating the independent components of neuroimaging time series via clustering and visualization. *Neuroimage* **22**: 1214–1222.
- Hollingshead A, Redlich F (1958). *Social Class and Mental Illness*. Wiley: New York.
- Ichikawa J, Ishii H, Bonaccorso S, Fowler WL, O'Laughlin IA, Meltzer HY (2001). 5-HT(2A) and D(2) receptor blockade increases cortical DA release via 5-HT(1A) receptor activation: a possible mechanism of atypical antipsychotic-induced cortical dopamine release. *J Neurochem* **76**: 1521–1531.
- Keefe RS, Bilder RM, Davis SM, Harvey PD, Palmer BW, Gold JM et al (2007). Neurocognitive effects of antipsychotic medications in patients with chronic schizophrenia in the CATIE Trial. *Arch Gen Psychiatry* **64**: 633–647.
- Kim DI, Manoach DS, Mathalon DH, Turner JA, Mannell M, Brown GG et al (2009). Dysregulation of working memory and default-mode networks in schizophrenia using independent component analysis, an fBIRN and MCIC study. *Hum Brain Mapp* **30**: 3795–3811.
- Li YO, Adali T, Calhoun VD (2007). Estimating the number of independent components for functional magnetic resonance imaging data. *Hum Brain Mapp* **28**: 1251–1266.
- Liang M, Zhou Y, Jiang T, Liu Z, Tian L, Liu H et al (2006). Widespread functional disconnectivity in schizophrenia with resting-state functional magnetic resonance imaging. *Neuroreport* **17**: 209–213.

- Longe O, Senior C, Rippon G (2009). The lateral and ventromedial prefrontal cortex work as a dynamic integrated system: evidence from fMRI connectivity analysis. *J Cogn Neurosci* 21: 141–154.
- Luk CH, Wallis JD (2009). Dynamic encoding of responses and outcomes by neurons in medial prefrontal cortex. *J Neurosci* 29: 7526–7539.
- Lundstrom BN, Ingvar M, Petersson KM (2005). The role of precuneus and left inferior frontal cortex during source memory episodic retrieval. *Neuroimage* 27: 824–834.
- Lynch JC (1980). The functional organization of posterior parietal association cortex. *Behav Brain Sci* 3: 485–534.
- Manoach DS, Halpern EF, Kramer TS, Chang Y, Goff DC, Rauch SL *et al* (2001). Test-retest reliability of a functional MRI working memory paradigm in normal and schizophrenic subjects. *Am J Psychiatry* 158: 955–958.
- Mason MF, Norton MI, Van Horn JD, Wegner DM, Grafton ST, Macrae CN (2007). Wandering minds: the default network and stimulus-independent thought. *Science* 315: 393–395.
- Mazoyer B, Zago L, Mellet E, Bricogne S, Etard O, Houde O *et al* (2001). Cortical networks for working memory and executive functions sustain the conscious resting state in man. *Brain Res Bull* 54: 287–298.
- Meyer-Lindenberg A, Poline JB, Kohn PD, Holt JL, Egan MF, Weinberger DR *et al* (2001). Evidence for abnormal cortical functional connectivity during working memory in schizophrenia. *Am J Psychiatry* 158: 1809–1817.
- Murphy K, Birn RM, Handwerker DA, Jones TB, Bandettini PA (2009). The impact of global signal regression on resting state correlations: are anti-correlated networks introduced? *Neuroimage* 44: 893–905.
- Nagano-Saito A, Liu J, Doyon J, Dagher A (2009). Dopamine modulates default mode network deactivation in elderly individuals during the Tower of London task. *Neurosci Lett* 458: 1–5.
- Oldfield RC (1971). The assessment and analysis of handedness: the Edinburgh inventory. *Neuropsychologia* 9: 97–113.
- Pomarol-Clotet E, Salvador R, Sarro S, Gomar J, Vila F, Martinez A *et al* (2008). Failure to deactivate in the prefrontal cortex in schizophrenia: dysfunction of the default mode network? *Psychol Med* 38: 1185–1193.
- Purdon SE, Jones BD, Stip E, Labelle A, Addington D, David SR *et al* (2000). Neuropsychological change in early phase schizophrenia during 12 months of treatment with olanzapine, risperidone, or haloperidol. The Canadian Collaborative Group for research in schizophrenia. *Arch Gen Psychiatry* 57: 249–258.
- Raichle ME, MacLeod AM, Snyder AZ, Powers WJ, Gusnard DA, Shulman GL (2001). A default mode of brain function. *Proc Natl Acad Sci USA* 98: 676–682.
- Rolls ET, Loh M, Deco G, Winterer G (2008). Computational models of schizophrenia and dopamine modulation in the prefrontal cortex. *Nat Rev Neurosci* 9: 696–709.
- Sambataro F, Murty VP, Callicott JH, Tan HY, Das S, Weinberger DR *et al* (2008). Age-related alterations in default mode network: impact on working memory performance. *Neurobiol Aging*. doi:10.1016/j.neurobiolaging.2008.05.022. (in press)
- Sato N, Sakata H, Tanaka YL, Taira M (2006). Navigation-associated medial parietal neurons in monkeys. *Proc Natl Acad Sci USA* 103: 17001–17006.
- Sohlagenhaut F, Wustenberg T, Schmack K, Dingas M, Wrase J, Koslowski M *et al* (2008). Switching schizophrenia patients from typical neuroleptics to olanzapine: effects on BOLD response during attention and working memory. *Eur Neuro* 18: 689–699.
- Seamans JK, Yang CR (2004). The principal features and mechanisms of dopamine modulation in the prefrontal cortex. *Prog Neurobiol* 74: 1–58.
- Smith SM, Fox PT, Miller KL, Glahn DC, Fox PM, Mackay CE *et al* (2009). Correspondence of the brain's functional architecture during activation and rest. *Proc Natl Acad Sci USA* 106: 13040–13045.
- Sonuga-Barke EJ, Castellanos FX (2007). Spontaneous attentional fluctuations in impaired states and pathological conditions: a neurobiological hypothesis. *Neurosci Biobehav Rev* 31: 977–986.
- Talairach J, Tournoux P (1988). *Co-Planar Stereotactic Atlas of the Human Brain*. Georg Thieme Verlag/Thieme Medical Publishers: Stuttgart-New York.
- Torrey EF (2007). Schizophrenia and the inferior parietal lobule. *Schizophr Res* 97: 215–225.
- Uddin LQ, Kelly AM, Biswal BB, Xavier Castellanos F, Milham MP (2009). Functional connectivity of default mode network components: correlation, anticorrelation, and causality. *Hum Brain Mapp* 30: 625–637.
- Weinberger DR, Berman KF, Suddath R, Torrey EF (1992). Evidence of dysfunction of a prefrontal-limbic network in schizophrenia: a magnetic resonance imaging and regional cerebral blood flow study of discordant monozygotic twins. *Am J Psychiatry* 149: 890–897.
- Weinberger DR, Berman KF, Zec RF (1986). Physiologic dysfunction of dorsolateral prefrontal cortex in schizophrenia. I. Regional cerebral blood flow evidence. *Arch Gen Psychiatry* 43: 114–124.
- Weissenbacher A, Kasess C, Gerstl F, Lanzenberger R, Moser E, Windischberger C (2009). Correlations and anticorrelations in resting-state functional connectivity MRI: a quantitative comparison of preprocessing strategies. *Neuroimage* 47: 1408–1416.
- Whitfield-Gabrieli S, Thermenos HW, Milanovic S, Tsuang MT, Faraone SV, McCarley RW *et al* (2009). Hyperactivity and hyperconnectivity of the default network in schizophrenia and in first-degree relatives of persons with schizophrenia. *Proc Natl Acad Sci USA* 106: 1279–1284.
- Zhou Y, Liang M, Tian L, Wang K, Hao Y, Liu H *et al* (2007). Functional disintegration in paranoid schizophrenia using resting-state fMRI. *Schizophr Res* 97: 194–205.

Supplementary Information accompanies the paper on the Neuropsychopharmacology website (<http://www.nature.com/npp>)