

SEARCHING FOR AN ANALOGUE OF ATR_0 IN THE WEIHRAUCH LATTICE

TAKAYUKI KIHARA, ALBERTO MARCONE, AND ARNO PAULY

Abstract. There are close similarities between the Weihrauch lattice and the zoo of axiom systems in reverse mathematics. Following these similarities has often allowed researchers to translate results from one setting to the other. However, amongst the *big five* axiom systems from reverse mathematics, so far ATR_0 has no identified counterpart in the Weihrauch degrees. We explore and evaluate several candidates, and conclude that the situation is complicated.

§1. Introduction. Reverse mathematics [42] is a program to find the sufficient and necessary axioms to prove theorems of mathematics (that can be formalized in second-order arithmetic). For this, a base system (RCA_0) is fixed, and then equivalences between theorems and certain benchmark axioms are proven. Sometimes, a careful reading of the original proof of the theorem reveals which of the benchmark axioms are used, and the main challenge is to show that the theorem indeed implies those axioms (hence the name *reverse* mathematics). A vast number of theorems turned out to be equivalent to one of only five systems: RCA_0 , WKL_0 , ACA_0 , ATR_0 , and $\Pi_1^1\text{-CA}_0$. While recently attention has shifted to theorems not equivalent to one of the *big five*, the big five still occupy a central role in the endeavour.

Computational metamathematics in the Weihrauch lattice starts with the observation that many theorems in analysis and other areas of mathematics have Π_2 -*gestalt*, i.e. are of the form $\forall x \in \mathbf{X} (Q(x) \rightarrow \exists y \in \mathbf{Y} P(x, y))$, and can hence be seen as computational tasks: Given some $x \in \mathbf{X}$ satisfying $Q(x)$, find a suitable witness $y \in \mathbf{Y}$. This task can also be viewed as a multivalued partial function $f : \subseteq \mathbf{X} \rightrightarrows \mathbf{Y}$, and thus the precise definition of Weihrauch reducibility (given in Section 2.2 below) deals with this kind of objects. Often, the task cannot be solved algorithmically (equivalently, the multivalued function is not computable). The research programme (as formulated by Gherardi and Marcone [20], Pauly [35, 37], and in particular Brattka and Gherardi [7, 6]) is to compare the degree of impossibility as follows: Assume we had a black box to solve the task for Theorem B. Can we solve the task for Theorem A using the black box exactly once? If so, then $A \leq_W B$, A is Weihrauch reducible to B.

Received November 30, 2018.

2020 *Mathematics Subject Classification*. Primary 03D78, Secondary 03D30, 03B30.

Key words and phrases. Weihrauch degrees, perfect set theorem, open and clopen determinacy, comparability of well-orders.

© 2020, Association for Symbolic Logic

0022-4812/20/8503-0007

DOI:10.1017/jsl.2020.12

As provability in RCA_0 is closely linked to computability, it is maybe not that surprising that very often, classification in reverse math can be translated easily into Weihrauch reductions.¹ While there are a number of obstacles for precise correspondence (see [24] for a detailed discussion), the resource-sensitivity of Weihrauch reductions might be the most obvious one: A proof in reverse mathematics can use a principle multiple times, a Weihrauch reduction uses its black box once. This obstacle does not apply to RCA_0 or WKL_0 classifications.

The analogue of RCA_0 are the computable principles, the analogue of WKL_0 is $\text{C}_{2^{\mathbb{N}}}$ (closed choice on Cantor space), and the analogue of ACA_0 is lim or finite iterations thereof. Theorems equivalent to $\Pi_1^1\text{-CA}_0$ have not yet been studied in the Weihrauch lattice, but an obvious analogue of $\Pi_1^1\text{-CA}_0$ is readily defined as the function which maps a countable sequence of trees to the characteristic function of the set of indices corresponding to well-founded trees. This leaves ATR_0 out of the big five, leading Marcone to initiate the search for an analogue in the Weihrauch lattice at a Dagstuhl meeting on Weihrauch reducibility [13].

Two candidates have been put forth as potential answers, $\text{UC}_{\mathbb{N}^{\mathbb{N}}}$ and $\text{C}_{\mathbb{N}^{\mathbb{N}}}$ (unique choice and closed choice on Baire space). We will examine some evidence for both of them, and show that the question is not as easily answered as those for the other big five. Our main focus is on three particular theorems equivalent to ATR_0 in reverse mathematics: Comparability of well orderings, open determinacy on Baire space² and the perfect tree theorem.

THEOREM (Comparability of well orderings). *If X and Y are well orderings over \mathbb{N} , then $|X| \leq |Y|$ or $|Y| \leq |X|$.*

THEOREM (Open determinacy). *Consider a two-player infinite sequential game with moves from \mathbb{N} . Let the first player have an open winning set. Then one player has a winning strategy.*

THEOREM (Perfect Tree Theorem). *If $T \subseteq \mathbb{N}^{<\mathbb{N}}$ is a tree, then either $[T]$ is countable or T has a perfect subtree.*

Structure of the paper. In Section 2 we recall the prerequisite notions about Weihrauch reducibility. While reverse mathematics serves as the motivation for this paper, its results are not invoked in our proofs, hence we do not expand on this area. In Section 3 we recall two Weihrauch degrees of central importance, unique choice $\text{UC}_{\mathbb{N}^{\mathbb{N}}}$ and closed choice $\text{C}_{\mathbb{N}^{\mathbb{N}}}$ on Baire space. We then prove some equivalences to those for variants of comprehension and separation principles. In Section 4, we re-examine the strength of a separation principle, which is shown to be equivalent to $\Sigma_1^1\text{-WKL}$, weak König's lemma for Σ_1^1 -trees (Theorem 4.3). The comparability of well orderings is studied in Section 5. We see two variants, one of which we prove to be equivalent to $\text{UC}_{\mathbb{N}^{\mathbb{N}}}$ (Theorem 5.5) whereas the other resists full classification (Question 5.8).

¹The reverse direction would also be possible, but as reverse mathematics is the older field, occurs seldom in practice.

²The version for Cantor space has been studied in the Weihrauch degrees by Le Roux and Pauly [30].

Open determinacy and the perfect tree theorem are investigated in Sections 6 and 7. Both principles are formulated as disjunctions, and the versions where we know in which case we are proven to be equivalent to $UC_{\mathbb{N}^{\mathbb{N}}}$ or $C_{\mathbb{N}^{\mathbb{N}}}$ in Section 6. The results about open determinacy can be seen as uniform versions of the study of the complexity of winning strategies in [2]. If no case is fixed, we arrive at Weihrauch degrees not previously studied. Some of their properties are exhibited in Section 7. Since the degrees studied in Section 7 are not very well behaved, we introduce the canonical principle $TC_{\mathbb{N}^{\mathbb{N}}}$, the total continuation of closed choice in Section 8. We prove that up to finite parallelization, it is equivalent to the two-sided versions of open determinacy and the perfect tree theorem, and show some additional properties of the degree. Some concluding remarks and open questions are found in Section 9.

The following illustrates the strength of key benchmark principles in this article:

$$UC_{\mathbb{N}^{\mathbb{N}}} <_w \Sigma_1^1\text{-WKL} <_w C_{\mathbb{N}^{\mathbb{N}}} <_w TC_{\mathbb{N}^{\mathbb{N}}} <_w \widehat{TC_{\mathbb{N}^{\mathbb{N}}}} <_w \Pi_1^1\text{-CA}.$$

§2. Background on represented spaces and Weihrauch degrees. For background on the theory of represented spaces we refer to [38], for an introduction to and survey of Weihrauch reducibility we point the reader to [12].

As usual in the area, we use angle brackets to denote a variety of pairing and coding functions, such as those from $\mathbb{N} \times \mathbb{N}$ to \mathbb{N} , from $\mathbb{N}^{<\mathbb{N}}$ to \mathbb{N} , and from $\mathbb{N}^{\mathbb{N}} \times \mathbb{N}^{\mathbb{N}}$, $(\mathbb{N}^{\mathbb{N}})^{<\mathbb{N}}$ and $(\mathbb{N}^{\mathbb{N}})^{\mathbb{N}}$ to $\mathbb{N}^{\mathbb{N}}$. The context provides information about the one actually employed in any given instance.

2.1. Represented spaces.

DEFINITION 2.1. A *represented space* \mathbf{X} is a set X together with a partial surjection $\delta_{\mathbf{X}} : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow X$. If $x \in X$, any element of $(\delta_{\mathbf{X}})^{-1}(x)$ is called a *name* or a *code* for x .

A partial function $F : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ is called a *realizer* of a function $f : \subseteq \mathbf{X} \rightarrow \mathbf{Y}$ between represented spaces, if $f(\delta_{\mathbf{X}}(p)) = \delta_{\mathbf{Y}}(F(p))$ holds for all $p \in \text{dom}(f \circ \delta_{\mathbf{X}})$. We denote F being a realizer of f by $F \vdash f$. We then call $f : \subseteq \mathbf{X} \rightarrow \mathbf{Y}$ *computable* (respectively *continuous*), iff it has a computable (respectively continuous) realizer.

Represented spaces can adequately model most spaces of interest in *everyday mathematics*. For our purposes, we only need a few specific spaces that we discuss in the following, as well as some constructions of hyperspaces.

The category of represented spaces and continuous functions is cartesian-closed, by virtue of the UTM-theorem. Thus, for any two represented spaces \mathbf{X}, \mathbf{Y} we have a represented space $\mathcal{C}(\mathbf{X}, \mathbf{Y})$ of continuous functions from \mathbf{X} to \mathbf{Y} . The expected operations involving $\mathcal{C}(\mathbf{X}, \mathbf{Y})$ (evaluation, composition, (un)currying) are all computable. Using the Sierpiński space \mathbb{S} with underlying set $\{\top, \perp\}$ and representation $\delta_{\mathbb{S}} : \mathbb{N}^{\mathbb{N}} \rightarrow \{\top, \perp\}$ defined via $\delta_{\mathbb{S}}(\perp)^{-1} = \{0^\omega\}$, we can then define the represented space $\mathcal{O}(\mathbf{X})$ of *open* subsets of \mathbf{X} by identifying a subset of \mathbf{X} with its (continuous) characteristic function into \mathbb{S} . Since countable *or* and binary *and* on \mathbb{S} are computable, so are countable union and binary intersection of open sets. The space $\mathcal{A}(\mathbf{X})$ of closed subsets is obtained by taking formal complements, i.e. the names for $A \in \mathcal{A}(\mathbf{X})$ are the same as the names of $X \setminus A \in \mathcal{O}(\mathbf{X})$ (i.e. we are using the negative information representation).

We indicate with \mathbf{Tr} the space of trees on \mathbb{N} represented in an obvious way via characteristic functions on the set of finite sequences. The computable map $[] : \mathbf{Tr} \rightarrow \mathcal{A}(\mathbb{N}^{\mathbb{N}})$ maps a tree to its set of infinite paths, and has a computable multivalued inverse. In other words, one can compute a code of a tree T from a code of a closed set $[T]$, and vice versa.

Given a represented space \mathbf{X} and $k \in \mathbb{N}$, using Borel codes, the collections $\Sigma_k^0(\mathbf{X})$ (respectively $\Pi_k^0(\mathbf{X})$) of Σ_k^0 (respectively Π_k^0) subsets of \mathbf{X} can be naturally viewed as a represented space, cf. [3, 22, 39]. Equivalently, we can use the jumps of \mathbb{S} to characterize these spaces. We find that \mathcal{A} and Π_1^0 (respectively \mathcal{O} and Σ_1^0) are identical.

The collection $\Sigma_1^1(\mathbf{X})$ of analytic subsets of \mathbf{X} can also be represented in a straightforward manner: p is a name of a Σ_1^1 set $S \subseteq \mathbf{X}$ iff p is a name of a closed set $P \subseteq \mathbb{N}^{\mathbb{N}} \times \mathbf{X}$ such that $S = \{x \in \mathbf{X} : (\exists g) (g, x) \in P\}$. Equivalently [40, Proposition 35], we can define the space $\mathbb{S}_{\Sigma_1^1}$ by letting it have the underlying set $\{\top, \perp\}$, and letting $p \in \mathbb{N}^{\mathbb{N}}$ be a name for \top iff the tree on \mathbb{N} coded by p is ill-founded; and then identify $\Sigma_1^1(\mathbf{X})$ with $\mathcal{C}(\mathbf{X}, \mathbb{S}_{\Sigma_1^1})$ (here $f \in \mathcal{C}(\mathbf{X}, \mathbb{S}_{\Sigma_1^1})$ represents the $\Sigma_1^1(\mathbf{X})$ set $f^{-1}(\top)$).

Again, the collection $\Pi_1^1(\mathbf{X})$ of coanalytic subsets of \mathbf{X} is represented in an obvious way by taking formal complements. We define the space $\mathbb{S}_{\Pi_1^1}$ with underlying set $\{\top, \perp\}$, so that $p \in \mathbb{N}^{\mathbb{N}}$ is a name for \top iff the tree on \mathbb{N} coded by p is well-founded.

We first check that basic operations on these represented spaces are well-behaved.

LEMMA 2.2. *The following operations are computable:*

1. $\vee, \wedge : \mathbb{S}_{\Sigma_1^1}^{\mathbb{N}} \rightarrow \mathbb{S}_{\Sigma_1^1}$.
2. $\exists : \Sigma_1^1(\mathbf{X}) \rightarrow \mathbb{S}_{\Sigma_1^1}$, mapping nonempty sets to \top and the empty set to \perp .
3. $\text{id}, \neg : \mathbb{S} \rightarrow \mathbb{S}_{\Sigma_1^1}$.

PROOF. 1. For \vee , we need to show that given a sequence of trees we can compute a tree that is ill-founded iff one of the contributing trees is. This can be done by simply joining them at the root. For \wedge , we need a tree that is ill-founded iff all them are. For that, we can take the product of the trees (e.g. as in [34]).

2. From $f \in \mathcal{C}(\mathbb{N}^{\mathbb{N}}, \mathbb{S}_{\Sigma_1^1})$ we can compute by type-conversion some $g : \mathbb{N}^{\mathbb{N}} \times \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{S}$ such that $f(p) = \top$ iff $\exists q \in \mathbb{N}^{\mathbb{N}} g(p, q) = \perp$. But then $\exists p \in \mathbb{N}^{\mathbb{N}} f(p) = \top \Leftrightarrow \exists (p, q) \in \mathbb{N}^{\mathbb{N}} g(p, q) = \perp$, and we are done.

3. For $\neg : \mathbb{S} \rightarrow \mathbb{S}_{\Sigma_1^1}$, given a name p for a point in \mathbb{S} let the tree T be defined by $w \in T$ iff $\forall n \leq |w| p(n) = 0$. For $\text{id} : \mathbb{S} \rightarrow \mathbb{S}_{\Sigma_1^1}$, we let T have only branches of the form $n0^\omega$, and such a branch is present iff $p(n) \neq 0$.

□

PROPOSITION 2.3. *The following operations are computable for any represented space \mathbf{X} and $k > 0$:*

1. $\Sigma_1^1(\mathbf{X})^{\mathbb{N}} \rightarrow \Sigma_1^1(\mathbf{X}), (A_n)_n \mapsto \bigcup_{n \in \mathbb{N}} A_n$ (countable union);
2. $\Sigma_1^1(\mathbf{X})^{\mathbb{N}} \rightarrow \Sigma_1^1(\mathbf{X}), (A_n)_n \mapsto \bigcap_{n \in \mathbb{N}} A_n$ (countable intersection);
3. $\Sigma_1^1(\mathbf{X} \times \mathbf{Y}) \rightarrow \Sigma_1^1(\mathbf{Y}), A \mapsto \{y \in \mathbf{Y} \mid \exists x \in \mathbf{X} (x, y) \in A\}$;

- 4. $\Sigma_k^0(\mathbf{X}) \rightarrow \Sigma_1^1(\mathbf{X}), \Pi_k^0(\mathbf{X}) \rightarrow \Sigma_1^1(\mathbf{X}), \Sigma_k^0(\mathbf{X}) \rightarrow \Pi_1^1(\mathbf{X}), \Pi_k^0(\mathbf{X}) \rightarrow \Pi_1^1(\mathbf{X})$ (inclusions);
- 5. $\Sigma_k^0(\mathbb{N}^{\mathbb{N}} \times \mathbf{X}) \rightarrow \Sigma_1^1(\mathbf{X}), \Pi_k^0(\mathbb{N}^{\mathbb{N}} \times \mathbf{X}) \rightarrow \Sigma_1^1(\mathbf{X})$, such that

$$B \mapsto A = \{x \in X : \exists g \in \mathbb{N}^{\mathbb{N}}(g, x) \in B\};$$

- 6. $\Sigma_k^0(\mathbb{N}^{\mathbb{N}} \times \mathbf{X}) \rightarrow \Pi_1^1(\mathbf{X}), \Pi_k^0(\mathbb{N}^{\mathbb{N}} \times \mathbf{X}) \rightarrow \Pi_1^1(\mathbf{X})$, such that

$$B \mapsto A = \{x \in X : \forall g \in \mathbb{N}^{\mathbb{N}}(g, x) \in B\};$$

- 7. $\Pi_1^0(\mathbb{N}^{\mathbb{N}} \times \mathbf{X}) \rightarrow \Pi_1^1(\mathbf{X})$, such that

$$C \mapsto A = \{x \in X : \exists! g \in \mathbb{N}^{\mathbb{N}}(g, x) \in C\}.$$

PROOF. (1–6) These all follow directly from Lemma 2.2 together with function composition.

(7) It is well-known that $a \in \mathbb{N}^{\mathbb{N}}$ is hyperarithmetical relative to $\{a\} \in \Pi_1^0(\mathbb{N}^{\mathbb{N}})$ (cf. Corollary 3.3 and accompanying remarks below). The section map $(x, C) \mapsto \{y \in \mathbb{N}^{\mathbb{N}} \mid (y, x) \in C\} : \mathbf{X} \times \Pi_1^0(\mathbb{N}^{\mathbb{N}} \times \mathbf{X}) \rightarrow \Pi_1^0(\mathbb{N}^{\mathbb{N}})$ is computable, see [38, Proposition 4.2(9)]. Thus, we find that

$$A = \{x \in \mathbf{X} \mid \exists y \in \text{HYP}(x) (y, x) \in C\} \cap \{x \in \mathbf{X} \mid \forall y, z((y, x), (z, x) \in C \rightarrow y = z)\}.$$

The first set on the right-hand side is Π_1^1 by Kleene’s HYP-quantification theorem [27, 28] (see also [41, Lemma III.3.1]); that is, the formula $\exists y \in \text{HYP}(x) P(x, y)$ means that there are natural numbers a, e such that $a \in \mathcal{O}^x$ (which represents an ordinal α) and the e -th real $\Phi_e(x^{(\alpha)})$ computable in the α th Turing jump of x satisfies $P(x, \Phi_e(x^{(\alpha)}))$, where \mathcal{O}^x is Kleene’s system of ordinal notations relative to x (which is a $\Pi_1^1(x)$ set), cf. [41]. This description is trivially Π_1^1 , uniformly relative to x and the complexity of P , so that we can actually compute the Π_1^1 set from C . The second set explicitly and uniformly defines a Π_1^1 set. The claim thus follows using that intersection is a computable operation on Π_1^1 sets from (2). \dashv

LEMMA 2.4. *Let \mathbf{X} be a represented space. Then the function $F : \bigsqcup_k \Pi_k^0(\mathbb{N}^{\mathbb{N}} \times \mathbf{X}) \rightarrow \Sigma_1^1(\mathbf{X})$ defined by*

$$B \mapsto A = \{x \in \mathbf{X} : \exists g \in \mathbb{N}^{\mathbb{N}}(g, x) \in B\},$$

is computable.

PROOF. Proposition 2.3(5) is typically proved by induction on k , and the inductive argument is uniform in k . Since (a name for) $B \in \bigsqcup_k \Pi_k^0(\mathbb{N}^{\mathbb{N}} \times \mathbf{X})$ includes the information about the k such that $B \in \Pi_k^0(\mathbb{N}^{\mathbb{N}} \times \mathbf{X})$, we can uniformly repeat k steps of the induction argument to obtain a name for $\{x \in \mathbf{X} : \exists g \in \mathbb{N}^{\mathbb{N}}(g, x) \in B\}$ as a $\Sigma_1^1(\mathbf{X})$ set. \dashv

We define the represented spaces **LO** and **WO** respectively of linear orderings and countable well orderings with domain contained in \mathbb{N} (thus **WO** is a subspace of **LO**) as follows: p is a name for the linear order (X, \preceq_X) with $X \subseteq \mathbb{N}$ if $p(\langle n, m \rangle) = 1$ if and only if $n \preceq_X m$. We often abuse notation by leaving \preceq_X implicit and writing $X \in \mathbf{LO}$.

We may assume without loss of generality that, for all $X \in \mathbf{LO}$, $0 \notin X$ (this will be useful in Definition 5.1 below). If $X \in \mathbf{LO}$ we use interchangeably $\mathbf{WO}(X)$ and $X \in \mathbf{WO}$. If $X \in \mathbf{WO}$ we indicate its order type by $|X|$. Given some tree $T \subseteq \mathbb{N}^{<\mathbb{N}}$, we define the Kleene–Brouwer ordering \preceq_{KB} on T as the transitive closure of $w \preceq_{\text{KB}} u$ if $w \sqsupseteq u$ and $un \preceq_{\text{KB}} um$ if $n \leq m$. Using the coding of finite strings we view (T, \preceq_{KB}) as a member of \mathbf{LO} .

OBSERVATION 2.5. *The map $\text{KB} : \mathbf{Tr} \rightarrow \mathbf{LO}$ mapping a tree to its Kleene–Brouwer ordering is computable. We have $\mathbf{WO}(\text{KB}(T))$ iff T is well-founded.*

We need a technical definition, which can be found in [42, Definition V.6.4], for some of our proofs related to well orderings.

DEFINITION 2.6 (double descent tree). If $X, Y \in \mathbf{LO}$ the double descent tree $\mathbf{T}(X, Y)$ is the set of all finite sequences of the form $\langle (m_0, n_0), (m_1, n_1), \dots, (m_{k-1}, n_{k-1}) \rangle \in \mathbb{N}^{<\mathbb{N}}$ such that

- $m_0, m_1, \dots, m_{k-1} \in X$ and $m_0 >_X m_1 >_X \dots >_X m_{k-1}$,
- $n_0, n_1, \dots, n_{k-1} \in Y$ and $n_0 >_Y n_1 >_Y \dots >_Y n_{k-1}$.

We define the linear ordering $X * Y = \text{KB}(\mathbf{T}(X, Y))$.

OBSERVATION 2.7. $(X, Y) \mapsto (X * Y) : \mathbf{LO} \times \mathbf{LO} \rightarrow \mathbf{LO}$ is computable.

With an abuse of notation, we use \mathbb{Q} and \mathbb{N} to denote respectively a computable presentation of the standard linear ordering of rational numbers and of the well ordering of natural numbers.

LEMMA 2.8. *Let $X, Y \in \mathbf{LO}$.*

1. *If $\mathbf{WO}(X)$ then $X * Y$ and $Y * X$ are well orderings.*
2. *If $\mathbf{WO}(X)$ and $\neg \mathbf{WO}(Y)$, then $|X| \leq |X * Y|$.*
3. *If $\mathbf{WO}(Y)$, then $|X * Y| \leq |\mathbb{Q} * Y|$.*

PROOF. The proofs of 1 and 2 can be found in Lemma V.6.5 of [42]. In order to prove 3, consider a function $g : X \rightarrow \mathbb{Q}$ such that, for all $x, x' \in X$,

- (a) $x <_X x' \rightarrow g(x) <_{\mathbb{Q}} g(x')$,
- (b) $x <_{\mathbb{N}} x' \rightarrow g(x) <_{\mathbb{N}} g(x')$.

It is easy to see that such a function exists. Define then $\hat{g} : (X * Y) \rightarrow (\mathbb{Q} * Y)$ by putting $\hat{g}(\langle (x_0, y_0), \dots, (x_{k-1}, y_{k-1}) \rangle) := \langle (g(x_0), y_0), \dots, (g(x_{k-1}), y_{k-1}) \rangle$. Property a. of g guarantees that \hat{g} is well-defined and property b. implies that \hat{g} respects the Kleene–Brouwer orderings of the double descent trees $X * Y$ and $\mathbb{Q} * Y$. \dashv

2.2. Weihrauch reducibility. Intuitively, f being Weihrauch reducible to g means that there is an otherwise computable procedure to solve f by invoking an oracle for g exactly once. We thus obtain a very fine-grained picture of the relative strength of partial multivalued functions. Consequently, a Weihrauch equivalence is a very strong result compared to other approaches that allow more generous access to the principle being reduced to.

DEFINITION 2.9 (Weihrauch reducibility). Let f, g be multivalued functions on represented spaces. Then f is said to be *Weihrauch reducible* to g , in symbols $f \leq_w g$, if there are computable functions $K, H : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ such that $(p \mapsto K(p, GH(p))) \vdash f$ for all $G \vdash g$.

If there are computable functions $K, H : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ such that $KGH \vdash f$ for all $G \vdash g$, then f is *strongly Weihrauch reducible* to g , in symbols $f \leq_{sW} g$.

The relations \leq_w, \leq_{sW} are reflexive and transitive. We use $\equiv_w (\equiv_{sW})$ to denote equivalence and by $<_w$ we denote strict reducibility. Both Weihrauch degrees [36] and strong Weihrauch degrees [17] form lattices, the former being distributive and the latter not (in general, Weihrauch degrees behave more naturally than strong Weihrauch degrees).

Rather than the lattice operations, we will use two kinds of products in this work: The parallel product $f \times g$ is just the usual cartesian product of (multivalued) functions, which is readily seen to induce an operation on (strong) Weihrauch degrees. We call f a *cylinder*, if $f \equiv_{sW} (\text{id}_{\mathbb{N}^{\mathbb{N}}} \times f)$, and note that for cylinders, Weihrauch reducibility and strong Weihrauch reducibility coincide.

The compositional product $f \star g$ satisfies that

$$f \star g \equiv_w \max_{\leq_w} \{f_1 \circ g_1 \mid f_1 \leq_w f \wedge g_1 \leq_w g\}$$

and thus is the hardest problem that can be realized using first g , then something computable, and finally f . The existence of the maximum is shown in [15]. Both products as well as the lattice-join can be interpreted as logical *and*, albeit with very different properties. The sequential product \star is not commutative, however, it is the only one that admits a matching implication [15, 23].

Two further (unary) operations on Weihrauch degrees are relevant for us, finite parallelization f^* and parallelization \hat{f} . The former has as input a finite tuple of instances to f and needs to solve all of them, the latter takes and solves a countable sequences of instances. Both operations are closure operators in the Weihrauch lattice. They can be used to relax the requirement of using the oracle only once, if so desired, by looking at the relevant quotient lattices.

In passing, we will refer to the third operation, the jump from [11] (studied further in [4], denoted by f'). We use $f^{(n)}$ to denote the result of applying the jump n -times. The jump only preserves strong Weihrauch degrees. The input to f' is a sequence converging (with unknown speed) to an input of f , the output is whatever f would output on the limit.

The well-studied Weihrauch degrees most relevant for us are unique closed choice and closed choice (on Baire space), to which we dedicate the following Section 3. Two other degrees we will refer to are $\text{LPO} : \mathbb{N}^{\mathbb{N}} \rightarrow \{0, 1\}$ and $\text{lim} : \subseteq (\mathbb{N}^{\mathbb{N}})^{\omega} \rightarrow \mathbb{N}^{\mathbb{N}}$. These are defined via $\text{LPO}(p) = 1$ iff $p = 0^{\omega}$, and $\text{lim}((p_i)_{i \in \mathbb{N}}) = \lim_{i \rightarrow \infty} p_i$. They are related by $\widehat{\text{LPO}} \equiv_w \text{lim}$. The importance of lim is found partially in the observation from [3] that lim is complete for Baire class 1 functions, and more generally, that $\text{lim}^{(n)}$ is complete for Baire class $n + 1$ functions.

§3. $\text{UC}_{\mathbb{N}^{\mathbb{N}}}$ and $\text{C}_{\mathbb{N}^{\mathbb{N}}}$. The two Weihrauch degrees of central importance for this paper are unique closed choice and closed choice (on Baire space). These are defined as follows:

DEFINITION 3.1. Given a represented space \mathbf{X} , let $C_{\mathbf{X}} : \subseteq \mathcal{A}(\mathbf{X}) \rightrightarrows \mathbf{X}$ be defined via $x \in C_{\mathbf{X}}(A)$ iff $x \in A$ (thus, $A \in \text{dom}(C_{\mathbf{X}})$ iff $A \neq \emptyset$). Let $\text{UC}_{\mathbf{X}}$ be the restriction of $C_{\mathbf{X}}$ to singletons.

In particular, $\text{UC}_{\mathbf{X}}$ is capable of finding an element of a given Π_1^0 singleton in \mathbf{X} . In [39] Pauly introduced the notion of iterating a Weihrauch degree f over a given countable ordinal, this is denoted by f^\dagger . It is then shown that:

THEOREM 3.2 ([39, Theorem 80]). $\text{UC}_{\mathbb{N}^{\mathbb{N}}} \equiv_W \text{lim}^\dagger$.

One can read the above result as a very uniform version of the famous classical result that the Turing downward closures of Π_1^0 singletons in $\mathbb{N}^{\mathbb{N}}$ exhausts the hyperarithmetical hierarchy (cf. [41, Corollary II.4.3]).

Remark: Seeing that ATR_0 asserts the existence of Turing jumps iterated along some countable ordinal and since lim is equivalent to the Turing jump, it may seem as if this theorem already establishes that $\text{UC}_{\mathbb{N}^{\mathbb{N}}}$ is the Weihrauch degree corresponding to ATR_0 . There is a significant difference here though in what is meant by countable ordinal: In lim^\dagger , the input includes a code for something which is an ordinal in the surrounding meta-theory. In particular, any computable ordinal can be used *for free*. For ATR_0 the notion of countable ordinal is that of the model used. For example, an ill-founded computable linear order without hyperarithmetical descending chains (Kleene, see [41, Chapter 3, Lemma 2.1]) counts as an ordinal in the ω -model HYP consisting exactly of hyperarithmetical sets, and a similar phenomenon may happen in non- β -models of ATR_0 . Things get worse if non- ω -models are considered: ATR_0 (indeed, any sound c.e. theory, of course) fails to prove well-foundedness of some computable ordinals.

Note that lim^\dagger roughly corresponds to a (uniform) hyperarithmetical reduction, and therefore Theorem 3.2, for instance, implies the following:

COROLLARY 3.3. *Whenever $\{a\} \in \mathcal{A}(\mathbb{N}^{\mathbb{N}})$ is computable, then $a \in \mathbb{N}^{\mathbb{N}}$ is hyperarithmetical.*

COROLLARY 3.4. *If $f \leq_W \text{UC}_{\mathbb{N}^{\mathbb{N}}}$ for $f : \subseteq \mathbb{N}^{\mathbb{N}} \rightrightarrows \mathbf{X}$, then for every $x \in \text{dom}(f)$, $f(x)$ contains some y hyperarithmetical relative to x .*

Corollary 3.3 is a well-known classical fact saying that every Π_1^0 singleton is hyperarithmetical. Indeed, Spector showed that every Σ_1^1 singleton is hyperarithmetical (cf. [41, Theorem I.1.6]). Thus, it is natural to ask whether choice from Σ_1^1 singletons has exactly the same strength as $\text{UC}_{\mathbb{N}^{\mathbb{N}}}$.

One can generalize Definition 3.1 to any $\Gamma \in \{\Sigma_k^i, \Pi_k^i, \Delta_k^i\}$ in a straightforward manner: Let $\Gamma\text{-C}_{\mathbf{X}} : \subseteq \Gamma(\mathbf{X}) \rightrightarrows \mathbf{X}$ be defined via $x \in \Gamma\text{-C}_{\mathbf{X}}(A)$ iff $x \in A$. In other words, any realizer of $\Gamma\text{-C}_{\mathbf{X}}$ sends a code of a Γ -definition of A to a name of an element of A . Let $\Gamma\text{-UC}_{\mathbf{X}}$ be the restriction of $\Gamma\text{-C}_{\mathbf{X}}$ to singletons. For instance, a realizer for Σ_1^1 -unique choice $\Sigma_1^1\text{-UC}_{\mathbb{N}^{\mathbb{N}}} : \subseteq \Sigma_1^1(\mathbb{N}^{\mathbb{N}}) \rightarrow \mathbb{N}^{\mathbb{N}}$ is a partial function which, given a Σ_1^1 -code of a singleton $\{x\} \subseteq \mathbb{N}^{\mathbb{N}}$, returns a name of its unique element x . We will see below (in Theorem 3.11) that $\Sigma_1^1\text{-UC}_{\mathbb{N}^{\mathbb{N}}} \equiv_W \text{UC}_{\mathbb{N}^{\mathbb{N}}}$.

We now explore the strength of $\text{C}_{\mathbb{N}^{\mathbb{N}}}$.

THEOREM 3.5 (Kleene [27]). *There exists computable nonempty $A \in \mathcal{A}(\mathbb{N}^{\mathbb{N}})$ containing no hyperarithmetical point.*

That is, there is a nonempty Π_1^0 set $A \subseteq \mathbb{N}^{\mathbb{N}}$ with no hyperarithmetical element. This shows that $C_{\mathbb{N}^{\mathbb{N}}}$ has a computable instance with no hyperarithmetical solution. Let $NHA : \mathbb{N}^{\mathbb{N}} \rightrightarrows \mathbb{N}^{\mathbb{N}}$ be defined via $q \in NHA(p)$ iff q is not hyperarithmetical relative to p .

COROLLARY 3.6. $NHA \not\leq_W UC_{\mathbb{N}^{\mathbb{N}}}$ but $NHA \leq_W C_{\mathbb{N}^{\mathbb{N}}}$.

We now get the separation between $UC_{\mathbb{N}^{\mathbb{N}}}$ and $C_{\mathbb{N}^{\mathbb{N}}}$.

COROLLARY 3.7. $UC_{\mathbb{N}^{\mathbb{N}}} <_W C_{\mathbb{N}^{\mathbb{N}}}$.

There are a number of variants of unique choice, comprehension and separation that are all equivalent to $UC_{\mathbb{N}^{\mathbb{N}}}$ w.r.t. Weihrauch reducibility. We explore some of these next:

DEFINITION 3.8 (Σ_1^1 -Separation). Let $\Sigma_1^1\text{-Sep} := \subseteq (\mathbf{Tr} \times \mathbf{Tr})^{\mathbb{N}} \rightrightarrows 2^{\mathbb{N}}$ be the multivalued function with $\text{dom}(\Sigma_1^1\text{-Sep}) = \{(S_n, T_n)_{n \in \mathbb{N}} : \forall n ([S_n] = \emptyset \vee [T_n] = \emptyset)\}$ that maps any sequence $(S_n, T_n)_{n \in \mathbb{N}}$ in the domain to the set

$$\{f \in 2^{\mathbb{N}} : \forall n (([S_n] \neq \emptyset \rightarrow f(n) = 0) \wedge ([T_n] \neq \emptyset \rightarrow f(n) = 1))\}.$$

One can introduce a similar multivalued function by directly using the space $\Sigma_1^1(\mathbb{N}) \times \Sigma_1^1(\mathbb{N})$ instead of $(\mathbf{Tr} \times \mathbf{Tr})^{\mathbb{N}}$ without affecting the Weihrauch degree.

DEFINITION 3.9 (Δ_1^1 -Comprehension). Let $\Delta_1^1\text{-CA} := \subseteq (\mathbf{Tr} \times \mathbf{Tr})^{\mathbb{N}} \rightarrow 2^{\mathbb{N}}$ be the restriction of $\Sigma_1^1\text{-Sep}$ to the set $\{(S_n, T_n)_{n \in \mathbb{N}} : \forall n ([S_n] = \emptyset \leftrightarrow [T_n] \neq \emptyset)\}$. Let $\Delta_1^1\text{-CA}^-$ be the restriction of $\Delta_1^1\text{-CA}$ to the set $\{(S_n, T_n)_{n \in \mathbb{N}} : \forall n |[S_n]| + |[T_n]| = 1\}$.

DEFINITION 3.10 (Weak Σ_1^1 -Comprehension). Let $\Sigma_1^1\text{-CA}^- := \mathbf{Tr}^{\mathbb{N}} \rightarrow 2^{\mathbb{N}}$ be the function with domain $\text{dom}(\Sigma_1^1\text{-CA}^-) = \{(T_n)_{n \in \mathbb{N}} : \forall n |[T_n]| \leq 1\}$ and that maps $(T_n)_{n \in \mathbb{N}}$ to the unique $f \in 2^{\mathbb{N}}$ such that $f(n) = 1 \leftrightarrow |[T_n]| = 1$ for all $n \in \mathbb{N}$.

THEOREM 3.11. *The following are strongly Weihrauch equivalent:*

1. $UC_{\mathbb{N}^{\mathbb{N}}}$.
2. $\Sigma_1^1\text{-}UC_{\mathbb{N}^{\mathbb{N}}}$.
3. $\Sigma_1^1\text{-Sep}$.
4. $\Delta_1^1\text{-CA}$.
5. $\Delta_1^1\text{-CA}^-$.
6. $\Sigma_1^1\text{-CA}^-$.

PROOF. $(\Sigma_1^1\text{-}UC_{\mathbb{N}^{\mathbb{N}}} \leq_{sW} UC_{\mathbb{N}^{\mathbb{N}}})$: The proof of [39, Theorem 80] implicitly contains a proof of $\Sigma_1^1\text{-}UC_{\mathbb{N}^{\mathbb{N}}} \leq_{sW} \lim^\dagger$ (in the last paragraph). It is clear that $\Sigma_1^1\text{-}UC_{\mathbb{N}^{\mathbb{N}}} \equiv_{sW} \widehat{\Sigma_1^1\text{-}UC_{\mathbb{N}^{\mathbb{N}}}}$ and that $\widehat{UC_{\mathbb{N}^{\mathbb{N}}}} \equiv_{sW} UC_{\mathbb{N}^{\mathbb{N}}}$, so the claim follows with Theorem 3.2.

An alternative proof can be obtained by noting that the proof of $\text{UC}_{\mathbb{N}^{\mathbb{N}}} \leq_{\text{sw}} \Delta_1^1\text{-CA}^-$ given below is readily adapted to show that $\Sigma_1^1\text{-UC}_{\mathbb{N}^{\mathbb{N}}} \leq_{\text{sw}} \Delta_1^1\text{-CA}$ instead, and use the reductions below.

$(\text{UC}_{\mathbb{N}^{\mathbb{N}}} \leq_{\text{sw}} \Sigma_1^1\text{-UC}_{\mathbb{N}^{\mathbb{N}}})$: Trivial, as $\text{id} : \Pi_1^0(\mathbb{N}^{\mathbb{N}}) \rightarrow \Sigma_1^1(\mathbb{N}^{\mathbb{N}})$ is computable by Proposition 2.3(4).

$(\Sigma_1^1\text{-Sep} \leq_{\text{sw}} \text{UC}_{\mathbb{N}^{\mathbb{N}}})$: By [39, Proposition 62 and Lemma 79]. An alternative proof can be obtained by combining Lemmata 5.6 and 5.7 below.

$(\Delta_1^1\text{-CA} \leq_{\text{sw}} \Sigma_1^1\text{-Sep})$: The former is a restriction of the latter.

$(\Delta_1^1\text{-CA}^- \leq_{\text{sw}} \Delta_1^1\text{-CA})$: The former is a restriction of the latter.

$(\text{UC}_{\mathbb{N}^{\mathbb{N}}} \leq_{\text{sw}} \Delta_1^1\text{-CA}^-)$: Let $\{f\}$ be a singleton of $\mathbb{N}^{\mathbb{N}}$ given via some tree T such that $[T] = \{f\}$. From T we compute the double-sequence of trees $(T_t^0, T_t^1)_{t \in \mathbb{N}^{<\mathbb{N}}}$ such that: for all $t \in \mathbb{N}^{<\mathbb{N}}$,

- $T_t^0 = \{s \in T : t \sqsubseteq s \vee s \sqsubseteq t\}$,
- $T_t^1 = \{s \in T : t \not\sqsubseteq s\}$.

Note that, for all $t \in \mathbb{N}^{<\mathbb{N}}$, exactly one between T_t^0 and T_t^1 is ill-founded. In fact, if $t \sqsubseteq f$ then $f \in [T_t^0]$ and, since T has only one path, T_t^1 is well-founded. Otherwise, if $t \not\sqsubseteq f$ then $f \in [T_t^1]$ and $[T_t^0] = \emptyset$. Hence, we even have that for all $t \in \mathbb{N}^{<\mathbb{N}}$, $|[T_t^0]| + |[T_t^1]| = 1$.

Since we can identify $\mathbb{N}^{<\mathbb{N}}$ with \mathbb{N} we can consider $g = \Delta_1^1\text{-CA}^-((T_t^0, T_t^1)_{t \in \mathbb{N}^{<\mathbb{N}}})$. For all $t \in \mathbb{N}^{<\mathbb{N}}$, $g(t) = 0 \iff [T_t^0] \neq \emptyset \iff t \sqsubseteq f$. Therefore, given $n \in \mathbb{N}$, to compute $f(n)$ it suffices to wait for the first $t \in \mathbb{N}^{n+1}$ such that $g(t) = 0$ and then put $f(n) = t(n)$. This concludes the proof.

$(\Delta_1^1\text{-CA}^- \leq_{\text{sw}} \Sigma_1^1\text{-CA}^-)$: For every $(T_n^0, T_n^1)_{n \in \mathbb{N}} \in \text{dom}(\Delta_1^1\text{-CA}^-)$ we have that $\Delta_1^1\text{-CA}^-((T_n^0, T_n^1)_{n \in \mathbb{N}}) = \Sigma_1^1\text{-CA}^-((T_n^1)_{n \in \mathbb{N}})$.

$(\Sigma_1^1\text{-CA}^- \leq_{\text{sw}} \Sigma_1^1\text{-UC}_{\mathbb{N}^{\mathbb{N}}})$: Let $(T_n)_{n \in \mathbb{N}}$ be a sequence of trees in $\text{dom}(\Sigma_1^1\text{-CA}^-)$. We claim that using $\Sigma_1^1\text{-UC}_{\mathbb{N}^{\mathbb{N}}}$ we are able to compute $f \in 2^{\mathbb{N}}$ such that:

$$\forall n(f(n) = 1 \leftrightarrow |[T_n]| = 1). \tag{1}$$

In fact, (1) is equivalent to

$$\forall n[(f(n) = 0 \vee \exists g(g \in [T_n])) \wedge (\neg \exists! g(g \in [T_n]) \vee f(n) = 1)],$$

which in turn is equivalent to

$$\forall n[\exists g(f(n) = 0 \vee g \in [T_n]) \wedge \neg \exists! g(g \in [T_n] \wedge f(n) = 0)]. \tag{2}$$

Now, for each n , we can uniformly compute from $(T_n)_{n \in \mathbb{N}}$ a name for

$$\left\{ (g, f) \in \mathbb{N}^{\mathbb{N}} \times \mathbb{N}^{\mathbb{N}} : f(n) = 0 \vee g \in [T_n] \right\}$$

as a closed subset of $\mathbb{N}^{\mathbb{N}} \times \mathbb{N}^{\mathbb{N}}$, which entails that we can uniformly compute from $(T_n)_{n \in \mathbb{N}}$ a name for

$$\left\{ f \in \mathbb{N}^{\mathbb{N}} : \exists g(f(n) = 0 \vee g \in [T_n]) \right\}$$

as a $\Sigma_1^1(\mathbb{N}^{\mathbb{N}})$ set for each $n \in \mathbb{N}$. Furthermore, for each $n \in \mathbb{N}$, we can uniformly compute from $(T_n)_{n \in \mathbb{N}}$ a name for

$$\left\{ (g, f) \in \mathbb{N}^{\mathbb{N}} \times \mathbb{N}^{\mathbb{N}} : g \in [T_n] \wedge f(n) = 0 \right\}$$

as a closed set and hence a name for

$$\left\{ f \in \mathbb{N}^{\mathbb{N}} : \neg \exists! g (g \in [T_n] \wedge f(n) = 0) \right\}$$

as a $\Sigma_1^1(\mathbb{N}^{\mathbb{N}})$ set by Proposition 2.3(7).

Finally, since the operations of finite and countable intersection of Σ_1^1 sets are computable, we are able to uniformly compute from $(T_n)_{n \in \mathbb{N}}$ a name (by Proposition 2.3(2)) for the $\Sigma_1^1(\mathbb{N}^{\mathbb{N}})$ singleton

$$\left\{ f \in 2^{\mathbb{N}} : \forall n [\exists g (f(n) = 0 \vee g \in [T_n]) \wedge \neg \exists! g (g \in [T_n] \wedge f(n) = 0)] \right\}.$$

Clearly, applying $\Sigma_1^1\text{-UC}_{\mathbb{N}^{\mathbb{N}}}$ to such set we obtain the unique f satisfying (1), which is exactly $\Sigma_1^1\text{-CA}^-((T_n)_n)$. ⊢

Arithmetical transfinite recursion. As mentioned above, the operation lim^\dagger from [39] is the ordinal-iteration of the map lim . Here, we will explore a direct encoding of arithmetical transfinite recursion as a Weihrauch degree, and give another proof of its equivalence with $\text{UC}_{\mathbb{N}^{\mathbb{N}}}$. Let us fix an effective enumeration $\langle \phi_n : n \in \mathbb{N} \rangle$ of all the computable functions $\phi : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$. Note that $\widehat{\text{LPO}}^{(k)}$ is a complete Σ_{k+2}^0 -computable function, and thus one can think of $\theta_n^k = \widehat{\text{LPO}}^{(k)} \circ \phi_n$ as the n^{th} Σ_{k+2}^0 -computable function. Instead, we could have used the n^{th} Σ_{k+2}^0 formula to define an equivalent notion.

DEFINITION 3.12 (Arithmetical transfinite recursion). Let $\text{ATR} : \subseteq 2^{\mathbb{N}} \times \mathbf{WO} \times \mathbb{N}^2 \rightarrow 2^{\mathbb{N}}$ be the function which maps each $(Z, X, (k, n)) \in 2^{\mathbb{N}} \times \mathbf{WO} \times \mathbb{N}^2$ to the set $Y \in 2^{\mathbb{N}}$ such that, for all $(y, j) \in \mathbb{N}^2$,

$$(y, j) \in Y \leftrightarrow j \in X \wedge y \in \theta_n^k(Y^j \oplus Z),$$

where $Y^j = \{ \langle y, i \rangle \in Y : i <_X j \}$.

Compare Definition 3.12 with ATR_0 in reverse mathematics, cf. [42, Definition V.2.4]. Note that our ATR is a *single-valued* function since, as mentioned in the first remark in this section, our X is truly well ordered, and therefore, we do not need to consider pseudo-hierarchies.

THEOREM 3.13. $\text{ATR} \equiv_{sW} \text{UC}_{\mathbb{N}^{\mathbb{N}}}$.

PROOF. By Lemmata 3.14, 3.15 below and Theorem 3.11. ⊢

The following is an analog of the classical reverse mathematical fact [42, Theorem V.5.1].

LEMMA 3.14. $\text{ATR}_{\leq_{sW}} \Sigma_1^1\text{-Sep}$.

PROOF. It is easy to see that $\Sigma_1^1\text{-Sep}$ is a cylinder and hence it suffices to show $\text{ATR}_{\leq_W} \Sigma_1^1\text{-Sep}$. Given $(Z, X, \langle k, n \rangle) \in 2^{\mathbb{N}} \times \mathbf{WO} \times \mathbb{N}^2$, we want to compute $\text{ATR}(Z, X, \langle k, n \rangle)$ as defined in Definition 3.12. For each $j \in X$ and $Y \in 2^{\mathbb{N}}$, let us consider the following formula:

$$H(Y, j) \equiv \forall \langle y, i \rangle \in \mathbb{N}^2 [\langle y, i \rangle \in Y \iff i <_X j \wedge y \in \theta_n^k(Y^i \oplus Z)].$$

Essentially, $H(Y, j)$ says that Y is the set $\{\langle y, i \rangle \in \text{ATR}(Z, X, \langle k, n \rangle) : i <_X j\}$. Using now H , we define the following two formulas for each $j, z \in \mathbb{N}$:

$$\begin{aligned} \varphi_0(j, z) &\equiv j \in X \wedge \exists Y \in 2^{\mathbb{N}} [H(Y, j) \wedge z \in \theta_n^k(Y^j \oplus Z)], \\ \varphi_1(j, z) &\equiv j \in X \wedge \exists Y \in 2^{\mathbb{N}} [H(Y, j) \wedge z \notin \theta_n^k(Y^j \oplus Z)]. \end{aligned}$$

Note that, for each $j \in X$ and $z \in \mathbb{N}$ we have $\varphi_0(j, z) \iff \langle z, j \rangle \in \text{ATR}(Z, X, \langle k, n \rangle)$.

Using the function F defined in Lemma 2.4 and the closure properties of Proposition 2.3, we are able to compute two names for the $\Sigma_1^1(\mathbb{N}^2)$ -sets A_0 and A_1 corresponding to the formulas φ_0 and φ_1 . Note that in this case the use of F is required and we cannot appeal to Proposition 2.3(5) because k is not fixed but is given with the input. It is easy to see that A_0 and A_1 are disjoint; hence one can ask $\Sigma_1^1\text{-Sep}$ to give us f separating A_0 from A_1 , which is clearly a solution of $\text{ATR}(Z, X, \langle k, n \rangle)$. Here are the details:

Since the names for A_0 and A_1 are $\Pi_1^0(\mathbb{N}^{\mathbb{N}} \times \mathbb{N}^2)$ -names, it is not difficult to see that we can build a double sequence of trees $(T_{\langle j, z \rangle}^0, T_{\langle j, z \rangle}^1)_{j, z \in \mathbb{N}}$ such that, for each $j \in \mathbb{N}$ and $z \in \mathbb{N}$,

- $\langle j, z \rangle \in A_0 \iff [T_{\langle j, z \rangle}^0] \neq \emptyset$,
- $\langle j, z \rangle \in A_1 \iff [T_{\langle j, z \rangle}^1] \neq \emptyset$.

Note that, if $j \notin X$ then for each $z \in \mathbb{N}$, $\neg\varphi_0(j, z)$ and $\neg\varphi_1(j, z)$, which means that $[T_{\langle j, z \rangle}^0] = [T_{\langle j, z \rangle}^1] = \emptyset$. If instead $j \in X$ we have, for each $z \in \mathbb{N}$, $\varphi_0(j, z) \iff \neg\varphi_1(j, z)$ which implies $[T_{\langle j, z \rangle}^0] \neq \emptyset \iff [T_{\langle j, z \rangle}^1] = \emptyset$. Therefore the double-sequence of trees $(T_{\langle j, z \rangle}^0, T_{\langle j, z \rangle}^1)_{j, z \in \mathbb{N}}$ belongs to the domain of $\Sigma_1^1\text{-Sep}$. So let $f \in \Sigma_1^1\text{-Sep}(T_{\langle j, z \rangle}^0, T_{\langle j, z \rangle}^1)_{j, n \in \mathbb{N}}$. Now we have, for each $j \in X$ and $z \in \mathbb{N}$, $f(j, z) = 0 \iff [T_{\langle j, z \rangle}^0] \neq \emptyset \iff \varphi_0(j, z) \iff \langle z, j \rangle \in \text{ATR}(Z, X, \langle k, n \rangle)$, i.e. we are able to compute $\text{ATR}(Z, X, \langle k, n \rangle) \in 2^{\mathbb{N}}$ using f .

Note that we are using the original input to test whether $j \in X$. ⊣

LEMMA 3.15. $\Delta_1^1\text{-CA}_{\leq_{sW}} \text{ATR}$.

PROOF. Let $(T_n^0, T_n^1)_{n \in \mathbb{N}} \in \text{dom}(\Delta_1^1\text{-CA})$, we want to compute $f \in 2^{\mathbb{N}}$ such that, for all $n \in \mathbb{N}$, $f(n) = 0 \iff [T_n^0] \neq \emptyset$. In order to apply ATR we have to specify a set parameter Z , a well ordering X and an arithmetical formula. The role of Z in this case will be played by $(T_n^0, T_n^1)_{n \in \mathbb{N}}$. The well ordering X is obtained as $\sum_{n \in \mathbb{N}} (\text{KB}(T_n^0) * \text{KB}(T_n^1)) + 1$ (which is a well ordering by Lemma 2.8(1)).

It remains to specify an arithmetical formula $\varphi(y, Y^j \oplus Z)$ which describes what to do at each step of the recursion. We read both Y^j and Z as coding a sequence of

pairs of trees. The idea is to eliminate at each step the leaves of all the trees in the sequence. Thus, $\varphi(y, Y^j \oplus Z)$ holds if either $Y^j = \emptyset$ and y codes a vertex with a child in Z , or y codes a vertex with a child in each tree from Y^j . This is easily verified to be an arithmetical formula, and hence can be coded as some θ_n^k .³

Finally, consider $Y = \text{ATR}((T_n^0, T_n^1)_n, X, \langle k, n \rangle)$, which is the set we obtain after repeating, along the well ordering X , the procedure of eliminating leaves from the trees T_n^0 and T_n^1 . Now, let fix n and consider $i \in \{0, 1\}$ such that T_n^i is well founded. Note that, in order to eliminate all the tree T_n^i , the recursion should be done at least over the ordinal rank(T_n^i). In our case, the recursion is done over X whose order type is greater than the order type of $\text{KB}(T_n^i)$ which in turn is greater than $\text{rank}(T_n^i)$, cf. Lemma 2.8(2). This means that Y does not contain any element of the tree T_n^i . This argument applies to each well founded tree in the sequence $(T_n^0, T_n^1)_n$, so we can know whether a tree in the sequence has a path or not simply by checking if its root is in Y . It is easy to see that this allows us to compute $\Delta_1^1\text{-CA}((T_n^0, T_n^1)_{n \in \mathbb{N}})$. ⊣

§4. Σ_1^1 -weak König’s lemma.

4.1. Σ_1^1 versus Π_1^1 . In this section, we focus on the following contrast between reverse mathematics and the Weihrauch lattice regarding Σ_1^1 and Π_1^1 -separation: On the one hand, in reverse mathematics, we have

$$\Pi_1^1\text{-SEP}_0 < \Sigma_1^1\text{-SEP}_0 \tag{3}$$

where $A < B$ indicates $\text{RCA}_0 \vdash B \rightarrow A$, but $\text{RCA}_0 \not\vdash A \rightarrow B$. On the other hand, in the Weihrauch lattice, we have

$$\Sigma_1^1\text{-Sep} <_w \Pi_1^1\text{-Sep}. \tag{4}$$

The former inequality (3) was proven by Montalbán [31] using Steel’s tagged tree forcing. The latter inequality (4) follows from the well-known fact in descriptive set theory that Σ_1^1 has the Δ_1^1 -separation property, while Π_1^1 does not (see also Lemma 4.4). It is not hard to explain the cause of the contrast between (3) and (4), namely the Spector–Gandy phenomenon.

Let \mathcal{M} be an ω -model, and let $(\Sigma_1^1)^\mathcal{M}$ be the collection of all subsets of ω which are Σ_1^1 -definable within \mathcal{M} , that is, $(\Sigma_1^1)^\mathcal{M} = \{\{n \in \omega : \mathcal{M} \models \varphi(n)\} : \varphi \in \Sigma_1^1\}$. We define $(\Pi_1^1)^\mathcal{M}$ analogously. Consider the ω -model HYP consisting of all hyperarithmetical reals. The Spector–Gandy theorem (cf. [41, Theorem III.3.5 and Lemma III.3.1] or [42, Theorems VIII.3.20 and VIII.3.27]) implies that

$$(\Sigma_1^1)^{\text{HYP}} = \Pi_1^1, \text{ and } (\Pi_1^1)^{\text{HYP}} = \Sigma_1^1.$$

The roles of Σ_1^1 and Π_1^1 are interchanged! We should always be careful about this role-exchange phenomenon of Σ_1^1 and Π_1^1 when comparing reverse math and computability theory. Of course, the notion of a β -model solves this role-exchange

³Similar ideas are found in the investigation of the Weihrauch degree of the *pruning derivative* of a tree in [34].

problem. To be precise, a β -model (see [42, Section VII]) is an ω -model \mathcal{M} satisfying the following condition:

$$(\Sigma_1^1)^{\mathcal{M}} = \Sigma_1^1, \text{ and } (\Pi_1^1)^{\mathcal{M}} = \Pi_1^1.$$

However, the notion of a β -model is obviously related to closed choice $C_{\mathbb{N}^{\mathbb{N}}}$: An ω -model \mathcal{M} is a β -model iff, for any $Z \in \mathcal{M}$ and nonempty $\Pi_1^0(Z)$ set $P \subseteq \mathbb{N}^{\mathbb{N}}$, some $\alpha \in P$ belongs to \mathcal{M} . Therefore, when studying principles weaker than $C_{\mathbb{N}^{\mathbb{N}}}$, we cannot work within the β -models.

Now, how should we interpret the reverse-mathematical Σ_1^1 -separation principle in our real universe? *The* right answer may not exist. It may be Π_1^1 -Sep or may be Σ_1^1 -Sep.

We have already examined the strength of the Σ_1^1 -separation principle Σ_1^1 -Sep. In this section, we will investigate the Π_1^1 -separation principle, Π_1^1 -Sep, in the Weihrauch lattice. In reverse mathematics, Montalbán [31] showed that the strength of the Π_1^1 -separation principle is strictly between $\Delta_1^1\text{-CA}_0$ and ATR_0 (⁴):

$$\Delta_1^1\text{-CA}_0 < \Pi_1^1\text{-SEP}_0 < \text{ATR}_0 \equiv \Sigma_1^1\text{-SEP}_0.$$

Moreover, $\Delta_1^1\text{-CA}_0$ and $\Pi_1^1\text{-SEP}_0$ are theories of hyperarithmetic analysis, that is, for every $Z \subseteq \omega$, $\text{HYP}(Z)$ is the least ω -model of that theory containing Z . On the other hand, $\text{HYP} \not\equiv \text{ATR}_0$. In contrast, we will see the following:

$$\text{UC}_{\mathbb{N}^{\mathbb{N}}} \equiv \Delta_1^1\text{-CA} \equiv \text{w ATR} \equiv \text{w } \Sigma_1^1\text{-Sep} < \text{w } \Pi_1^1\text{-Sep} < \text{w } C_{\mathbb{N}^{\mathbb{N}}}.$$

4.2. The strength of Σ_1^1 -weak König’s lemma. The principle of Π_1^0 -separation was studied already in the precursor works by Weihrauch [44], and Weak König’s Lemma (aka closed choice on Cantor space) was a focus in the earliest work on Weihrauch reducibility in the modern understanding [20, 7, 5]. Here, we explore their higher-level analogues.

Let Π_1^1 -Sep be the following partial multivalued function: Given Π_1^1 -codes of sets $A, B \subseteq \mathbb{N}$, if A and B are disjoint, then return a set $C \subseteq \mathbb{N}$ separating A from B , that is, $A \subseteq C$ and $B \cap C = \emptyset$. To be more precise:

DEFINITION 4.1. Let $\Pi_1^1\text{-Sep} : \subseteq \Pi_1^1(\mathbb{N}) \times \Pi_1^1(\mathbb{N}) \rightrightarrows 2^{\mathbb{N}}$ be such that $C \in \Pi_1^1\text{-Sep}(A, B)$ iff C separates A from B , where $(A, B) \in \text{dom}(\Pi_1^1\text{-Sep})$ iff $A \cap B = \emptyset$.

We also consider Σ_1^1 -weak König’s lemma Σ_1^1 -WKL: Given a Σ_1^1 -code of a set $T \subseteq 2^{<\omega}$, if T is an infinite binary tree, then return a path through T . Formally speaking:

DEFINITION 4.2. Let $\Sigma_1^1\text{-WKL} : \subseteq \Sigma_1^1(2^{<\omega}) \rightrightarrows 2^{\mathbb{N}}$ be such that $p \in \Sigma_1^1\text{-WKL}(T)$ iff p is an infinite path through T , where $T \in \text{dom}(\Sigma_1^1\text{-WKL})$ iff T is an infinite binary tree.

While Σ_1^1 -WKL appears as a Σ_1^1 -version of closed choice on Cantor space, it is not equivalent to Σ_1^1 -choice on $2^{\mathbb{N}}$ (nor, equivalently, closed choice on $\mathbb{N}^{\mathbb{N}}$). Instead, it is

⁴Actually, Montalbán showed that Π_1^1 -separation is strictly weaker than Σ_1^1 -AC.

equivalent to the parallelization $\widehat{\Sigma_1^1-C_2}$ of Σ_1^1 choice on the discrete space $\mathbf{2} = \{0, 1\}$. We will show the following.

THEOREM 4.3. $UC_{\mathbb{N}^{\mathbb{N}}} <_W \widehat{\Sigma_1^1-C_2} \equiv_W \Pi_1^1\text{-Sep} \equiv_W \Sigma_1^1\text{-WKL} <_W \widehat{\Sigma_1^1-C_{\mathbb{N}}} \leq_W C_{\mathbb{N}^{\mathbb{N}}}$.

We will use the following fundamental notion in HYP-theory. A Π_1^1 -norm on a Π_1^1 set $P \subseteq \mathbb{N}$ is a map $\varphi : \mathbb{N} \rightarrow \omega_1^{CK} \cup \{\infty\}$ such that $P = \{n : \varphi(n) < \infty\}$ and that the following relations \leq_φ and $<_\varphi$ are Π_1^1 :

$$\begin{aligned} a \leq_\varphi b &\iff \varphi(a) < \infty \text{ and } \varphi(a) \leq \varphi(b), \\ a <_\varphi b &\iff \varphi(a) < \infty \text{ and } \varphi(a) < \varphi(b). \end{aligned}$$

It is well-known that every Π_1^1 set admits a Π_1^1 -norm (in an effective manner): Consider a many-one reduction from a Π_1^1 set P to the set WO of well orderings. We will explore the uniform complexity of this kind of stage comparison principle in Section 5.

One can easily separate unique choice on $\mathbb{N}^{\mathbb{N}}$ and the Π_1^1 -separation principle by considering the *diagonally nonhyperarithmetical* functions, which is a HYP version of DNC_2 (known as diagonally noncomputable functions). A very basic fact in HYP-theory is the existence of a computable enumeration $(\psi_e)_{e \in \mathbb{N}}$ of all partial Π_1^1 functions on \mathbb{N} . For instance, let ψ_e be a standard Π_1^1 -uniformization of the e^{th} Π_1^1 set $P_e \subseteq \mathbb{N} \times \mathbb{N}$, that is, $\psi_e(n)$ is an element in the n^{th} section of P_e attaining the smallest φ -value if it exists, where φ is a Π_1^1 -norm on P_e .

LEMMA 4.4. $UC_{\mathbb{N}^{\mathbb{N}}} <_W \Pi_1^1\text{-Sep}$.

PROOF. To see that $UC_{\mathbb{N}^{\mathbb{N}}} \leq_W \Pi_1^1\text{-Sep}$, note that $UC_{\mathbb{N}^{\mathbb{N}}} \equiv_W \Delta_1^1\text{-CA}$ by Theorem 3.11, and $\Delta_1^1\text{-CA} \leq_W \Pi_1^1\text{-Sep}$ is straightforward. For the separation, let $(\psi_e)_{e \in \mathbb{N}}$ be an enumeration of all partial Π_1^1 functions on \mathbb{N} as above. For $i < 2$, consider $P_i = \{e \in \mathbb{N} : \psi_e(e) \downarrow = i\}$. Clearly P_i is Π_1^1 , and $P_0 \cap P_1 = \emptyset$. It is easy to see that there is no Δ_1^1 set separating P_0 and P_1 . \dashv

The proof of Lemma 4.4 motivates us to introduce the following multivalued function $\Pi_1^1\text{-DNC}_2 : 2^{\mathbb{N}} \rightrightarrows 2^{\mathbb{N}}$: Given an oracle X , return a two-valued X -diagonally nonhyperarithmetical function f , that is, $f \in \Pi_1^1\text{-DNC}_2(X)$ iff, whenever $\psi_e^X(e) \downarrow$, $f(e) \neq \psi_e^X(e)$, where $(\psi_e^X)_{e \in \mathbb{N}}$ is a canonical enumeration of all partial $\Pi_1^1(X)$ functions on \mathbb{N} . The following is an analog of the well-known fact that every DNC_2 -function has a PA-degree.

PROPOSITION 4.5. $\Pi_1^1\text{-Sep} \equiv_W \Pi_1^1\text{-DNC}_2$.

PROOF. Let P_0 and P_1 be disjoint Π_1^1 sets. Clearly there is e such that $n \in P_i$ iff $\psi_e(n) \downarrow = i$. By the recursion theorem, one can uniformly find a computable function r such that $\psi_{r(n)}(r(n)) \simeq \psi_e(n)$. Let f be a diagonally nonhyperarithmetical function. If $f(r(n)) = i$ then $\psi_{r(n)}(r(n)) \simeq \psi_e(n) \neq i$, which implies $n \notin P_i$. Therefore, $S = \{n : f(r(n)) = 1\}$ separates P_0 from P_1 . This argument is easily relativizable uniformly. The converse direction is also clear. \dashv

Using a Π_1^1 -norm, one can show $\Sigma_1^1\text{-WKL} \equiv_W \Pi_1^1\text{-Sep}$ by modifying the usual proof of the well-known equivalence between WKL and $\Sigma_1^0\text{-Sep}$.

LEMMA 4.6. $\Sigma_1^1\text{-WKL} \equiv_W \Pi_1^1\text{-Sep} \equiv_W \widehat{\Sigma_1^1\text{-C}_2}$.

PROOF. By a straightforward modification of the usual proof of $\Sigma_1^0\text{-Sep} \equiv_W \widehat{\text{C}_2}$, it is easy to see that $\Pi_1^1\text{-Sep} \equiv_W \widehat{\Sigma_1^1\text{-C}_2}$ holds. It is also clear that $\Pi_1^1\text{-Sep} \leq_W \Sigma_1^1\text{-WKL}$. Thus, it suffices to show that $\Sigma_1^1\text{-WKL} \leq_W \Pi_1^1\text{-Sep}$.

Given a Σ_1^1 -tree $T \subseteq 2^{<\omega}$, let $\text{Ext}_T \subseteq 2^{<\omega}$ be the set of all extendible nodes of T . Clearly, its complement $\neg\text{Ext}_T = 2^{<\omega} \setminus \text{Ext}_T$ is Π_1^1 , and thus admits a Π_1^1 -norm φ (we need to get φ in a uniform way, but it is straightforward). Consider the Π_1^1 set $P_i = \{\sigma : \sigma \frown i <_\varphi \sigma \frown (1-i)\}$ for each $i < 2$. Obviously, $P_0 \cap P_1 = \emptyset$. We claim that

$$\sigma \in \text{Ext}_T \text{ and } \sigma \notin P_j \implies \sigma \frown j \in \text{Ext}_T.$$

If $\sigma \notin P_j$ then $\sigma \frown j \not<_\varphi \sigma \frown (1-j)$, that is, either $\varphi(\sigma \frown j) = \infty$ or $\varphi(\sigma \frown (1-j)) \leq \varphi(\sigma \frown j)$ holds. If the former holds then we must have $\sigma \frown j \in \text{Ext}_T$. If $\varphi(\sigma \frown j) < \infty$, then we must have $\varphi(\sigma \frown (1-j)) = \infty$ since $\sigma \in \text{Ext}_T$ implies that $\sigma \frown i \in \text{Ext}_T$ for some $i < 2$. By the latter condition, $\infty = \varphi(\sigma \frown (1-j)) \leq \varphi(\sigma \frown j)$; hence $\varphi(\sigma \frown j)$ must be ∞ . In any case, we have $\varphi(\sigma \frown j) = \infty$, which means that $\sigma \frown j \in \text{Ext}_T$. This verifies the above claim.

Let S be such that $P_0 \subseteq S$ and $S \cap P_1 = \emptyset$. Let σ_0 be the empty string, and put $\sigma_{n+1} = \sigma_n \frown S(\sigma_n)$. Then, by the above claim, we have $\sigma_n \in \text{Ext}_T$ for any n , and therefore $\bigcup_n \sigma_n \in [T]$. One can easily relativize this argument uniformly. \dashv

LEMMA 4.7. $\Sigma_1^1\text{-WKL} <_W \widehat{\Sigma_1^1\text{-C}_\mathbb{N}}$.

PROOF. By Lemma 4.6, we have $\Sigma_1^1\text{-WKL} \leq_W \widehat{\Sigma_1^1\text{-C}_\mathbb{N}}$. It remains to show that $\widehat{\Sigma_1^1\text{-C}_\mathbb{N}} \not\leq_W \Sigma_1^1\text{-WKL}$. It is easy to see that $\Sigma_1^1\text{-WKL}$ is a cylinder, and hence it suffices to show that $\widehat{\Sigma_1^1\text{-C}_\mathbb{N}} \not\leq_{sW} \Sigma_1^1\text{-WKL}$.

We first show the following claim: Let $T \subseteq 2^{<\omega}$ be a Σ_1^1 tree, and Φ a Turing functional such that for every $x \in [T]$, Φ^x is total. Then there exists a Δ_1^1 function $h : \mathbb{N} \rightarrow \mathbb{N}$ majorizing $n \mapsto \Phi^x(n)$ for every $x \in [T]$.

Let $g : \mathbb{N} \rightarrow \mathbb{N}$ be a function such that for any n , if $|\sigma| = g(n)$ then either $\sigma \notin \text{Ext}_T$ or $\Phi^\sigma(n) \downarrow$. This condition is clearly Π_1^1 , and by compactness, g is total. Hence, g is a total Π_1^1 function, and thus actually Δ_1^1 . Then define $h(n) = \max\{\Phi^\sigma(n) : |\sigma| = g(n) \text{ and } \Phi^\sigma(n) \downarrow\}$. Clearly h is Δ_1^1 and $\Phi^x(n) \leq h(n)$ for any $x \in [T]$. This verifies the claim.

Let $(\psi_e)_{e \in \omega}$ be a computable enumeration of partial Π_1^1 functions on \mathbb{N} . Let S_e be the set of all k such that

$$(\forall n \leq e)(\psi_n(e) \downarrow \implies \psi_n(e) < k).$$

Clearly S_e is Σ_1^1 and cofinite. Then every element of $S = \prod_e S_e$ dominates all Δ_1^1 functions. If $\widehat{\Sigma_1^1\text{-C}_\mathbb{N}} \leq_{sW} \Sigma_1^1\text{-WKL}$ then we must have a Σ_1^1 -tree $T \subseteq 2^{<\omega}$ whose paths compute uniformly an element of S , which is impossible by the above claim. \dashv

Recall that $A \star B$ denotes the sequential composition of A and B , cf. [15], that is, a function attaining the greatest Weihrauch degree among $\{g \circ f : g \leq_w A \text{ and } f \leq_w B\}$.

PROPOSITION 4.8. $\Sigma_1^1\text{-WKL} \star \Sigma_1^1\text{-WKL} \equiv_W \Sigma_1^1\text{-WKL}$.

PROOF. This is a modification of the independent choice theorem from [5]. We can assume that the inputs to $\Sigma_1^1\text{-WKL} \star \Sigma_1^1\text{-WKL}$ are a computable function f , $z \in 2^{\mathbb{N}}$ as well as (relativizable) Σ_1^1 trees S and T . Then, $\{x \oplus y : x \in [S^z] \text{ and } y \in [T^{f(z,x)}]\}$ is a Σ_1^1 closed set, and any of its elements is a solution to $\Sigma_1^1\text{-WKL} \star \Sigma_1^1\text{-WKL}$. \dashv

There is a natural principle between $\text{UC}_{\mathbb{N}^{\mathbb{N}}}$ and $\Sigma_1^1\text{-WKL}$. Let us define $\Sigma_1^1\text{-weak weak König's lemma}$ $\Sigma_1^1\text{-WWKL}$ as follows: Given a Σ_1^1 set $T \subseteq 2^{<\omega}$, if T is an infinite binary tree and if $[T]$ has a positive measure, then return a path through T . This is in analogy to the usual weak König's lemma, whose Weihrauch degree was studied in [14, 8, 10].

Note that Hjorth and Nies (see [33, Chapter 9.2]) showed that there is a Σ_1^1 -closed set consisting of Π_1^1 -Martin-Löf random reals. Indeed, the proof shows that $\Pi_1^1\text{-MLR}$ is Weihrauch reducible to $\Sigma_1^1\text{-WWKL}$, where $\Pi_1^1\text{-MLR}$ is a multivalued functions representing Π_1^1 -Martin-Löf randomness, which is introduced in a straightforward manner. We also have $\text{WKL} \not\leq_w \Sigma_1^1\text{-WWKL}$ since the Turing upward closure of any nontrivial separating class has measure zero (cf. [25, Theorem 5.3]). We show that, even if we enhance $\Sigma_1^1\text{-WWKL}$ by adding a hyperarithmetical power, its strength is strictly weaker than $\Sigma_1^1\text{-WKL}$:

THEOREM 4.9. $\text{UC}_{\mathbb{N}^{\mathbb{N}}} <_W \text{UC}_{\mathbb{N}^{\mathbb{N}}} \star \Sigma_1^1\text{-WWKL} <_W \Sigma_1^1\text{-WKL}$.

PROOF. The inequality $\text{UC}_{\mathbb{N}^{\mathbb{N}}} <_W \text{UC}_{\mathbb{N}^{\mathbb{N}}} \star \Sigma_1^1\text{-WWKL}$ is obvious since no Π_1^1 -Martin-Löf random real is hyperarithmetical. Moreover, by Proposition 4.8, we have $\text{UC}_{\mathbb{N}^{\mathbb{N}}} \star \Sigma_1^1\text{-WWKL} \leq_w \Sigma_1^1\text{-WKL}$. Suppose for the sake of contradiction that $\Sigma_1^1\text{-WKL} \leq_w \text{UC}_{\mathbb{N}^{\mathbb{N}}} \star \Sigma_1^1\text{-WWKL}$. Then, for any Σ_1^1 closed set S , there are a Σ_1^1 closed set P of positive measure and a Π_1^1 function $f : P \rightarrow S$, so that $f(x) \leq_h x$ for any $x \in P$.

In particular, assume that S is the set of all $\Pi_1^1\text{-DNC}_2$ functions, and let P and f be as above. It is known that x is Π_1^1 -random iff x is Δ_1^1 -random and $\omega_1^{\text{CK},x} = \omega_1^{\text{CK}}$ (see [33, Theorem 9.3.9]). Since there are conull many Π_1^1 -random reals, $Q = \{x \in P : \omega_1^{\text{CK},x} = \omega_1^{\text{CK}}\}$ also has positive measure. Given $x \in Q$, there is an ordinal $\alpha < \omega_1^{\text{CK},x} = \omega_1^{\text{CK}}$ such that $f(x) \leq_T x \oplus \emptyset^{(\alpha)}$ (cf. [16, Lemma 4.2] and [1, Section 2.3.2]). As in [25, Theorem 5.3], it is easy to see that the $\emptyset^{(\alpha)}$ -Turing upward closure, $S_\alpha = \{z : h \leq_T z \oplus \emptyset^{(\alpha)} \text{ for some } h \in S\}$, of S has measure zero for any computable ordinal α . Hence, $\hat{S} = \bigcup \{S_\alpha : \alpha < \omega_1^{\text{CK}}\}$ is also null. Our previous argument shows that $Q \subseteq \hat{S}$, however $\mu(\hat{S}) = 0$ contradicts $\mu(Q) > 0$. \dashv

QUESTION 4.10 ([9]). $\widehat{\Sigma_1^1\text{-C}_{\mathbb{N}^{\mathbb{N}}}} <_W \text{C}_{\mathbb{N}^{\mathbb{N}}}$?

§5. Comparability of well orderings. Two statements which are equivalent to ATR_0 in the context of reverse mathematics are comparability of well orderings and weak comparability of well orderings ([42, Theorem V.6.8] and [19]). These involve two kinds of effective witnesses that one well ordering is shorter than another: strong comparison maps and order preserving maps.

DEFINITION 5.1. If $X, Y \in \mathbf{WO}$ then we say that $f : \mathbb{N} \rightarrow \mathbb{N}$ is a *strong comparison map* between X and Y , in symbols $f : X \leq_s Y$, if the following conditions hold:

- $\forall n(n \notin X \rightarrow f(n) = 0)$,
- $\forall n, m \in X(n \leq_X m \leftrightarrow f(n) \leq_Y f(m))$,
- $\forall n \in X \forall k \in Y(k \leq_Y f(n) \rightarrow \exists m \in X f(m) = k)$.

In other words, f is an order embedding of X into Y whose image is an initial segment of Y .

DEFINITION 5.2 (Comparability of well orderings). Let $\text{CWO} : \mathbf{WO} \times \mathbf{WO} \rightarrow \mathbb{N}^{\mathbb{N}}$ be the function that maps any pair (X, Y) of countable well orderings to the unique $f \in \mathbb{N}^{\mathbb{N}}$ such that $f : X \leq_s Y$ or $f : Y + 1 \leq_s X$.

The use of $Y + 1$ in the previous definition makes sure that f is unique even when X and Y are isomorphic.

DEFINITION 5.3. If $X, Y \in \mathbf{LO}$ we say that $f : \mathbb{N} \rightarrow \mathbb{N}$ is an *order preserving map* between X and Y , in symbols $f : X \leq Y$, if the following conditions hold:

- $\forall n(n \notin X \rightarrow f(n) = 0)$,
- $\forall n, m \in X(n \leq_X m \leftrightarrow f(n) \leq_Y f(m))$.

DEFINITION 5.4 (Weak comparability of well orderings). Let $\text{WCWO} : \mathbf{WO} \times \mathbf{WO} \rightrightarrows \mathbb{N}^{\mathbb{N}}$ be the multivalued function that maps any pair (X, Y) of countable well orderings to the set $\{f \in \mathbb{N}^{\mathbb{N}} : (f : X \leq Y) \vee (f : Y \leq X)\}$.

The following classifies the Weihrauch degree of comparability of well orderings:

THEOREM 5.5. $\text{UC}_{\mathbb{N}^{\mathbb{N}}} \equiv_{sW} \text{CWO}$.

PROOF. By Lemmata 5.6 and 5.7 below. ⊣

LEMMA 5.6. $\text{CWO} \leq_{sW} \text{UC}_{\mathbb{N}^{\mathbb{N}}}$.

PROOF. If $X, Y \in \mathbf{WO}$, the conjunction of the three conditions in Definition 5.1 is a Π_2^0 formula with X, Y and f as free variables. In particular, a name for the Π_2^0 set $\{f\} = \text{CWO}(X, Y)$ is computable from X and Y . Then, since $\text{UC}_{\mathbb{N}^{\mathbb{N}}} \equiv_{sW} \Pi_2^0\text{-UC}_{\mathbb{N}^{\mathbb{N}}}$ by Theorem 3.11 and Proposition 2.3, we can use the second one to obtain f . ⊣

LEMMA 5.7. $\Sigma_1^1\text{-Sep} \leq_{sW} \text{CWO}$.

PROOF. We follow essentially the proof of Theorem V.6.8 in [42]. The only modification concerns the definition of the well orderings U and V , for which the original proof uses the Σ_1^1 bounding principle.

So, let $(S_n, T_n)_{n \in \omega}$ be a double-sequence of trees in $\text{dom}(\Sigma_1^1\text{-Sep})$. Without loss of generality we assume that for all $n \in \mathbb{N}$, S_n and T_n are nonempty. We can build the corresponding double-sequence of linear orderings $(X_n, Y_n)_n$ such that, for all n , $X_n = \text{KB}(S_n)$ and $Y_n = \text{KB}(T_n)$. Note that, since $(S_n, T_n)_n \in \text{dom}(\Sigma_1^1\text{-Sep})$, we have

$$\forall n(\mathbf{WO}(X_n) \vee \mathbf{WO}(Y_n)). \tag{5}$$

Consider $U = \sum_{n \in \mathbb{N}}(\mathbb{Q} * Y_n) * X_n$, which by (5) and by Lemma 2.8.1 is a well ordering. We claim that the following holds:

$$\forall X \in \mathbf{LO} \forall n(\neg \mathbf{WO}(X_n) \rightarrow |X * Y_n| < |U|). \tag{6}$$

In fact, let $X \in \mathbf{LO}$ and n be such that $\neg \mathbf{WO}(X_n)$. Then by (5) we have $\mathbf{WO}(Y_n)$, which means that $X * Y_n$ is also a well ordering. Furthermore, by 3 and 2 of Lemma 2.8, we have $|X * Y_n| \leq |\mathbb{Q} * Y_n| \leq |(\mathbb{Q} * Y_n) * X_n| < |U|$.

For all $n \in \mathbb{N}$, define $Z_n = (U + X_n) * Y_n$. By (6) and by 1 and 2 of Lemma 2.8 we have, for all $n \in \mathbb{N}$,

$$\neg \mathbf{WO}(X_n) \rightarrow |Z_n| < |U|, \tag{7}$$

$$\neg \mathbf{WO}(Y_n) \rightarrow |U| < |Z_n|. \tag{8}$$

Finally, consider $V = U + \sum_{n \in \mathbb{N}} Z_n$ and define the well orderings

- $Z = \sum_{n \in \mathbb{N}}(Z_n + V \cdot \mathbb{N})$,
- $W = \sum_{n \in \mathbb{N}}(V + V \cdot \mathbb{N})$.

Note that all the well orderings we defined so far, in particular Z and W , are computable from the double-sequence $(X_n, Y_n)_n$. In the construction of V we can also use a special mark for its least element. Furthermore, we can code Z in such a way that, if $x \in Z_n + V \cdot \mathbb{N}$, for some $n \in \mathbb{N}$, then we are able to compute whether x belongs to Z_n or to the first copy of V , and in the second case, whether x belongs to the copy of U contained in V . Similar assumptions can be made for the construction of W .

Let now $f = \text{CWO}(Z, W)$ be the comparing map between Z and W . Since $|Z_n + V \cdot \mathbb{N}| = |V + V \cdot \mathbb{N}|$ for all n , we have $|Z| = |W|$ and f is the isomorphism of Z onto W . In particular, for each $n \in \mathbb{N}$, f induces an isomorphism f_n of $Z_n + V \cdot \mathbb{N}$ onto $V + V \cdot \mathbb{N}$. Define $g \in 2^{\mathbb{N}}$ by $g(n) = 0$ if and only if the image of Z_n under f_n is a strict initial segment of U , i.e. $|Z_n| < |U|$. This can be done computably by checking whether f_n maps the first element of the first copy of V in $Z_n + V \cdot \mathbb{N}$ to U or not. Then, recalling the definition of $(X_n, Y_n)_n$, if $[S_n] \neq \emptyset$ then $\neg \mathbf{WO}(X_n)$ and, by (7), $|Z_n| < |U|$ so that $g(n) = 0$. Similarly, if $[T_n] \neq \emptyset$ then, by (8), $|U| \leq |Z_n|$ so that $g(n) = 1$. ⊣

The Weihrauch degree of weak comparability of well orderings, however, has eluded our classification attempts:

QUESTION 5.8. *Does $\text{WCWO} \equiv_W \text{UC}_{\mathbb{N}^{\mathbb{N}}}$?*

Recently, Jun Le Goh [21] obtained a positive answer to our question.

§6. The one-sided versions of PTT and open determinacy. Both the perfect tree theorem and open determinacy have at its core a disjunction $A \vee B$ which is not to be read constructively. A typical approach to formulate these as computational tasks is to view these as implications $\neg A \Rightarrow B$ or $\neg B \Rightarrow A$. In this section, we explore these variants.

Recall that a tree is perfect if every node has at least two incomparable extensions. In particular, every perfect tree is pruned. The perfect tree theorem states that every tree with uncountably many paths has a perfect subtree and leads to the following two problems: The first problem is given a closed set A which has no perfect subset (that simply means that A is countable), and has to show its countability, that is, to enumerate all elements of A . We consider two variants of this task, depending on what exactly is meant by *listing*. The weak version contains no information about the cardinality, the strong version does. The second problem is more direct: it asks to find a perfect subset of a given tree with uncountably many paths.

DEFINITION 6.1. $\text{wList} : \subseteq \mathcal{A}(\mathbb{N}^{\mathbb{N}}) \rightrightarrows (\mathbb{N}^{\mathbb{N}})^{\omega}$ maps a countable set A to some $\langle b_0 p_0, b_1 p_1, \dots \rangle$ such that $A = \{p_i \mid b_i = 1\}$. $\text{List} : \subseteq \mathcal{A}(\mathbb{N}^{\mathbb{N}}) \rightrightarrows (\mathbb{N}^{\mathbb{N}})^{\omega}$ maps a countable set A to some $n \langle p_0, p_1, \dots \rangle$ such that either $n = 0, p_i \neq p_j$ for $i \neq j$ and $A = \{p_i \mid i \in \mathbb{N}\}$; or $n > 0, |A| = n - 1$ and $A = \{p_i \mid i < n - 1\}$.

DEFINITION 6.2. $\text{PTT}_1 : \subseteq \mathbf{Tr} \rightrightarrows \mathbf{Tr}$ maps T such that $[T]$ is uncountable to some perfect $T' \subseteq T$.

We start by reporting a result originating from discussion during the Dagstuhl seminar on Weihrauch reducibility [13], in particular including a contribution by Brattka:

PROPOSITION 6.3. $\text{PTT}_1 \equiv_W \mathbf{C}_{\mathbb{N}^{\mathbb{N}}}$.

PROOF. For $\mathbf{C}_{\mathbb{N}^{\mathbb{N}}} \leq_W \text{PTT}_1$, note that from $A \in \mathcal{A}(\mathbb{N}^{\mathbb{N}})$ we can compute a tree T such that $[T] = A \times \mathbb{N}^{\mathbb{N}}$. If A is nonempty, then $[T]$ is uncountable. Given some perfect subtree T' of T , we can compute a path through T' and hence through T . By projecting, we obtain a point in A .

For $\text{PTT}_1 \leq_W \mathbf{C}_{\mathbb{N}^{\mathbb{N}}}$, call a function $\lambda : \mathbb{N}^{<\mathbb{N}} \rightarrow \mathbb{N}$ a modulus of perfectness for T , if $v \in T$ implies that there are incomparable $u, w \in [0, \lambda(v)]^{\lambda(v)}$ with $vu, vw \in T$. A nonempty tree has a modulus of perfectness iff it is perfect, and given T the set

$$\{(T', \lambda) \in \mathbf{Tr} \times \mathbb{N}^{(\mathbb{N}^{<\mathbb{N}})} \mid \emptyset \neq T' \subseteq T \wedge \lambda \text{ is a modulus of perfectness for } T'\}$$

is closed, and nonempty for $[T]$ uncountable by the perfect tree theorem. Taking into account that $\mathbf{Tr} \times \mathbb{N}^{(\mathbb{N}^{<\mathbb{N}})}$ is computably isomorphic to $\mathbb{N}^{\mathbb{N}}$, we can thus apply $\mathbf{C}_{\mathbb{N}^{\mathbb{N}}}$ and project to obtain a perfect subtree of T . ⊣

6.1. Listing the points in a countable set. We now examine the strength of the contrapositive of the perfect tree theorem PTT_1 , which is List in our setting as explained above.

THEOREM 6.4. $\text{wList} \equiv_W \text{List} \equiv_W \mathbf{UC}_{\mathbb{N}^{\mathbb{N}}}$.

The main ingredient of our proof is a variant of the Cantor–Bendixson decomposition, designed in such a way that it can be carried out in a Borel way. This modified version works as the usual one for countable sets, but can differ for uncountable ones.⁵ If u and w are finite words on \mathbb{N} , $u \sqsubseteq w$ means that u is a prefix of w .

DEFINITION 6.5. A one-step mCB-certificate of $A \in \mathcal{A}(\mathbb{N}^{\mathbb{N}})$ consists of

- (a) A prefix-independent⁶ sequence $(w_i)_{i \in \mathbb{N}}$ of finite words ordered in a canonical way,
- (b) A sequence of bits $(b_i)_{i \in \mathbb{N}}$ which are not all 0,
- (c) A sequence of points $(p_i)_{i \in \mathbb{N}}$

subject to the following constraints:

- 1. If $b_i = 1$, then $p_i \in A \cap w_i \mathbb{N}^{\mathbb{N}}$.
- 2. If $b_i = 0$, then $\forall p \in \text{HYP}(A) p \notin A \cap w_i \mathbb{N}^{\mathbb{N}}$ and $p_i = 0^\omega$.
- 3. $\forall p, q \in \text{HYP}(A) (p \in A \cap w_i \mathbb{N}^{\mathbb{N}} \wedge q \in A \cap w_i \mathbb{N}^{\mathbb{N}} \Rightarrow p = q)$.
- 4. If $w_i \not\sqsubseteq w$ for all $i \in \mathbb{N}$, then $\exists p, q \in A \cap w \mathbb{N}^{\mathbb{N}} p \neq q$.

For a one-step mCB-certificate for A , its residue is $A \setminus \bigcup_{i \in \mathbb{N}} w_i \mathbb{N}^{\mathbb{N}}$.

DEFINITION 6.6. A global mCB-certificate for $A \in \mathcal{A}(\mathbb{N}^{\mathbb{N}})$ is indexed by some initial $I \subseteq \mathbb{N}$ (which may be empty). It consists of a sequence $(c_i)_{i \in I}$ of one-step mCB-certificates such that there exists a linear ordering $\sqsubseteq \subseteq I \times I$ with minimum 0 (if nonempty), such that c_0 is a one-step mCB-certificate for A , for each $n \in I \setminus \{0\}$, c_n is an mCB-certificate for $\bigcap_{i \sqsubseteq n} A_i$, where A_i is the residue of c_i ; and $\forall p \in \text{HYP}(A) p \notin A \cap \bigcap_{i \in I} A_i$.

LEMMA 6.7. The set of global mCB-certificates of A is uniformly Σ_1^1 in A .

PROOF. This is almost immediate from the definition, besides the quantification over HYP. That this is unproblematic follows from Kleene’s HYP-quantification theorem [27, 28] (the converse of the Spector–Gandy theorem). \dashv

LEMMA 6.8. For nonempty nonperfect $A \in \mathcal{A}(\mathbb{N}^{\mathbb{N}})$, A has a one-step mCB-certificate such that its residue is equal to its Cantor–Bendixson derivative. If all points in A are hyperarithmetical relative to A , then A has a unique one-step mCB-certificate.

PROOF. Let (q_j) be the finite or infinite list of isolated points in A , and let (u_j) be the shortest prefix such that $A \cap u_j \mathbb{N}^{\mathbb{N}} = \{q_j\}$. It follows from Corollary 3.3 applied to $A \cap u_j \mathbb{N}^{\mathbb{N}}$ that each q_j is hyperarithmetical relative to A . Let (v_k) be the list of shortest prefixes such that $A \cap v_k \mathbb{N}^{\mathbb{N}} = \emptyset$, excluding those extending some u_j . Now the sequence (w_i) is obtained such that $\{w_i\} = \{u_j\} \cup \{v_k\}$, subject to the canonical ordering condition. If $w_i = v_k$, then $b_i = 0$ and $p_i = 0^\omega$, if $w_i = u_j$ then $b_i = 1$ and $p_i = q_j$.

⁵Kreisel has shown that computable $A \in \mathcal{A}(\mathbb{N}^{\mathbb{N}})$ may have uncomputable Cantor–Bendixson rank [29]. As any total function from $\mathbb{N}^{\mathbb{N}}$ into the countable ordinals that is effectively Borel is dominated by a computable function (the Spector Σ_1^1 -boundedness principle, cf. [39]), this implies that the Cantor–Bendixson decomposition cannot be done in a Borel way.

⁶Meaning that $w_i \sqsubset w_j$ never holds.

It is immediate that the construction satisfies Conditions (1, 2, 3, and 4) and that the residue sees exactly the isolated points removed, i.e. is the Cantor–Bendixson derivative of A . It remains to argue that the mCB-certificate constructed as such is unique if all points in A are hyperarithmetical relative to A (this is a classic result, of course). As the choice of b_i and p_i was uniquely determined by the sequence (w_i) , we only need to prove that there is no alternative sequence (w'_i) . As no w_i can satisfy the conclusion of Condition (4), we know that for each w_i there exists some w'_i with $w'_i \sqsubseteq w_i$.

Assume that $w'_i \sqsubset w_i$ for some i . If $b_i = 1$, then w_i was chosen minimal under the constraint that $A \cap w_i \mathbb{N}^{\mathbb{N}}$ is a singleton, $A \cap w'_i$ contains at least two points, which are both hyperarithmetical. Hence, w'_i fails Condition (3). If $b_i = 0$, then $w'_i \mathbb{N}^{\mathbb{N}} \cap A = \emptyset$ contradicts the choice of v_k as shortest prefix, $|w'_i \mathbb{N}^{\mathbb{N}} \cap A| = 1$ contradicts the choice of u_j as shortest prefix of an isolated point in A , and $|w'_i \mathbb{N}^{\mathbb{N}} \cap A| \geq 2$ again violates Condition (3). Hence we know that all (w_i) must appear as some (w'_i) .

Assume that there is some w occurring as a w'_i but not as a w_i . As the (w'_i) are prefix-free, w is not an extension of some w_i . Hence, Condition (4) for the (w_i) implies that $|A \cap w \mathbb{N}^{\mathbb{N}}| \geq 2$. But as all points in A are hyperarithmetical, this shows that neither the conclusion of Condition (2) nor that of Condition (3) can be satisfied for $w'_i = w$, and we have obtained the desired contradiction. \dashv

COROLLARY 6.9. *If $A \in \mathcal{A}(\mathbb{N}^{\mathbb{N}})$ is countable, then A has a unique global mCB-certificate, the p_i for $b_i = 1$ occurring in some one-step mCB-certificate list all points in A , and the order type of the implied linear ordering is the Cantor–Bendixson rank of A plus 1.*

PROOF OF THEOREM 6.4. That $\text{UC}_{\mathbb{N}^{\mathbb{N}}} \leq_w \text{wList}$ is simple: Any instance of the former is an instance of the latter, and from a list repeating a single element, we can recover that element. For the other direction, we show $\text{wList} \leq_w \Sigma_1^1\text{-UC}_{\mathbb{N}^{\mathbb{N}}}$ instead and invoke Theorem 3.11. By Lemma 6.7 the set of global mCB-certificates of $A \in \mathcal{A}(\mathbb{N}^{\mathbb{N}})$ is computable as a Σ_1^1 -set from A , and by Corollary 6.9 this is a singleton for countable A . We can distinguish whether the global mCB-certificate uses an empty or nonempty linear order. In the former case, the set is empty, and in the latter case, we can compute a list of all points in A .

Again, $\text{wList} \leq_w \text{List}$ is trivial. For the reverse direction, we observe that $\text{List} \leq_w \text{UC}_{\mathbb{N}^{\mathbb{N}}} \star \text{wList}$, since $\text{UC}_{\mathbb{N}^{\mathbb{N}}}$ more than suffices to extract the required additional information from an unstructured list. We then use the preceding result and $\text{UC}_{\mathbb{N}^{\mathbb{N}}} \equiv_w \text{UC}_{\mathbb{N}^{\mathbb{N}}} \star \text{UC}_{\mathbb{N}^{\mathbb{N}}}$ from [5]. \dashv

Regarding the nonuniform aspect, it is known that every countable Π_1^0 (indeed Σ_1^1) set $A \subseteq \mathbb{N}^{\mathbb{N}}$ consists only of hyperarithmetical elements [41, Theorem III.6.2]. Theorem 6.4 concludes that every countable Π_1^0 set $A \subseteq \mathbb{N}^{\mathbb{N}}$ admits a hyperarithmetical enumeration. Combining Proposition 6.3 (and Gandy’s basis theorem [41, Corollary III.1.5]) and Theorem 6.4, we indeed get the following:

COROLLARY 6.10. *For any computable tree $T \subseteq \omega^{<\omega}$, either T has a hyperlow perfect subtree or there is a hyperarithmetical enumeration of all infinite paths through T .*

Listing on Cantor space. We have seen that for subsets of Baire space, it makes no difference whether we intend to list all points of a countable set or all points of a finite set. We briefly explore the corresponding versions for Cantor space. Let $\text{List}_{2^{\mathbb{N}}, < \omega} : \subseteq \mathcal{A}(2^{\mathbb{N}}) \rightrightarrows (2^{\mathbb{N}})^*$ denote the problem to produce a tuple of all elements of a finite closed subset of $2^{\mathbb{N}}$ (i.e. $(p_0, \dots, p_{n-1}) \in \text{List}_{2^{\mathbb{N}}, < \omega}(A)$ iff $A = \{p_i \mid i < n\}$). Let $\text{wList}_{2^{\mathbb{N}}, \leq \omega} : \subseteq \mathcal{A}(2^{\mathbb{N}}) \rightrightarrows (2^{\mathbb{N}})^{\omega}$ denote the problem to list all elements of a nonempty countable closed subset of $2^{\mathbb{N}}$ (i.e. $(p_i)_{i \in \mathbb{N}} \in \text{wList}_{2^{\mathbb{N}}, \leq \omega}(A)$ iff $\{p_i \mid i \in \mathbb{N}\} = A$). Note that $\text{List}_{2^{\mathbb{N}}, < \omega}$ is not a restriction of $\text{wList}_{2^{\mathbb{N}}, \leq \omega}$, since finite tuples and lists with finite range have distinct properties. We will in fact show in Corollary 6.15 that these two multivalued functions are incomparable with respect to Weihrauch reducibility.

PROPOSITION 6.11. $\text{List}_{2^{\mathbb{N}}, < \omega} \equiv_W \mathbf{\Pi}_2\text{-C}_{\mathbb{N}}$.

PROOF. To see that $\text{List}_{2^{\mathbb{N}}, < \omega} \leq_W \mathbf{\Pi}_2\text{-C}_{\mathbb{N}}$, note that we can guess a finite partition of $2^{\mathbb{N}}$ into clopens A_0, \dots, A_n such that $|A \cap A_i| = 1$ for input A and any i . Verifying a correct partition is $\mathbf{\Pi}_2^0$ (because $A \cap A_i \neq \emptyset$ and $|A \cap A_i| \leq 1$ are respectively a $\mathbf{\Pi}_1^0$ and a $\mathbf{\Pi}_2^0$ condition), and given a correct partition, we can compute the listing since $\text{UC}_{2^{\mathbb{N}}}$ is computable.

For the other direction, note that we can view $\mathbf{\Pi}_2\text{-C}_{\mathbb{N}}$ as the following task: Given $(p_0, p_1, \dots) \in (2^{\mathbb{N}})^{\omega}$ with the promise that if $|\{j \mid p_i(j) = 1\}| = \infty$ then $|\{j \mid p_{i+1}(j) = 1\}| = \infty$, and that there exists some i with $|\{j \mid p_i(j) = 1\}| = \infty$, find such an i (for details, see [9]). We now construct $A \in 2^{\mathbb{N}}$ as follows: For each i , keep track of an auxiliary variable k_i , which is initially 0. Start enumerating all $0^{(i,k)}1$ into the complement of A except the $0^{(i,k_i)}1$. Also enumerate all 0^i1^s0 . Whenever we read another 1 in p_i , we do enumerate $0^{(i,k_i)}1$, and set the new k_i to be the least k such that $0^{(i,k)}1$ has not been enumerated yet.

Whenever $|\{j \mid p_i(j) = 1\}| < \infty$ for some i , then k_i will eventually remain constant. The resulting set A will be of the form $\{0^{\omega}\} \cup \{0^{(i,k_i)}1^{\omega} \mid i \in I\}$ where I is the finite set of nonsolutions. Having a finite listing of A lets us easily pick some solution. \dashv

As a corollary one can see that every finite $\mathbf{\Pi}_1^0$ subset of $2^{\mathbb{N}}$ admits a computable listing uniformly in $\mathbf{0}''$, and the complexity $\mathbf{0}''$ is optimal: If a function f sends an index (i.e. a Gödel number) of a $\mathbf{\Pi}_1^0$ set $P \subseteq 2^{\mathbb{N}}$ to an index of a computable listing of elements of P whenever P is finite, then f must compute $\mathbf{0}''$.

PROPOSITION 6.12. $\text{wList}_{2^{\mathbb{N}}, \leq \omega} \equiv_W \widehat{\text{wList}_{2^{\mathbb{N}}, \leq \omega}} \leq_W \text{UC}_{\mathbb{N}^{\mathbb{N}}} \equiv_W \mathbf{\Pi}_2\text{-C}_{\mathbb{N}} \star \text{wList}_{2^{\mathbb{N}}, \leq \omega}$.

PROOF. To note that $\text{wList}_{2^{\mathbb{N}}, \leq \omega}$ is parallelizable, observe that we can effectively join countably many trees along a comb, and the set of paths of the result is essentially the disjoint union of the original paths. The second reduction follows from the obvious embedding of $2^{\mathbb{N}}$ into $\mathbb{N}^{\mathbb{N}}$ as a closed set and Theorem 6.4. For the third reduction, note that we can embed $\mathbb{N}^{\mathbb{N}}$ as a $\mathbf{\Pi}_2^0$ -subspace B into $2^{\mathbb{N}}$ such that $2^{\mathbb{N}} \setminus B$ is countable. Given some singleton $A \in \mathcal{A}(\mathbb{N}^{\mathbb{N}})$, we can compute some countable $\bar{A} \in \mathcal{A}(2^{\mathbb{N}})$ such that $\bar{A} \cap B$ is the image of A under that embedding. If we have a list of all points in \bar{A} , we can then use $\mathbf{\Pi}_2^0\text{-C}_{\mathbb{N}}$ to pick the one in B . That the third reduction is an equivalence follows from the second, the observation that $\mathbf{\Pi}_2^0\text{-C}_{\mathbb{N}} \leq_W \text{UC}_{\mathbb{N}^{\mathbb{N}}}$ and $\text{UC}_{\mathbb{N}^{\mathbb{N}}} \star \text{UC}_{\mathbb{N}^{\mathbb{N}}} \equiv_W \text{UC}_{\mathbb{N}^{\mathbb{N}}}$ (cf. [5]). \dashv

PROPOSITION 6.13. $\lim \leq_W \text{wList}_{2^{\mathbb{N}}, \leq \omega}$.

PROOF. Consider the map $\text{id} : \mathcal{A}(\mathbb{N}) \rightarrow \mathcal{O}(\mathbb{N})$ translating an enumeration of a complement of a set to an enumeration of the set. Studied under the name EC in [43], it is known to be equivalent to \lim . Now from $A \in \mathcal{A}(\mathbb{N})$ we can compute $\{0^\omega\} \cup \{0^n 1^\omega \mid n \in A\} \in \mathcal{A}(2^{\mathbb{N}})$. From any list of the elements of the latter set, we can then compute $A \in \mathcal{O}(\mathbb{N})$. \dashv

PROPOSITION 6.14. *The following are equivalent for single-valued $f : \subseteq \mathbf{X} \rightarrow \mathbb{N}^{\mathbb{N}}$ where \mathbf{X} is a represented space:*

1. $f \leq_W \lim$;
2. $f \leq_W \text{wList}_{2^{\mathbb{N}}, \leq \omega}$.

PROOF. Proposition 6.13 entails that 1. implies 2.

To see that 2. implies 1., consider some single-valued $f : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ with $f \leq_W \text{wList}_{2^{\mathbb{N}}, \leq \omega}$. So from any $p \in \text{dom}(f)$, we can compute some countable $A_p \in \mathcal{A}(2^{\mathbb{N}})$, and from any enumeration of the points in A_p together with p we can compute $f(p)$ via some computable K . We will argue that having access to a pruned tree T with $[T] = A_p$ suffices to compute $f(p)$, and note that pruning a binary tree is equivalent to \lim (see e.g. [34]). Let us assume that there are prefixes w_0, \dots, w_n in the pruned tree such that K upon reading p and w_0, \dots, w_n outputs some prefix w . Then there is some enumeration q_0, q_1, \dots of points in A_p such that w_0, \dots, w_n are prefixes of q_0, \dots, q_n , hence w is a prefix of $f(p)$. Conversely, for any fixed enumeration q_0, q_1, \dots of points in A_p and desired prefix length m of $f(p)$ there is some $k \in \mathbb{N}$ such that K outputs $f(p)_{\leq m}$ after having read no more than the k -length prefixes of q_i for $i \leq k$. Moreover, each $(q_i)_{\leq k}$ occurs in the pruned tree T . Thus, having access to T lets us compute longer and longer prefixes of $f(p)$, and since f is single-valued, this suffices to compute $f(p)$. \dashv

In particular, $A \subseteq \mathbb{N}$ is computable from all listings of some countable Π_1^0 set $P \subseteq 2^{\mathbb{N}}$ iff A is $\mathbf{0}'$ -computable. On the other hand, there is no computable ordinal α such that $\mathbf{0}^{(\alpha)}$ computes a listing of any countable Π_1^0 subset of $2^{\mathbb{N}}$.

COROLLARY 6.15. $\text{List}_{2^{\mathbb{N}}, < \omega} \not\leq_W \text{wList}_{2^{\mathbb{N}}, \leq \omega}$ and $\text{wList}_{2^{\mathbb{N}}, \leq \omega} \not\leq_W \text{List}_{2^{\mathbb{N}}, < \omega}$.

PROOF. For the first claim, it is known that $\Pi_2^0\text{-C}_{\mathbb{N}} \equiv_W \Pi_2^0\text{-UC}_{\mathbb{N}}$ [9]. (Sketch: Take $(p_i)_{i \in \mathbb{N}}$ as in Proposition 6.11, and then put $\hat{p}_{i,s}(n) = 1$ iff $p_i(n) = 1$ and $p_j(t) = 0$ for all $j < i$ and $s \leq t < n$. It is easy to see that there is a unique i, s such that $|\{n \mid \hat{p}_{i,s}(n) = 1\}| = \infty$, and then $|\{n \mid p_i(n) = 1\}| = \infty$.) Then observe that $\Pi_2^0\text{-UC}_{\mathbb{N}}$ is single-valued, and that \lim is Σ_2^0 -computable while $\Pi_2^0\text{-C}_{\mathbb{N}}$ is not. The claim then follows by Proposition 6.14.

The second claim follows from the observation that any solution of a (computable) instance of $\Pi_2^0\text{-C}_{\mathbb{N}}$ must be computable, while \lim has computable instances without computable solutions. \dashv

COROLLARY 6.16. $\text{wList}_{2^{\mathbb{N}}, \leq \omega} <_W \text{wList}_{2^{\mathbb{N}}, \leq \omega} \star \text{wList}_{2^{\mathbb{N}}, \leq \omega} \star \text{wList}_{2^{\mathbb{N}}, \leq \omega} \equiv_W \text{UC}_{\mathbb{N}^{\mathbb{N}}}$.

PROOF. In Proposition 6.13 we have shown that $\lim \leq_W \text{List}_{2^{\mathbb{N}}, \leq \omega}$, which implies $\Pi_2^0\text{-C}_{\mathbb{N}} \leq_W \lim \star \lim \leq_W \text{wList}_{2^{\mathbb{N}}, \leq \omega} \star \text{wList}_{2^{\mathbb{N}}, \leq \omega}$; hence the assertion follows from Proposition 6.12 and $\text{UC}_{\mathbb{N}^{\mathbb{N}}} \star \text{UC}_{\mathbb{N}^{\mathbb{N}}} \equiv_W \text{UC}_{\mathbb{N}^{\mathbb{N}}}$. The strictness follows from Proposition 6.14 since $\text{UC}_{\mathbb{N}^{\mathbb{N}}}$ is single-valued and $\text{UC}_{\mathbb{N}^{\mathbb{N}}} \not\leq_W \lim$. \dashv

QUESTION 6.17. *Does $\text{wList}_{2^{\mathbb{N}}, \leq \omega} \star \text{wList}_{2^{\mathbb{N}}, \leq \omega} \equiv_W \text{UC}_{\mathbb{N}^{\mathbb{N}}}$ hold?*

The feature that $\text{wList}_{2^{\mathbb{N}}, \leq \omega}$ is not closed under composition itself, but that the hierarchy of more and more compositions stabilizes at a finite level, seems surprising for a *natural* degree. A similar observation was made before regarding the degree of finding Nash equilibria in bimatrix games [26].

6.2. Finding winning strategies. We now move on to the complexity of finding winning strategies in open Gale–Stewart games. In formulating the corresponding multivalued functions, we implicitly code strategies in sequential games into Baire space elements.

DEFINITION 6.18. $\text{FindWS}_{\Sigma} : \subseteq \mathcal{O}(\mathbb{N}^{\mathbb{N}}) \rightrightarrows \mathbb{N}^{\mathbb{N}}$ ($\text{FindWS}_{\Pi} : \subseteq \mathcal{O}(\mathbb{N}^{\mathbb{N}}) \rightrightarrows \mathbb{N}^{\mathbb{N}}$) maps an open game where Player 2 (Player 1) has no winning strategy to a winning strategy for Player 1 (Player 2). Likewise, FindWS_{Δ} maps a clopen game where Player 2 has no winning strategy to a winning strategy for Player 1. Here a name for a clopen set consists of two names for open sets which are one the complement of the other.

On the one hand, the difficulty of finding a winning strategy for a closed player is the same as the closed choice on Baire space.

PROPOSITION 6.19. $\text{FindWS}_{\Pi} \equiv_W \text{C}_{\mathbb{N}^{\mathbb{N}}}$.

PROOF. For $\text{C}_{\mathbb{N}^{\mathbb{N}}} \leq_W \text{FindWS}_{\Pi}$, note that we can turn any $A \in \mathcal{A}(\mathbb{N}^{\mathbb{N}})$ into a Σ_1^0 game where Player 1’s moves do not matter, and Player 2 wins iff his moves form a point $p \in A$.

For $\text{FindWS}_{\Pi} \leq_W \text{C}_{\mathbb{N}^{\mathbb{N}}}$, note that given a Player 2 strategy τ and the Σ_1^0 winning condition $W \subseteq \mathbb{N}^{\mathbb{N}}$ we can compute a tree $T_{W, \tau}$ describing the options available to Player 1: Essentially, the strategies σ winning against τ correspond to finite paths in $T_{W, \tau}$ ending in a leaf, whereas strategies σ' losing against τ correspond to infinite paths through $T_{W, \tau}$. Thus, τ is a winning strategy for Player 2 iff $T_{W, \tau}$ is a pruned tree, i.e. a tree without any leaves. Let $\lambda : \mathbb{N}^* \rightarrow \mathbb{N}$ be a witness of prunedness of T iff $\forall v \in T \ v\lambda(v) \in T$. If Player 2 has a winning strategy for the game W , then the set

$$\{(\tau, \lambda) \mid \lambda \text{ is a witness of prunedness for } T_{W, \tau}\}$$

is a nonempty closed set computable from W , and projecting a member of it yields a winning strategy for Player 2. \dashv

On the other hand, the difficulty of finding a winning strategy for an open/clopen player is the same as the unique choice on Baire space. In the case of clopen games, we even get full determinacy defined as follows:

DEFINITION 6.20. $\text{Det}_{\Delta} : \Delta_1^0(\mathbb{N}^{\mathbb{N}}) \rightrightarrows \mathbb{N}^{\mathbb{N}} \times \mathbb{N}^{\mathbb{N}}$ maps a clopen game W to a pair of strategies σ, τ such that either σ is winning for Player 1 or τ is winning for Player 2 (i.e. a Nash equilibrium).

THEOREM 6.21. $\text{FindWS}_\Delta \equiv_W \text{Det}_\Delta \equiv_W \text{FindWS}_\Sigma \equiv_W \text{UC}_{\mathbb{N}^{\mathbb{N}}}$.

We will prove Theorem 6.21 using the following lemmata.

LEMMA 6.22. $\text{FindWS}_\Sigma \leq_W \Sigma_1^1\text{-UC}_{\mathbb{N}^{\mathbb{N}}}$.

PROOF. Let T be a tree describing the complement of some open set, the payoff for Player 1. Fix some strategy σ of Player 1. We understand this to prescribe the action even at positions made impossible by σ itself. For any $v \in \mathbb{N}^*$ where Player 1 moves, consider the trees T_i^v describing the options available to Player 2 if the game starts at v , Player 1 plays i and otherwise follows σ . σ is a winning strategy iff for any v compatible with σ we find that $T_{\sigma(v)}^v$ is well-founded. Only $\Sigma_1^1\text{-UC}_{\mathbb{N}^{\mathbb{N}}}$ is available here while a lot of strategies may exist. We overcome this difficulty by considering the optimal strategy, that is, the one that minimizes the rank of $T_{\sigma(v)}^v$.

Let v be a position where Player 1 moves. A certificate of optimality for σ at v describes maps preserving \sqsubset from $T_{\sigma(v)}^v$ to $T_i^v \setminus \{\lambda\}$ (here λ denotes the empty sequence) for every $i < \sigma(v)$, and maps preserving \sqsubset from $T_{\sigma(v)}^v$ to T_j^v for every $j > \sigma(v)$. The set of strategies σ and corresponding certificates of optimality for all positions is a closed set computable from the game.

If we fix partial strategies of all proper extensions of v such that Player 1 can win from v , then there is a unique action of Player 1 at v such that extending the strategy to v admits a certificate of optimality. It follows that if Player 1 has a winning strategy, then there is a unique strategy admitting a certificate of optimality at all compatible positions; and this strategy is winning. We can compute this using $\Sigma_1^1\text{-UC}_{\mathbb{N}^{\mathbb{N}}}$. \dashv

COROLLARY 6.23. $\text{FindWS}_\Sigma \leq_W \text{UC}_{\mathbb{N}^{\mathbb{N}}}$.

PROOF. By Lemma 6.22 and Theorem 3.11. \dashv

LEMMA 6.24. $\text{Det}_\Delta \leq_W \text{FindWS}_\Delta$.

PROOF. Given a Δ_1^0 -game G , we can compute the derived Δ_1^0 -game G' where the first player can decide whether to play G as Player 1, or as Player 2, and then proceed a play of a chosen side. Thus, Player 1 can definitely win G' , and a winning strategy of Player 1 in G' tells us who wins G and how. \dashv

LEMMA 6.25. $\widehat{\text{FindWS}}_\Delta \leq_W \text{FindWS}_\Delta$.

PROOF. Given a sequence G_0, G_1, \dots of Δ_1^0 -games all won by Player 1, we combine them into a single Δ_1^0 game where Player 2 first chooses n , and then the players play G_n . Player 1 wins the combined game, and any winning strategy in that game yields in the obvious way winning strategies for every G_i . \dashv

Let \mathbb{S}_B denote the space of Borel-truth values (cf. [22, 39]). Roughly speaking, if p is a Borel code of a Borel subset A of the singleton space $\{\bullet\}$, then we think of p as a name of \top (\perp , resp.) iff $A \neq \emptyset$ ($A = \emptyset$, resp.); if p is not a Borel code, p is not in the domain of the representation.

LEMMA 6.26. $(id : \mathbb{S}_B \rightarrow \mathbf{2}) \leq_W \text{Det}_\Delta$.

PROOF. A Borel code can be viewed as a well-founded tree whose even-levels (odd-levels, resp.) consist of \exists -vertices (\forall -vertices, resp.) and leaves are labeled by either \top or \perp (corresponding to either $\{\bullet\}$ or \emptyset) [22, 39]. We can turn a \mathbb{S}_B -name into a Δ_1^0 -game by letting Player 1 control the \exists -vertices, Player 2 the \forall -vertices, make the \top -leaves winning for Player 1 and the \perp -leaves losing. Then Player 1 has a winning strategy iff the value of the root is \top . Given a Nash equilibrium (σ, τ) we can compute the leaf reached by the induced play, and find it to be equal to the truth value of the root. \dashv

PROOF OF THEOREM 6.21. As shown in [39, Theorem 80], $UC_{\mathbb{N}^{\mathbb{N}}} \leq_W (id : \widehat{\mathbb{S}_B} \rightarrow \mathbf{2})$. By Lemma 6.26, the latter is reducible to $\widehat{\text{Det}}_\Delta$. This is reducible to FindWS_Δ by Lemma 6.24, which in turn reduces to FindWS_Δ by Lemma 6.25. $\text{FindWS}_\Delta \leq_W \text{Det}_\Delta$ is trivial, and so is $\text{FindWS}_\Delta \leq_W \text{FindWS}_\Sigma$. $\text{FindWS}_\Sigma \leq_W UC_{\mathbb{N}^{\mathbb{N}}}$ follows by Corollary 6.23. \dashv

As in the case of the perfect tree theorem (Corollary 6.10), the results in this section can be viewed as a refinement of the following known result [2]:

COROLLARY 6.27. *For any open game, either the open player has a hyperarithmetical winning strategy or the closed player has a hyperlow winning strategy.*

§7. The two-sided versions of PTT and open determinacy. Rather than demanding a promise about the case of the theorem we are in, we could alternatively consider the task completely uniformly. As distinguishing the two cases is a Π_1^1 -complete question (cf. the well-known equation $\exists \Sigma_1^0 = \Pi_1^1$), the fully uniform task should **not** include the information in which case we are. A priori, since we considered two versions of listing, we also have the two corresponding version of the two-sided perfect tree theorem. We are left with the following formulations:

DEFINITION 7.1. $w\text{PTT}_2 : \mathbf{Tr} \rightrightarrows \mathbf{Tr} \times \mathbb{N}^{\mathbb{N}}$ has $(T', \langle b_0p_0, b_1p_1, b_2p_2, \dots \rangle) \in w\text{PTT}_2(T)$ iff one of the following holds:

- T' is a perfect subtree of T ;
- $[T] = \{p_i \mid b_i \neq 0\}$.

DEFINITION 7.2. $\text{PTT}_2 : \mathbf{Tr} \rightrightarrows \mathbf{Tr} \times \mathbb{N}^{\mathbb{N}}$ has $(T', n \langle p_0, p_1, p_2, \dots \rangle) \in \text{PTT}_2(T)$ iff one of the following holds:

- T' is a perfect subtree of T ;
- $n = 0, p_i \neq p_j$ for $i \neq j$ and $[T] = \{p_i \mid i \in \mathbb{N}\}$;
- $n > 0, |[T]| = n - 1$ and $[T] = \{p_i \mid i < n - 1\}$.

DEFINITION 7.3. $\text{Det}_\Sigma : \mathcal{O}(\mathbb{N}^{\mathbb{N}}) \rightrightarrows \mathbb{N}^{\mathbb{N}} \times \mathbb{N}^{\mathbb{N}}$ maps an open game W to a pair of strategies σ, τ such that either σ is winning for Player 1 or τ is winning for Player 2.

These variants are strictly harder than the nonuniform ones (which are Weihrauch reducible to $C_{\mathbb{N}^{\mathbb{N}}}$ by the results of Section 6). To see that, let $\chi_{\Pi_1^1} : \mathbb{N}^{\mathbb{N}} \rightarrow \mathbf{2}$ be the characteristic function of a Π_1^1 -complete set. Since the single-valued functions between computable Polish spaces which are Weihrauch reducible to $C_{\mathbb{N}^{\mathbb{N}}}$ are exactly

those that are effectively Borel measurable [5, Theorem 7.7], and $\chi_{\Pi_1^1}$ is not such, we have $\chi_{\Pi_1^1} \not\leq_W C_{\mathbb{N}^{\mathbb{N}}}$.

OBSERVATION 7.4. $\chi_{\Pi_1^1} \leq_W \text{LPO}' \star \text{wPTT}_2$ and $\chi_{\Pi_1^1} \leq_W \text{LPO} \star \text{Det}_{\Sigma}$.

PROOF. Deciding whether $[T]$ is uncountable and who wins a Σ_1^0 -game are Π_1^1/Σ_1^1 -complete decision problems. Given trees T' and T , we can use LPO' to decide whether or not T' is a perfect subtree of T . Given a Nash equilibrium (σ, τ) of a Σ_1^0 -game, we can compute the induced play and then use LPO to decide who wins that play—and this is the same player that has a winning strategy in the game. \dashv

COROLLARY 7.5. $C_{\mathbb{N}^{\mathbb{N}}} <_W \text{wPTT}_2 \leq_W \text{PTT}_2$ and $C_{\mathbb{N}^{\mathbb{N}}} <_W \text{Det}_{\Sigma}$.

PROOF. Using the fact that $C_{\mathbb{N}^{\mathbb{N}}}$ is closed under composition [5, Corollary 7.6] we have $\chi_{\Pi_1^1} \not\leq_W C_{\mathbb{N}^{\mathbb{N}}} \equiv_W \text{LPO} \star C_{\mathbb{N}^{\mathbb{N}}} \equiv_W \text{LPO}' \star C_{\mathbb{N}^{\mathbb{N}}}$. \dashv

In particular, we find that $\text{FindWS}_{\Sigma} <_W \text{Det}_{\Sigma}$ and $\text{FindWS}_{\Pi} <_W \text{Det}_{\Sigma}$. Thus, knowing who wins a Σ_1^0 -game makes it strictly easier to find a Nash equilibrium. This is in contrast to Δ_1^0 -games (as seen in Theorem 6.21), as well as to games on Cantor space with winning sets in the difference hierarchy over Σ_1^0 (cf. [30]). Knowing who wins the game allows for constructions such as the one used in Lemma 6.25 to conclude that finding a winning strategy is parallelizable (i.e. $\widehat{\text{FindWS}}_{\Sigma} \equiv_W \text{FindWS}_{\Sigma}$ and $\widehat{\text{FindWS}}_{\Pi} \equiv_W \text{FindWS}_{\Pi}$). We will see in Corollary 7.13 below that this is not just an obstacle for the proof strategy, but that the result differs for Det_{Σ} .

If then else. As we have seen, many theorems equivalent to ATR_0 are described as *dichotomy*-type theorems: Exactly one of A or B holds. Thus, it is natural to consider the following if-then-else problem for a given dichotomy A xor B : Provide two descriptions (α, β) trying to verify A and B simultaneously. If A is true, then α is a correct proof validating A ; or else β is a correct proof of B , where we do not need to know which one is correct. We formalize this idea as follows.

A space of truth values is just a represented space \mathbb{B} with underlying set $\{\top, \perp\}$.

DEFINITION 7.6. Let \mathbb{B} be a space of truth values. For $f : \subseteq \mathbf{X} \rightrightarrows \mathbf{Y}$ and $g : \subseteq \mathbf{A} \rightrightarrows \mathbf{B}$, we define

$$[\text{if } \mathbb{B} \text{ then } f \text{ else } g] : \subseteq \mathbb{B} \times \mathbf{X} \times \mathbf{A} \rightrightarrows \mathbf{Y} \times \mathbf{B}$$

via $(b, x_0, x_1) \in \text{dom}([\text{if } \mathbb{B} \text{ then } f \text{ else } g])$ iff $b = \top$ and $x_0 \in \text{dom}(f)$ or $b = \perp$ and $x_1 \in \text{dom}(g)$, and $(y_0, y_1) \in [\text{if } \mathbb{B} \text{ then } f \text{ else } g](b, x_0, x_1)$ iff $b = \top$ and $y_0 \in f(x_0)$ or $b = \perp$ and $y_1 \in g(x_1)$.

Note that the degree of $[\text{if } \mathbb{B} \text{ then } f \text{ else } g]$ depends on the precise choice of spaces for domain and codomains involved, beyond what matters for where f and g are actually defined and are taking their range. In particular, $[\text{if } \mathbb{B} \text{ then } f \text{ else } g]$ is not an operation on Weihrauch degrees.⁷

⁷Let \mathbf{X} be the represented space of the noncomputable elements of $\mathbb{N}^{\mathbb{N}}$, and $f : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ the restriction of $\text{id}_{\mathbb{N}^{\mathbb{N}}}$ to the noncomputable elements ($\text{id}_{\mathbf{X}}$ and f are the same function, but defined on

The upper bound. Let \mathbb{S}_{Σ_1} be the space of truth values where p is a name for \top iff p codes an ill-founded tree, and a name for \perp iff it codes a well-founded tree.

In the proofs of Propositions 6.3 and 6.19, we constructed closed sets containing information over the perfect subtrees or the winning strategies of Player 2 respectively. In particular, by testing whether these are empty or not, we can decide in which case we are, and obtain the answer in \mathbb{S}_{Σ_1} . Thus, by combining Proposition 6.3 and Theorem 6.4, respectively Proposition 6.19 and Theorem 6.21, we obtain the following:

COROLLARY 7.7. $\text{PTT}_2 \leq_W [\text{if } \mathbb{S}_{\Sigma_1} \text{ then } C_{\mathbb{N}^{\mathbb{N}}} \text{ else } UC_{\mathbb{N}^{\mathbb{N}}}]$.

COROLLARY 7.8. $\text{Det}_{\Sigma} \leq_W [\text{if } \mathbb{S}_{\Sigma_1} \text{ then } C_{\mathbb{N}^{\mathbb{N}}} \text{ else } UC_{\mathbb{N}^{\mathbb{N}}}]$.

As $UC_{\mathbb{N}^{\mathbb{N}}} \leq_W C_{\mathbb{N}^{\mathbb{N}}}$, it follows that $[\text{if } \mathbb{S}_{\Sigma_1} \text{ then } C_{\mathbb{N}^{\mathbb{N}}} \text{ else } UC_{\mathbb{N}^{\mathbb{N}}}] \leq_W C_{\mathbb{N}^{\mathbb{N}}} \star \chi_{\Pi_1^1}$. In particular, the difference between $[\text{if } \mathbb{S}_{\Sigma_1} \text{ then } C_{\mathbb{N}^{\mathbb{N}}} \text{ else } UC_{\mathbb{N}^{\mathbb{N}}}]$ and $C_{\mathbb{N}^{\mathbb{N}}}$ disappears if we move from Weihrauch reducibility to computable reducibility. It follows immediately that Gandy’s basis theorem applies to Det_{Σ} : Every Σ_1^0 -game has a Nash equilibrium that is hyperlow relative to the game.

Idempotency. We can show a kind of absorption result for the if-then-else construction. Recall that NHA asks for an output that is not hyperarithmetical relative to the input.

PROPOSITION 7.9. *Let g have a hyperarithmetical point p in its codomain. If we have $f \times \text{NHA} \leq_W [\text{if } \mathbb{B} \text{ then } g \text{ else } UC_{\mathbb{N}^{\mathbb{N}}}]$, then $f \leq_W g$.*

PROOF. Any $x \in \text{dom}(f)$ is provided in the form of some name p_x , which is a valid input to NHA. If some $(x, p_x) \in \text{dom}(f \times \text{NHA})$ were mapped to some (\perp, a, A) via the reduction, then $A = \{q\}$ where q is hyperarithmetical in p_x . Then (p, q) is a valid output of $[\text{if } \mathbb{B} \text{ then } g \text{ else } UC_{\mathbb{N}^{\mathbb{N}}}]$, but we cannot compute a solution to $\text{NHA}(p_x)$ from (p, q) .

Thus, every (x, p_x) gets mapped to (\top, a_x, A) such that from $b \in g(a_x)$ we can compute $y \in f(x)$ (since (b, z) for any z , say (b, \emptyset) , is a solution to the instance (\top, a_x, A)). This provides the claimed reduction $f \leq_W g$. ←

By Corollaries 7.5, 7.8, and 7.7, and Proposition 7.9 we get the following:

COROLLARY 7.10. $w\text{PTT}_2 \times \text{NHA} \not\leq_W [\text{if } \mathbb{S}_{\Sigma_1} \text{ then } C_{\mathbb{N}^{\mathbb{N}}} \text{ else } UC_{\mathbb{N}^{\mathbb{N}}}]$.

COROLLARY 7.11. $\text{Det}_{\Sigma} \times \text{NHA} \not\leq_W [\text{if } \mathbb{S}_{\Sigma_1} \text{ then } C_{\mathbb{N}^{\mathbb{N}}} \text{ else } UC_{\mathbb{N}^{\mathbb{N}}}]$.

Using the corollaries above in conjunction with Corollary 3.6, we obtain:

different spaces); then $\text{id}_X \equiv_W f$, yet $[\text{if } \mathbb{S} \text{ then } f \text{ else } \text{id}_{\mathbb{N}^{\mathbb{N}}}] \not\leq_W [\text{if } \mathbb{S} \text{ then } \text{id}_X \text{ else } \text{id}_{\mathbb{N}^{\mathbb{N}}}]$ because the former has computable inputs while the latter does not.

COROLLARY 7.12. $\text{wPTT}_2 \times C_{\mathbb{N}^{\mathbb{N}}} \not\leq_W \text{PTT}_2$ and hence $\text{wPTT}_2 \times \text{wPTT}_2 \not\leq_W \text{PTT}_2$.

COROLLARY 7.13. $\text{Det}_{\Sigma} \times C_{\mathbb{N}^{\mathbb{N}}} \not\leq_W \text{Det}_{\Sigma}$ and hence $\text{Det}_{\Sigma} \times \text{Det}_{\Sigma} \not\leq_W \text{Det}_{\Sigma}$.

Products with $\text{UC}_{\mathbb{N}^{\mathbb{N}}}$. While we just saw that Det_{Σ} , PTT_2 and $[\text{if } \mathbb{S}_{\Sigma}^1]$ then $C_{\mathbb{N}^{\mathbb{N}}}$ else $\text{UC}_{\mathbb{N}^{\mathbb{N}}}$ are not closed under products with $C_{\mathbb{N}^{\mathbb{N}}}$, the situation for products with $\text{UC}_{\mathbb{N}^{\mathbb{N}}}$ is different:

PROPOSITION 7.14. $\text{UC}_{\mathbb{N}^{\mathbb{N}}} \times [\text{if } \mathbb{B} \text{ then } C_{\mathbb{N}^{\mathbb{N}}} \text{ else } \text{UC}_{\mathbb{N}^{\mathbb{N}}}] \equiv_W [\text{if } \mathbb{B} \text{ then } C_{\mathbb{N}^{\mathbb{N}}} \text{ else } \text{UC}_{\mathbb{N}^{\mathbb{N}}}]$ for any space of truth values \mathbb{B} .

PROOF. Let $\{a\}, b \in \mathbb{B}$, A, B be the input to $\text{UC}_{\mathbb{N}^{\mathbb{N}}} \times [\text{if } \mathbb{B} \text{ then } C_{\mathbb{N}^{\mathbb{N}}} \text{ else } \text{UC}_{\mathbb{N}^{\mathbb{N}}}]$. We can use $[\text{if } \mathbb{B} \text{ then } C_{\mathbb{N}^{\mathbb{N}}} \text{ else } \text{UC}_{\mathbb{N}^{\mathbb{N}}}]$ on $b, \{a\} \times A$ and $\{a\} \times B$, as $\{a\} \times A$ is nonempty iff A is, and $\{a\} \times B$ is a singleton iff B is. We will receive as output $(\langle p, x \rangle, \langle q, y \rangle)$ such that $\langle x, y \rangle$ is a valid output to $[\text{if } \mathbb{B} \text{ then } C_{\mathbb{N}^{\mathbb{N}}} \text{ else } \text{UC}_{\mathbb{N}^{\mathbb{N}}}] (b, A, B)$, and at least one of p and q is a . Let us write $p_{\leq n}$ for the prefix of p of length $n + 1$. We have that, if $p_{\leq n} = q_{\leq n}$, then $p_{\leq n} = a_{\leq n}$, and if $p_{\leq n} \neq q_{\leq n}$, then either $p \notin \{a\}$ or $q \notin \{a\}$, hence we can compute a from p, q , and $\{a\}$. \dashv

PROPOSITION 7.15. $\text{UC}_{\mathbb{N}^{\mathbb{N}}} \times \text{PTT}_2 \equiv_W \text{PTT}_2$.

PROOF. Let $(\{a\}, T)$ be the input to $\text{UC}_{\mathbb{N}^{\mathbb{N}}} \times \text{PTT}_2$. From this input we can build a tree T_0 such that $[T_0] = \{a\} \times (\{0^\omega\} \cup 1[T])$ (notice that $|[T_0]| = |[T]| + 1$). $\text{PTT}_2(T_0)$ yields a tree T' and a sequence $n(\langle q_0, t_0p_0 \rangle, \langle q_1, t_1p_1 \rangle, \dots)$.

We first explain how to compute the sequence part of $\text{PTT}_2(T)$. If $n = 1$, or $n = 0$ and more than one t_i is 0, or $n > 1$ and more than one t_i for $i < n - 1$ is 0, then the sequence is not listing $[T_0]$ (because $[T_0] \neq \emptyset$ and $(a, 0^\omega)$ is the only member of $[T_0]$ whose second component starts with 0), which implies that $[T_0]$, and hence $[T]$, was uncountable. In this case, we can just output some arbitrary sequence. Otherwise let p'_i be the sequence consisting of the odd digits of p_i . If $n = 0$, we output $0\langle p'_{i_0}, p'_{i_1}, \dots \rangle$ where the i_k are the (all but one) indices such that $t_i \neq 0$ (in this way, if $(\langle q_0, t_0p_0 \rangle, \langle q_1, t_1p_1 \rangle, \dots)$ lists injectively $[T_0]$, our output lists injectively $[T]$). To achieve the same result when $n > 1$ we output $(n - 1)\langle p'_{i_0}, p'_{i_1}, \dots \rangle$ where we are omitting the (at most one) $i < n - 1$ such that $t_i = 0$.

To compute the tree part of $\text{PTT}_2(T)$, starting from T' we obtain a tree T'' as follows: On the first three levels (corresponding to the first two digits of a and the control bit), go down some arbitrary edge in T' . Then alternate adding all children of the present vertices into T'' , and passing down some arbitrary edge. If T' is perfect, then so is T'' , and moreover, $T'' \subseteq T$ in that case.

We need also to compute a . To produce a possible candidate, we attempt to compute the left-most branch q of T' . If we ever reach a leaf (which never happens if T' is perfect), then we continue q by constant 0. In any case, let q' be the even digits of q : if T' is a perfect subtree of T_0 then $a = q'$. On the other hand, if $(q_0, t_0p_0), (q_1, t_1p_1), \dots$ lists $[T_0]$ then $a = q_0$. Thus $a = q_0$ or $a = q'$. As in the proof of Proposition 7.14 it follows that we can compute a from q_0, q' and $\{a\}$. \dashv

PROPOSITION 7.16. $\text{UC}_{\mathbb{N}^{\mathbb{N}}} \times \text{Det}_{\Sigma} \equiv_W \text{Det}_{\Sigma}$.

PROOF. By Theorem 6.21, we have $UC_{\mathbb{N}^{\mathbb{N}}} \leq_w \text{FindWS}_{\Delta}$, i.e. we can compute a Δ_1^0 -game G'_1 from $\{a\}$ such that Player 1 wins G'_1 , and from a winning strategy of Player 1 in G'_1 we can compute a . Let G'_2 be the game with the roles of Player 1 and Player 2 exchanged, which is still Δ_1^0 . Now we construct a Σ_1^0 game G'' from a Σ_1^0 -game G , and from G'_1 and G'_2 .

The players start playing G and G'_2 in parallel. If Player 2 wins both of these, he wins in G'' . Else, if he loses one of them (which would happen at some finite time), the players proceed to play G'_1 , and whoever wins G'_1 wins G'' . W.l.o.g. we assume that Player 2 can choose to lose G right at the start of G'' .

Since by assumption Player 2 has a winning strategy in G'_2 , and Player 1 has a winning strategy in G'_1 , the winning strategies of Player 2 are exactly those that consists of playing winning strategies in G and G'_2 simultaneously. On the other hand, Player 1 can win the game for sure only by first playing a winning strategy in G (and arbitrarily in G'_2), followed by a winning strategy in G'_1 .

From a Nash equilibrium of the whole game we thus obtain a Nash equilibrium in G by considering how the players play in G . Furthermore, we consider how Player 1 plays in the copy of G'_1 played when Player 2 loses in G right at the start of G'' , and how Player 2 plays in G'_2 , and compute two candidates q_0, q_1 for a from that. As in the proof of Proposition 7.14, we can then compute a from $\{a\}$, q_0 and q_1 . \dashv

Here the difference between $w\text{PTT}_2$ and PTT_2 is revealed, as the former is more sensitive to products. We recall that a Weihrauch degree is called *fractal*, if it has a representative $f : \subseteq \mathbb{N}^{\mathbb{N}} \rightrightarrows \mathbb{N}^{\mathbb{N}}$ such that for any $w \in \mathbb{N}^{<\mathbb{N}}$ such that $w^{\mathbb{N}^{\mathbb{N}}} \cap \text{dom}(f) \neq \emptyset$ it holds that $f|_{w^{\mathbb{N}^{\mathbb{N}}}} \equiv_w f$. Most of the degrees considered in this articles are fractals, including $w\text{PTT}_2$.

PROPOSITION 7.17. *If f is a fractal and $\text{LPO} \times f \leq_w w\text{PTT}_2$, then $f \leq_w C_{\mathbb{N}^{\mathbb{N}}}$.*

PROOF. W.l.o.g. assume that $f : \subseteq \mathbb{N}^{\mathbb{N}} \rightrightarrows \mathbb{N}^{\mathbb{N}}$ witnesses its own fractality.

Fix a reduction of $\text{LPO} \times f$ to $w\text{PTT}_2$ and let K_1 be the computable function that transforms the output of $w\text{PTT}_2$ and the original input of $\text{LPO} \times f$ into the answer to the LPO -instance. We distinguish the following cases:

1. There exists $0^n, w \in \mathbb{N}^{<\mathbb{N}}$, a finite tree T , and a finite prefix of a list $\langle 0q_0, 0q_1, 0q_2, \dots \rangle$ such that K_1 provides its answer upon reading those (as input for LPO , input for f , first and second component of the output of $w\text{PTT}_2$, in that order).

Then by fixing the input to LPO to something consistent with 0^n and incompatible with the answer provided, we can make sure that the reduction needs to avoid the prefix to be valid for any input to f extending w . But this can only be achieved by making the input to $w\text{PTT}_2$ having uncountable body and not having T as prefix of any perfect subtree. This means in particular that we are dealing with an input to PTT_1 . As f is a fractal, restricting to those of its inputs extending w does not decrease its Weihrauch degree, and we conclude $f \leq_w C_{\mathbb{N}^{\mathbb{N}}}$.

2. For no $0^n, w \in \mathbb{N}^{<\mathbb{N}}$, finite tree T , and finite prefix of a list $\langle 0q_0, 0q_1, 0q_2, \dots \rangle$, K_1 provides its answer upon reading those.

If we fix the LPO-input to be 0^ω , we see that to ensure that K_1 behaves correctly, the list-component of the output of wPTT_2 must actually list some elements. This can only be guaranteed if the input to wPTT_2 is a tree with countable nonempty body, i.e. is already in the domain of List. We thus conclude $f \leq_W \text{List} \equiv_W \text{UC}_{\mathbb{N}^{\mathbb{N}}}$ (by Theorem 6.4) and, a fortiori, $f \leq_W \text{C}_{\mathbb{N}^{\mathbb{N}}}$. \dashv

COROLLARY 7.18. $\text{LPO} \times \text{wPTT}_2 \not\leq_W \text{wPTT}_2$.

COROLLARY 7.19. $\text{wPTT}_2 <_W \text{PTT}_2$.

PROOF. By contrasting Corollary 7.18 and Proposition 7.15. \dashv

We shall see that wPTT_2 is still closed under some nontrivial products. For that, let $\text{NON} : 2^{\mathbb{N}} \rightrightarrows 2^{\mathbb{N}}$ be defined via $q \in \text{NON}(p)$ iff $q \not\leq_T p$; i.e. NON is the function corresponding to the theorem asserting the existence of sets noncomputable in any given set.

PROPOSITION 7.20. $\text{NON} \times \text{wPTT}_2 \leq_W \text{wPTT}_2$.

PROOF. Fix a Turing functional Φ such that for every $p \in 2^{\mathbb{N}}$, Φ^p is an injective enumeration of p' , the Turing jump of p . Let $\hat{p} \in \mathbb{N}^{\mathbb{N}}$ be such that for every n we have that $\hat{p}(n) = 0$ implies $n \notin p'$ and $\hat{p}(n) > 0$ implies $\Phi^p(p(n) - 1) = n$. Then \hat{p} is Turing equivalent to p' and hence $\hat{p} \not\leq_T p$.

Notice that the function from $2^{\mathbb{N}}$ to $\mathcal{A}(\mathbb{N}^{\mathbb{N}})$ which sends p to $\{\hat{p}\}$ is computable. Therefore, from $(p, A) \in 2^{\mathbb{N}} \times \mathcal{A}(\mathbb{N}^{\mathbb{N}})$ we can compute $\{\hat{p}\} \times (\{0^\omega\} \cup 1A) \in \mathcal{A}(\mathbb{N}^{\mathbb{N}})$. From any solution to $\text{wPTT}_2(\{\hat{p}\} \times (\{0^\omega\} \cup 1A))$ we can compute a solution to $\text{wPTT}_2(A)$ with the argument of the first part of the proof of Proposition 7.15. Moreover, any solution to $\text{wPTT}_2(\{\hat{p}\} \times (\{0^\omega\} \cup 1A))$ is $\geq_T \hat{p}$, and hence solves $\text{NON}(p)$. \dashv

In [18], products with LPO and NON are used to separate Weihrauch degrees in a similar fashion.

§8. $\text{TC}_{\mathbb{N}^{\mathbb{N}}}$ —a candidate for ATR_0 ? Our separation proofs of principles like Det_{Σ} and PTT_2 from $\text{C}_{\mathbb{N}^{\mathbb{N}}}$ relied on being able to transform an arbitrary closed subset into an input for the former, with specified behaviour occurring only for nonempty closed sets. We can capture this using the notion of *total continuation* of closed choice on $\mathbb{N}^{\mathbb{N}}$:

DEFINITION 8.1. Let $\text{TC}_{\mathbb{N}^{\mathbb{N}}} : \mathcal{A}(\mathbb{N}^{\mathbb{N}}) \rightrightarrows \mathbb{N}^{\mathbb{N}}$ be defined via $p \in \text{TC}_{\mathbb{N}^{\mathbb{N}}}(A)$ iff $A \neq \emptyset \Rightarrow p \in A$.

In the same vein, we can define the total continuation of other choice principles. The computable compactness of $2^{\mathbb{N}}$ yields $\text{TC}_{2^{\mathbb{N}}} \equiv_W \text{C}_{2^{\mathbb{N}}}$. The principle $\text{TC}_{\mathbb{N}}$ was studied in [32].

- PROPOSITION 8.2.
1. $\text{C}_{\mathbb{N}^{\mathbb{N}}} <_W \text{TC}_{\mathbb{N}^{\mathbb{N}}}$;
 2. $\text{TC}_{\mathbb{N}^{\mathbb{N}}} <_W \text{LPO} \times \text{TC}_{\mathbb{N}^{\mathbb{N}}}$;
 3. If $\text{NON} \times f \leq_W \text{TC}_{\mathbb{N}^{\mathbb{N}}}$, then $f \leq_W \text{C}_{\mathbb{N}^{\mathbb{N}}}$;
 4. $\text{TC}_{\mathbb{N}^{\mathbb{N}}} <_W \text{wPTT}_2$;

5. $\text{TC}_{\mathbb{N}^{\mathbb{N}}} <_W \text{Det}_{\Sigma}$;
6. $[\text{if } \mathbb{S}_{\Sigma_1} \text{ then } C_{\mathbb{N}^{\mathbb{N}}} \text{ else } \text{UC}_{\mathbb{N}^{\mathbb{N}}}] <_W \text{TC}_{\mathbb{N}^{\mathbb{N}}} \times C_{\mathbb{N}^{\mathbb{N}}}$.

PROOF. 1. The reduction is trivial. Separation follows from $\text{LPO} \star C_{\mathbb{N}^{\mathbb{N}}} \equiv_W C_{\mathbb{N}^{\mathbb{N}}}$ and $\chi_{\Pi_1^1} \leq_W \text{LPO} \star \text{TC}_{\mathbb{N}^{\mathbb{N}}}$ (the latter is straightforward because LPO can check whether the output of $\text{TC}_{\mathbb{N}^{\mathbb{N}}}(A)$ belongs to A).

2. Again, the reduction is trivial. For the separation, assume that $\text{LPO} \times \text{TC}_{\mathbb{N}^{\mathbb{N}}} \leq_W \text{TC}_{\mathbb{N}^{\mathbb{N}}}$ via computable H, K_1, K_2 . Recall that $\text{LPO}(r) = 1$ iff $r = 0^\omega$. Consider the input 0^ω for LPO and $\mathbb{N}^{\mathbb{N}} \in \mathcal{A}(\mathbb{N}^{\mathbb{N}})$ (coded as some name t) for $\text{TC}_{\mathbb{N}^{\mathbb{N}}}$ on the left. There has to be some $p \in \mathbb{N}^{\mathbb{N}}$ such that $K_1(0^\omega, t, p) = 1$. By continuity, we find that $K_1(0^k q, t_{\leq k} t', p) = 1$ for sufficiently large k and arbitrary q, t' .

For any $A \in \mathcal{A}(\mathbb{N}^{\mathbb{N}})$ we can compute some name of the form $t_{\leq k} t'$. Now consider what happens if the inputs on the left are $0^k 1^\omega$ and some $t_{\leq k} t'$: If $H(0^k 1^\omega, t_{\leq k} t')$ ever returns a name for the empty set, then p is a valid solution to $\text{TC}_{\mathbb{N}^{\mathbb{N}}}$ on the right. But then K_1 will answer incorrectly 1. Thus, $H(0^k 1^\omega, t_{\leq k} t')$ never returns a name for the empty set. But then we obtain a reduction $\text{TC}_{\mathbb{N}^{\mathbb{N}}} \leq_W C_{\mathbb{N}^{\mathbb{N}}}$, contradicting (1).

3. As $\text{TC}_{\mathbb{N}^{\mathbb{N}}}(\emptyset)$ has computable solutions, the reduction $\text{NON} \times f \leq_W \text{TC}_{\mathbb{N}^{\mathbb{N}}}$ already has to be a reduction to $C_{\mathbb{N}^{\mathbb{N}}}$.
4. The reduction given in Proposition 6.3 works for this, by using the following observation: given $A \in \mathcal{A}(\mathbb{N}^{\mathbb{N}})$, $T \in \mathbf{Tr}$ such that $[T] = A \times \mathbb{N}^{\mathbb{N}}$ and $(T', \langle b_0 p_0, b_1 p_1, \dots \rangle) \in \text{PTT}_2(T)$, if we realize that T' is not pruned (which can happen only if $A = \emptyset$) we can continue our output with 0^ω .

Strictness follows by (3), Proposition 7.20 and Corollary 7.5.

5. The reduction given in Proposition 6.19 works for this, by using the following observation: if $A = \emptyset$ then Player 1 has a winning strategy in the Σ_1^0 game we constructed (in fact, any strategy for 1 is winning), however following the strategy for 2 provided by Det_{Σ} we find an element of $\text{TC}_{\mathbb{N}^{\mathbb{N}}}(A)$.

Strictness follows by (2), Proposition 7.16 and Corollary 7.5.

6. The arguments used to establish Lemma 6.7 or 6.22 show that the total continuation $\text{TUC}_{\mathbb{N}^{\mathbb{N}}}$ of $\text{UC}_{\mathbb{N}^{\mathbb{N}}}$ (i.e. the total multivalued function defined on $\mathcal{A}(\mathbb{N}^{\mathbb{N}})$ which extends $\text{UC}_{\mathbb{N}^{\mathbb{N}}}$ and is defined as $\mathbb{N}^{\mathbb{N}}$ on nonsingletons) is reducible to $C_{\mathbb{N}^{\mathbb{N}}}$. For example, given an arbitrary closed $A \subseteq \mathbb{N}^{\mathbb{N}}$ we can compute the nonempty Σ_1^1 set of the mCB-certificates of A and, choosing an element in it, compute the list of the elements of A whenever A is a countable, and in particular a singleton.

Thus, we can consider $\text{TC}_{\mathbb{N}^{\mathbb{N}}} \times \text{TUC}_{\mathbb{N}^{\mathbb{N}}}$ in place of $\text{TC}_{\mathbb{N}^{\mathbb{N}}} \times C_{\mathbb{N}^{\mathbb{N}}}$. Given some input b, A, B to $[\text{if } \mathbb{S}_{\Sigma_1} \text{ then } C_{\mathbb{N}^{\mathbb{N}}} \text{ else } \text{UC}_{\mathbb{N}^{\mathbb{N}}}]$ we ignore b , we feed A to $\text{TC}_{\mathbb{N}^{\mathbb{N}}}$, and B to $\text{TUC}_{\mathbb{N}^{\mathbb{N}}}$. Any resulting output pair is a valid output to $[\text{if } \mathbb{S}_{\Sigma_1} \text{ then } C_{\mathbb{N}^{\mathbb{N}}} \text{ else } \text{UC}_{\mathbb{N}^{\mathbb{N}}}]$.

To see that $\text{TC}_{\mathbb{N}^{\mathbb{N}}} \times C_{\mathbb{N}^{\mathbb{N}}} \not\equiv_W [\text{if } \mathbb{S}_{\Sigma_1} \text{ then } C_{\mathbb{N}^{\mathbb{N}}} \text{ else } \text{UC}_{\mathbb{N}^{\mathbb{N}}}]$ first notice that $\text{TC}_{\mathbb{N}^{\mathbb{N}}} \times C_{\mathbb{N}^{\mathbb{N}}} \times C_{\mathbb{N}^{\mathbb{N}}} \equiv_W \text{TC}_{\mathbb{N}^{\mathbb{N}}} \times C_{\mathbb{N}^{\mathbb{N}}}$. On the other hand, we have

$$[\text{if } \mathbb{S}_{\Sigma_1} \text{ then } C_{\mathbb{N}^{\mathbb{N}}} \text{ else } \text{UC}_{\mathbb{N}^{\mathbb{N}}}] \times C_{\mathbb{N}^{\mathbb{N}}} \not\leq_W [\text{if } \mathbb{S}_{\Sigma_1} \text{ then } C_{\mathbb{N}^{\mathbb{N}}} \text{ else } \text{UC}_{\mathbb{N}^{\mathbb{N}}}] :$$

otherwise, since by Corollaries 7.7 and 3.6 we have $\text{PTT}_2 \leq_W [\text{if } \mathbb{S}_{\Sigma_1} \text{ then } C_{\mathbb{N}^{\mathbb{N}}} \text{ else } \text{UC}_{\mathbb{N}^{\mathbb{N}}}]$ and $\text{NHA} \leq_W C_{\mathbb{N}^{\mathbb{N}}}$, we would have $\text{PTT}_2 \times \text{NHA} \leq_W [\text{if } \mathbb{S}_{\Sigma_1} \text{ then } C_{\mathbb{N}^{\mathbb{N}}} \text{ else } \text{UC}_{\mathbb{N}^{\mathbb{N}}}]$

else $\text{UC}_{\mathbb{N}^{\mathbb{N}}}$] and Proposition 7.9 would imply $\text{PTT}_2 \leq_w \text{C}_{\mathbb{N}^{\mathbb{N}}}$, against Corollary 7.5. \dashv

COROLLARY 8.3. $\text{PTT}_2^* \equiv_w \text{Det}_{\Sigma}^* \equiv_w \text{TC}_{\mathbb{N}^{\mathbb{N}}}^*$.

PROOF. $\text{TC}_{\mathbb{N}^{\mathbb{N}}}^* \leq_w \text{PTT}_2^*$ is immediate from Proposition 8.2(4). On the other hand we have

$$\text{PTT}_2^* \leq_w [\text{if } \mathbb{S}_{\Sigma_1} \text{ then } \text{C}_{\mathbb{N}^{\mathbb{N}}} \text{ else } \text{UC}_{\mathbb{N}^{\mathbb{N}}}]^* \leq_w (\text{TC}_{\mathbb{N}^{\mathbb{N}}} \times \text{C}_{\mathbb{N}^{\mathbb{N}}})^* \leq_w \text{TC}_{\mathbb{N}^{\mathbb{N}}}^*,$$

using Corollary 7.7 and Proposition 8.2(6). \dashv

The argument for Det_{Σ}^* is similar. \dashv

It is reasonable to expect a Weihrauch degree corresponding to an axiom system from reverse mathematics to be closed under finite parallelization. For candidates for WKL_0 or ACA_0 this happens inherently. Here, we might need to demand it explicitly, and thus consider the degree $\text{TC}_{\mathbb{N}^{\mathbb{N}}}^*$ rather than any directly defined one to be one of the most promising candidates.

A potentially convenient way to think about the separation between $\text{C}_{\mathbb{N}^{\mathbb{N}}}$ and $\text{TC}_{\mathbb{N}^{\mathbb{N}}}$ is in terms of translations between truth values. $\text{TC}_{\mathbb{N}^{\mathbb{N}}}$ allows us to treat a single Π_1^1 -set as an open set, whereas $\text{C}_{\mathbb{N}^{\mathbb{N}}}$ cannot even bridge the gap from Σ_1^1 to Borel.

PROPOSITION 8.4. $(\text{id} : \mathbb{S}_{\Pi_1^1} \rightarrow \mathbb{S}) \leq_w \text{TC}_{\mathbb{N}^{\mathbb{N}}}$, but $(\text{id} : \mathbb{S}_{\Sigma_1^1} \rightarrow \mathbb{S}_B) \not\leq_w \text{C}_{\mathbb{N}^{\mathbb{N}}}$.

PROOF. For the reduction, we observe that $A = \emptyset$ iff $p \notin A$ for some $p \in \text{TC}_{\mathbb{N}^{\mathbb{N}}}(A)$.

For the nonreduction, we recall that $\text{id} : \mathbb{S}_B \rightarrow \mathbf{2} \leq_w \text{UC}_{\mathbb{N}^{\mathbb{N}}}$ was shown in [39, Lemma 79], and that $\text{UC}_{\mathbb{N}^{\mathbb{N}}} \star \text{C}_{\mathbb{N}^{\mathbb{N}}} \equiv_w \text{C}_{\mathbb{N}^{\mathbb{N}}}$ as shown in [5, Theorem 7.3]. Thus, assuming the reduction would hold, we would even have that $(\text{id} : \mathbb{S}_{\Pi_1^1} \rightarrow \mathbf{2}) \leq_w \text{C}_{\mathbb{N}^{\mathbb{N}}}$, which contradicts [5, Theorem 7.7] because the unique realizer of $\text{id} : \mathbb{S}_{\Pi_1^1} \rightarrow \mathbf{2}$ is not effectively Borel measurable. \dashv

Next, we shall see that the additional computational power of $\text{TC}_{\mathbb{N}^{\mathbb{N}}}$ (even of its parallelization $\widehat{\text{TC}}_{\mathbb{N}^{\mathbb{N}}}$) over $\text{UC}_{\mathbb{N}^{\mathbb{N}}}$ concerns only multivalued problems.

THEOREM 8.5. *The following are equivalent for single-valued $f : \subseteq \mathbf{X} \rightarrow \mathbb{N}^{\mathbb{N}}$ where \mathbf{X} is a represented space:*

1. $f \leq_w \text{UC}_{\mathbb{N}^{\mathbb{N}}}$;
2. $f \leq_w \widehat{\text{TC}}_{\mathbb{N}^{\mathbb{N}}}$.

PROOF. That 1 implies 2 is trivial. For the other direction, we first argue that it suffices to consider single-valued $f : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow \{0, 1\}$. Then we show that for single-valued $f : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow \{0, 1\}$, $f \leq_{sw} \widehat{\text{TC}}_{\mathbb{N}^{\mathbb{N}}}$ implies $f \leq_w \Delta_1^1\text{-CA}$ and invoke Theorem 3.11.

Let $\delta_{\mathbf{X}}$ be the representation of \mathbf{X} . For $f : \subseteq \mathbf{X} \rightarrow \mathbb{N}^{\mathbb{N}}$, consider the map $F : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow \{0, 1\}$ where $F(nmp) = 1$ if $f(\delta_{\mathbf{X}}(p))(n) = m$ and $F(nmp) = 0$ otherwise, provided $p \in \text{dom}(f\delta_{\mathbf{X}})$. Now it holds that $F \leq_w f \leq_w \widehat{F}$ (the latter reduction holds because f

is single-valued). As $UC_{\mathbb{N}^{\mathbb{N}}}$ is parallelizable, $F \leq_w UC_{\mathbb{N}^{\mathbb{N}}}$ is equivalent to $\widehat{F} \leq_w UC_{\mathbb{N}^{\mathbb{N}}}$ and hence $f \leq_w UC_{\mathbb{N}^{\mathbb{N}}}$.

For the second claim, we can start from a strong Weihrauch reduction because $\widehat{TC_{\mathbb{N}^{\mathbb{N}}}}$ is a cylinder. Assume that $f : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow \{0, 1\}$ and $f \leq_{sW} \widehat{TC_{\mathbb{N}^{\mathbb{N}}}}$ via computable K, H . The outer reduction witness K essentially consists of two open sets $U^0, U^1 \in \mathcal{O}((\mathbb{N}^{\mathbb{N}})^{\mathbb{N}})$, while the inner reduction witness H gives us for each $p \in \mathbb{N}^{\mathbb{N}}$ a sequence $(A_n(p))_{n \in \mathbb{N}}$ of closed sets. For $S \subseteq \mathbb{N}$ and $U \in \mathcal{O}((\mathbb{N}^{\mathbb{N}})^{\mathbb{N}})$, let $\pi_S(U)$ denote the projection of U to the components in S . Now we find that:

$$f(p) = b \Leftrightarrow \forall S \subseteq \mathbb{N} \prod_{n \in S} A_n(p) \subseteq \pi_S(U^b).$$

(Notice that $\prod_{n \in \mathbb{N}} A_n(p) \subseteq U^b$ does not imply $f(p) = b$ in general because some of the $A_n(p)$ could be empty.) This is a Π_1^1 -condition. Since exactly one of $f(p) = 0$ and $f(p) = 1$ holds, we thus have a valid instance for Δ_1^1 -CA. \dashv

In particular, $\widehat{TC_{\mathbb{N}^{\mathbb{N}}}}$ does not reach the level of Π_1^1 -CA₀.

COROLLARY 8.6. $\chi_{\Pi_1^1} \not\leq_w \widehat{TC_{\mathbb{N}^{\mathbb{N}}}}$.

§9. Open questions and discussion. The results reported in Section 7 immediately lead to three interlinked questions, which unfortunately we have been unable to resolve so far:

QUESTION 9.1. Does $\text{Det}_{\Sigma} \equiv_w [\text{if } \mathbb{S}_{\Sigma_1} \text{ then } C_{\mathbb{N}^{\mathbb{N}}} \text{ else } UC_{\mathbb{N}^{\mathbb{N}}}]?$

QUESTION 9.2. Does $\text{PTT}_2 \equiv_w [\text{if } \mathbb{S}_{\Sigma_1} \text{ then } C_{\mathbb{N}^{\mathbb{N}}} \text{ else } UC_{\mathbb{N}^{\mathbb{N}}}]?$

QUESTION 9.3. How do PTT_2 and Det_{Σ} relate?

We would expect that other theorems equivalent to ATR_0 (e.g. open Ramsey) exhibit similar behaviour, i.e. a nonconstructive disjunction between cases equivalent to $C_{\mathbb{N}^{\mathbb{N}}}$ and $UC_{\mathbb{N}^{\mathbb{N}}}$ respectively. Proving any reductions between the two-sided versions of these theorems could be very illuminating. Until then, we might have to settle for classifications in the Weihrauch lattice up to $*$, and strive to understand better the degree $TC_{\mathbb{N}^{\mathbb{N}}}^*$.

Brattka has also raised the question whether the strong two-sided versions, which return an answer on the applicable case together with a witness, are worthwhile studying. It seems conceivable that finding reductions here would be easier. Up to $*$, these problems would have the degree $TC_{\mathbb{N}^{\mathbb{N}}}^* \times \chi_{\Pi_1^1}^*$. Would this be an acceptable candidate for an ATR_0 -equivalent, or is this degree too close to Π_1^1 -CA₀?

Given that $TC_{\mathbb{N}^{\mathbb{N}}}^*$ is not closed under composition, one could make the case that $TC_{\mathbb{N}^{\mathbb{N}}}^{\diamond}$ (its closure under generalized register machines, cf. [32]) is the better candidate. Note that $TC_{\mathbb{N}^{\mathbb{N}}}^{\diamond} \equiv_w (TC_{\mathbb{N}^{\mathbb{N}}} \times \chi_{\Pi_1^1})^{\diamond}$, so the distinction between the weak and strong two-sided versions of the theorems would disappear here. How well justified this step would be in particular depends on whether there exists a natural

theorem equivalent to ATR_0 in reverse mathematics where ATR_0 is actually used in a sequential way, i.e. a theorem naturally associated with a Weihrauch degree not reducible to $TC_{\mathbb{N}\mathbb{N}}^*$.

Acknowledgments. In the earlier stages of this research Marcone collaborated with Andrea Cettolo, and some of the proofs were obtained jointly with him. Pauly began working on this project while being a visiting fellow at the Isaac Newton Institute for Mathematical Sciences in the programme “Mathematical, Foundational and Computational Aspects of the Higher Infinite.” He thanks Vasco Brattka, Jun Le Goh, Luca San Mauro, and Richard Shore for inspiring conversations. The research project benefitted from discussion at the Dagstuhl seminars 15392 “Measuring the Complexity of Computational Content: Weihrauch Reducibility and Reverse Analysis” and 18361 “Measuring the Complexity of Computational Content: From Combinatorial Problems to Analysis.”

Kihara’s research was partially supported by JSPS KAKENHI Grant 17H06738, 15H03634, and the JSPS Core-to-Core Program (A. Advanced Research Networks). Marcone’s research was partially supported by PRIN 2012 Grant “Logica, Modelli e Insiemi” and by the departmental PRID funding “HiWei—The higher levels of the Weihrauch hierarchy.”

Pauly has received funding from the European Union’s Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 731143, *Computing with Infinite Data*.

REFERENCES

- [1] L. BIENVENU, N. GREENBERG, and B. MONIN, *Continuous higher randomness*. *Journal of Mathematical Logic*, vol. 17 (2017), no. 1, p. 53.
- [2] A. BLASS, *Complexity of winning strategies*. *Discrete Mathematics*, vol. 3 (1972), pp. 295–300.
- [3] V. BRATTKA, *Effective borel measurability and reducibility of functions*. *Mathematical Logic Quarterly*, vol. 51 (2005), pp. 19–44.
- [4] ———, *A Galois connection between turing jumps and limits*. *Logical Methods in Computer Science*, vol. 14 (2018), no. 3, pp. 1–37.
- [5] V. BRATTKA, M. DE BRECHT, and A. PAULY, *Closed choice and a uniform low basis theorem*. *Annals of Pure and Applied Logic*, vol. 163 (2012), no. 8, pp. 968–1008.
- [6] V. BRATTKA and G. GHERARDI, *Effective choice and boundedness principles in computable analysis*. *Bulletin of Symbolic Logic*, vol. 17 (2011), pp. 73–117.
- [7] ———, *Weihrauch degrees, omniscience principles and weak computability*, this JOURNAL, vol. 76 (2011), pp. 143–176.
- [8] V. BRATTKA, G. GHERARDI, and R. HÖLZL, *Probabilistic computability and choice*. *Information and Computation*, vol. 242 (2015), pp. 249–286.
- [9] V. BRATTKA, G. GHERARDI, R. HÖLZL, H. NOBREGA, and A. PAULY, *Borel choice*, in preparation.
- [10] V. BRATTKA, G. GHERARDI, R. HÖLZL, and A. PAULY, *The Vitali covering theorem in the Weihrauch lattice*. *Computability and Complexity: Essays Dedicated to Rodney G. Downey on the Occasion of His 60th Birthday* (A. Day, M. Fellows, N. Greenberg, B. Khossainov, A. Melnikov, and F. Rosamond, editors), Springer International Publishing, Cham, 2017, pp. 188–200.
- [11] V. BRATTKA, G. GHERARDI, and A. MARCONE, *The Bolzano-Weierstrass Theorem is the jump of Weak König’s Lemma*. *Annals of Pure and Applied Logic*, vol. 163 (2012), no. 6, pp. 623–625.
- [12] V. BRATTKA, G. GHERARDI, and A. PAULY, *Weihrauch complexity in computable analysis*. Preprint, 2017. <http://arxiv.org/abs/1707.03202>.
- [13] V. BRATTKA, A. KAWAMURA, A. MARCONE, and A. PAULY, *Measuring the complexity of computational content (Dagstuhl Seminar 15392)*. *Dagstuhl Reports*, vol. 5 (2016), no. 9, pp. 77–104.

- [14] V. BRAITKA and A. PAULY, *Computation with advice*. *Electronic Proceedings in Theoretical Computer Science*, vol. 24 (2010), pp. 41–55
- [15] ———, *On the algebraic structure of Weihrauch degrees*. *Logical Methods in Computer Science*, vol. 14 (2018), no. 4, pp. 1–36
- [16] C. T. CHONG and L. YU, *Randomness in the higher setting*, this JOURNAL, vol. 80 (2015), no. 4, pp. 1131–1148.
- [17] D. DZHAFAROV, *Joins in the strong Weihrauch degrees*. *Mathematical Research Letters*, vol. 26 (2019), no. 3, pp. 749–767.
- [18] D. D. DZHAFAROV, J. LE GOH, D. R. HIRSCHFELDT, L. PATEY, and A. PAULY, *Ramsey's theorem and products in the Weihrauch degrees*. Preprint, 2018. <https://arxiv.org/abs/1804.10968>.
- [19] H. M. FRIEDMAN and J. L. HIRST, *Weak comparability of well orderings and reverse mathematics*. *Annals of Pure and Applied Logic*, vol. 47 (1990), pp. 11–29.
- [20] G. GHERARDI and A. MARCONE, *How incomputable is the separable Hahn-Banach theorem?* *Notre Dame Journal of Formal Logic*, vol. 50 (2009), no. 4, pp. 393–425.
- [21] J. L. GOH, *Embeddings between well-orderings: computability-theoretic reductions*. *Annals of Pure and Applied Logic*, vol. 171 (2020), no. 6, art. 102789.
- [22] V. GREGORIADES, T. KISPÉTER, and A. PAULY, *A comparison of concepts from computable analysis and effective descriptive set theory*. *Mathematical Structures in Computer Science*, vol. 27 (2017), vol. 8, pp. 1414–1436.
- [23] K. HIGUCHI and A. PAULY, *The degree-structure of Weihrauch-reducibility*. *Logical Methods in Computer Science*, vol. 9 (2013), no. 2, pp. 1–17.
- [24] D. R. HIRSCHFELDT, *Slicing the Truth: On the Computability Theoretic and Reverse Mathematical Analysis of Combinatorial Principles*, World Scientific, Hackensack, NJ, 2014.
- [25] C. G. JOCKUSCH, JR. and R. I. SOARE, Π_1^0 classes and degrees of theories. *Transactions of the American Mathematical Society*, vol. 173 (1972), pp. 33–56.
- [26] T. KIHARA and A. PAULY, *Dividing by zero – How bad is it, really?* *41st International Symposium on Mathematical Foundations of Computer Science (MFCS 2016)* (P. Faliszewski, A. Muscholl, and R. Niedermeier, editors), Leibniz International Proceedings in Informatics (LIPIcs), volume 58, Schloss Dagstuhl, Dagstuhl, 2016, pp. 58:1–58:14.
- [27] S. C. KLEENE, *Hierarchies of number-theoretic predicates*. *Bulletin of the American Mathematical Society*, vol. 61 (1955), pp. 193–213.
- [28] ———, *Quantification of number-theoretic functions*. *Compositio Mathematica*, vol. 14 (1959), pp. 23–40.
- [29] G. KREISEL, *Analysis of the Cantor-Bendixson theorem by means of the analytic hierarchy*. *Bulletin de l'Académie Polonaise des Sciences. Série des Sciences Mathématiques, Astronomiques et Physiques*, vol. 7 (1959), pp. 621–626.
- [30] S. LE ROUX and A. PAULY, *Weihrauch degrees of finding equilibria in sequential games*. *Evolving Computability* (A. Beckmann, V. Mitran, and M. Soskova, editors), Lecture Notes in Computer Science, volume 9136, Springer, Cham, 2015, pp. 246–257.
- [31] A. MONTALBÁN, *On the Π_1^1 -separation principle*. *Mathematical Logic Quarterly*, vol. 54 (2008), no. 6, pp. 563–578.
- [32] E. NEUMANN and A. PAULY, *A topological view on algebraic computations models*. *Journal of Complexity*, vol. 44 (2018), pp. 1–22.
- [33] A. NIES, *Computability and Randomness*, Oxford Logic Guides, volume 51, Oxford, Oxford University Press, 2009.
- [34] H. NOBREGA and A. PAULY, *Game characterizations and lower cones in the Weihrauch degrees*. *Logical Methods in Computer Science*, vol. 15 (2019), no. 3, pp. 11.1–11.29.
- [35] A. PAULY, *How incomputable is finding Nash equilibria?* *Journal of Universal Computer Science*, vol. 16 (2010), no. 18, pp. 2686–2710.
- [36] ———, *On the (semi)lattices induced by continuous reducibilities*. *Mathematical Logic Quarterly*, vol. 56 (2010), no. 5, pp. 488–502.
- [37] ———, *Computable metamathematics and its application to game theory*. Ph.D. thesis, University of Cambridge, 2012.
- [38] ———, *On the topological aspects of the theory of represented spaces*. *Computability*, vol. 5 (2016), no. 2, pp. 159–180.

- [39] ———, *Computability on the space of countable ordinals*, this JOURNAL, accepted for publication. <http://arxiv.org/abs/1501.00386>.
- [40] A. PAULY and M. DE BRECHT, *Towards synthetic descriptive set theory: An instantiation with represented spaces*. Preprint, 2013. <http://arxiv.org/abs/1307.1850>.
- [41] G. E. SACKS, *Higher Recursion Theory*, Perspectives in Mathematical Logic, vol. 2, Springer-Verlag, Berlin, 1990.
- [42] S. SIMPSON, *Subsystems of Second Order Arithmetic*, Perspectives in Logic, Cambridge University Press, Cambridge, 2009.
- [43] T. VON STEIN, *Vergleich nicht konstruktiv lösbarer Probleme in der Analysis*, *Diplomarbeit*, Fachbereich Informatik, FernUniversität Hagen, 1989.
- [44] K. WEIHRAUCH, *The degrees of discontinuity of some translators between representations of the real numbers*, *Informatik Berichte*, vol. 129, FernUniversität Hagen, Hagen, 1992.

DEPARTMENT OF MATHEMATICAL INFORMATICS
NAGOYA UNIVERSITY
NAGOYA, JAPAN

E-mail: kihara@i.nagoya-u.ac.jp

DIPARTIMENTO DI SCIENZE MATEMATICHE, INFORMATICHE E FISICHE,
UNIVERSITÀ DI UDINE
UDINE, ITALY

E-mail: alberto.marcone@uniud.it

DEPARTMENT OF COMPUTER SCIENCE
SWANSEA UNIVERSITY
SWANSEA, UK

and

DEPARTMENT OF COMPUTER SCIENCE
UNIVERSITY OF BIRMINGHAM
BIRMINGHAM, UK

E-mail: Arno.M.Pauly@gmail.com